

**BUKTI KORESPONDENSI**  
**ARTIKEL JURNAL INTERNASIONAL BEREPUTASI**

Judul artikel : New emerging entomopathogenic fungi isolated from soil in South Sumatra (Indonesia) and their filtrate and conidial insecticidal activity against *Spodoptera litura*


Jurnal : Biodiversitas

Penulis: Siti Herlinda, Riski Anwar Efendi, Radix Suharjo, Hasbi, Arum Setiawan, Elfita, Marieska Verawaty

**Bukti korespondensi**

No.	Perihal	Tanggal
1.	Bukti Konfirmasi submit paper dan full paper yang disubmit	20 September 2020
2.	Bukti konfirmasi review pertama dan hasil revisi pertama	7 Oktober 2020
3.	Bukti konfirmasi paper accepted, uncorrected Proof dan hasil koreksi penulis	9 Oktober 2020
4.	Bukti tagihan untuk penerbitan artikel	10 Oktober 2020

**1. Bukti Konfirmasi submit paper dan full paper yang disubmit**



**COVERING LETTER**

Dear **Editor-in-Chief**,

I herewith enclosed a research article,

**Title:**

New emerging entomopathogenic fungi isolated from soil in South Sumatra (Indonesia) and their filtrate and conidial insecticidal activity against *Spodoptera litura*

**Author(s) name:**

Siti Herlinda<sup>1,2,♥</sup>, Riski Anwar Efendi<sup>1</sup>, Radix Suharjo<sup>3</sup>, Hasbi<sup>2,4</sup>, Arum Setiawan<sup>5</sup>, Elfita<sup>6</sup>, Marieska Verawaty<sup>5</sup>

#### Address

(Fill in your institution's name and address, your personal cellular phone and email)

<sup>1</sup>Department of Plant Pests and Diseases, Faculty of Agriculture, Universitas Sriwijaya, Indralaya 30662, Indonesia

♥Email: [sitiherlinda@unsri.ac.id](mailto:sitiherlinda@unsri.ac.id)

<sup>2</sup>Research Center for Sub-optimal Lands (PUR-PLSO), Universitas Sriwijaya. Palembang 30139, Indonesia

<sup>3</sup>Department of Plant Protection, Faculty of Agriculture, Universitas Lampung. Jl. Prof. Sumantri Brojonegoro No. 1 Bandar Lampung 35145, Lampung, Indonesia

<sup>4</sup>Agricultural Engineering Study Program, Faculty of Agriculture, Universitas Sriwijaya, Indralaya 30662, Indonesia

<sup>5</sup>Department of Biology, Faculty of Mathematics and Natural Sciences, Universitas Sriwijaya, Indralaya 30662, Indonesia

<sup>6</sup>Department of Chemistry, Faculty of Mathematics and Natural Sciences, Universitas Sriwijaya, Indralaya 30662, Indonesia

#### For possibility publication on the journal:

(fill in *Biodiversitas* or *Nusantara Bioscience* or mention the others)

*Biodiversitas*

#### Novelty:

(state your claimed novelty of the findings versus current knowledge)

This study highlights the finding that there are two new emerging entomopathogenic fungal species, *Penicillium citrinum* and *Talaromyces diversus* found in South Sumatra soil. The both fungi have higher insecticidal activity than other previously entomopathogenic fungal species, *Beauveria bassiana* and *Metarhizium anisopliae* found in the South Sumatra soil.

#### Statements:

This manuscript has not been published and is not under consideration for publication to any other journal or any other type of publication (including web hosting) either by me or any of my co-authors.

Author(s) has been read and agree to the Ethical Guidelines.

#### List of five potential reviewers

(Fill in names of five potential reviewers **that agree to review your manuscript** and their **email** addresses. He/she should have Scopus ID and come from different institution with the authors; and from at least three different countries)

1. Dr. Hari Purnono, Department of Plant Pests and Diseases, Faculty of Agriculture, Universitas Jember, Indonesia, e-mail: [haripurnomo.faperta@unej.ac.id](mailto:haripurnomo.faperta@unej.ac.id)
2. Prof. Itji Diana Daud, Department of Plant Pests and Diseases, Faculty of Agriculture, Universitas Hasanuddin University, Makassar, Indonesia, email: [itfir@yahoo.com](mailto:itfir@yahoo.com)
3. Dr. Eka Candra Lina, Department of Plant Pests and Diseases, Faculty of Agriculture, Universitas Andalas, Padang, Indonesia, e-mail: [ekacandra222@gmail.com](mailto:ekacandra222@gmail.com)
4. Prof. Abu Hassan Ahmad, School of Biological Sciences, Universiti Sains Malaysia, 11800 Penang, Malaysia, e-mail: [stegoculex@gmail.com](mailto:stegoculex@gmail.com)
5. Dr. Buyung A. R. Hadi, International Rice Research Institute, Phnom Penh, Cambodia, e-mail: [b.hadi@irri.org](mailto:b.hadi@irri.org) and [buyung.hadi@sdstate.edu](mailto:buyung.hadi@sdstate.edu)

#### Place and date:

Palembang, 20 September 2020

#### Sincerely yours,

(fill in your name, no need scanned autograph)

Siti Herlinda

# New emerging entomopathogenic fungi isolated from soil in South Sumatra (Indonesia) and their filtrate and conidial insecticidal activity against *Spodoptera litura*

SITI HERLINDA<sup>1,2,\*</sup>, RISKI ANWAR EFENDI<sup>1</sup>, RADIX SUHARJO<sup>3</sup>, HASBI<sup>2,4</sup>, ARUM SETIAWAN<sup>5</sup>, ELFITA<sup>6</sup>,  
MARIESKA VERAWATY<sup>5</sup>

<sup>1</sup>Department of Plant Pests and Diseases, Faculty of Agriculture, Universitas Sriwijaya, Indralaya 30662, Indonesia

\*Email: [sitiherlinda@unsri.ac.id](mailto:sitiherlinda@unsri.ac.id)

<sup>2</sup>Research Center for Sub-optimal Lands (PUR-PLSO), Universitas Sriwijaya, Palembang 30139, Indonesia

<sup>3</sup>Department of Plant Protection, Faculty of Agriculture, Universitas Lampung, Jl. Prof. Sumantri Brojonegoro No. 1 Bandar Lampung 35145, Lampung, Indonesia

<sup>4</sup>Agricultural Engineering Study Program, Faculty of Agriculture, Universitas Sriwijaya, Indralaya 30662, Indonesia

<sup>5</sup>Department of Biology, Faculty of Mathematics and Natural Sciences, Universitas Sriwijaya, Indralaya 30662, Indonesia

<sup>6</sup>Department of Chemistry, Faculty of Mathematics and Natural Sciences, Universitas Sriwijaya, Indralaya 30662, Indonesia

Manuscript received: 20 September 2020. Revision accepted: .....

**Abstract.** Exploring entomopathogenic fungi (EPF) which are more toxic to *S. litura* larvae still needs to be carried out. This study was performed to confirm identity of *Beauveria bassiana* and *Metarhizium anisopliae* using sequence analysis of Internal Transcribed Spacer (ITS) region as well as the two unknown isolates of EPF and to investigate their filtrate and conidial insecticidal activity against *S. litura* larvae. The EPF species were identified based on morphological characters and molecular analysis. Their culture filtrate and conidial bioassays were performed on *S. litura* larvae. The species found were *Penicillium citrinum* (BKbTp), *Talaromyces diversus* (MSwTp1), *B. bassiana* (BSwTd4), and *M. anisopliae* (MSwTp3) and have been deposited in the GenBank. All EPF species have a high insecticidal activity against *S. litura* larvae. The highest larvae mortality caused by filtrate was found on *P. citrinum* (98.67%) and not significantly different from those of *T. diversus* (96%), but significantly different from *B. bassiana* (86.67%) and *M. anisopliae* (82.67%). The highest larvae mortality caused by the conidia was also occurred to *P. citrinum* (98.67%) which was not significantly different from those of *T. diversus* (97.33%), *B. bassiana* (86.67%), and *M. anisopliae* (86.67%). The LT<sub>50</sub> caused by *T. diversus* filtrate was the shortest (6.19 days) and not significantly different from those caused by *P. citrinum* (6.62 days). The shortest LT<sub>50</sub> caused by conidia was found on *P. citrinum* (5.45 days). Therefore, the conidia and filtrate of *P. citrinum*, a new emerging entomopathogenic fungus, has the highest insecticidal activity. *P. citrinum* has the potential to be developed into commercial mycoinsecticides.

**Key words:** *Beauveria bassiana*; leaf eating caterpillar; *Metarhizium anisopliae*; *Penicillium citrinum*; *Talaromyces diversus*

**Abbreviations** (if any): -

**Running title:** New emerging entomopathogenic fungi from South Sumatra

## INTRODUCTION

South Sumatra, Indonesia has a stretch of land ranging from lowlands to high ones with varied topography. This topographic variation can grow many species of plants, for example vegetables and the food crops. Vegetables that can adapt to various land topography in South Sumatra include chili (*Capsicum* spp.), mustard greens (*Brassica juncea*), kale (*Ipomoea reptans*), and spinach (*Amaranthus tricolor*) (Siaga et al. 2019). The main constraints of vegetable cultivation, especially chili pests are fruit flies (*Bactrocera dorsalis*) (Syamsudin et al. 2019), sap sucker (Thrips) (Johari et al. 2016), and leaf eating caterpillar (*Spodoptera litura*) (Tompe et al. 2020).

*S. litura* is a polyphagous insect pest that attacks many plant species from various families (Tompe et al. 2020), like chili, mustard greens, water spinach, and spinach. The yield loss of chili caused by this pest reaches 50% every year (Vijayalakshmi et al. 2016). The first instar of *S. litura* larvae eats the leaves in groups and skeletonize them, while the 2nd instar up to the last instar spread and eat individually leaves, flowers and pistil chili (Tompe et al. 2020). In addition, *S. litura* is a key pest for the leafy vegetables, such as mustard greens, water spinach, and spinach because the leaves that have been eaten become no longer valuable (Vijayalakshmi et al. 2016; Tompe et al. 2020). Since the public awareness on the good health and environmental quality is increasing, it is needed to reduce on using synthetic pesticides and improving application of eco-friendly control strategies for *S. litura*. One of the methods is applying the entomopathogenic fungi (EPF).

Some EPF species have been shown to be effective in killing *S. litura*, for example, *Beauveria bassiana* cultured in liquid media has been shown to kill 98% of *S. litura* larvae (Ayudya et al. 2019; Gustianingtyas et al. 2020). *Metarhizium anisopliae*

was reportedly also effective in killing *S. litura* larvae (Sowmya et al., 2017; Yang et al., 2019; Gustianingtyas et al. 2020) and can kill 96% of *S. litura* larvae (Gustianingtyas et al. 2020). Furthermore, *Metarhizium rileyi* also effectively kills *S. litura* larvae (Liu et al. 2017). In addition to being effective in killing the insect pests, EPF can also be integrated with entomopathogenic viruses, such as nucleopolyhedroviruses (NPV) (Souza et al. 2019) and botanical insecticides (Hernandez-Trejo et al. 2019) and does not harm parasitoids (Ngangambe and Mwatawala 2020).

Four isolates of EPF have been isolated from several kinds of plant rhizosphere in lowland to high land of South Sumatra. On the basis of morphological characters, the two isolates have been identified as *B. bassiana* and *M. anisopliae* (Safitri et al. 2018; Sumikarsih et al. 2019; Gustianingtyas et al. 2020) and their filtrates were proven to be effectively killed up to 98% of *S. litura* larvae (Ayudya et al. 2019; Gustianingtyas et al. 2020). The other two EPF, until now, have not been fully revealed on molecular identifications and their filtrate and conidial capability to cause mortality on *S. litura*. This research aimed to confirm identity of *Beauveria bassiana* and *Metarhizium anisopliae* using sequence analysis of Internal Transcribed Spacer (ITS) region as well as the two unknown isolates of EPF and to investigate their filtrate and conidial insecticidal activity against *S. litura* larvae.

## MATERIALS AND METHODS

### Exploring EPF

The EPF exploration was carried out on soil near rooting for oil palm, rice, maize and vegetables in various locations (Table 1) ranging from lowlands to highlands of South Sumatra using the method of Safitri et al. (2018). Those locations chosen have objective to obtain the fungi that had adapted in the soils. Morphological identifications were carried out in 2018 and 2019, and identified based on the molecular analysis was conducted in April 2020.

The EPF exploration method modified those of Anwar et al. (2015), namely using a bait method with *Tenebrio molitor* larvae as insect bait which was fed in the soil sample. The soil sampling used the method of Anwar et al. (2015) by digging the soil using surface soil sampler ( $\varnothing$  20 mm) as deep as 10 cm around the rhizosphere plant according to the method of Thaochan and Sausa-Ard (2017) and 5 points were taken to collect 1 kg of soil. Then, the soil sample was put into a plastic bag and labeled with information about the height of the location, type of commodity, and date of collection.

The soil sample was first cleaned from the roots and sieved with a 10 mesh sieve. Then, it was put into a plastic tray (32 x 25 x 5 cm<sup>3</sup>) each containing as much as 1 kg. The soil was then moistened with sterile distilled water with soil moisture exceeding 20% according to the method of Chen et al. (2014). Then, 20 newly molted 3rd-instar larvae of *T. molitor* were placed at the bottom of the tray and the larvae were sprinkled with the soil to reach 3 cm in thickness. Then, the tray containing the soil was covered with a black cloth and sprayed with distilled water to maintain the soil moisture. The larvae were infested in the soil for 7 days to give time for the entomopathogenic fungal conidia to infect them, after which the dead larvae infected with entomopathogenic fungi were grown in the Sabouraud Dextrose Agar (SDA, Merck) medium.

### Isolation and identification of EPF

The EPF infecting the larvae of *T. molitor* which was characterized by its body was overgrown with EPF mycelia, dry and contracted, then isolated and purified using the method of Youssef (2014). The isolation of EPF from the larvae's body surface began with the surface sterilization by modifying the method of Sharma et al. (2018) by rinsing with 1% of NaOCl for 1 minute, then rinsing with 100 mL of sterile distilled water for 3 times. The larvae sterilization was carried out to obtain EPF inside the haemocoel and cuticle larvae and prevent water fungi from being obtained. The sterilized larvae were grown in SDA media and incubated for 2 days. The growing fungus was purified in the SDA media until after getting pure isolates. The pure EPF was identified based on the molecular analysis.

### Fungal identification

Morphological observations were carried out based on microscopic and macroscopic characteristics of the EPF. The macroscopic observation of the fungi was the color and shape of colony, whereas the microscopic observation was the shape and size of conidia and conidiophores.

The EPF were identified based on the molecular analysis at the Laboratory of Agricultural Biotechnology (accredited according to the ISO/IEC 17025 standard), Department of Plant Protection, Faculty of Agriculture, University of Lampung, Indonesia.

### DNA extraction and PCR amplification

The DNA extraction was performed on the mycelium of 7 days old fungi using InstaGene™ Matrix (BioRad) based on the company recommendation. Amplification was carried out by Sensoquest Thermal Cycler Machine (Germany). PCR amplification on Internal Transcribed Spacer (ITS) region was conducted using primer ITS1 and ITS4 (Table 2) (White et al. 1990). The PCR was carried out with total volume of 25  $\mu$ L involving 12.5  $\mu$ L Master Mix (Red Mix) (bioline), 1  $\mu$ L of 10  $\mu$ M of primer ITS 1 and ITS 4, 2  $\mu$ L DNA and 8.5  $\mu$ L of sterile distilled water. DNA amplification consist of 1 cycle of the initiation at 95°C for 5 min, 30 cycles of denaturation at 95°C for 1 minute, primer annealing at 52°C for 1 min, primer extension at 72°C for 1 minute, and one cycle of elongation at 72°C for 5 minutes. The PCR results were run using 0.5% agarose gel in 20 mL of 1x buffer Tris-Boric Acid-EDTA (TBE) (1<sup>st</sup> Base Malaysia) with 1  $\mu$ L Ethidium Bromide (EtBr 10 mg

mL<sup>-1</sup>). The electrophoresis was performed by 1x TBE buffer at 50 volts for 70 minutes. The results were envisioned by DigiDoc UV transilluminator (UVP, USA).

**Table 1.** Locations where the EPF found in South Sumatra, Indonesia

Fungal species (codes)	Village, District/City	Ecosystem	Crop	GPS (coordinat)	Altitude (m)	GenBank Acc. No.
<i>Penicillium citrinum</i> (BKbTp)	Talang Patai, Pagar Alam	Highland soil	Cabbage	S 04°02.161' E 103°10.484'	175	MT448730
<i>Talaromyces diversus</i> (MSwTp1)	Talang Patai, Pagar Alam	Highland soil	Mustard	S 03°50.174', E 103°31.293'	193	MT448731
<i>Beauveria bassiana</i> (BSwTd4)	Talang Dabok, Ogan Komerling Ilir	Peat soil	Oil palm	S 03°23.570' E 104°51.498'	24	MT448732
<i>Metarhizium anisopliae</i> (MSwTp3)	Talang Patai, Pagar Alam	Highland soil	Mustard	S 03°50.174', E 103°31.293'	193	MT448733

**Table 2.** Primer used for amplification in this study

Target of amplification	Primer name	Sequence (5'-3')	Product size (bp)	Reference
Internal Transcribe Spacer Region	ITS1 ITS4	TCC GTA GGT GAA CCT TGC GG TCC TCC GCT TAT TGA TAT GC	600	(White et al. 1990)

The PCR product was emitted to 1<sup>st</sup> Base Malaysia for sequencing. The sequencing results were analyzed with Bio Edit program ver. 7.2.6 for windows. The result was then submitted to Basic Local Alignment Search Tool (BLAST) (<https://blast.ncbi.nlm.nih.gov/Blast.cgi>) to find their potential identity. The phylogenetic tree was developed by Mega 7 for Windows (Kumar et al. 2016) with Unweighted-pair Group Method with Arithmetic means (UPGMA). Reference strains employed in this research were obtained from NCBI (<https://www.ncbi.nlm.nih.gov/>).

### ***Spodoptera litura* mass rearing**

*S. litura* larvae used in the insecticide activity test was collected from the vegetable fields around the campus of Universitas Sriwijaya and the farmers' vegetable fields that were not applied with synthetic insecticides. The larvae were taken to the laboratory and maintained in plastic containers (Ø 25 cm, height 30 cm) topped with gauze that has been washed thoroughly using detergent. Next, pesticide-free water spinach (*Ipomoea aquatic*) was put into the container for the first instar feed of *S. litura*. The second instar was fed mulberry leaves (*Morus alba*) planted in the experimental field of the Faculty of Agriculture, Universitas Sriwijaya, Indonesia and not applied with pesticides.

Every day the larvae feed was replaced with fresh new feed. The larvae entering the pupae stage were transferred into plastic containers (Ø 10 cm, height 15 cm) whose top was exposed and the base was sprinkled with sifted soil (10 mesh) and already sterilized in an oven for 1 hour at 100°C. The thickness of the soil put in the container was 3 cm. This plastic container containing pupae was put into a gauze cage (30 x 30 x 30 cm<sup>3</sup>) already tapped with 10 water spinach sticks for laying eggs of the adults. The hatched eggs were transferred into a plastic cylinder and fed with water spinach which was replaced every day. The larvae used for toxicity testing were the second instar of the third offspring (F3) or afterwards.

### **Producing of EPF mycoinsecticides**

The EPF already identified molecularly, namely *P. citrinum*, *T. diversus*, *B. bassiana*, and *M. anisopliae* were made liquid formulation mycoinsecticides. The mycoinsecticides made of four types, each of which had active ingredients *P. citrinum*, *T. diversus*, *B. bassiana*, and *M. anisopliae* and for control the sterile distilled water was used. The making of mycoinsecticides modified the method of which Mascarin et al. (2015) began with the cultivation of EPF in solid media of Sabouraud Dextrose Agar (SDA, Merck) incubated for 7 days, and then the fungal cultures that grew were transferred to the liquid media of Sabouraud Dextrose Broth (SDB, Merck). The broth fungal cultures were incubated while being shaken using a shaker for 7 days at a speed of 120 rpm. The broth fungal cultures from SDB with a minimum density of 1x10<sup>9</sup> conidia mL<sup>-1</sup> were used for the active ingredient mycoinsecticides. The mycoinsecticides were made liquid formulation containing a mixture of 600 mL of the broth fungal cultures, plus 400 mL of carrier from shrimp shell flour compost extract, 10 mL of vegetable oil, and 10 mL of sterile distilled water solution of 0.04% polyoxyethyl-189 ene sorbitan mono-oleate (Tween®). The shrimp skin flour compost extract was made using the method of Suwandi et al. (2012). After that, the formulation was stored for 30 days before it was applied to obtain stable and non-contaminant mycoinsecticides. The mycoinsecticides activity was tested in the form of EPF culture and conidia filtrate.

### **Culture filtrate preparation of the mycoinsecticides.**

Culture filtrates containing raw secondary metabolites have high insecticidal activity (Gustianingtyas et al., 2020) and to get the culture filtrate that contains the raw secondary metabolites according to Soesanto et al. (2019) can be done by filtering fungal broth or liquid mycoinsecticides. The culture filtrate of liquid mycoinsecticides was produced from 2 stages of filtering using the method of Ayudya et al. (2019). The first stage, 100 mL liquid mycoinsecticides was filtered using Whatman filter paper no. 42 and cotton coated with a thickness of 1 cm. The filtration culture produced from the first stage filter was then

sucked up to 10 mL using a sterile hypodermic syringe to be filtered in the second stage. The second stage filtering used a syringe filter (0.45 µm pore size, Sartorius). The method of installing the syringe filter used the method of Alhusein et al. (2016) by placing it at the base of the needle, by the way the needle was removed first from the syringe, then the syringe filter was installed at the base of the needle. After the syringe filter was installed at the base of the needle, the needle was re-fitted to the syringe. The syringe containing 10 mL was pressed so that it released the culture filtrate that escaped from the syringe filter. The 100 mL of mycoinsecticide produced  $\pm$  70 mL of culture filtrate. According to Soesanto et al. (2019), this second filtering culture filtrate was the raw secondary metabolites. Next, the culture filtrate was measured by its insecticidal activity by taking as much as 1 mL to be dropped on a piece of chili leaves.

#### Bioassays of culture filtrate and conidia of EPF

Bioassays of culture filtrate from mycoinsecticides of *P. citrinum*, *T. diversus*, *B. bassiana*, and *M. anisopliae* were carried out by means of chili leaves already dropped with 1 mL of culture filtrate. As a control, it was used sterile distilled water which was dripped on chili leaves. Then the leaves were dried before putting 25 second-instar *S. litura* larvae which had been fasted for 2 hours. The second instar larvae were let to eat the leaves for 6 hours which had been dropped by the culture filtrate. After 6 hours the larvae were transferred into a plastic cylinder topped with gauze (Ø 25 cm, height 30 cm) containing 15-30 pieces of chili leaves (depending on the instar) which were clean and not dripped with culture filtrate. To maintain the moisture of chili leaves, the stalk was wrapped in cotton wool moistened with sterile distilled water. Every day chili leaves were replaced with new fresh leaves. The weight of the leaves eaten was measured according to the method of Ayudya et al. (2019) and recorded daily. Every day the dead larvae were recorded up to 12 days after the application.

Bioassays of conidia from mycoinsecticides of *P. citrinum*, *T. diversus*, *B. bassiana*, and *M. anisopliae* were carried out by means of 25 second instar *S. litura* larvae dripped topically 1 mL of conidia suspension of EPF with a concentration of  $1 \times 10^6$  conidia mL<sup>-1</sup>. Then, the larvae were fed 15-30 pieces of chili leaves. As a control it was used sterile distilled water. The weight of leaves eaten by larvae and weight of the larvae were recorded daily. Every day the dead larvae were recorded up to 12 days after the application.

The number of dead larvae was used to calculate mortality and the median Lethal Time (LT<sub>50</sub>). The weight of the chili leaves eaten and the body weight of the larvae were measured every day from the first day to 12 days after the application. The measurement of the eaten leaf weight was conducted to determine whether the EPF affected the larvae appetite, whereas the measurement of larvae body weight was carried out to determine whether or not the EPF influenced the larvae growth.


#### Data Analysis

The data differences among the treatments for edible leaf weight, larvae weight, larvae mortality, and LT<sub>50</sub> were analyzed using Analysis of Variance (ANOVA) if there were differences, then further tests were carried out using the 5% Tukey's Honestly Significant Difference (HSD). LT<sub>50</sub> data were obtained from probit analysis. All calculations were conducted using software of SAS University Edition 2.7 9.4 M5.

## RESULTS AND DISCUSSION

#### Species of EPF


There were 4 EPF species found from lowland and highland soils in South Sumatra and the results of molecular identification showed that they were *P. citrinum*, *T. diversus*, *B. bassiana*, and *M. anisopliae*. The macroscopic and microscopic appearance of the four fungal species showed obvious differences (Figure 1 and 2). The colony characteristics of *P. citrinum* on SDA medium are initially white, the older the fungus, the color turned to bluish green with a white margin, round conidia with a diameter of 2.01-2.28 µm. The characteristics matched the description written by Mathew et al. (2010) stating that *P. citrinum* has globose green granular conidia. *T. diversus* has a colony characteristic generally in the form of microcolony scattering in the medium; the color was dark green with yellow margins. According to Visagie and Jacobs (2012), the size of *T. diversus* colony ranged from 2-4 mm and sometimes only microcolony and produced yellow colored mycelia at colony peripheral zones. In this study, *T. diversus* was more difficult to culture than the other three species and showed poor growth of colony on SDA. *B. bassiana* had a colony that was white but when it got older it was yellowish or turbid white and *B. bassiana* colonized SDA faster than the other three species. Conidia of *B. bassiana* was one-celled and the same circular shape as the observation of Safitri et al. (2018). *M. anisopliae* had a colony which was also white initially similar to *B. bassiana* colony but the older the fungus the colony turned greenish white and finally dark green. Conidia of *M. anisopliae* was 6.42-6.76 µm in size. The found characteristics of the colony and conidia of *M. anisopliae* were the same as those described by Lopes et al. (2018) and Thaochan and Sausa-Ard (2017) stating that the colony is more than 5 days old, the conidia changes from white to green indicating that the conidia is matured with the conidial mass bluish olive and has cylindrical, single and one-celled, hyaline conidia.



A B

C D


**Figure 1.** Colony of *Penicillium citrinum* (A), *Talaromyces diversus* (B), *Beauveria bassiana* (C), and *Metarhizium anisopliae*



**Figure 2.** Conidia of *Penicillium citrinum* (A), *Talaromyces diversus* (B), *Beauveria bassiana* (C), and *Metarhizium anisopliae* (D)

The result of BLAST search revealed that BKbTp showed 99.38% of similarity with *P. citrinum* strain NFML\_CH33\_5BbE.1 (Acc. No. KM458810.1), MSwTp1 showed 99.05% of similarity with *T. diversus* strain DTO\_13116 (Acc. No. KJ775700.1), BSwtD4 showed 99.65% of similarity with *B. bassiana* isolate GZMS-28 (Acc. No. KT715480.1) and MSwTp3 exhibited 99.82% of similarity with *M. anisopliae* isolate HT008 (Acc. No. MH512951.1). The result of BLAST search corresponded to those of phylogenetic tree analysis. The phylogenetic tree (Figure 3) was constructed using Mega 7 for Windows (Kumar et al. 2016) with Unweighted-pair Group Method with Arithmetic means (UPGMA). Based on the phylogenetic tree, BKbTp was placed in the group of *P. citrinum* strain NFML\_CH33\_5BbE.1 (Acc. No. KM458810.1), and SL1 (Acc. No. MN735924.1). MSwTp1 was in the group of *T. diversus* strain DTO\_13116 (Acc. No. KJ775700.1) and strain SCZ5-2 (Acc. No. KY445783.1). BSwtD4 was in the group of *B. bassiana* isolate CGAIPFBS-012 (Acc. No. KY495188.1),

isolate 08F04 (Acc. No. KF359944.1), and isolate GZMS-28 (Acc. No. KT715480.1). Meanwhile, MSwTp3 was in the group of *M. anisopliae* HT008 (Acc. no. MH512951.1), A1 (Acc. No. KX806656.1), and IPPM010202 (Acc. No. KY437678.1). Thus, based on the morphological characters and molecular analysis, it confirmed that the identity of the four isolates were *P. citrinum* (BKbTp), *T. diversus* (MSwTp1), *B. bassiana* (BSwTd4), and *M. anisopliae* (MSwTp3). The isolates have been deposited in the GenBank and the accession number were MT448730 (BKbTp) (<https://www.ncbi.nlm.nih.gov/nuccore/MT448730>), MT448731 (MSwTp1) (<https://www.ncbi.nlm.nih.gov/nuccore/MT448731>), MT448732 (BSwTd4) (<https://www.ncbi.nlm.nih.gov/nuccore/MT448732>), and MT448733 (MSwTp3) (<https://www.ncbi.nlm.nih.gov/nuccore/MT448733>).



**Figure 3.** Dendrogram created on the basis of sequence analysis of ITS region using primer ITS1 and ITS4 with UPGMA method developed by MEGA7 for windows ■ = fungal isolates used in this study



**EPF insecticidal activity against *Spodoptera litura***

The EPF insecticidal activity against *S. litura* larvae is stated from the data of the effect of EPF on the weight reduction of edible chili leaves, body weight of larvae, mortality, and length of time of death. On day 1 (1 x 24 hours) after the larvae were treated with the filtrate culture of the EPF, the weight of the leaves eaten by *S. litura* larvae did not show any significant difference compared to control (sterile distilled water), but after that the weight of the leaves eaten by the larvae was more lower than control (Table 3). For the larvae treated with conidia of the EPF on day 1 (1 x 24 hours) and day 2 (2 x 24 hours), the weight of the eaten leaves did not show any significant difference compared to control, on the following day the weight of the eaten leaves by the larvae was significantly lower than those of the control.

**Table 3.** Mean of leaf weight eaten by *Spodoptera litura* larvae treated with culture filtrates and conidia of mycoinsecticides from *Penicillium citrinum*, *Talaromyces diversus*, *Beauveria bassiana*, and *Metarhizium anisopliae*

Fungal species	Leaf weight eaten by larvae (g larvae <sup>-1</sup> day <sup>-1</sup> ) on observation for 12 days											
	1	2	3	4	5	6	7	8	9	10	11	12
<b>Culture filtrate</b>												
<i>P. citrinum</i>	0.10	0.26 a	0.40 a	0.51 a	0.62 a	0.73 a	0.82 a	0.96 a	1.08 a	1.17 a	1.22 a	1.24 a
<i>T. diversus</i>	0.14	0.35 a	0.46 a	0.55 a	0.65 a	0.77 a	0.86 a	0.99 a	1.09 a	1.18 a	1.23 a	1.25 a
<i>B. bassiana</i>	0.17	0.36 a	0.47 a	0.58 a	0.74 a	0.94 a	1.08 a	1.22 a	1.35 ab	1.45 ab	1.53 ab	1.59 b
<i>M. anisopliae</i>	0.17	0.36 ab	0.53 a	0.65 a	0.78 a	0.97 a	1.11 a	1.26 a	1.42 b	1.55 b	1.65 ab	1.77 b
Control	0.24	0.56 b	0.94 b	1.32 b	1.74 b	2.33 b	2.91 b	3.45 b	4.02 c	4.65 c	5.04 c	5.58 c
F-value	1.96 <sup>ns</sup>	6.86*	10.28*	19.63*	44.96*	76.82*	124.79*	131.90*	203.70*	263.92*	281.95*	324.89*
P value	0.19	0.011	0.0031	0.00034	0.000016	0.000020	0.000031	0.000025	0.000004	0.000001	0.000012	0.000070
HSD value	-	0.10	0.14	0.15	0.12	0.12	0.12	0.13	0.117	0.115	0.119	0.122
<b>Conidia</b>												
<i>P. citrinum</i>	0.45	0.76	1.13 a	1.50 a	1.84 a	2.00 a	2.28 a	2.47 a	2.62 a	2.74 a	2.85 a	2.92 a
<i>T. diversus</i>	0.45	0.84	1.17 a	1.50 a	1.87 a	2.15 a	2.51 a	2.71 a	2.94 a	3.04 a	3.19 a	3.28 a
<i>B. bassiana</i>	0.44	0.76	1.09 a	1.43 a	1.76 a	2.07 a	2.40 a	2.59 a	2.87 a	3.01 a	3.21 a	3.45 a
<i>M. anisopliae</i>	0.34	0.70	1.08 a	1.39 a	1.78 a	2.10 a	2.35 a	2.54 a	2.72 a	2.86 a	3.01 a	3.23 a
Control	0.56	1.28	1.97 b	2.64 b	3.33 b	3.99 b	4.99 b	5.73 b	6.57 b	7.49 b	8.32 b	9.16 b
F-value	1.78 <sup>ns</sup>	3.58 <sup>ns</sup>	7.14 *	18.26 *	16.06 *	17.38 *	14.98 *	14.07 *	17.72 *	23.27 *	26.81 *	33.23 *
P value	0.23	0.06	0.01	0.0004	0.0007	0.0005	0.0011	0.0011	0.0005	0.0002	0.0001	0.00005
HSD value			0.23	0.18	0.23	0.26	0.35	0.41	0.42	0.42	0.43	0.42


Note: ns = not significantly different; \* = significantly different; values within a column followed by the same letters were not significantly different at P < 0.05 according to Tukey's HSD test



**Figure 4.** The symptoms on chili leaves eaten by *Spodoptera litura* larvae treated with culture filtrates of mycoinsecticides and control

The symptoms of leaf infested with larvae treated with filtrate or conidia culture showed less damage, whereas the control leaves left only the bone of the leaf (Figure 4 and 5). The decreased appetite in leaves treated with the culture filtrate of the EPF was faster than in leaves treated with the conidia of the EPF. All tested species of the EPF significantly

23 reduced the weight of leaves eaten by larvae compared to controls (Table 3). Thus, all species of the EPF (*P. citrinum*, *T.*  
 24 *diversus*, *B. bassiana*, and *M. anisopliae*) were effective in reducing the appetite of *S. litura* larvae.  
 25



26  
 27 **Figure 5.** The symptoms on chili leaves eaten by *Spodoptera litura* larvae treated with conidia of mycoinsecticides and control  
 28  
 29  
 30

31 **Table 4.** Mean of weight of *Spodoptera litura* larvae treated with culture filtrates and conidia of mycoinsecticides from *Penicillium*  
 32 *citrinum*, *Talaromyces diversus*, *Beauveria bassiana*, and *Metarhizium anisopliae*

Fungal species	Larvae weight (g larvae <sup>-1</sup> ) on observation for 12 days											
	1	2	3	4	5	6	7	8	9	10	11	12
<b>Culture filtrate</b>												
<i>P. citrinum</i>	0.06	0.09 a	0.12 a	0.15 a	0.19 a	0.22 a	0.25 a	0.29 a	0.34 ab	0.40 a	0.48 ab	0.50 a
<i>T. diversus</i>	0.06	0.09 a	0.12 a	0.15 a	0.18 a	0.22 a	0.27 a	0.31 a	0.38 b	0.46 b	0.53 b	0.58 b
<i>B. bassiana</i>	0.06	0.09 a	0.12 a	0.15 a	0.18 a	0.21 a	0.25 a	0.29 a	0.33 a	0.39 a	0.45 a	0.51 a
<i>M. anisopliae</i>	0.06	0.09 a	0.12 a	0.15 a	0.18 a	0.22 a	0.26 a	0.31 a	0.36 ab	0.41 ab	0.47 a	0.53 ab
Control	0.07	0.11 b	0.16 b	0.21 b	0.28 b	0.34 b	0.42 b	0.50 b	0.58 c	0.68 c	0.78 c	0.89 c
F-value	2.91 <sub>ns</sub>	6.31*	12.24*	21.07*	36.38*	52.94*	74.25*	109.18*	97.99*	96.07*	107.63*	40.12*
P value	0.09	0.014	0.0017	0.00026	0.000035	0.000085	0.00023	0.00052	0.00079	0.00085	0.00055	0.00024
HSD value	-	0.01	0.01	0.02	0.02	0.02	0.02	0.02	0.024	0.03	0.028	0.053
<b>Conidia</b>												
<i>P. citrinum</i>	0.12	0.18 ab	0.23 a	0.29 a	0.35 a	0.41 a	0.47 a	0.53 a	0.59 a	0.65 a	0.73 a	0.75 a
<i>T. diversus</i>	0.13	0.19 ab	0.25 ab	0.32 a	0.39 a	0.46 a	0.52 a	0.59 a	0.66 a	0.74 a	0.83 a	0.85 a
<i>B. bassiana</i>	0.11	0.17 a	0.24 a	0.31 a	0.38 a	0.45 a	0.50 a	0.57 a	0.64 a	0.72 a	0.81 a	0.90 a
<i>M. anisopliae</i>	0.12	0.18 ab	0.25 ab	0.31 a	0.38 a	0.45 a	0.52 a	0.59 a	0.66 a	0.74 a	0.82 a	0.91 a
Control	0.12	0.20 b	0.27 b	0.36 b	0.45 b	0.55 b	0.65 b	0.77 b	0.90 b	1.03 b	1.18 b	1.33 b
F-value	3.37 <sub>ns</sub>	4.07*	6.49*	7.49*	9.23*	12.46*	15.82*	20.61*	27.40*	32.25*	36.85*	64.25*
P value	0.07	0.04	0.01	0.01	0.004	0.002	0.001	0.0003	0.0001	0.00006	0.00003	0.00004
HSD value	-	0.01	0.01	0.02	0.028	0.03	0.04	0.04	0.05	0.05	0.05	0.05

33 Note: ns = not significantly different; \* = significantly different; values within a column followed by the same letters were not  
 34 significantly different at P < 0.05 according to Tukey's HSD test  
 35

36 The decreased appetite of *S. litura* larvae treated with culture filtrate or conidia EPF was one of the stages of the larvae  
 37 to be ill due to the treatment with EPF. In culture filtrate experiments, on day 2 the larvae appetite already decreased and  
 38 was faster on day 1 than the trial using the EPF conidia. In culture filtrate experiments, the decreased appetite was faster  
 39 due to the raw secondary metabolite content of larvae consumed from the first day since the EPF culture filtrate was  
 40 exposed, whereas in conidia treatment it required conidia time to enter the haemolymph and secrete secondary metabolites.  
 41 The process of being ill due to the EPF started with the conidia attached to the cuticle larvae, and then the conidia ruptures  
 42 to form germ tubes to penetrate the cuticle (Fernandes et al. 2007). The infection started when germ tubes were able to  
 43 penetrate the insect cuticle and the ability to infect was a determinant of the level of virulence EPF (Altre and Vandenberg  
 44 2001). After germ tubes reach the haemocoel, they produced specific infection hyphae originating at appressoria (El-

45 Ghany 2015). Then the hyphae spread to the haemolymph and developed to produce blastospores. The blastospores then  
 46 produced the secondary metabolites and enzymes that could disrupt normal cell metabolism (Mancillas-Paredes et al.  
 47 2019). The penetration stages of germ tubes to blastospores produced secondary metabolites and the enzymes occurred in  
 48 the experiments using conidia, whereas in culture filtrate experiments this stage did not occur because the two stages of  
 49 filtration were already carried out to separate conidia from the filtrate. Therefore, the process of the larvae starting to be ill  
 50 in the culture filtrate was faster than the conidia experiment. The next process, the toxins from secondary metabolites  
 51 began to kill the host insect but the death of the host insect was not only by the toxin but also by the mechanical damage of  
 52 the fungal penetration into the body of the insect (El-Ghany 2015).

53 The reduction of the weight of the eaten affected the weight of the larvae. At 1 x 24 hours after the larvae were treated  
 54 with culture filtrate or EPF conidia, the weight of the *S. litura* larvae did not show any significant difference, starting on  
 55 the second day the larvae weight began to be lower than that of the control (Table 4), as well as the third day up to the 12  
 56 day the larvae weight was lower than that of the control. All the tested species of the EPF decreased significantly the  
 57 larvae body weight compared to that of the control. Consequently, the larvae treated with mycoinsecticides containing the  
 58 culture filtrate or EPF conidia could reduce the larvae appetite to make their body weight decrease.

59 Percentage of the mortality or mortality of the larvae caused by the EPF culture filtrate showed significant differences  
 60 when compared to that of the control (Table 5). All EPF species were toxic to *S. litura* larvae, both their culture filtrate and  
 61 conidia. The highest larvae mortality due to culture filtrate was found in the treatment of *P. citrinum* (98.67%) and not  
 62 significantly different from the larvae mortality by *T. diversus* (96%), but significantly different from the larvae mortality  
 63 by *B. bassiana* (86.67%) and *M. anisopliae* (82.67%). The LT<sub>50</sub> larvae treated with culture filtrate *T. diversus* was the  
 64 shortest (6.19 days) and not significantly different from the LT<sub>50</sub> larvae treated with *P. citrinum* (6.62 days), but  
 65 significantly different from the LT<sub>50</sub> larvae treated with *B. bassiana* (7.55 days) and *M. anisopliae* (7.60 days). Thus, the  
 66 most toxic EPF species against *S. litura* larvae was *P. citrinum*. The larvae mortality due to the EPF conidia showed  
 67 significant differences when compared to that of the control (Table 5). All of the EPF species were pathogenic to *S. litura*  
 68 larvae. The larvae mortality caused by the conidia was highest in *P. citrinum* (98.67%) which was not significantly  
 69 different from the larvae mortality by *T. diversus* (97.33%), *B. bassiana* (86.67%), and *M. anisopliae* (86.67%). The LT<sub>50</sub>  
 70 of the larvae treated with the conidia of *P. citrinum* was the shortest (5.45 days) and not significantly different from the  
 71 LT<sub>50</sub> of larvae treated with *T. diversus* (6.55 days), but it was significantly different from the LT<sub>50</sub> of larvae treated with *B.*  
 72 *bassiana* (8.88 days) and *M. anisopliae* (7.70 days). Thus, the most pathogenic EPF species against *S. litura* larvae was *P.*  
 73 *citrinum*.

74  
75  
76 **Table 5.** Mean of larvae mortality and LT<sub>50</sub> of *Spodoptera litura* larvae treated with culture filtrates and conidia of mycoinsecticides  
 77 from *Penicillium citrinum*, *Talaromyces diversus*, *Beauveria bassiana*, and *Metarhizium anisopliae*

Fungal species	Mortality ± SE (%)	LT <sub>50</sub> ± SE (days)
<b>Culture filtrate</b>		
<i>P. citrinum</i>	98.67±1.09 c	6.62±0.135 a
<i>T. diversus</i>	96.00±1.89 c	6.19±0.072 a
<i>B. bassiana</i>	86.67±2.88 b	7.55±0.154 a
<i>M. anisopliae</i>	82.67±1.09 b	7.60±0.203 b
Control	0.00±0.00 a	-
F-value	187.26*	13.12*
P value	0.00000006	0.00480044
HSD value	11.38	0.87
<b>Conidia</b>		
<i>P. citrinum</i>	98.67 ± 1.09 b	5.45 ± 0.232 a
<i>T. diversus</i>	97.33 ± 2.18 b	6.55 ± 0.081 ab
<i>B. bassiana</i>	86.67 ± 1.09 b	8.88 ± 0.420 c
<i>M. anisopliae</i>	86.67 ± 1.09 b	7.70 ± 0.404 bc
Control	0.00 ± 0.00 a	-
F-value	156.31*	29.38*
P value	0.00000013	0.00055
HSD value	12.70	1.24


78 Note: \* = significantly different; values within a column followed by the same letters were not significantly different at P < 0.05  
 79 according to Tukey's HSD test  
 80

81 Both culture filtrate and conidia of EPF in this study had significantly higher insecticidal activity than the control, and  
 82 the highest insecticidal activity was found in *P. citrinum*. *P. citrinum* had the highest insecticidal activity due to its ability  
 83 to infect the larvae faster shown by its LT<sub>50</sub> value which was the shortest compared to the other species. Altre and  
 84 Vandenberg (2001) state that the most pathogenic EPF species due to the fungus has germ tubes that are able to penetrate  
 85 the insect cuticle faster than other fungal species and have more toxins and enzymes that disrupt normal cell metabolism in  
 86 insects. In addition, the color of the *P. citrinum* culture filtrate was the darkest compared to the color of the culture filtrate

87 of the other three EPF species. The darker color of the EPF culture filtrate proved to be more toxic than the lighter color of  
88 the culture filtrate (Ayudya et al., 2019; Gustianingtyas et al., 2020) because the darker the color of the fungus culture  
89 filtrate the higher the concentration of the secondary metabolites (Luo et al., 2017). The *P. citrinum* culture filtrate  
90 contains secondary metabolites and is toxic to insects, for example patulin which can kill *Culex quinquefasciatus* larvae  
91 up to 100% (Maketon et al., 2014). *P. citrinum* are reported to be effective in killing insect pests (Foo et al., 2017;  
92 Maketon et al., 2014) and mite pests (Mazid et al., 2016).


93 The other EPF species were also toxic because they had enzymes or secondary metabolites that could kill the host  
94 insects, for example *B. bassiana* produced the enzymes of proteolytic and chitinolytic that could disrupt normal cell  
95 metabolism (Mancillas-Paredes et al., 2019). The secondary metabolite produced by *B. bassiana* containing the  
96 bassianolone B toxic to the *T. molitor* larvae (Luo et al., 2017). *M. anisopliae* produces secondary metabolites, including  
97 destruxins toxic to insect pests (Borisade et al., 2016). So far there has been no information on the secondary metabolites  
98 or enzymes produced by *T. diversus* that kills insects. At present there is only general information stating that *T. diversus*  
99 can kill insects (Visagie et al., 2009). For this reason, in the future it is necessary to study to identify the secondary  
100 metabolites or *T. diversus* enzymes that can kill insect pests.

101 The *S. litura* larvae treated with EPF culture or conidia showed the same behavior, i.e., they ate less and less and got  
102 lazy to move, while the control not treated with the EPF continued to eat greedily and actively moved. After 3 days the  
103 larvae given culture filtrate began to change integument color that was previously bright green to dull and began to find  
104 dead larvae and the same symptoms occurred in larvae treated with EPF conidia. Four days after the larvae were treated  
105 with culture filtrate or EPF conidia, the larvae's body began to shrink, wrinkle, get odorless and dull black, and eventually  
106 died (Figure 6 and 7). After 2-3 days the larvae died, the symptoms in the larvae integument died due to the conidia  
107 covered by mycelia and the conidia appeared, whereas in larvae that died due to the culture filtrate there was no mycelia  
108 and conidia. Thus, the symptoms of ill *S. litura* larvae caused by the culture filtrate were slightly different from the  
109 symptoms caused by the EPF conidia. The symptoms of sick larvae due to the culture of this filtrate were the same as the  
110 symptoms described by Ayudya et al. (2019) and Gustianingtyas et al. (2020), namely the larvae got shrunk, wrinkled, dry,  
111 odorless, and the integument was not covered by EPF conidia. The symptoms of sick insects caused by EPF conidia in this  
112 study were the same as those described by Bugti et al. (2018) and Sumikarsih et al. (2019), that is, the insect's body got  
113 smaller, wrinkled, dry, odorless, and the integument was covered in mycelia and EPF conidia.



114  
115  
116  
117  
118  
119  
120  
121  
122  
123  
124  
125  
126  
127  
128  
129  
130  
131  
132  
133  
134  
135  
136  
137  
138  
139  
140  
141  
142  
143 **Figure 6.** Morphology of *Spodoptera litura* larvae of healthy larvae of control (right) and dead larvae (left) treated with culture  
144 filtrate of of mycoinsecticides from *Penicillium citrinum* (A), *Talaromyces diversus* (B), *Beauveria bassiana* (C), and *Metarhizium*  
145 *anisopliae* (D)  
146

147  
148  
149  
150  
151  
152  
153  
154  
155  
156  
157  
158  
159  
160  
161  
162  
163  
164  
165  
166  
167  
168  
169  
170  
171  
172  
173  
174  
175  
176  
177  
178  
179  
180  
181  
182  
183  
184  
185  
186  
187  
188  
189  
190  
191  
192  
193  
194  
195  
196  
197  
198  
199  
200  
201  
202  
203  
204  
205  
206



**Figure 7.** Morphology of *Spodoptera litura* larvae of healthy larvae of control (right) and dead larvae (left) treated with conidia of mycoinsecticides from *Penicillium citrinum* (A), *Talaromyces diversus* (B), *Beauveria bassiana* (C), and *Metarhizium anisopliae* (D)

The results of bioassays of culture filtrate and EPF conidia showed that both culture filtrate and conidia had high insecticidal activity. However, for the future development of mycoinsecticides, the use of culture filtrate without conidia as the active ingredient of mycoinsecticide needs to be further investigated. The fungus culture filtrate is relatively more stable than the culture that still has conidia and can be stored for a longer time and remains toxic, besides the culture filtrate does not clog the sprayer nozzle when sprayed (Soesanto et al. 2019).

The EPF species found in the soils of South Sumatra were *P. citrinum*, *T. diversus*, *B. bassiana*, and *M. anisopliae*. All species have a high insecticidal activity against *S. litura* larvae and the most toxic species is *P. citrinum*. The filtrate and conidia of the *P. citrinum* are toxic. *P. citrinum*, a new emerging entomopathogenic fungus, has the highest insecticidal activity and the highest potential to be developed into commercial mycoinsecticides.

#### ACKNOWLEDGEMENTS

This research was funded by the scheme of Applied Research (Riset Terapan) with budget year of 2020 according to Directorate of Research and Community Service (*Direktorat Riset dan Pengabdian Masyarakat*), Ministry of Research and Technology/National Research and Innovation Agency of Republic Indonesia, Grant/Contract Number: SP DIPA-042.06.1.401516/2020.

#### REFERENCES

- Alhusein N, Scott J, Kasprzyk-Hordern B, Bolhuis A. 2016. Development of a filter to prevent infections with spore-forming bacteria in injecting drug users. *Harm Reduct. J.* 13: 1–8. <https://doi.org/10.1186/s12954-016-0122-1>
- Altre JA, Vandenberg JD. 2001. Factors influencing the infectivity of isolates of *Paecilomyces fumosoroseus* against diamondback moth, *Plutella xylostella*. *J. Invertebr. Pathol.* 78: 31–36. <https://doi.org/10.1006/jjpa.2001.5043>
- Anwar W, Khan SN, Aslam M, Haider MS, Shahid AA, Ali M. 2015. Exploring fungal flora associated with insects of cotton agroecological zones of Punjab, Pakistan. *Pakistan Entomol.* 37: 27–31.
- Ayudya DR, Herlinda S, Suwandi S. 2019. Insecticidal activity of culture filtrates from liquid medium of *Beauveria bassiana* isolates from South Sumatra (Indonesia) wetland soil against larvae of *Spodoptera litura*. *Biodiversitas* 20: 2101–2109. <https://doi.org/10.13057/biodiv/d200802>
- Borisade OA, Medina A, Magan N. 2016. Interacting temperature and water activity modulate production of destruxin a by *Metarhizium anisopliae* on galleria larvae-modified agar based media invitro. *West African J. Appl. Ecol.* 24: 31–42.
- Bugti GA, Bin W, Na C, Feng LH. 2018. Pathogenicity of *Beauveria bassiana* strain 202 against sap-sucking insect pests. *Plant Prot. Sci.* 54: 111–117. <https://doi.org/10.17221/45/2017-PPS>
- Chen Z, Xu L, Yang F, Ji G, Yang J, Wang J. 2014. Efficacy of *Metarhizium anisopliae* isolate MAX-2 from Shangri-la, China under desiccation stress. *BBMC Microbiol.* 14, 1–8. <https://doi.org/http://www.biomedcentral.com/1471-2180/14/4>

207 El-Ghany TMA. 2015. Entomopathogenic Fungi and their Role in Biological Control. Biology Department Faculty of Science Jazan University KSA,  
208 Cairo. <https://doi.org/10.4172/978-1-63278-065-2-66>

209 Fernandes EKK, Rangel DEN, Moraes AM, Bittencourt VREP, Roberts DW. 2007. Variability in tolerance to UV-B radiation among *Beauveria* spp.  
210 isolates. J Invertebr Pathol 96: 237–243. <https://doi.org/10.1016/j.jip.2007.05.007>

211 Foo K, Seelan JSS, Dawood MM. 2017. Microfungi associated with *Pteroptyx bearni* (Coleoptera: Lampyridae) eggs and larvae from Kawang River,  
212 Sabah (Northern Borneo). Insects 8: 1–12. <https://doi.org/10.3390/insects8030066>

213 Gustianingtyas M, Herlinda S, Suwandi, Suparman, Hamidson H, Hasbi, Setiawan A, Verawaty M, Elfita, Arsi. 2020. Toxicity of entomopathogenic  
214 fungal culture filtrate of lowland and highland soil of South Sumatra (Indonesia) against *Spodoptera litura* larvae. Biodiversitas 21: 1839–1849.  
215 <https://doi.org/10.13057/biodiv/d210510>

216 Hernandez-Trejo A, Estrada-Drouaillet B, López-Santillán, J, Rios-Velasco C, Varela-Fuentes S, Rodríguez-Herrera R, Osorio-Hernández E. 2019. In  
217 vitro evaluation of native entomopathogenic fungi and neem (*Azadirachta indica*) extracts on *Spodoptera frugiperda*. Int. J. Exp. Bot. 4: 75–84.  
218 <https://doi.org/10.1037/0033-2909.126.1.78>

219 Johari A, Herlinda S, Irsan C, Pujiastuti Y. 2016. Phenomenon of Thrips (Thysanoptera) attack on chili plant (*Capsicum annuum*). Am. J. Agric. Biol.  
220 Sci. 11: 103–109. <https://doi.org/10.3844/ajabssp.2016.103.109>

221 Kumar S, Stecher G, Tamura K. 2016. MEGA7: Molecular evolutionary genetics analysis version 7.0 for bigger data sets. Mol. Biol. Evol. 33: 1870–  
222 1874. <https://doi.org/10.1093/molbev/msw054>

223 Liu SF, Wang GJ, Nong XQ, Liu B, Wang MM, Li SL, Cao GC, Zhang ZH. 2017. Entomopathogen *Metarhizium anisopliae* promotes the early  
224 development of peanut root. Plant Prot. Sci. 53: 101–107. <https://doi.org/10.17221/49/2016-PPS>

225 Lopes RB, Souza DA, Rocha LFN, Montalva C, Luz C, Humber RA, Faria M. 2018. *Metarhizium alvesii* sp. nov.: A new member of the *Metarhizium*  
226 *anisopliae* species complex. J. Invertebr. Pathol. 151: 165–168. <https://doi.org/10.1016/j.jip.2017.12.001>

227 Luo Z, Ren H, Mousa JJ, Rangel DEN, Zhang Y, Bruner SD, Keyhani NO. 2017. The PacC transcription factor regulates secondary metabolite  
228 production and stress response, but has only minor effects on virulence in the insect pathogenic fungus *Beauveria bassiana*. Environ. Microbiol. 19:  
229 788–802. <https://doi.org/10.1111/1462-2920.13648>

230 Maketon M, Amnuaykanjanasin A, Kaysornpug A. 2014. A rapid knockdown effect of *Penicillium citrinum* for control of the mosquito *Culex*  
231 *quinquefasciatus* in Thailand. World J. Microbiol. Biotechnol. 30: 727–736. <https://doi.org/10.1007/s11274-013-1500-4>

232 Mancillas-Paredes J, Hernández-Sánchez H, Jaramillo-Flores ME, García-Gutiérrez C. 2019. Proteases and chitinases induced in *Beauveria bassiana*  
233 during infection by *Zabrotes subfasciatus*. Southwest. Entomol 44: 125–137. <https://doi.org/10.3958/059.044.0114>

234 Mascarin GM, Jackson MA, Kobori NN, Behle RW, Delalibera Júnior Í. 2015. Liquid culture fermentation for rapid production of desiccation tolerant  
235 blastospores of *Beauveria bassiana* and *Isaria fumosorosea* strains. J. Invertebr. Pathol. 127: 11–20. <https://doi.org/10.1016/j.jip.2014.12.001>

236 Mathew AJ, Jayachandran K, Mathew J. 2010. Endophytic *Penicillium citrinum* Thom. from *Scoparia dulcis* Linn. Indian J. Microbiol. 50: 99–102.  
237 <https://doi.org/10.1007/s12088-010-0065-2>

238 Mazid S, Kalita JC, Rajkhowa RC. 2016. Biocontrol potential of *Penicillium citrinum* and *Penicillium chrysogenum* against red spider mite, *Oligonychus*  
239 *caffaeae* Nietner infesting tea. J. Entomol. Res. <https://doi.org/10.5958/0974-4576.2016.00008.6>

240 Ngangambe MH, Mwatawala MW. 2020. Effects of entomopathogenic fungi (EPFs) and cropping systems on parasitoids of fall armyworm (*Spodoptera*  
241 *frugiperda*) on maize in eastern central, Tanzania. Biocontrol Sci. Technol. 30: 1–13. <https://doi.org/10.1080/09583157.2020.1726878>

242 Safitri A, Herlinda S, Setiawan A. 2018. Entomopathogenic fungi of soils of freshwater swamps, tidal lowlands, peatlands, and highlands of South  
243 Sumatra, Indonesia. Biodiversitas 19: 2365–2373. <https://doi.org/10.13057/biodiv/d190647>

244 Sharma L, Oliveira I, Torres L, Marques G. 2018. Entomopathogenic fungi in Portuguese vineyards soils: suggesting a ‘Galleria-Tenebrio-bait method’  
245 as bait-insects *Galleria* and *Tenebrio* significantly underestimate the respective recoveries of *Metarhizium (robertsii)* and *Beauveria (bassiana)*.  
246 MycoKeys 23: 1–23. <https://doi.org/10.3897/mycokeys.38.26970>

247 Siaga E, Lakitan B, Hasbi H, Bernas SM, Widuri LI, Kartika K. 2019. Floating seedbed for preparing rice seedlings under unpredictable flooding  
248 occurrence at tropical riparian wetland. Bulg. J. Agric. Sci. 25: 326–336.

249 Soesanto L, Mugiastuti E, Manan A. 2019. Raw secondary metabolites application of two *Trichoderma harzianum* isolates towards vascular streak  
250 dieback on cocoa seedlings. Pelita Perkeb. 35, 22–32. <https://doi.org/10.22302/iccri.jur.pelitaperkebunan.v35i1.346>

251 Souza ML, Sanches MM, Souza DAde, Faria M, Espinel-Correal C, Sihler W, Lopes RB. 2019. Within-host interactions of *Metarhizium rileyi* strains  
252 and nucleopolyhedroviruses in *Spodoptera frugiperda* and *Anticarsia gemmatilis* (Lepidoptera: Noctuidae). J. Invertebr. Pathol. 162: 10–18.  
253 <https://doi.org/10.1016/j.jip.2019.01.006>

254 Sowmya G, Ramanagouda SH, Jayappa J. 2017. Lethal and sub lethal effects of indigenous isolates of *Metarhizium anisopliae* (Metchnikoff) Sorokin  
255 against *Spodoptera litura* (Fabricius). Entomol. Zool. Stud. 5, 1154–1160.

256 Sumikarsih E, Herlinda S, Pujiastuti Y. 2019. Conidial density and viability of *Beauveria bassiana* isolates from Java and Sumatra and their virulence  
257 against *Nilaparvata lugens* at different temperatures. Agrivita J. Agric. Sci. 41: 335–349. <http://doi.org/10.17503/agrivita.v41i2.2105>

258 Suwandi, Ammar M, Chandra I. 2012. Application of extract compost increased yield and suppressed the diseases of ratoon rice crop in tidal swamp of  
259 Banyuasin Regency. J. Suboptimal Lands 1: 116–122. [Indonesia] <https://doi.org/10.33230/JLSO.1.2.2012.15>

260 Syamsudin TS, Faizal A, Kirana, R. 2019. Dataset on antixenosis and antibiosis of chili fruit by fruit fly (*Bactrocera dorsalis*) infestation Tati. Data Br.  
261 23: 1–4. <https://doi.org/10.1016/j.dib.2019.103758>

262 Thaochan N, Sausa-Ard W. 2017. Occurrence and effectiveness of indigenous *Metarhizium anisopliae* against adults *Zeugodacus cucurbitae* (Coquillett)  
263 (Diptera: Tephritidae) in Southern Thailand. Songklanakarin J. Sci. Technol. 39: 325–334. <https://doi.org/10.14456/sjst-psu.2017.35>

264 Tompe AA, Hole UB, Kulkarni SR, Chaudhari CS, Chavan SK. 2020. Studies on seasonal incidence of leaf eating caterpillar, *Spodoptera litura* (Fab.)  
265 infesting capsicum under polyhouse condition. J. Entomol. Zool. Stud. 8: 761–764.

266 Vijayalakshmi P, Vijayalakshmi T, Naidu LN. 2016. Evaluation of certain insecticide molecules against chillipod borer, *Spodoptera litura* in Andhra  
267 Pradesh. J. Res. ANGRAU 44: 26–30.

268 Visagie CM, Jacobs K. 2012. Three new additions to the genus *Talaromyces* isolated from Atlantis sandveld fynbos soils. Persoonia 28: 14–24.  
269 <https://doi.org/10.3767/003158512X632455>

270 Visagie CM, Roets F, Jacobs K. 2009. A new species of *Penicillium*, *P. ramulosum* sp. nov., from the natural environment. Mycologia 101: 888–895.  
271 <https://doi.org/10.3852/08-149>

272 White T, Bruns T, Lee S, Taylor J. 1990. Amplification and direct sequencing of fungal ribosomal RNA genes for phylogenetics, in: Innis, M.A.,  
273 Gelfand, D.H., Sninsky, J.J., White, T.J. (Eds.), PCR Protocols: A Guide to Methods and Applications. Academic Press Inc., New York.

274 Yang H, Qin CS, Chen YM, Zhang GY, Dong LH, Wan SQ. 2019. Persistence of *Metarhizium* (Hypocreales: Clavicipitaceae) and *Beauveria bassiana*  
275 (Hypocreales: Clavicipitaceae) in tobacco soils and potential as biocontrol agents of *Spodoptera litura* (Lepidoptera: Noctuidae). Environ. Entomol.  
276 48: 147–155. <https://doi.org/10.1093/ee/nvy161>

277 Youssef NA. 2014. Effect of certain entomopathogenic fungi and nematode on the desert locust *Schistocerca gregaria* (Forsk.). Ann. Agric. Sci. 59,  
278 125–131. <https://doi.org/10.1016/j.aos.2014.06.017>

279

## SUBMISSION CHECKLIST

Ensure that the following items are present:

The first corresponding author must be accompanied with contact details:

**Give mark (X)**

• E-mail address	X
• Full postal address (incl street name and number (location), city, postal code, state/province, country)	X
• Phone and facsimile numbers (incl country phone code)	X


All necessary files have been uploaded, and contain:

• Keywords	X
• Running titles	X
• All figure captions	X
• All tables (incl title and note/description)	X

Further considerations

• Manuscript has been “spell & grammar-checked” Better, if it is revised by a professional science editor or a native English speaker	X
• References are in the correct format for this journal	X
• All references mentioned in the Reference list are cited in the text, and vice versa	X
• Colored figures are only used if the information in the text may be losing without those images	X
• Charts (graphs and diagrams) are drawn in black and white images; use shading to differentiate	X

## 2. Bukti konfirmasi review pertama dan hasil revisi pertama



## New emerging entomopathogenic fungi isolated from soil in South Sumatra (Indonesia) and their filtrate and conidial insecticidal activity against *Spodoptera litura*

**Abstract.** This study was carried out to confirm identity of *Beauveria bassiana* and *Metarhizium anisopliae* using sequence analysis of Internal Transcribed Spacer (ITS) region as well as the two unknown isolates of EPF and to investigate their filtrate and conidial insecticidal activity against *S. litura* larvae. The EPF species were identified based on morphological characters and molecular analysis. Their culture filtrate and conidial bioassays were performed on *S. litura* larvae. The species found were *Penicillium citrinum* (BKbTp), *Talaromyces diversus* (MSwTp1), *B. bassiana* (BSwTd4), and *M. anisopliae* (MSwTp3) and have been deposited in the GenBank. All EPF species have a high insecticidal activity against *S. litura* larvae. The highest larvae mortality caused by filtrate was found on *P. citrinum* (98.67%) and not significantly different from those of *T. diversus* (96%), but significantly different from *B. bassiana* (86.67%) and *M. anisopliae* (82.67%). The highest larvae mortality caused by the conidia was also occurred to *P. citrinum* (98.67%) which was not significantly different from those of *T. diversus* (97.33%), *B. bassiana* (86.67%), and *M. anisopliae* (86.67%). The  $LT_{50}$  caused by *T. diversus* filtrate was the shortest (6.19 days) and not significantly different from those caused by *P. citrinum* (6.62 days). The shortest  $LT_{50}$  caused by conidia was found on *P. citrinum* (5.45 days). Therefore, the conidia and filtrate of *P. citrinum*, a new emerging entomopathogenic fungus, has the highest insecticidal activity. *P. citrinum* has the potential to be developed into commercial mycoinsecticides.

**Key words:** *Beauveria bassiana*; leaf eating caterpillar; *Metarhizium anisopliae*; *Penicillium citrinum*; *Talaromyces diversus*

**Abbreviations** (if any): -



## INTRODUCTION

South Sumatra, Indonesia has a stretch with varying topography from the height of the land to the high ground. This topographic variation can develop many species of plants, for example vegetables and the food crops. Vegetables that can adapt to various land topography in South Sumatra include chili (*Capsicum* spp.), mustard greens (*Brassica juncea*), kale (*Ipomoea reptans*), and spinach (*Amaranthus tricolor*) (Siaga et al. 2019). The main constraints of vegetable cultivation, especially chili pests are fruit flies (*Bactrocera dorsalis*) (Syamsudin et al. 2019), sap sucker (Thrips) (Johari et al. 2016), and leaf eating caterpillar (*Spodoptera litura*) (Tompe et al. 2020).

*S. litura* is a polyphagous insect pest that attacks many plant species from various families (Tompe et al. 2020), like chili, mustard greens, water spinach, and spinach. The yield loss of chili caused by this pest reaches 50% every year (Vijayalakshmi et al. 2016). The first instar of *S. litura* larvae eats the leaves in groups and skeletonize them, while the second instar extend to the last instar spread and eat individually leaves, flowers and pistil chili (Tompe et al. 2020). In addition, *S. litura* is a key pest for the leafy vegetables, such as mustard greens, water spinach, and spinach because the leaves that have been eaten become no longer valuable (Vijayalakshmi et al. 2016; Tompe et al. 2020). Since the public awareness on the good health and environmental quality is increasing, there is a need to reduce the use of synthetic pesticides and improving application of eco-friendly control strategies for *S. litura*. One of the methods is applying the entomopathogenic fungi (EPF).

Some EPF species have been shown to be effective in killing *S. litura*, for example, *Beauveria bassiana* cultured in liquid media has been shown to kill 98% of *S. litura* larvae (Ayudya et al. 2019; Gustianingtyas et al. 2020). *Metarhizium anisopliae* was reportedly also effective in killing *S. litura* larvae (Sowmya et al., 2017; Yang et al., 2019; Gustianingtyas et al. 2020) and can kill 96% of *S. litura* larvae (Gustianingtyas et al. 2020). Furthermore, *Metarhizium rileyi* also effectively kills *S. litura* larvae (Liu et al. 2017). In addition to being effective in killing the insect pests, EPF can also be integrated with entomopathogenic viruses, such as nucleopolyhedroviruses (NPV) (Souza et al. 2019) and botanical insecticides (Hernandez-Trejo et al. 2019) and does not harm parasitoids (Ngangambe and Mwatawala 2020).

Four isolates of EPF have been isolated from rhizosphere of several kinds of plant in lowland to high land of South Sumatra. On the basis of morphological characters, the two isolates have been identified as *B. bassiana* and *M. anisopliae* (Safitri et al. 2018; Sumikarsih et al. 2019; Gustianingtyas et al. 2020) and their filtrates were proven to be effectively killed up to 98% of *S. litura* larvae (Ayudya et al. 2019; Gustianingtyas et al. 2020). The other two EPF, until now, have not been fully revealed on molecular identity and their filtrate and conidial capability to cause mortality on *S. litura*. This research aimed to confirm identity of *Beauveria bassiana* and *Metarhizium anisopliae* using sequence analysis of Internal Transcribed Spacer (ITS) region as well as the two unknown isolates of EPF and to investigate their filtrate and conidial insecticidal activity against *S. litura* larvae.

## MATERIALS AND METHODS

### Exploring EPF

The EPF exploration was carried out on soil near root of oil palm, rice, maize and vegetables in various locations (Table 1) ranging from lowland to highland of South Sumatra using the method of Safitri et al. (2018). Those locations chosen have objective to obtain the fungi that had adapted in the soils. The morphological identifications were carried out in 2018 and 2019, and their identities were confirmed by molecular analysis. The EPF exploration method modified those of Anwar et al. (2015), namely using a bait method with *Tenebrio molitor* larvae as insect bait which was fed in the soil sample. The soil sampling used the method of Anwar et al. (2015) by digging the soil using surface soil sampler ( $\varnothing$  20 mm) as deep as 10 cm around the rhizosphere plant according to the method of Thaochan and Sausa-Ard (2017) and 5 points were taken to collect 1 kg of soil. Then, the soil sample was put into a plastic bag and labeled with information about the height of the location, type of commodity, and date of collection.

The soil sample was first cleaned from the roots and sieved with a 10 mesh sieve. Then, it was put into a plastic tray (32 x 25 x 5 cm<sup>3</sup>) each containing as much as 1 kg. The soil was then moistened with sterile distilled water with soil moisture exceeding 20% according to the method of Chen et al. (2014). Then, 20 newly molted 3rd-instar larvae of *T. molitor* were placed at the bottom of the tray and the larvae were sprinkled with the soil to reach 3 cm in thickness. Then, the tray containing the soil was covered with a black cloth and sprayed with distilled water to maintain the soil moisture. The larvae were infested in the soil for 7 days to give time for the entomopathogenic fungal conidia to infect them, after which the dead larvae infected with entomopathogenic fungi were grown in the Sabouraud Dextrose Agar (SDA, Merck) medium.

### Isolation and identification of EPF

The EPF infecting the larvae of *T. molitor* which was characterized by its body was overgrown with EPF mycelia, dry and contracted, then isolated and purified using the method of Youssef (2014). The isolation of EPF from the larvae's body surface began with the surface sterilization by modifying the method of Sharma et al. (2018) by rinsing with 1% of NaOCl for 1 minute, then rinsing with 100 mL of sterile distilled water for 3 times. The larvae sterilization was carried out to obtain EPF inside the

haemocoel and cuticle larvae and prevent water fungi from being obtained. The sterilized larvae were grown in SDA media and incubated for 2 days. The growing fungus was purified in the SDA media until after getting pure isolates. The pure EPF was identified based on the molecular analysis.

### Fungal identification

Morphological observations were carried out based on microscopic and macroscopic characteristics of the EPF. The macroscopic observation of the fungi was based on the color and shape of colony, whereas the microscopic observation was based on the shape and size of conidia and conidiophores.

The EPF were identified based on the molecular analysis at the Laboratory of Agricultural Biotechnology (accredited according to the ISO/IEC 17025 standard), Department of Plant Protection, Faculty of Agriculture, University of Lampung, Indonesia.

### DNA extraction and PCR amplification

The DNA extraction was performed on the mycelium of 7 days old fungi using InstaGene™ Matrix (BioRad). Amplification was carried out by Sensoquest Thermal Cycler Machine (Germany). PCR amplification on Internal Transcribed Spacer (ITS) region was conducted using primer ITS1 and ITS4 (Table 2) (White et al. 1990). The PCR was carried out with total volume of 25 µL involving 12.5 µL Master Mix (Red Mix) (bioline), 1 µL of 10 µM of primer ITS 1 and ITS 4, 2 µL DNA and 8.5 µL of sterile distilled water. DNA amplification consist of 1 cycle of the initiation at 95°C for 5 min, 30 cycles of denaturation at 95°C for 1 minute, primer annealing at 52°C for 1 min, primer extension at 72°C for 1 minute, and one cycle of elongation at 72°C for 5 minutes. The PCR results were run using 0.5% agarose gel in 20 mL of 1x buffer Tris-Boric Acid-EDTA (TBE) (1<sup>st</sup> Base Malaysia) with 1 µL Ethidium Bromide (EtBr 10 mg mL<sup>-1</sup>). The electrophoresis was performed by 1x TBE buffer at 50 volts for 70 minutes. The results were visualized by DigiDoc UV transilluminator (UVP, USA).

**Table 1.** Locations of EPF in South Sumatra, Indonesia

Fungal species (codes)	Village, District/City	Ecosystem	Crop	GPS (coordinat)	Altitude (m)	GenBank Acc. No.
<i>Penicillium citrinum</i> (BKbTp)	Talang Patai, Pagar Alam	Highland soil	Cabbage	S 04°02.161' E 103°10.484'	175	MT448730
<i>Talaromyces diversus</i> (MSwTp1)	Talang Patai, Pagar Alam	Highland soil	Mustard	S 03°50.174', E 103°31.293'	193	MT448731
<i>Beauveria bassiana</i> (BSwTd4)	Talang Dabok, Ogan Komering Ilir	Peat soil	Oil palm	S 03°23.570' E 104°51.498'	24	MT448732
<i>Metarhizium anisopliae</i> (MSwTp3)	Talang Patai, Pagar Alam	Highland soil	Mustard	S 03°50.174', E 103°31.293'	193	MT448733

**Table 2.** Primer used for amplification

Target of amplification	Primer name	Sequence (5'-3')	Product size (bp)	Reference
Internal Transcribe Spacer Region	ITS1	TCC GTA GGT GAA CCT TGC GG	600	(White et al. 1990)
	ITS4	TCC TCC GCT TAT TGA TAT GC		

The PCR product was emitted to 1<sup>st</sup> Base Malaysia for sequencing. The sequencing results were analyzed with Bio Edit program ver. 7.2.6 for windows. The result was then submitted to Basic Local Alignment Search Tool (BLAST) (<https://blast.ncbi.nlm.nih.gov/Blast.cgi>) to find their potential identity. The phylogenetic tree was developed by Mega 7 for Windows (Kumar et al. 2016) with Unweighted-pair Group Method with Arithmetic means (UPGMA). Reference strains employed in this research were obtained from NCBI (<https://www.ncbi.nlm.nih.gov/>).

### *Spodoptera litura* mass rearing

*S. litura* larvae used in the insecticide activity test was collected from the vegetable fields around the campus of Universitas Sriwijaya and those vegetable fields that were not sprayed with synthetic insecticides. The larvae were taken to the laboratory and maintained in plastic containers (Ø 25 cm, height 30 cm) topped with gauze. that has been washed thoroughly using detergent. Next, pesticide-free spinach (*Ipomoea aquatic*) was put into the containers for the first instar feed of *S. litura*. The second instar was fed mulberry leaves (*Morus alba*) planted in the experimental field of the Faculty of Agriculture, Universitas Sriwijaya, Indonesia and not sprayed with pesticides.

Every day the larvae feed was replaced with fresh new feed. The larvae entering the pupae stage were transferred into plastic containers (Ø 10 cm, height 15 cm) whose top was exposed and the base was sprinkled with sifted soil (10 mesh). and already sterilized in an oven for 1 hour at 100°C. The thickness of the soil in the container was 3 cm. This plastic container containing pupae was put into a gauze cage (30 x 30 x 30 cm<sup>3</sup>) already tapped with 10 water spinach sticks for laying eggs to the adults. The hatched eggs were transferred into a plastic cylinder and fed with water spinach which was replaced every day. The larvae used for toxicity testing were the second instar of the third offspring (F3) or afterwards.

### Producing of EPF mycoinsecticides

The EPF already identified molecularly, namely *P. citrinum*, *T. diversus*, *B. bassiana*, and *M. anisopliae* were made liquid formulation mycoinsecticides. The mycoinsecticides made of four types, each of which had active ingredients *P. citrinum*, *T. diversus*, *B. bassiana*, and *M. anisopliae* and sterile distilled water was used for control. The mycoinsecticides were made by the method of Mascarin et al. (2015). In this, EPF was cultivated in solid media of Sabouraud Dextrose Agar (SDA, Merck), incubated for 7 days, and then the fungal cultures that grew were transferred to the liquid media of Sabouraud Dextrose Broth (SDB, Merck). The broth fungal cultures were incubated using a shaker for 7 days at 120 rpm. The broth fungal cultures from SDB with a minimum density of  $1 \times 10^9$  conidia  $\text{mL}^{-1}$  were used for the active ingredient mycoinsecticides. The mycoinsecticides were made liquid formulation containing a mixture of 600 mL of the broth fungal cultures, 400 mL of carrier from shrimp shell flour compost extract, 10 mL of vegetable oil, and 10 mL of sterile distilled water solution of 0.04% polyoxyethyl-189 ene sorbitan mono-oleate (Tween®). The shrimp skin flour compost extract was made using the method of Suwandi et al. (2012). After that, the formulation was stored for 30 days before it was applied to obtain stable and non-contaminant mycoinsecticides. The mycoinsecticides activity was tested in the form of EPF culture and conidia filtrate.

#### **Culture filtrate preparation of the mycoinsecticides.**

The presence of raw secondary metabolites in culture filtrate was performed according to Soesanto et al. (2019) can be done by filtering fungal broth or liquid mycoinsecticides. The culture filtrate of liquid mycoinsecticides was produced from 2 stages of filtering using the method of Ayudya et al. (2019). The first stage, 100 mL liquid mycoinsecticides was filtered using Whatman filter paper no. 42 and cotton with a thickness of 1 cm. The filtration culture produced from the first stage filter was then sucked up to 10 mL using a sterile hypodermic syringe to be filtered in the second stage. The second stage filtering used a syringe filter (0.45  $\mu\text{m}$  pore size, Sartorius). The method of installing the syringe filter used the method of Alhusein et al. (2016) by placing it at the base of the needle, by the way the needle was removed first from the syringe, then the syringe filter was installed at the base of the needle. After the syringe filter was installed at the base of the needle, the needle was re-fitted to the syringe. The syringe containing 10 mL was pressed so that it released the culture filtrate that escaped from the syringe filter. The 100 mL of mycoinsecticide produced  $\pm 70$  mL of culture filtrate. According to Soesanto et al. (2019), this second filtering culture filtrate was the raw secondary metabolites. Next, the culture filtrate was measured by its insecticidal activity by taking as much as 1 mL to be dropped on a piece of chili leaves.

#### **Bioassays of culture filtrate and conidia of EPF**

Bioassays of culture filtrate from mycoinsecticides of *P. citrinum*, *T. diversus*, *B. bassiana*, and *M. anisopliae* were carried out on chili leaves sprayed with 1 mL of culture filtrate. As a control, sterile distilled water was dripped on chili leaves. Then the leaves were dried before putting 25 second-instar *S. litura* larvae which were fasted for 2 hours. The second instar larvae were let to eat the leaves for 6 hours which had been dropped by the culture filtrate. After 6 hours the larvae were transferred into a plastic cylinder topped with gauze ( $\varnothing$  25 cm, height 30 cm) containing 15-30 pieces of chili leaves (depending on the instar) which were clean and not dripped with culture filtrate. To maintain the moisture of chili leaves, the stalk was wrapped in cotton wool moistened with sterile distilled water. Every day chili leaves were replaced with new fresh leaves. The weight of the leaves eaten was measured according to the method of Ayudya et al. (2019) and recorded daily. Every day the dead larvae were recorded up to 12 days after the application.

Bioassays of conidia from mycoinsecticides of *P. citrinum*, *T. diversus*, *B. bassiana*, and *M. anisopliae* were carried out by means of 25 second instar *S. litura* larvae dripped topically 1 mL of conidia suspension of EPF with a concentration of  $1 \times 10^6$  conidia  $\text{mL}^{-1}$ . Then, the larvae were fed 15-30 pieces of chili leaves. As a control it was used sterile distilled water. The weight of leaves eaten by larvae and weight of the larvae were recorded daily. Every day the dead larvae were recorded up to 12 days after the application.

The number of dead larvae was used to calculate mortality and the median Lethal Time ( $\text{LT}_{50}$ ). The weight of the chili leaves eaten and the body weight of the larvae were measured every day from the first day to 12 days after the application. The measurement of the eaten leaf weight was conducted to determine whether the EPF affected the larvae appetite, whereas the measurement of larvae body weight was carried out to determine whether or not the EPF influenced the larvae growth.

#### **Data Analysis**


The data differences among the treatments for edible leaf weight, larvae weight, larvae mortality, and  $\text{LT}_{50}$  were analyzed using Analysis of Variance (ANOVA) if there were differences, then further tests were carried out using the 5% Tukey's Honestly Significant Difference (HSD).  $\text{LT}_{50}$  data were obtained from probit analysis. All calculations were conducted using software of SAS University Edition 2.7 9.4 M5.

## **RESULTS AND DISCUSSION**

#### **Species of EPF**

There were 4 EPF species namely, *P. citrinum*, *T. diversus*, *B. bassiana*, and *M. anisopliae* found from lowland and highland soils in South Sumatra. The macroscopic and microscopic appearance of the four fungal species showed obvious differences (Figure 1 and 2). The colony characteristics of *P. citrinum* on SDA medium were initially white, the older the fungus, the color turned to bluish green with a white margin, round conidia with a diameter of 2.01-2.28  $\mu\text{m}$ . The characteristics


matched the description **described** by Mathew et al. (2010) stating that *P. citrinum* has globose green granular conidia. *T. diversus* has a colony characteristic generally in the form of microcolony scattering in the medium; the color was dark green with yellow margins. According to Visagie and Jacobs (2012), the size of *T. diversus* colony ranged from 2-4 mm and sometimes only microcolony and produced yellow colored mycelia at colony peripheral zones. In this study, *T. diversus* was more difficult to culture than the other three species and showed poor growth of colony on SDA. *B. bassiana* had a colony that was white but when it got older it was yellowish or turbid white and *B. bassiana* colonized SDA faster than the other three species. Conidia of *B. bassiana* was one-celled and the same circular shape as the observation of Safitri et al. (2018). *M. anisopliae* had a colony which was also white initially similar to *B. bassiana* colony but the older the fungus the colony turned greenish white and finally dark green. Conidia of *M. anisopliae* was 6.42-6.76  $\mu\text{m}$  in size. The found characteristics of the colony and conidia of *M. anisopliae* were the same as those described by Lopes et al. (2018) and Thaochan and Sausa-Ard (2017) stating that the colony is more than 5 days old, the conidia changes from white to green indicating that the conidia was matured with the conidial mass bluish olive and has cylindrical, single and one-celled, hyaline conidia.



A B

C D


**Figure 1.** Colony of *Penicillium citrinum* (A), *Talaromyces diversus* (B), *Beauveria bassiana* (C), and *Metarhizium anisopliae*



**Figure 2.** Conidia of *Penicillium citrinum* (A), *Talaromyces diversus* (B), *Beauveria bassiana* (C), and *Metarhizium anisopliae* (D)

The result of BLAST search revealed that BKbTp showed 99.38% of similarity with *P. citrinum* strain NFML\_CH33\_5BbE.1 (Acc. No. KM458810.1), MSwTp1 showed 99.05% of similarity with *T. diversus* strain DTO\_13116 (Acc. No. KJ775700.1), BSwTd4 showed 99.65% of similarity with *B. bassiana* isolate GZMS-28 (Acc. No. KT715480.1) and MSwTp3 exhibited 99.82% of similarity with *M. anisopliae* isolate HT008 (Acc. No. MH512951.1). The result of BLAST search corresponded to those of phylogenetic tree analysis. The phylogenetic tree (Figure 3) was constructed using Mega 7 for Windows (Kumar et al. 2016) with Unweighted-pair Group Method with Arithmetic means (UPGMA). Based on the phylogenetic tree, BKbTp was placed in the group of *P. citrinum* strain NFML\_CH33\_5BbE.1 (Acc. No. KM458810.1), and SL1 (Acc. No. MN735924.1). MSwTp1 was in the group of *T. diversus* strain DTO\_13116 (Acc. No. KJ775700.1) and strain SCZ5-2 (Acc. No. KY445783.1). BSwTd4 was in the group of *B. bassiana* isolate CGAIPFBS-012 (Acc. No. KY495188.1),

isolate 08F04 (Acc. No. KF359944.1), and isolate GZMS-28 (Acc. No. KT715480.1). Meanwhile, MSwTp3 was in the group of *M. anisopliae* HT008 (Acc. no. MH512951.1), A1 (Acc. No. KX806656.1), and IPPM010202 (Acc. No. KY437678.1). Thus, based on the morphological characters and molecular analysis, it confirmed that the identity of the four isolates were *P. citrinum* (BKbTp), *T. diversus* (MSwTp1), *B. bassiana* (BSwTd4), and *M. anisopliae* (MSwTp3). The isolates have been deposited in the GenBank and the accession number were MT448730 (BKbTp) (<https://www.ncbi.nlm.nih.gov/nuccore/MT448730>), MT448731 (MSwTp1) (<https://www.ncbi.nlm.nih.gov/nuccore/MT448731>), MT448732 (BSwTd4) (<https://www.ncbi.nlm.nih.gov/nuccore/MT448732>), and MT448733 (MSwTp3) (<https://www.ncbi.nlm.nih.gov/nuccore/MT448733>).



**Figure 3.** Dendrogram created on the basis of sequence analysis of ITS region using primer ITS1 and ITS4 with UPGMA method developed by MEGA7 for windows ■ = fungal isolates used in this study

**EPF insecticidal activity against *Spodoptera litura***

The EPF insecticidal activity against *S. litura* larvae was stated from the data of the effect of EPF on the weight reduction of edible chili leaves, body weight of larvae, mortality, and length of time of death. On day 1 (1 x 24 hours) after the larvae were treated with the filtrate culture of the EPF, the weight of the leaves eaten by *S. litura* larvae did not show any significant difference compared to control (sterile distilled water), but after that the weight of the leaves eaten by the larvae was more lower than control (Table 3). For the larvae treated with conidia of the EPF on day 1 (1 x 24 hours) and day 2 (2 x 24 hours), the weight of the eaten leaves did not show any significant difference compared to control, on the following day the weight of the eaten leaves by the larvae was significantly lower than those of the control.

**Table 3.** Mean of leaf weight eaten by *Spodoptera litura* larvae treated with culture filtrates and conidia of mycoinsecticides from *Penicillium citrinum*, *Talaromyces diversus*, *Beauveria bassiana*, and *Metarhizium anisopliae*

Fungal species	Leaf weight eaten by larvae (g larvae <sup>-1</sup> day <sup>-1</sup> ) on observation for 12 days											
	1	2	3	4	5	6	7	8	9	10	11	12
<b>Culture filtrate</b>												
<i>P. citrinum</i>	0.10	0.26 a	0.40 a	0.51 a	0.62 a	0.73 a	0.82 a	0.96 a	1.08 a	1.17 a	1.22 a	1.24 a
<i>T. diversus</i>	0.14	0.35 a	0.46 a	0.55 a	0.65 a	0.77 a	0.86 a	0.99 a	1.09 a	1.18 a	1.23 a	1.25 a
<i>B. bassiana</i>	0.17	0.36 a	0.47 a	0.58 a	0.74 a	0.94 a	1.08 a	1.22 a	1.35 ab	1.45 ab	1.53 ab	1.59 b
<i>M. anisopliae</i>	0.17	0.36 ab	0.53 a	0.65 a	0.78 a	0.97 a	1.11 a	1.26 a	1.42 b	1.55 b	1.65 ab	1.77 b
Control	0.24	0.56 b	0.94 b	1.32 b	1.74 b	2.33 b	2.91 b	3.45 b	4.02 c	4.65 c	5.04 c	5.58 c
F-value	1.96 <sup>ns</sup>	6.86*	10.28*	19.63*	44.96*	76.82*	124.79*	131.90*	203.70*	263.92*	281.95*	324.89*
P value	0.19	0.011	0.0031	0.00034	0.000016	0.000020	0.000031	0.000025	0.000044	0.000016	0.000012	0.000070
HSD value	-	0.10	0.14	0.15	0.12	0.12	0.12	0.13	0.117	0.115	0.119	0.122
<b>Conidia</b>												
<i>P. citrinum</i>	0.45	0.76	1.13 a	1.50 a	1.84 a	2.00 a	2.28 a	2.47 a	2.62 a	2.74 a	2.85 a	2.92 a
<i>T. diversus</i>	0.45	0.84	1.17 a	1.50 a	1.87 a	2.15 a	2.51 a	2.71 a	2.94 a	3.04 a	3.19 a	3.28 a
<i>B. bassiana</i>	0.44	0.76	1.09 a	1.43 a	1.76 a	2.07 a	2.40 a	2.59 a	2.87 a	3.01 a	3.21 a	3.45 a
<i>M. anisopliae</i>	0.34	0.70	1.08 a	1.39 a	1.78 a	2.10 a	2.35 a	2.54 a	2.72 a	2.86 a	3.01 a	3.23 a
Control	0.56	1.28	1.97 b	2.64 b	3.33 b	3.99 b	4.99 b	5.73 b	6.57 b	7.49 b	8.32 b	9.16 b
F-value	1.78 <sup>ns</sup>	3.58 <sup>ns</sup>	7.14 *	18.26 *	16.06 *	17.38 *	14.98 *	14.07 *	17.72 *	23.27 *	26.81 *	33.23 *
P value	0.23	0.06	0.01	0.0004	0.0007	0.0005	0.0011	0.0011	0.0005	0.0002	0.0001	0.00005
HSD value			0.23	0.18	0.23	0.26	0.35	0.41	0.42	0.42	0.43	0.42

Note: ns = not significantly different; \* = significantly different; values within a column followed by the same letters were not significantly different at P < 0.05 according to Tukey's HSD test



**Figure 4.** The symptoms on chili leaves eaten by *Spodoptera litura* larvae treated with culture filtrates of mycoinsecticides and control

The symptoms of leaf infested with larvae treated with filtrate or conidia culture showed less damage, whereas, the control leaves left only the bone of the leaf (Figure 4 and 5). The decreased appetite in leaves treated with the culture filtrate of the EPF was faster than in leaves treated with the conidia of the EPF. All tested species of the EPF significantly reduced the weight of leaves eaten by larvae compared to controls (Table 3). Thus, all species of the EPF (*P. citrinum*, *T. diversus*, *B. bassiana*, and *M. anisopliae*) were effective in reducing the appetite of *S. litura* larvae.



**Figure 5.** The symptoms on chili leaves eaten by *Spodoptera litura* larvae treated with conidia of mycoinsecticides and control

**Table 4.** Mean of weight of *Spodoptera litura* larvae treated with culture filtrates and conidia of mycoinsecticides from *Penicillium citrinum*, *Talaromyces diversus*, *Beauveria bassiana*, and *Metarhizium anisopliae*

Fungal species	Larvae weight (g larvae <sup>-1</sup> ) on observation for 12 days											
	1	2	3	4	5	6	7	8	9	10	11	12
<b>Culture filtrate</b>												
<i>P. citrinum</i>	0.06	0.09 a	0.12 a	0.15 a	0.19 a	0.22 a	0.25 a	0.29 a	0.34 ab	0.40 a	0.48 ab	0.50 a
<i>T. diversus</i>	0.06	0.09 a	0.12 a	0.15 a	0.18 a	0.22 a	0.27 a	0.31 a	0.38 b	0.46 b	0.53 b	0.58 b
<i>B. bassiana</i>	0.06	0.09 a	0.12 a	0.15 a	0.18 a	0.21 a	0.25 a	0.29 a	0.33 a	0.39 a	0.45 a	0.51 a
<i>M. anisopliae</i>	0.06	0.09 a	0.12 a	0.15 a	0.18 a	0.22 a	0.26 a	0.31 a	0.36 ab	0.41 ab	0.47 a	0.53 ab
Control	0.07	0.11 b	0.16 b	0.21 b	0.28 b	0.34 b	0.42 b	0.50 b	0.58 c	0.68 c	0.78 c	0.89 c
F-value	2.91 <sub>ns</sub>	6.31*	12.24*	21.07*	36.38*	52.94*	74.25*	109.18*	97.99*	96.07*	107.63*	40.12*
P value	0.09	0.014	0.0017	0.00026	0.000035	0.000085	0.00023	0.0000052	0.0000079	0.0000085	0.0000055	0.000024
HSD value	-	0.01	0.01	0.02	0.02	0.02	0.02	0.02	0.024	0.03	0.028	0.053
<b>Conidia</b>												
<i>P. citrinum</i>	0.12	0.18 ab	0.23 a	0.29 a	0.35 a	0.41 a	0.47 a	0.53 a	0.59 a	0.65 a	0.73 a	0.75 a
<i>T. diversus</i>	0.13	0.19 ab	0.25 ab	0.32 a	0.39 a	0.46 a	0.52 a	0.59 a	0.66 a	0.74 a	0.83 a	0.85 a
<i>B. bassiana</i>	0.11	0.17 a	0.24 a	0.31 a	0.38 a	0.45 a	0.50 a	0.57 a	0.64 a	0.72 a	0.81 a	0.90 a
<i>M. anisopliae</i>	0.12	0.18 ab	0.25 ab	0.31 a	0.38 a	0.45 a	0.52 a	0.59 a	0.66 a	0.74 a	0.82 a	0.91 a
Control	0.12	0.20 b	0.27 b	0.36 b	0.45 b	0.55 b	0.65 b	0.77 b	0.90 b	1.03 b	1.18 b	1.33 b
F-value	3.37 <sub>ns</sub>	4.07*	6.49*	7.49*	9.23*	12.46*	15.82*	20.61*	27.40*	32.25*	36.85*	64.25*
P value	0.07	0.04	0.01	0.01	0.004	0.002	0.001	0.0003	0.0001	0.00006	0.00003	0.00004
HSD value	-	0.01	0.01	0.02	0.028	0.03	0.04	0.04	0.05	0.05	0.05	0.05

Note: ns = not significantly different; \* = significantly different; values within a column followed by the same letters were not significantly different at  $P < 0.05$  according to Tukey's HSD test

The decreased appetite of *S. litura* larvae treated with culture filtrate or conidia EPF was one of the stages of the larvae to be ill due to the treatment with EPF. In culture filtrate experiments, the larvae appetite already decreased on 2<sup>nd</sup> day and was faster on day 1<sup>st</sup> than the trial using the EPF conidia. In culture filtrate experiments, the decreased appetite was faster due to the raw secondary metabolite content of larvae consumed from the first day since the EPF culture filtrate was exposed, whereas in conidia treatment it required conidia time to enter the haemolymph and secrete secondary metabolites. The process of being ill due to the EPF started with the conidia attached to the cuticle larvae, and then the conidia ruptures



to form germ tubes to penetrate the cuticle (Fernandes et al. 2007). The infection started when germ tubes were able to penetrate the insect cuticle and the ability to infect was a determinant of the level of virulence EPF (Altre and Vandenberg 2001). After germ tubes reach the haemocoel, they produced specific infection hyphae originating at appressoria (El-Ghany 2015). Then the hyphae spread to the haemolymph and developed to produce blastospores. The blastospores then produced the secondary metabolites and enzymes that could disrupt normal cell metabolism (Mancillas-Paredes et al. 2019). The penetration stages of germ tubes to blastospores produced secondary metabolites and the enzymes occurred in the experiments using conidia, whereas in culture filtrate experiments this stage did not occur because the two stages of filtration were already carried out to separate conidia from the filtrate. Therefore, the process of the larvae starting to be ill in the culture filtrate was faster than the conidia experiment. The next process, the toxins from secondary metabolites began to kill the host insect but the death of the host insect was not only by the toxin but also by the mechanical damage of the fungal penetration into the body of the insect (El-Ghany 2015).

The reduction of the weight of the eaten affected the weight of the larvae. At 1 x 24 hours after the larvae were treated with culture filtrate or EPF conidia, the weight of the *S. litura* larvae did not show any significant difference, starting on the second day the larvae weight began to be lower than that of the control (Table 4), as well as the third day up to the 12 day the larvae weight was lower than that of the control. All the tested species of the EPF decreased significantly the larvae body weight compared to that of the control. Consequently, the larvae treated with mycoinsecticides containing the culture filtrate or EPF conidia could reduce the larvae appetite to make their body weight decrease.

Percentage of the mortality or mortality of the larvae caused by the EPF culture filtrate showed significant differences when compared to that of the control (Table 5). All EPF species were toxic to *S. litura* larvae, both their culture filtrate and conidia. The highest larvae mortality due to culture filtrate was found in the treatment of *P. citrinum* (98.67%) and not significantly different from the larvae mortality by *T. diversus* (96%), but significantly different from the larvae mortality by *B. bassiana* (86.67%) and *M. anisopliae* (82.67%). The  $LT_{50}$  larvae treated with culture filtrate *T. diversus* was the shortest (6.19 days) and not significantly different from the  $LT_{50}$  larvae treated with *P. citrinum* (6.62 days), but significantly different from the  $LT_{50}$  larvae treated with *B. bassiana* (7.55 days) and *M. anisopliae* (7.60 days). Thus, the most toxic EPF species against *S. litura* larvae was *P. citrinum*. The larvae mortality due to the EPF conidia showed significant differences when compared to that of the control (Table 5). All of the EPF species were pathogenic to *S. litura* larvae. The larvae mortality caused by the conidia was highest in *P. citrinum* (98.67%) which was not significantly different from the larvae mortality by *T. diversus* (97.33%), *B. bassiana* (86.67%), and *M. anisopliae* (86.67%). The  $LT_{50}$  of the larvae treated with the conidia of *P. citrinum* was the shortest (5.45 days) and not significantly different from the  $LT_{50}$  of larvae treated with *T. diversus* (6.55 days), but it was significantly different from the  $LT_{50}$  of larvae treated with *B. bassiana* (8.88 days) and *M. anisopliae* (7.70 days). Thus, the most pathogenic EPF species against *S. litura* larvae was *P. citrinum*.

**Table 5.** Mean of larvae mortality and  $LT_{50}$  of *Spodoptera litura* larvae treated with culture filtrates and conidia of mycoinsecticides from *Penicillium citrinum*, *Talaromyces diversus*, *Beauveria bassiana*, and *Metarhizium anisopliae*

Fungal species	Mortality $\pm$ SE (%)	$LT_{50} \pm$ SE (days)
<b>Culture filtrate</b>		
<i>P. citrinum</i>	98.67 $\pm$ 1.09 c	6.62 $\pm$ 0.135 a
<i>T. diversus</i>	96.00 $\pm$ 1.89 c	6.19 $\pm$ 0.072 a
<i>B. bassiana</i>	86.67 $\pm$ 2.88 b	7.55 $\pm$ 0.154 a
<i>M. anisopliae</i>	82.67 $\pm$ 1.09 b	7.60 $\pm$ 0.203 b
Control	0.00 $\pm$ 0.00 a	-
F-value	187.26*	13.12*
P value	0.00000006	0.00480044
HSD value	11.38	0.87
<b>Conidia</b>		
<i>P. citrinum</i>	98.67 $\pm$ 1.09 b	5.45 $\pm$ 0.232 a
<i>T. diversus</i>	97.33 $\pm$ 2.18 b	6.55 $\pm$ 0.081 ab
<i>B. bassiana</i>	86.67 $\pm$ 1.09 b	8.88 $\pm$ 0.420 c
<i>M. anisopliae</i>	86.67 $\pm$ 1.09 b	7.70 $\pm$ 0.404 bc
Control	0.00 $\pm$ 0.00 a	-
F-value	156.31*	29.38*
P value	0.00000013	0.00055
HSD value	12.70	1.24


Note: \* = significantly different; values within a column followed by the same letters were not significantly different at  $P < 0.05$  according to Tukey's HSD test

Both culture filtrate and conidia of EPF in this study had significantly higher insecticidal activity than the control, and the highest insecticidal activity was found in *P. citrinum*. *P. citrinum* had the highest insecticidal activity due to its ability to infect the larvae faster shown by its  $LT_{50}$  value which was the shortest compared to the other species. Altre and


Vandenberg (2001) state that the most pathogenic EPF species due to the fungus has germ tubes that are able to penetrate the insect cuticle faster than other fungal species and have more toxins and enzymes that disrupt normal cell metabolism in insects. In addition, the color of the *P. citrinum* culture filtrate was the darkest compared to the color of the culture filtrate of the other three EPF species. The darker color of the EPF culture filtrate proved to be more toxic than the lighter color of the culture filtrate (Ayudya et al., 2019; Gustianingtyas et al., 2020) because the darker the color of the fungus culture filtrate the higher the concentration of the secondary metabolites (Luo et al., 2017). The *P. citrinum* culture filtrate contains secondary metabolites and is toxic to insects, for example patulin which can kill *Culex quinquefasciatus* larvae up to 100% (Maketon et al., 2014). *P. citrinum* are reported to be effective in killing insect pests (Foo et al., 2017; Maketon et al., 2014) and mite pests (Mazid et al., 2016).

The other EPF species were also toxic because they had enzymes or secondary metabolites that could kill the host insects, for example *B. bassiana* produced the enzymes of proteolytic and chitinolytic that could disrupt normal cell metabolism (Mancillas-Paredes et al., 2019). The secondary metabolite produced by *B. bassiana* containing the bassianolone B toxic to the *T. molitor* larvae (Luo et al., 2017). *M. anisopliae* produces secondary metabolites, including destruxins toxic to insect pests (Borisade et al., 2016). So far there has been no information on the secondary metabolites or enzymes produced by *T. diversus* that kills insects. At present there is only general information stating that *T. diversus* can kill insects (Visagie et al., 2009). For this reason, in the future it is necessary to study to identify the secondary metabolites or *T. diversus* enzymes that can kill insect pests.

The *S. litura* larvae treated with EPF culture or conidia showed the same behavior, i.e., they ate less and less and got lazy to move, while the control not treated with the EPF continued to eat greedily and actively moved. After 3 days the larvae given culture filtrate began to change integument color that was previously bright green to dull and began to find dead larvae and the same symptoms occurred in larvae treated with EPF conidia. Four days after the larvae were treated with culture filtrate or EPF conidia, the larvae's body began to shrink, wrinkle, get odorless and dull black, and eventually died (Figure 6 and 7). After 2-3 days the larvae died, the symptoms in the larvae integument died due to the conidia covered by mycelia and the conidia appeared, whereas in larvae that died due to the culture filtrate there was no mycelia and conidia. Thus, the symptoms of ill *S. litura* larvae caused by the culture filtrate were slightly different from the symptoms caused by the EPF conidia. The symptoms of sick larvae due to the culture of this filtrate were the same as the symptoms described by Ayudya et al. (2019) and Gustianingtyas et al. (2020), namely the larvae got shrunk, wrinkled, dry, odorless, and the integument was not covered by EPF conidia. The symptoms of sick insects caused by EPF conidia in this study were the same as those described by Bugti et al. (2018) and Sumikarsih et al. (2019), that is, the insect's body got smaller, wrinkled, dry, odorless, and the integument was covered in mycelia and EPF conidia.



**Figure 6.** Morphology of *Spodoptera litura* larvae of healthy larvae of control (right) and dead larvae (left) treated with culture filtrate of of mycoinsecticides from *Penicillium citrinum* (A), *Talaromyces diversus* (B), *Beauveria bassiana* (C), and *Metarhizium anisopliae* (D)



**Figure 7.** Morphology of *Spodoptera litura* larvae of healthy larvae of control (right) and dead larvae (left) treated with conidia of mycoinsecticides from *Penicillium citrinum* (A), *Talaromyces diversus* (B), *Beauveria bassiana* (C), and *Metarhizium anisopliae* (D)

The results of bioassays of culture filtrate and EPF conidia showed that both culture filtrate and conidia had high insecticidal activity. However, for the future development of mycoinsecticides, the use of culture filtrate without conidia as the active ingredient of mycoinsecticide needs to be further investigated. The fungus culture filtrate is relatively more stable than the culture that still has conidia and can be stored for a longer time and remains toxic, besides the culture filtrate does not clog the sprayer nozzle when sprayed (Soesanto et al. 2019).

The EPF species found in the soils of South Sumatra were *P. citrinum*, *T. diversus*, *B. bassiana*, and *M. anisopliae*. All species have a high insecticidal activity against *S. litura* larvae and the most toxic species is *P. citrinum*. The filtrate and conidia of the *P. citrinum* are toxic. *P. citrinum*, a new emerging entomopathogenic fungus, has the highest insecticidal activity and the highest potential to be developed into commercial mycoinsecticides.

#### ACKNOWLEDGEMENTS

This research was funded by the scheme of Applied Research (Riset Terapan) with budget year of 2020 according to Directorate of Research and Community Service (*Direktorat Riset dan Pengabdian Masyarakat*), Ministry of Research and Technology/National Research and Innovation Agency of Republic Indonesia, Grant/Contract Number: SP DIPA-042.06.1.401516/2020.


#### REFERENCES

- Alhusein N, Scott J, Kasprzyk-Hordern B, Bolhuis A. 2016. Development of a filter to prevent infections with spore-forming bacteria in injecting drug users. *Harm Reduct. J.* 13: 1–8. <https://doi.org/10.1186/s12954-016-0122-1>
- Altre JA, Vandenberg JD. 2001. Factors influencing the infectivity of isolates of *Paecilomyces fumosoroseus* against diamondback moth, *Plutella xylostella*. *J. Invertebr. Pathol.* 78: 31–36. <https://doi.org/10.1006/jipa.2001.5043>
- Anwar W, Khan SN, Aslam M, Haider MS, Shahid AA, Ali M. 2015. Exploring fungal flora associated with insects of cotton agroecological zones of Punjab, Pakistan. *Pakistan Entomol.* 37: 27–31.
- Ayudya DR, Herlinda S, Suwandi S. 2019. Insecticidal activity of culture filtrates from liquid medium of *Beauveria bassiana* isolates from South Sumatra (Indonesia) wetland soil against larvae of *Spodoptera litura*. *Biodiversitas* 20: 2101–2109. <https://doi.org/10.13057/biodiv/d200802>
- Borisade OA, Medina A, Magan N. 2016. Interacting temperature and water activity modulate production of destruxin a by *Metarhizium anisopliae* on galleria larvae-modified agar based media invitro. *West African J. Appl. Ecol.* 24: 31–42.
- Bugti GA, Bin W, Na C, Feng LH. 2018. Pathogenicity of *Beauveria bassiana* strain 202 against sap-sucking insect pests. *Plant Prot. Sci.* 54: 111–117. <https://doi.org/10.17221/45/2017-PPS>
- Chen Z, Xu L, Yang F, Ji G, Yang J, Wang J. 2014. Efficacy of *Metarhizium anisopliae* isolate MAX-2 from Shangri-la, China under desiccation stress. *BBMC Microbiol.* 14, 1–8. <https://doi.org/http://www.biomedcentral.com/1471-2180/14/4>

- El-Ghany TMA. 2015. Entomopathogenic Fungi and their Role in Biological Control. Biology Department Faculty of Science Jazan University KSA, Cairo. <https://doi.org/10.4172/978-1-63278-065-2-66>
- Fernandes EKK, Rangel DEN, Moraes AM, Bittencourt VREP, Roberts DW. 2007. Variability in tolerance to UV-B radiation among *Beauveria* spp. isolates. *J Invertebr Pathol* 96: 237–243. <https://doi.org/10.1016/j.jip.2007.05.007>
- Foo K, Seelan JSS, Dawood MM. 2017. Microfungi associated with *Pteroptyx bearni* (Coleoptera: Lampyridae) eggs and larvae from Kawang River, Sabah (Northern Borneo). *Insects* 8: 1–12. <https://doi.org/10.3390/insects8030066>
- Gustianingtyas M, Herlinda S, Suwandi, Suparman, Hamidson H, Hasbi, Setiawan A, Verawaty M, Elfita, Arsi. 2020. Toxicity of entomopathogenic fungal culture filtrate of lowland and highland soil of South Sumatra (Indonesia) against *Spodoptera litura* larvae. *Biodiversitas* 21: 1839–1849. <https://doi.org/10.13057/biodiv/d210510>
- Hernandez-Trejo A, Estrada-Drouaillet B, López-Santillán, J, Rios-Velasco C, Varela-Fuentes S, Rodríguez-Herrera R, Osorio-Hernández E. 2019. In vitro evaluation of native entomopathogenic fungi and neem (*Azadirachta indica*) extracts on *Spodoptera frugiperda*. *Int. J. Exp. Bot.* 4: 75–84. <https://doi.org/10.1037/0033-2909.126.1.78>
- Johari A, Herlinda S, Irsan C, Pujiastuti Y. 2016. Phenomenon of Thrips (Thysanoptera) attack on chili plant (*Capsicum annuum*). *Am. J. Agric. Biol. Sci.* 11: 103–109. <https://doi.org/10.3844/ajabssp.2016.103.109>
- Kumar S, Stecher G, Tamura K. 2016. MEGA7: Molecular evolutionary genetics analysis version 7.0 for bigger data sets. *Mol. Biol. Evol.* 33: 1870–1874. <https://doi.org/10.1093/molbev/msw054>
- Liu SF, Wang GJ, Nong XQ, Liu B, Wang MM, Li SL, Cao GC, Zhang ZH. 2017. Entomopathogen *Metarhizium anisopliae* promotes the early development of peanut root. *Plant Prot. Sci.* 53: 101–107. <https://doi.org/10.17221/49/2016-PPS>
- Lopes RB, Souza DA, Rocha LFN, Montalva C, Luz C, Humber RA, Faria M. 2018. *Metarhizium alvesii* sp. nov.: A new member of the *Metarhizium anisopliae* species complex. *J. Invertebr. Pathol.* 151: 165–168. <https://doi.org/10.1016/j.jip.2017.12.001>
- Luo Z, Ren H, Mousa JJ, Rangel DEN, Zhang Y, Bruner SD, Keyhani NO. 2017. The PacC transcription factor regulates secondary metabolite production and stress response, but has only minor effects on virulence in the insect pathogenic fungus *Beauveria bassiana*. *Environ. Microbiol.* 19: 788–802. <https://doi.org/10.1111/1462-2920.13648>
- Maketon M, Amnuaykanjanasin A, Kaysorngup A. 2014. A rapid knockdown effect of *Penicillium citrinum* for control of the mosquito *Culex quinquefasciatus* in Thailand. *World J. Microbiol. Biotechnol.* 30: 727–736. <https://doi.org/10.1007/s11274-013-1500-4>
- Mancillas-Paredes J, Hernández-Sánchez H, Jaramillo-Flores ME, García-Gutiérrez C. 2019. Proteases and chitinases induced in *Beauveria bassiana* during infection by *Zabrotes subfasciatus*. *Southwest. Entomol* 44: 125–137. <https://doi.org/10.3958/059.044.0114>
- Mascarin GM, Jackson MA, Kobori NN, Behle RW, Delalibera Júnior Í. 2015. Liquid culture fermentation for rapid production of desiccation tolerant blastospores of *Beauveria bassiana* and *Isaria fumosorosea* strains. *J. Invertebr. Pathol.* 127: 11–20. <https://doi.org/10.1016/j.jip.2014.12.001>
- Mathew AJ, Jayachandran K, Mathew J. 2010. Endophytic *Penicillium citrinum* Thom. from *Scoparia dulcis* Linn. *Indian J. Microbiol.* 50: 99–102. <https://doi.org/10.1007/s12088-010-0065-2>
- Mazid S, Kalita JC, Rajkhowa RC. 2016. Biocontrol potential of *Penicillium citrinum* and *Penicillium chrysogenum* against red spider mite, *Oligonychus coffeae* Nietner infesting tea. *J. Entomol. Res.* <https://doi.org/10.5958/0974-4576.2016.00008.6>
- Ngangambe MH, Mwatawala MW. 2020. Effects of entomopathogenic fungi (EPFs) and cropping systems on parasitoids of fall armyworm (*Spodoptera frugiperda*) on maize in eastern central, Tanzania. *Biocontrol Sci. Technol.* 30: 1–13. <https://doi.org/10.1080/09583157.2020.1726878>
- Safitri A, Herlinda S, Setiawan A. 2018. Entomopathogenic fungi of soils of freshwater swamps, tidal lowlands, peatlands, and highlands of South Sumatra, Indonesia. *Biodiversitas* 19: 2365–2373. <https://doi.org/10.13057/biodiv/d190647>
- Sharma L, Oliveira I, Torres L, Marques G. 2018. Entomopathogenic fungi in Portuguese vineyards soils: suggesting a ‘Galleria-Tenebrio-bait method’ as bait-insects *Galleria* and *Tenebrio* significantly underestimate the respective recoveries of *Metarhizium (robertsii)* and *Beauveria (bassiana)*. *MycKeys* 23: 1–23. <https://doi.org/10.3897/mycokeys.38.26970>
- Siaga E, Lakitan B, Hasbi H, Bernas SM, Widuri LI, Kartika K. 2019. Floating seedbed for preparing rice seedlings under unpredictable flooding occurrence at tropical riparian wetland. *Bulg. J. Agric. Sci.* 25: 326–336.
- Soesanto L, Mugiastuti E, Manan A. 2019. Raw secondary metabolites application of two *Trichoderma harzianum* isolates towards vascular streak dieback on cocoa seedlings. *Pelita Perkeb.* 35, 22–32. <https://doi.org/10.22302/iccri.jur.pelitaperkebunan.v35i1.346>
- Souza ML, Sanches MM, Souza DAd, Faria M, Espinel-Correal C, Sihler W, Lopes RB. 2019. Within-host interactions of *Metarhizium rileyi* strains and nucleopolyhedroviruses in *Spodoptera frugiperda* and *Anticarsia gemmatilis* (Lepidoptera: Noctuidae). *J. Invertebr. Pathol.* 162: 10–18. <https://doi.org/10.1016/j.jip.2019.01.006>
- Sowmya G, Ramanagouda SH, Jayappa J. 2017. Lethal and sub lethal effects of indigenous isolates of *Metarhizium anisopliae* (Metchnikoff) Sorokin against *Spodoptera litura* (Fabricius). *Entomol. Zool. Stud.* 5, 1154–1160.
- Sumikarsih E, Herlinda S, Pujiastuti Y. 2019. Conidial density and viability of *Beauveria bassiana* isolates from Java and Sumatra and their virulence against *Nilaparvata lugens* at different temperatures. *Agrivita J. Agric. Sci.* 41: 335–349. <http://doi.org/10.17503/agrivita.v41i2.2105>
- Suwandi, Ammar M, Chandra I. 2012. Application of extract compost increased yield and suppressed the diseases of ratoon rice crop in tidal swamp of Banyuasin Regency. *J. Suboptimal Lands* 1: 116–122. [Indonesia] <https://doi.org/10.33230/JLSO.1.2.2012.15>
- Syamsudin TS, Faizal A, Kirana, R. 2019. Dataset on antixenosis and antibiosis of chili fruit by fruit fly (*Bactrocera dorsalis*) infestation Tati. *Data Br.* 23: 1–4. <https://doi.org/10.1016/j.dib.2019.103758>
- Thaochan N, Sausa-Ard W. 2017. Occurrence and effectiveness of indigenous *Metarhizium anisopliae* against adults *Zeugodacus cucurbitae* (Coquillett) (Diptera: Tephritidae) in Southern Thailand. *Songklanakarin J. Sci. Technol.* 39: 325–334. <https://doi.org/10.14456/sjst-psu.2017.35>
- Tompe AA, Hole UB, Kulkarni SR, Chaudhari CS, Chavan SK. 2020. Studies on seasonal incidence of leaf eating caterpillar, *Spodoptera litura* (Fab.) infesting capsicum under polyhouse condition. *J. Entomol. Zool. Stud.* 8: 761–764.
- Vijayalakshmi P, Vijayalakshmi T, Naidu LN. 2016. Evaluation of certain insecticide molecules against chillipod borer, *Spodoptera litura* in Andhra Pradesh. *J. Res. ANGRAU* 44: 26–30.
- Visagie CM, Jacobs K. 2012. Three new additions to the genus *Talaromyces* isolated from Atlantis sandveld fynbos soils. *Persoonia* 28: 14–24. <https://doi.org/10.3767/003158512X632455>
- Visagie CM, Roets F, Jacobs K. 2009. A new species of *Penicillium*, *P. ramulosum* sp. nov., from the natural environment. *Mycologia* 101: 888–895. <https://doi.org/10.3852/08-149>
- White T, Bruns T, Lee S, Taylor J. 1990. Amplification and direct sequencing of fungal ribosomal RNA genes for phylogenetics, in: Innis, M.A., Gelfand, D.H., Sninsky, J.J., White, T.J. (Eds.), *PCR Protocols: A Guide to Methods and Applications*. Academic Press Inc., New York.
- Yang H, Qin CS, Chen YM, Zhang GY, Dong LH, Wan SQ. 2019. Persistence of *Metarhizium* (Hypocreales: Clavicipitaceae) and *Beauveria bassiana* (Hypocreales: Clavicipitaceae) in tobacco soils and potential as biocontrol agents of *Spodoptera litura* (Lepidoptera: Noctuidae). *Environ. Entomol.* 48: 147–155. <https://doi.org/10.1093/ee/nvy161>
- Youssef NA. 2014. Effect of certain entomopathogenic fungi and nematode on the desert locust *Schistocerca gregaria* (Forsk.). *Ann. Agric. Sci.* 59, 125–131. <https://doi.org/10.1016/j.aos.2014.06.017>



## 3. Bukti konfirmasi paper accepted, uncorrected Proof dan hasil koreksi penulis



# New emerging entomopathogenic fungi isolated from soil in South Sumatra (Indonesia) and their filtrate and conidial insecticidal activity against *Spodoptera litura*

SITI HERLINDA<sup>1,2,\*</sup>, RISKI ANWAR EFENDI<sup>1</sup>, RADIX SUHARJO<sup>3</sup>, HASBI<sup>2,4</sup>, ARUM SETIAWAN<sup>5</sup>, ELFITA<sup>6</sup>,  
MARIESKA VERAWATY<sup>5</sup>

<sup>1</sup>Department of Plant Pests and Diseases, Faculty of Agriculture, Universitas Sriwijaya, Indralaya 30662, Indonesia

\*Email: sitiherlinda@unsri.ac.id

<sup>2</sup>Research Center for Sub-optimal Lands (PUR-PLSO), Universitas Sriwijaya, Palembang 30139, Indonesia

<sup>3</sup>Department of Plant Protection, Faculty of Agriculture, Universitas Lampung, Jl. Prof. Sumantri Brojonegoro No. 1 Bandar Lampung 35145, Lampung, Indonesia

<sup>4</sup>Agricultural Engineering Study Program, Faculty of Agriculture, Universitas Sriwijaya, Indralaya 30662, Indonesia

<sup>5</sup>Department of Biology, Faculty of Mathematics and Natural Sciences, Universitas Sriwijaya, Indralaya 30662, Indonesia

<sup>6</sup>Department of Chemistry, Faculty of Mathematics and Natural Sciences, Universitas Sriwijaya, Indralaya 30662, Indonesia

Manuscript received: xxx. Revision accepted: xxx October 2020.

**Abstract.** Herlinda S, Efendi RA, Suharjo R, Hasbi, Setiawan A, Elfita, Verawaty M. 2020. *New emerging entomopathogenic fungi isolated from soil in South Sumatra (Indonesia) and their filtrate and conidial insecticidal activity against Spodoptera litura.* *Biodiversitas* 21: xxxx. This study was carried out to confirm identity of *Beauveria bassiana* and *Metarhizium anisopliae* using sequence analysis of Internal Transcribed Spacer (ITS) region as well as the two unknown isolates of EPF and to investigate their filtrate and conidial insecticidal activity against *S. litura* larvae. The EPF species were identified based on morphological characters and molecular analysis. Their culture filtrate and conidial bioassays were performed on *S. litura* larvae. The species found were *Penicillium citrinum* (BKbTp), *Talaromyces diversus* (MSwTp1), *B. bassiana* (BSwTd4), and *M. anisopliae* (MSwTp3) and have been deposited in the GenBank. All EPF species have a high insecticidal activity against *S. litura* larvae. The highest larvae mortality caused by filtrate was found on *P. citrinum* (98.67%) and not significantly different from those of *T. diversus* (96%), but significantly different from *B. bassiana* (86.67%) and *M. anisopliae* (82.67%). The highest larvae mortality caused by the conidia was also occurred to *P. citrinum* (98.67%) which was not significantly different from those of *T. diversus* (97.33%), *B. bassiana* (86.67%), and *M. anisopliae* (86.67%). The LT<sub>50</sub> caused by *T. diversus* filtrate was the shortest (6.19 days) and not significantly different from those caused by *P. citrinum* (6.62 days). The shortest LT<sub>50</sub> caused by conidia was found on *P. citrinum* (5.45 days). Therefore, the conidia and filtrate of *P. citrinum*, a new emerging entomopathogenic fungus, has the highest insecticidal activity. *P. citrinum* has the potential to be developed into commercial mycoinsecticides.

**Keywords:** *Beauveria bassiana*; leaf eating caterpillar; *Metarhizium anisopliae*; *Penicillium citrinum*; *Talaromyces diversus*

## INTRODUCTION

South Sumatra, Indonesia has a stretch with varying topography from the height of the land to the high ground. This topographic variation can develop many species of plants, for example vegetables and the food crops. Vegetables that can adapt to various land topography in South Sumatra include chili (*Capsicum* spp.), mustard greens (*Brassica juncea*), kale (*Ipomoea reptans*), and spinach (*Amaranthus tricolor*) (Siaga et al. 2019). The main constraints of vegetable cultivation, especially chili pests are fruit flies (*Bactrocera dorsalis*) (Syamsudin et al. 2019), sap sucker (Thrips) (Johari et al. 2016), and leaf eating caterpillar (*Spodoptera litura*) (Tompe et al. 2020).

*S. litura* is a polyphagous insect pest that attacks many plant species from various families (Tompe et al. 2020), like chili, mustard greens, water spinach, and spinach. The yield loss of chili caused by this pest reaches 50% every year (Vijayalakshmi et al. 2016). The first instar of *S. litura* larvae eats the leaves in groups and skeletonize them, while

the second instar extend to the last instar spread and eat individually leaves, flowers and pistil chili (Tompe et al. 2020). In addition, *S. litura* is a key pest for the leafy vegetables, such as mustard greens, water spinach, and spinach because the leaves that have been eaten become no longer valuable (Vijayalakshmi et al. 2016; Tompe et al. 2020). Since the public awareness on the good health and environmental quality is increasing, there is a need to reduce the use of synthetic pesticides and improving application of eco-friendly control strategies for *S. litura*. One of the methods is applying the entomopathogenic fungi (EPF).

Some EPF species have been shown to be effective in killing *S. litura*, for example, *Beauveria bassiana* cultured in liquid media has been shown to kill 98% of *S. litura* larvae (Ayudya et al. 2019; Gustianingtyas et al. 2020). *Metarhizium anisopliae* was reportedly also effective in killing *S. litura* larvae (Sowmya et al. 2017; Yang et al. 2019; Gustianingtyas et al. 2020) and can kill 96% of *S. litura* larvae (Gustianingtyas et al. 2020). Furthermore,

*Metarhizium rileyi* also effectively kills *S. litura* larvae (Liu et al. 2017). In addition to being effective in killing the insect pests, EPF can also be integrated with entomopathogenic viruses, such as nucleopolyhedroviruses (NPV) (Souza et al. 2019) and botanical insecticides (Hernandez-Trejo et al. 2019) and does not harm parasitoids (Ngangambe and Mwatawala 2020).

Four isolates of EPF have been isolated from rhizosphere of several kinds of plant in lowland to high land of South Sumatra. On the basis of morphological characters, the two isolates have been identified as *B. bassiana* and *M. anisopliae* (Safitri et al. 2018; Sumikarsih et al. 2019; Gustianingtyas et al. 2020) and their filtrates were proven to be effectively killed up to 98% of *S. litura* larvae (Ayudya et al. 2019; Gustianingtyas et al. 2020). The other two EPF, until now, have not been fully revealed on molecular identity and their filtrate and conidial capability to cause mortality on *S. litura*. This research aimed to confirm identity of *Beauveria bassiana* and *Metarhizium anisopliae* using sequence analysis of Internal Transcribed Spacer (ITS) region as well as the two unknown isolates of EPF and to investigate their filtrate and conidial insecticidal activity against *S. litura* larvae.

## MATERIALS AND METHODS

### Exploring EPF

The EPF exploration was carried out on soil near root of oil palm, rice, maize and vegetables in various locations (Table 1) ranging from lowland to highland of South Sumatra using the method of Safitri et al. (2018). Those locations chosen have objective to obtain the fungi that had adapted in the soils. The morphological identifications were carried out in 2018 and 2019, and their identities were confirmed by molecular analysis in February 2020. The EPF exploration method modified those of Anwar et al. (2015), namely using a bait method with *Tenebrio molitor* larvae as insect bait which was fed in the soil sample. The soil sampling used the method of Anwar et al. (2015) by digging the soil using surface soil sampler ( $\varnothing$  20 mm) as deep as 10 cm around the rhizosphere plant according to the method of Thaochan and Sausa-Ard (2017) and 5 points were taken to collect 1 kg of soil. Then, the soil sample was put into a plastic bag and labeled with information about the height of the location, type of commodity, and date of collection.

The soil sample was first cleaned from the roots and sieved with a 10 mesh sieve. Then, it was put into a plastic tray (32 x 25 x 5 cm<sup>3</sup>) each containing as much as 1 kg. The soil was then moistened with sterile distilled water with soil moisture exceeding 20% according to the method of Chen et al. (2014). Then, 20 newly molted 3rd-instar larvae of *T. molitor* were placed at the bottom of the tray and the larvae were sprinkled with the soil to reach 3 cm in thickness. Then, the tray containing the soil was covered with a black cloth and sprayed with distilled water to maintain the soil moisture. The larvae were infested in the soil for 7 days to give time for the entomopathogenic fungal conidia to infect them, after which the dead larvae infected with entomopathogenic fungi were grown in the Sabouraud Dextrose Agar (SDA, Merck) medium.

### Isolation and identification of EPF

The EPF infecting the larvae of *T. molitor* which was characterized by its body was overgrown with EPF mycelia, dry and contracted, then isolated and purified using the method of Youssef (2014). The isolation of EPF from the larvae's body surface began with the surface sterilization by modifying the method of Sharma et al. (2018) by rinsing with 1% of NaOCl for 1 minute, then rinsing with 100 mL of sterile distilled water for 3 times. The larvae sterilization was carried out to obtain EPF inside the haemocoel and cuticle larvae and prevent water fungi from being obtained. The sterilized larvae were grown in SDA media and incubated for 2 days. The growing fungus was purified in the SDA media until after getting pure isolates. The pure EPF was identified based on the molecular analysis.

### Fungal identification

Morphological observations were carried out based on microscopic and macroscopic characteristics of the EPF. The macroscopic observation of the fungi was based on the color and shape of colony, whereas the microscopic observation was based on the shape and size of conidia and conidiophores.

The EPF were identified based on the molecular analysis in February 2020 at the Laboratory of Agricultural Biotechnology (accredited according to the ISO/IEC 17025 standard), Department of Plant Protection, Faculty of Agriculture, Universitas Lampung, Indonesia.

**Table 1.** Locations of EPF in South Sumatra, Indonesia

Fungal species (codes)	Village, District/City	Ecosystem	Crop	GPS (coordinat)	Altitude (m)	GenBank Acc. No.
<i>Penicillium citrinum</i> (BKbTp)	Talang Patai, Pagar Alam	Highland soil	Cabbage	S 04°02.161' E 103°10.484'	175	MT448730
<i>Talaromyces diversus</i> (MSwTp1)	Talang Patai, Pagar Alam	Highland soil	Mustard	S 03°50.174', E 103°31.293'	193	MT448731
<i>Beauveria bassiana</i> (BSwTd4)	Talang Dabok, Ogan Komering Ilir	Peat soil	Oil palm	S 03°23.570' E 104°51.498'	24	MT448732
<i>Metarhizium anisopliae</i> (MSwTp3)	Talang Patai, Pagar Alam	Highland soil	Mustard	S 03°50.174', E 103°31.293'	193	MT448733



**Table 2.** Primer used for amplification

Target of amplification	Primer name	Sequence (5'-3')	Product size (bp)	Reference
Internal Transcribe	ITS1	TCC GTA GGT GAA CCT TGC GG	600	(White et al. 1990)
Spacer Region	ITS4	TCC TCC GCT TAT TGA TAT GC		

### DNA extraction and PCR amplification

The DNA extraction was performed on the mycelium of 7 days old fungi using InstaGene™ Matrix (BioRad). Amplification was carried out by Sensoquest Thermal Cycler Machine (Germany). PCR amplification on Internal Transcribed Spacer (ITS) region was conducted using primer ITS1 and ITS4 (Table 2) (White et al. 1990). The PCR was carried out with total volume of 25 µL involving 12.5 µL Master Mix (Red Mix) (bioline), 1 µL of 10 µM of primer ITS 1 and ITS 4, 2 µL DNA and 8.5 µL of sterile distilled water. DNA amplification consist of 1 cycle of the initiation at 95°C for 5 min, 30 cycles of denaturation at 95°C for 1 minute, primer annealing at 52°C for 1 min, primer extension at 72°C for 1 minute, and one cycle of elongation at 72°C for 5 minutes. The PCR results were run using 0.5% agarose gel in 20 mL of 1x buffer Tris-Boric Acid-EDTA (TBE) (1<sup>st</sup> Base Malaysia) with 1 µL Ethidium Bromide (EtBr 10 mg mL<sup>-1</sup>). The electrophoresis was performed by 1x TBE buffer at 50 volts for 70 minutes. The results were visualized by DigiDoc UV transilluminator (UVP, USA).

The PCR product was emitted to 1<sup>st</sup> Base Malaysia for sequencing. The sequencing results were analyzed with Bio Edit program ver. 7.2.6 for windows. The result was then submitted to Basic Local Alignment Search Tool (BLAST) (<https://blast.ncbi.nlm.nih.gov/Blast.cgi>) to find their potential identity. The phylogenetic tree was developed by Mega 7 for Windows (Kumar et al. 2016) with Unweighted-pair Group Method with Arithmetic means (UPGMA). Reference strains employed in this research were obtained from NCBI (<https://www.ncbi.nlm.nih.gov/>).

### *Spodoptera litura* mass rearing

*S. litura* larvae used in the insecticide activity test was collected from the vegetable fields around the campus of Universitas Sriwijaya and those vegetable fields that were not sprayed with synthetic insecticides. The larvae were taken to the laboratory and maintained in plastic containers (Ø 25 cm, height 30 cm) topped with gauze. that has been washed thoroughly using detergent. Next, pesticide-free spinach (*Ipomoea aquatic*) was put into the containers for the first instar feed of *S. litura*. The second instar was fed mulberry leaves (*Morus alba*) planted in the experimental field of the Faculty of Agriculture, Universitas Sriwijaya, Indonesia and not sprayed with pesticides.

Every day the larvae feed was replaced with fresh new feed. The larvae entering the pupae stage were transferred into plastic containers (Ø 10 cm, height 15 cm) whose top was exposed and the base was sprinkled with sifted soil (10 mesh). and already sterilized in an oven for 1 hour at 100°C. The thickness of the soil in the container was 3 cm.

This plastic container containing pupae was put into a gauze cage (30 x 30 x 30 cm<sup>3</sup>) already tapped with 10 water spinach sticks for laying eggs to the adults. The hatched eggs were transferred into a plastic cylinder and fed with water spinach which was replaced every day. The larvae used for toxicity testing were the second instar of the third offspring (F3) or afterwards.

### Producing of EPF mycoinsecticides

The EPF already identified molecularly, namely *P. citrinum*, *T. diversus*, *B. bassiana*, and *M. anisopliae* were made liquid formulation mycoinsecticides. The mycoinsecticides made of four types, each of which had active ingredients *P. citrinum*, *T. diversus*, *B. bassiana*, and *M. anisopliae* and sterile distilled water was used for control. The mycoinsecticides were made by the method of Mascarin et al. (2015). In this, EPF was cultivated in solid media of Sabouraud Dextrose Agar (SDA, Merck), incubated for 7 days, and then the fungal cultures that grew were transferred to the liquid media of Sabouraud Dextrose Broth (SDB, Merck). The broth fungal cultures were incubated using a shaker for 7 days at 120 rpm. The broth fungal cultures from SDB with a minimum density of 1x10<sup>9</sup> conidia mL<sup>-1</sup> were used for the active ingredient mycoinsecticides. The mycoinsecticides were made liquid formulation containing a mixture of 600 mL of the broth fungal cultures, 400 mL of carrier from shrimp shell flour compost extract, 10 mL of vegetable oil, and 10 mL of sterile distilled water solution of 0.04% polyoxyethyl-189 ene sorbitan mono-oleate (Tween®). The shrimp skin flour compost extract was made using the method of Suwandi et al. (2012). After that, the formulation was stored for 30 days before it was applied to obtain stable and non-contaminant mycoinsecticides. The mycoinsecticides activity was tested in the form of EPF culture and conidia filtrate.

### Culture filtrate preparation of the mycoinsecticides.

The presence of raw secondary metabolites in culture filtrate was performed according to Soesanto et al. (2019) can be done by filtering fungal broth or liquid mycoinsecticides. The culture filtrate of liquid mycoinsecticides was produced from 2 stages of filtering using the method of Ayudya et al. (2019). The first stage, 100 mL liquid mycoinsecticides was filtered using Whatman filter paper no. 42 and cotton with a thickness of 1 cm. The filtration culture produced from the first stage filter was then sucked up to 10 mL using a sterile hypodermic syringe to be filtered in the second stage. The second stage filtering used a syringe filter (0.45 µm pore size, Sartorius). The method of installing the syringe filter used the method of Alhusein et al. (2016) by placing it at the base of the needle, by the way the needle was removed

first from the syringe, then the syringe filter was installed at the base of the needle. After the syringe filter was installed at the base of the needle, the needle was re-fitted to the syringe. The syringe containing 10 mL was pressed so that it released the culture filtrate that escaped from the syringe filter. The 100 mL of mycoinsecticide produced  $\pm$  70 mL of culture filtrate. According to Soesanto et al. (2019), this second filtering culture filtrate was the raw secondary metabolites. Next, the culture filtrate was measured by its insecticidal activity by taking as much as 1 mL to be dropped on a piece of chili leaves.

#### Bioassays of culture filtrate and conidia of EPF

Bioassays of culture filtrate from mycoinsecticides of *P. citrinum*, *T. diversus*, *B. bassiana*, and *M. anisopliae* were carried out on chili leaves sprayed with 1 mL of culture filtrate. As a control, sterile distilled water was dripped on chili leaves. Then the leaves were dried before putting 25 second-instar *S. litura* larvae which were fasted for 2 hours. The second instar larvae were let to eat the leaves for 6 hours which had been dropped by the culture filtrate. After 6 hours the larvae were transferred into a plastic cylinder topped with gauze ( $\varnothing$  25 cm, height 30 cm) containing 15-30 pieces of chili leaves (depending on the instar) which were clean and not dripped with culture filtrate. To maintain the moisture of chili leaves, the stalk was wrapped in cotton wool moistened with sterile distilled water. Every day chili leaves were replaced with new fresh leaves. The weight of the leaves eaten was measured according to the method of Ayudya et al. (2019) and recorded daily. Every day the dead larvae were recorded up to 12 days after the application.

Bioassays of conidia from mycoinsecticides of *P. citrinum*, *T. diversus*, *B. bassiana*, and *M. anisopliae* were carried out by means of 25 second instar *S. litura* larvae dripped topically 1 mL of conidia suspension of EPF with a

concentration of  $1 \times 10^6$  conidia mL<sup>-1</sup>. Then, the larvae were fed 15-30 pieces of chili leaves. As a control it was used sterile distilled water. The weight of leaves eaten by larvae and weight of the larvae were recorded daily. Every day the dead larvae were recorded up to 12 days after the application.

The number of dead larvae was used to calculate mortality and the median Lethal Time (LT<sub>50</sub>). The weight of the chili leaves eaten and the body weight of the larvae were measured every day from the first day to 12 days after the application. The measurement of the eaten leaf weight was conducted to determine whether the EPF affected the larvae appetite, whereas the measurement of larvae body weight was carried out to determine whether or not the EPF influenced the larvae growth.


#### Data analysis

The data differences among the treatments for edible leaf weight, larvae weight, larvae mortality, and LT<sub>50</sub> were analyzed using Analysis of Variance (ANOVA) if there were differences, then further tests were carried out using the 5% Tukey's Honestly Significant Difference (HSD). LT<sub>50</sub> data were obtained from probit analysis. All calculations were conducted using software of SAS University Edition 2.7 9.4 M5.


## RESULTS AND DISCUSSION

#### Species of EPF

There were 4 EPF species namely, *P. citrinum*, *T. diversus*, *B. bassiana*, and *M. anisopliae* found from lowland and highland soils in South Sumatra. The macroscopic and microscopic appearance of the four fungal species showed obvious differences (Figure 1 and 2).



**Figure 1.** Colony of *Penicillium citrinum* (A), *Talaromyces diversus* (B), *Beauveria bassiana* (C), and *Metarhizium anisopliae*




**Figure 2.** Conidia of *Penicillium citrinum* (A), *Talaromyces diversus* (B), *Beauveria bassiana* (C), and *Metarhizium anisopliae* (D)

The colony characteristics of *P. citrinum* on SDA medium were initially white, the older the fungus, the color turned to bluish green with a white margin, round conidia with a diameter of 2.01-2.28  $\mu\text{m}$ . The characteristics matched the description described by Mathew et al. (2010) stating that *P. citrinum* has globose green granular conidia. *T. diversus* has a colony characteristic generally in the form of microcolony scattering in the medium; the color was dark green with yellow margins. According to Visagie and Jacobs (2012), the size of *T. diversus* colony ranged from 2-4 mm and sometimes only microcolony and produced yellow colored mycelia at colony peripheral zones. In this study, *T. diversus* was more difficult to culture than the other three species and showed poor growth of colony on SDA. *B. bassiana* had a colony that was white but when it got older it was yellowish or turbid white and *B. bassiana* colonized SDA faster than the other three species. Conidia of *B. bassiana* was one-celled and the same circular shape as the observation of Safitri et al. (2018). *M. anisopliae* had a colony which was also white initially similar to *B. bassiana* colony but the older the fungus the colony turned greenish white and finally dark green. Conidia of *M. anisopliae* was 6.42-6.76  $\mu\text{m}$  in size. The found characteristics of the colony and conidia of *M. anisopliae* were the same as those described by Lopes et al. (2018) and Thaochan and Sausa-Ard (2017) stating that the colony is more than 5 days old, the conidia changes from white to green indicating that the conidia was matured with the conidial mass bluish olive and has cylindrical, single and one-celled, hyaline conidia.

The result of BLAST search revealed that BKbTp showed 99.38% of similarity with *P. citrinum* strain NFML\_CH33\_5BbE.1 (Acc. No. KM458810.1), MSwTp1

showed 99.05% of similarity with *T. diversus* strain DTO\_13116 (Acc. No. KJ775700.1), BSwtD4 showed 99.65% of similarity with *B. bassiana* isolate GZMS-28 (Acc. No. KT715480.1) and MSwTp3 exhibited 99.82% of similarity with *M. anisopliae* isolate HT008 (Acc. No. MH512951.1). The result of BLAST search corresponded to those of phylogenetic tree analysis. The phylogenetic tree (Figure 3) was constructed using Mega 7 for Windows (Kumar et al. 2016) with Unweighted-pair Group Method with Arithmetic means (UPGMA). Based on the phylogenetic tree, BKbTp was placed in the group of *P. citrinum* strain NFML\_CH33\_5BbE.1 (Acc. No. KM458810.1), and SL1 (Acc. No. MN735924.1). MSwTp1 was in the group of *T. diversus* strain DTO\_13116 (Acc. No. KJ775700.1) and strain SCZ5-2 (Acc. No. KY445783.1). BSwtD4 was in the group of *B. bassiana* isolate CGAIPFBS-012 (Acc. No. KY495188.1), isolate 08F04 (Acc. No. KF359944.1), and isolate GZMS-28 (Acc. No. KT715480.1). Meanwhile, MSwTp3 was in the group of *M. anisopliae* HT008 (Acc. no. MH512951.1), A1 (Acc. No. KX806656.1), and IPPM010202 (Acc. No. KY437678.1). Thus, based on the morphological characters and molecular analysis, it confirmed that the identity of the four isolates were *P. citrinum* (BKbTp), *T. diversus* (MSwTp1), *B. bassiana* (BSwtD4), and *M. anisopliae* (MSwTp3). The isolates have been deposited in the GenBank and the accession number were MT448730 (BKbTp) (<https://www.ncbi.nlm.nih.gov/nuccore/MT448730>), MT448731 (MSwTp1) (<https://www.ncbi.nlm.nih.gov/nuccore/MT448731>), MT448732 (BSwtD4) (<https://www.ncbi.nlm.nih.gov/nuccore/MT448732>), and MT448733 (MSwTp3) (<https://www.ncbi.nlm.nih.gov/nuccore/MT448733>).



**Figure 3.** Dendrogram created on the basis of sequence analysis of ITS region using primer ITS1 and ITS4 with UPGMA method developed by MEGA7 for windows ■ = fungal isolates used in this study

### EPF insecticidal activity against *Spodoptera litura*

The EPF insecticidal activity against *S. litura* larvae was stated from the data of the effect of EPF on the weight reduction of edible chili leaves, body weight of larvae, mortality, and length of time of death. On day 1 (1 x 24 hours) after the larvae were treated with the filtrate culture of the EPF, the weight of the leaves eaten by *S. litura* larvae did not show any significant difference compared to control (sterile distilled water), but after that the weight of the leaves eaten by the larvae was more lower than control (Table 3). For the larvae treated with conidia of the EPF on day 1 (1 x 24 hours) and day 2 (2 x 24 hours), the weight of the eaten leaves did not show any significant difference compared to control, on the following day the weight of the

eaten leaves by the larvae was significantly lower than those of the control.

The symptoms of leaves infested with larvae treated with filtrate or conidia culture showed less damage, whereas, the control leaves left only the bone of the leaf (Figures 4 and 5). The decreased appetite in leaves treated with the culture filtrate of the EPF was faster than in leaves treated with the conidia of the EPF. All tested species of the EPF significantly reduced the weight of leaves eaten by larvae compared to controls (Table 3). Thus, all species of the EPF (*P. citrinum*, *T. diversus*, *B. bassiana*, and *M. anisopliae*) were effective in reducing the appetite of *S. litura* larvae.



**Table 3.** Mean of leaf weight eaten by *Spodoptera litura* larvae treated with culture filtrates and conidia of mycoinsecticides from *Penicillium citrinum*, *Talaromyces diversus*, *Beauveria bassiana*, and *Metarhizium anisopliae*

Fungal species	Leaf weight eaten by a larvae (g day <sup>-1</sup> ) on observation for 12 days											
	1	2	3	4	5	6	7	8	9	10	11	12
<b>Culture filtrate</b>												
<i>P. citrinum</i>	0.10	0.26 a	0.40 a	0.51 a	0.62 a	0.73 a	0.82a	0.96 a	1.08 a	1.17 a	1.22 a	1.24 a
<i>T. diversus</i>	0.14	0.35 a	0.46 a	0.55 a	0.65 a	0.77 a	0.86 a	0.99 a	1.09 a	1.18 a	1.23 a	1.25 a
<i>B. bassiana</i>	0.17	0.36 a	0.47 a	0.58 a	0.74 a	0.94 a	1.08 a	1.22 a	1.35 ab	1.45 ab	1.53 ab	1.59 b
<i>M. anisopliae</i>	0.17	0.36ab	0.53 a	0.65 a	0.78 a	0.97 a	1.11 a	1.26 a	1.42 b	1.55 b	1.65 ab	1.77 b
Control	0.24	0.56 b	0.94 b	1.32 b	1.74 b	2.33 b	2.91 b	3.45 b	4.02 c	4.65 c	5.04 c	5.58 c
F-value	1.96 <sup>ns</sup>	6.86*	10.28*	19.63*	44.96*	76.82*	124.79*	131.90*	203.70*	263.92*	281.95*	324.89*
P value	0.19	0.011	0.0031	0.00034	0.000016	0.000020	0.0000031	0.0000025	0.00000044	0.00000016	0.00000012	0.0000070
HSD value	-	0.10	0.14	0.15	0.12	0.12	0.12	0.13	0.117	0.115	0.119	0.122
<b>Conidia</b>												
<i>P. citrinum</i>	0.45	0.76	1.13 a	1.50 a	1.84 a	2.00 a	2.28 a	2.47 a	2.62 a	2.74 a	2.85 a	2.92 a
<i>T. diversus</i>	0.45	0.84	1.17 a	1.50 a	1.87 a	2.15 a	2.51 a	2.71 a	2.94 a	3.04 a	3.19 a	3.28 a
<i>B. bassiana</i>	0.44	0.76	1.09 a	1.43 a	1.76 a	2.07 a	2.40 a	2.59 a	2.87 a	3.01 a	3.21 a	3.45 a
<i>M. anisopliae</i>	0.34	0.70	1.08 a	1.39 a	1.78 a	2.10 a	2.35 a	2.54 a	2.72 a	2.86 a	3.01 a	3.23 a
Control	0.56	1.28	1.97 b	2.64 b	3.33 b	3.99 b	4.99 b	5.73 b	6.57 b	7.49 b	8.32 b	9.16 b
F-value	1.78 <sup>ns</sup>	3.58 <sup>ns</sup>	7.14 *	18.26 *	16.06 *	17.38 *	14.98 *	14.07 *	17.72 *	23.27 *	26.81 *	33.23 *
P value	0.23	0.06	0.01	0.0004	0.0007	0.0005	0.001	0.0011	0.0005	0.0002	0.0001	0.00005
HSD value			0.23	0.18	0.23	0.26	0.35	0.41	0.42	0.42	0.43	0.42


Note: ns = not significantly different; \* = significantly different; values within a column followed by the same letters were not significantly different at P < 0.05 according to Tukey's HSD test

**Table 4.** Mean of larvae weight of *Spodoptera litura* treated with culture filtrates and conidia of mycoinsecticides from *Penicillium citrinum*, *Talaromyces diversus*, *Beauveria bassiana*, and *Metarhizium anisopliae*


Fungal species	Weight of larvae (g larvae <sup>-1</sup> ) on observation for 12 days											
	1	2	3	4	5	6	7	8	9	10	11	12
<b>Culture filtrate</b>												
<i>P. citrinum</i>	0.06	0.09 a	0.12 a	0.15 a	0.19 a	0.22 a	0.25 a	0.29 a	0.34 ab	0.40 a	0.48 ab	0.50 a
<i>T. diversus</i>	0.06	0.09 a	0.12 a	0.15 a	0.18 a	0.22 a	0.27 a	0.31 a	0.38 b	0.46 b	0.53 b	0.58 b
<i>B. bassiana</i>	0.06	0.09 a	0.12 a	0.15 a	0.18 a	0.21 a	0.25 a	0.29 a	0.33 a	0.39 a	0.45 a	0.51 a
<i>M. anisopliae</i>	0.06	0.09 a	0.12 a	0.15 a	0.18 a	0.22 a	0.26 a	0.31 a	0.36 ab	0.41 ab	0.47 a	0.53 ab
Control	0.07	0.11 b	0.16 b	0.21 b	0.28 b	0.34 b	0.42 b	0.50 b	0.58 c	0.68 c	0.78 c	0.89 c
F-value	2.91 <sup>ns</sup>	6.31*	12.24*	21.07*	36.38*	52.94*	74.25*	109.18*	97.99*	96.07*	107.63*	40.12*
P value	0.09	0.014	0.0017	0.00026	0.000035	0.0000085	0.0000023	0.00000052	0.00000079	0.00000085	0.00000055	0.000024
HSD value	-	0.01	0.01	0.02	0.02	0.02	0.02	0.02	0.024	0.03	0.028	0.053
<b>Conidia</b>												
<i>P. citrinum</i>	0.12	0.18 ab	0.23 a	0.29 a	0.35 a	0.41 a	0.47 a	0.53 a	0.59 a	0.65 a	0.73 a	0.75 a
<i>T. diversus</i>	0.13	0.19 ab	0.25 ab	0.32 a	0.39 a	0.46 a	0.52 a	0.59 a	0.66 a	0.74 a	0.83 a	0.85 a
<i>B. bassiana</i>	0.11	0.17 a	0.24 a	0.31 a	0.38 a	0.45 a	0.50 a	0.57 a	0.64 a	0.72 a	0.81 a	0.90 a
<i>M. anisopliae</i>	0.12	0.18 ab	0.25 ab	0.31 a	0.38 a	0.45 a	0.52 a	0.59 a	0.66 a	0.74 a	0.82 a	0.91 a
Control	0.12	0.20 b	0.27 b	0.36 b	0.45 b	0.55 b	0.65 b	0.77 b	0.90 b	1.03 b	1.18 b	1.33 b

F-value	3.37 <sup>ns</sup>	4.07*	6.49*	7.49*	9.23*	12.46*	15.82*	20.61*	27.40*	32.25*	36.85*	64.25*
P value	0.07	0.04	0.01	0.01	0.004	0.002	0.001	0.0003	0.0001	0.00006	0.00003	0.000004
HSD value	-	0.01	0.01	0.02	0.028	0.03	0.04	0.04	0.05	0.05	0.05	0.05

Note: ns = not significantly different; \* = significantly different; values within a column followed by the same letters were not significantly different at  $P < 0.05$  according to Tukey's HSD test.



**Figure 4.** The symptoms on chili leaves eaten by *Spodoptera litura* larvae treated with culture filtrates of mycoinsecticides and control.



**Figure 5.** The symptoms on chili leaves eaten by *Spodoptera litura* larvae treated with conidia of mycoinsecticides and control

The decreased appetite of *S. litura* larvae treated with culture filtrate or conidia EPF was one of the stages of the larvae to be ill due to the treatment with EPF. In culture filtrate experiments, the larvae appetite already decreased on 2<sup>nd</sup> day and was faster on day 1<sup>st</sup> than the trial using the EPF conidia. In culture filtrate experiments, the decreased appetite was faster due to the raw secondary metabolite content of larvae consumed from the first day since the EPF culture filtrate was exposed, whereas in conidia treatment it required conidia time to enter the haemolymph and secrete secondary metabolites. The process of being ill due to the EPF started with the conidia attached to the cuticle larvae, and then the conidia ruptures to form germ tubes to penetrate the cuticle (Fernandes et al. 2007). The infection started when germ tubes were able to penetrate the insect cuticle and the ability to infect was a determinant of the level of virulence EPF (Altre and Vandenberg 2001). After germ tubes reach the haemocoel, they produced specific infection hyphae originating at appressoria (El-Ghany 2015). Then the hyphae spread to the haemolymph and developed to produce blastospores. The blastospores then produced the secondary metabolites and enzymes that could disrupt normal cell metabolism (Mancillas-Paredes et

al. 2019). The penetration stages of germ tubes to blastospores produced secondary metabolites and the enzymes occurred in the experiments using conidia, whereas in culture filtrate experiments this stage did not occur because the two stages of filtration were already carried out to separate conidia from the filtrate. Therefore, the process of the larvae starting to be ill in the culture filtrate was faster than the conidia experiment. The next process, the toxins from secondary metabolites began to kill the host insect but the death of the host insect was not only by the toxin but also by the mechanical damage of the fungal penetration into the body of the insect (El-Ghany 2015).

The reduction of the weight of the eaten affected the weight of the larvae. At 1 x 24 hours after the larvae were treated with culture filtrate or EPF conidia, the weight of the *S. litura* larvae did not show any significant difference, starting on the second day the larvae weight began to be lower than that of the control (Table 4), as well as the third day up to the 12 day the larvae weight was lower than that of the control. All the tested species of the EPF decreased significantly the larvae body weight compared to that of the control. Consequently, the larvae treated with mycoinsecticides containing the culture filtrate or **EPF**



conidia could reduce the larvae appetite to make their body weight decrease.

**Table 5.** Mean of larvae mortality and LT<sub>50</sub> of *Spodoptera litura* larvae treated with culture filtrates and conidia of mycoinsecticides from *Penicillium citrinum*, *Talaromyces diversus*, *Beauveria bassiana*, and *Metarhizium anisopliae*

Fungal species	Mortality ± SE (%)	LT <sub>50</sub> ± SE (days)
<b>Culture filtrate</b>		
<i>P. citrinum</i>	98.67±1.09 c	6.62±0.135 a
<i>T. diversus</i>	96.00±1.89 c	6.19±0.072 a
<i>B. bassiana</i>	86.67±2.88 b	7.55±0.154 a
<i>M. anisopliae</i>	82.67±1.09 b	7.60±0.203 b
Control	0.00±0.00 a	-
F-value	187.26*	13.12*
P value	0.00000006	0.00480044
HSD value	11.38	0.87
<b>Conidia</b>		
<i>P. citrinum</i>	98.67 ± 1.09 b	5.45 ± 0.232 a
<i>T. diversus</i>	97.33 ± 2.18 b	6.55 ± 0.081 ab
<i>B. bassiana</i>	86.67 ± 1.09 b	8.88 ± 0.420 c
<i>M. anisopliae</i>	86.67 ± 1.09 b	7.70 ± 0.404 bc
Control	0.00 ± 0.00 a	-
F-value	156.31*	29.38*
P value	0.00000013	0.00055
HSD value	12.70	1.24

Note: \* = significantly different; values within a column followed by the same letters were not significantly different at P < 0.05 according to Tukey's HSD test.


Percentage of the mortality or mortality of the larvae caused by the EPF culture filtrate showed significant differences when compared to that of the control (Table 5). All EPF species were toxic to *S. litura* larvae, both their culture filtrate and conidia. The highest larvae mortality due to culture filtrate was found in the treatment of *P. citrinum* (98.67%) and not significantly different from the larvae mortality by *T. diversus* (96%), but significantly different from the larvae mortality by *B. bassiana* (86.67%) and *M. anisopliae* (82.67%). The LT<sub>50</sub> larvae treated with culture filtrate *T. diversus* was the shortest (6.19 days) and not significantly different from the LT<sub>50</sub> larvae treated with *P. citrinum* (6.62 days), but significantly different from the LT<sub>50</sub> larvae treated with *B. bassiana* (7.55 days) and *M. anisopliae* (7.60 days). Thus, the most toxic EPF species against *S. litura* larvae was *P. citrinum*. The larvae mortality due to the EPF conidia showed significant differences when compared to that of the control (Table 5). All of the EPF species were pathogenic to *S. litura* larvae. The larvae mortality caused by the conidia was highest in *P. citrinum* (98.67%) which was not significantly different from the larvae mortality by *T. diversus* (97.33%), *B. bassiana* (86.67%), and *M. anisopliae* (86.67%). The LT<sub>50</sub> of the larvae treated with the conidia of *P. citrinum* was the shortest (5.45 days) and not significantly different from the LT<sub>50</sub> of larvae treated with *T. diversus* (6.55 days), but it was significantly different from the LT<sub>50</sub> of larvae treated with *B. bassiana* (8.88 days) and *M. anisopliae* (7.70 days).

Thus, the most pathogenic EPF species against *S. litura* larvae was *P. citrinum*.


Both culture filtrate and conidia of EPF in this study had significantly higher insecticidal activity than the control, and the highest insecticidal activity was found in *P. citrinum*. *P. citrinum* had the highest insecticidal activity due to its ability to infect the larvae faster shown by its LT<sub>50</sub> value which was the shortest compared to the other species. Altre and Vandenberg (2001) state that the most pathogenic EPF species due to the fungus has germ tubes that are able to penetrate the insect cuticle faster than other fungal species and have more toxins and enzymes that disrupt normal cell metabolism in insects. In addition, the color of the *P. citrinum* culture filtrate was the darkest compared to the color of the culture filtrate of the other three EPF species. The darker color of the EPF culture filtrate proved to be more toxic than the lighter color of the culture filtrate (Ayudya et al. 2019; Gustianingtyas et al., 2020) because the darker the color of the fungus culture filtrate the higher the concentration of the secondary metabolites (Luo et al., 2017). The *P. citrinum* culture filtrate contains secondary metabolites and is toxic to insects, for example patulin which can kill *Culex quinquefasciatus* larvae up to 100% (Maketon et al. 2014). *P. citrinum* are reported to be effective in killing insect pests (Foo et al. 2017; Maketon et al. 2014) and mite pests (Mazid et al. 2016).

The other EPF species were also toxic because they had enzymes or secondary metabolites that could kill the host insects, for example *B. bassiana* produced the enzymes of proteolytic and chitinolytic that could disrupt normal cell metabolism (Mancillas-Paredes et al. 2019). The secondary metabolite produced by *B. bassiana* containing the bassianolone B toxic to the *T. molitor* larvae (Luo et al. 2017). *M. anisopliae* produces secondary metabolites, including destruxins toxic to insect pests (Borisade et al. 2016). So far there has been no information on the secondary metabolites or enzymes produced by *T. diversus* that kills insects. At present there is only general information stating that *T. diversus* can kill insects (Visagie et al. 2009). For this reason, in the future it is necessary to study to identify the secondary metabolites or *T. diversus* enzymes that can kill insect pests.

The *S. litura* larvae treated with EPF culture or conidia showed the same behavior, i.e., they ate less and less and got lazy to move, while the control not treated with the EPF continued to eat greedily and actively moved. After 3 days the larvae given culture filtrate began to change integument color that was previously bright green to dull and began to find dead larvae and the same symptoms occurred in larvae treated with EPF conidia. Four days after the larvae were treated with culture filtrate or EPF conidia, the larvae's body began to shrink, wrinkle, get odorless and dull black, and eventually died (Figure 6 and 7). After 2-3 days the larvae died, the symptoms in the larvae integument died due to the conidia covered by mycelia and the conidia appeared, whereas in larvae that died due to the culture filtrate there was no mycelia and conidia.



**Figure 6.** Morphology of *Spodoptera litura* larvae of healthy larvae of control (right) and dead larvae (left) treated with culture filtrate of of mycoinsecticides from *Penicillium citrinum* (A), *Talaromyces diversus* (B), *Beauveria bassiana* (C), and *Metarhizium anisopliae* (D)



**Figure 7.** Morphology of *Spodoptera litura* larvae of healthy larvae of control (right) and dead larvae (left) treated with conidia of mycoinsecticides from *Penicillium citrinum* (A), *Talaromyces diversus* (B), *Beauveria bassiana* (C), and *Metarhizium anisopliae* (D)

Thus, the symptoms of ill *S. litura* larvae caused by the culture filtrate were slightly different from the symptoms caused by the EPF conidia. The symptoms of sick larvae due to the culture of this filtrate were the same as the symptoms described by Ayudya et al. (2019) and Gustianingtyas et al. (2020), namely the larvae got shrunk, wrinkled, dry, odorless, and the integument was not covered by EPF conidia. The symptoms of sick insects caused by EPF conidia in this study were the same as those described by Bugti et al. (2018) and Sumikarsih et al. (2019), that is, the insect's body got smaller, wrinkled, dry, odorless, and the integument was covered in mycelia and EPF conidia.

The results of bioassays of culture filtrate and EPF conidia showed that both culture filtrate and conidia had high insecticidal activity. However, for the future development of mycoinsecticides, the use of culture filtrate without conidia as the active ingredient of mycoinsecticide needs to be further investigated. The fungus culture filtrate is relatively more stable than the culture that still has conidia and can be stored for a longer time and remains toxic, besides the culture filtrate does not clog the sprayer nozzle when sprayed (Soesanto et al. 2019).

The conclusion of this research is that the EPF species found in the soils of South Sumatra were *P. citrinum*, *T. diversus*, *B. bassiana*, and *M. anisopliae*. All species have a high insecticidal activity against *S. litura* larvae and the most toxic species is *P. citrinum*. The filtrate and conidia of the *P. citrinum* are toxic. *P. citrinum*, a new emerging entomopathogenic fungus, has the highest insecticidal activity and the highest potential to be developed into commercial mycoinsecticides.

### ACKNOWLEDGEMENTS


This research was funded by the scheme of Applied Research (Riset Terapan) with budget year of 2020 according to Directorate of Research and Community Service (*Direktorat Riset dan Pengabdian Masyarakat*), Ministry of Research and Technology/National Research and Innovation Agency of Republic Indonesia, Grant/Contract Number: SP DIPA-042.06.1.401516/2020.

### REFERENCES

- Alhusein N, Scott J, Kasprzyk-Hordern B, Bolhuis A. 2016. Development of a filter to prevent infections with spore-forming bacteria in injecting drug users. *Harm Reduct. J.* 13: 1-8. DOI: 10.1186/s12954-016-0122-1
- Altre JA, Vandenberg JD. 2001. Factors influencing the infectivity of isolates of *Paecilomyces fumosoroseus* against diamondback moth, *Plutella xylostella*. *J. Invertebr. Pathol.* 78: 31-36. DOI: 10.1006/jipa.2001.5043
- Anwar W, Khan SN, Aslam M, Haider MS, Shahid AA, Ali M. 2015. Exploring fungal flora associated with insects of cotton agroecological zones of Punjab, Pakistan. *Pakistan Entomol.* 37: 27-31.
- Ayudya DR, Herlinda S, Suwandi S. 2019. Insecticidal activity of culture filtrates from liquid medium of *Beauveria bassiana* isolates from South Sumatra (Indonesia) wetland soil against larvae of *Spodoptera litura*. *Biodiversitas* 20: 2101-2109. DOI: 10.13057/biodiv/d200802
- Borisade OA, Medina A, Magan N. 2016. Interacting temperature and water activity modulate production of destruxin A by *Metarhizium anisopliae* on galleria larvae-modified agar based media invitro. *West African J. Appl. Ecol.* 24: 31-42.
- Bugti GA, Bin W, Na C, Feng LH. 2018. Pathogenicity of *Beauveria bassiana* strain 202 against sap-sucking insect pests. *Plant Prot. Sci.* 54: 111-117. DOI: 10.17221/45/2017-PPS
- Chen Z, Xu L, Yang F, Ji G, Yang J, Wang J. 2014. Efficacy of *Metarhizium anisopliae* isolate MAX-2 from Shangri-la, China under desiccation stress. *BBMC Microbiol* 14: 1-8. DOI: <http://www.biomedcentral.com/1471-2180/14/4>
- El-Ghany TMA. 2015. Entomopathogenic Fungi and their Role in Biological Control. Biology Department Faculty of Science Jazan University KSA, Cairo. DOI: 10.4172/978-1-63278-065-2-66
- Fernandes EKK, Rangel DEN, Moraes AM, Bittencourt VREP, Roberts DW. 2007. Variability in tolerance to UV-B radiation among *Beauveria* spp. isolates. *J Invertebr Pathol* 96: 237-243. DOI: 10.1016/j.jip.2007.05.007
- Foo K, Seelan JSS, Dawood MM. 2017. Microfungi associated with *Pteroptyx bearni* (Coleoptera: Lampyridae) eggs and larvae from Kawang River, Sabah (Northern Borneo). *Insects* 8: 1-12. DOI: 10.3390/insects8030066
- Gustianingtyas M, Herlinda S, Suwandi, Suparman, Hamidson H, Hasbi, Setiawan A, Verawaty M, Elfita, Arsi. 2020. Toxicity of entomopathogenic fungal culture filtrate of lowland and highland soil of South Sumatra (Indonesia) against *Spodoptera litura* larvae. *Biodiversitas* 21: 1839-1849. DOI: 10.13057/biodiv/d210510
- Hernandez-Trejo A, Estrada-Drouaillet B, López-Santillán, J, Rios-Velasco C, Varela-Fuentes S, Rodríguez-Herrera R, Osorio-Hernández E. 2019. In vitro evaluation of native entomopathogenic fungi and neem (*Azadirachta indica*) extracts on *Spodoptera frugiperda*. *Int. J. Exp. Bot.* 4: 75-84. DOI: .1037//0033-2909.126.1.78
- Johari A, Herlinda S, Irsan C, Pujiastuti Y. 2016. *Phenomenon of Thrips* (Thysanoptera) attack on chili plant (*Capsicum annum*). *Am. J. Agric. Biol. Sci.* 11: 103-109. DOI: 10.3844/ajabssp.2016.103.109
- Kumar S, Stecher G, Tamura K. 2016. MEGA7: Molecular evolutionary genetics analysis version 7.0 for bigger data sets. *Mol. Biol. Evol.* 33: 1870-1874. DOI: 10.1093/molbev/msw054
- Liu SF, Wang GJ, Nong XQ, Liu B, Wang MM, Li SL, Cao GC, Zhang ZH. 2017. Entomopathogen *Metarhizium anisopliae* promotes the early development of peanut root. *Plant Prot. Sci.* 53: 101-107. DOI: 10.17221/49/2016-PPS
- Lopes RB, Souza DA, Rocha LFN, Montalva C, Luz C, Humber RA, Faria M. 2018. *Metarhizium alvesii* sp. nov.: A new member of the *Metarhizium anisopliae* species complex. *J. Invertebr. Pathol.* 151: 165-168. DOI: 10.1016/j.jip.2017.12.001
- Luo Z, Ren H, Mousa JJ, Rangel DEN, Zhang Y, Bruner SD, Keyhani NO. 2017. The PacC transcription factor regulates secondary metabolite production and stress response, but has only minor effects on virulence in the insect pathogenic fungus *Beauveria bassiana*. *Environ. Microbiol.* 19: 788-802. DOI: 10.1111/1462-2920.13648

- Maketon M, Amnuaykanjanasin A, Kaysorngup A. 2014. A rapid knockdown effect of *Penicillium citrinum* for control of the mosquito *Culex quinquefasciatus* in Thailand. *World J. Microbiol. Biotechnol.* 30: 727–736. DOI: 10.1007/s11274-013-1500-4
- Mancillas-Paredes J, Hernández-Sánchez H, Jaramillo-Flores ME, García-Gutiérrez C. 2019. Proteases and chitinases induced in *Beauveria bassiana* during infection by *Zabrotes subfasciatus*. *Southwest. Entomol* 44: 125-137. DOI: 10.3958/059.044.0114
- Mascarin GM, Jackson MA, Kobori NN, Behle RW, Delalibera Júnior Í. 2015. Liquid culture fermentation for rapid production of desiccation tolerant blastospores of *Beauveria bassiana* and *Isaria fumosorosea* strains. *J. Invertebr. Pathol.* 127: 11-20. DOI: 10.1016/j.jip.2014.12.001
- Mathew AJ, Jayachandran K, Mathew J. 2010. Endophytic *Penicillium citrinum* Thom. from *Scoparia dulcis* Linn. *Indian J. Microbiol.* 50: 99–102. DOI: 10.1007/s12088-010-0065-2
- Mazid S, Kalita JC, Rajkhowa RC. 2016. Biocontrol potential of *Penicillium citrinum* and *Penicillium chrysogenum* against red spider mite, *Oligonychus coffeae* Nietner infesting tea. *J. Entomol. Res.* DOI: 10.5958/0974-4576.2016.00008.6
- Ngangambe MH, Mwatawala MW. 2020. Effects of entomopathogenic fungi (EPFs) and cropping systems on parasitoids of fall armyworm (*Spodoptera frugiperda*) on maize in eastern central, Tanzania. *Biocontrol Sci. Technol.* 30: 1-13. DOI: 10.1080/09583157.2020.1726878
- Safitri A, Herlinda S, Setiawan A. 2018. Entomopathogenic fungi of soils of freshwater swamps, tidal lowlands, peatlands, and highlands of South Sumatra, Indonesia. *Biodiversitas* 19: 2365-2373. DOI: 10.13057/biodiv/d190647
- Sharma L, Oliveira I, Torres L, Marques G. 2018. Entomopathogenic fungi in Portuguese vineyards soils: suggesting a ‘Galleria-Tenebrio-bait method’ as bait-insects *Galleria* and *Tenebrio* significantly underestimate the respective recoveries of *Metarhizium (robertsii)* and *Beauveria (bassiana)*. *MycKeys* 23: 1-23. DOI: 10.3897/mycokeys.38.26970
- Siaga E, Lakitan B, Hasbi H, Bernas SM, Widuri LI, Kartika K. 2019. Floating seedbed for preparing rice seedlings under unpredictable flooding occurrence at tropical riparian wetland. *Bulg. J. Agric. Sci.* 25: 326-336.
- Soesanto L, Mugiastuti E, Manan A. 2019. Raw secondary metabolites application of two *Trichoderma harzianum* isolates towards vascular streak dieback on cocoa seedlings. *Pelita Perkeb.* 35: 22-32. DOI: 10.22302/iccri.jur.pelitaperkebunan.v35i1.346
- Souza ML, Sanches MM, Souza DAde, Faria M, Espinel-Correal C, Sihler W, Lopes RB. 2019. Within-host interactions of *Metarhizium rileyi* strains and nucleopolyhedroviruses in *Spodoptera frugiperda* and *Anticarsia gemmatilis* (Lepidoptera: Noctuidae). *J. Invertebr. Pathol.* 162: 10-18. DOI: 10.1016/j.jip.2019.01.006
- Sowmya G, Ramanagouda SH, Jayappa J. 2017. Lethal and sub lethal effects of indigenous isolates of *Metarhizium anisopliae* (Metchnikoff) Sorokin against *Spodoptera litura* (Fabricius). *Entomol. Zool. Stud.* 5: 1154-1160.
- Sumikarsih E, Herlinda S, Pujastuti Y. 2019. Conidial density and viability of *Beauveria bassiana* isolates from Java and Sumatra and their virulence against *Nilaparvata lugens* at different temperatures. *Agrivita J. Agric. Sci.* 41: 335-349. <http://doi.org/10.17503/agrivita.v41i2.2105>
- Suwandi, Ammar M, Chandra I. 2012. Application of extract compost increased yield and suppressed the diseases of ratoon rice crop in tidal swamp of Banyuwasin Regency. *J. Suboptimal Lands* 1: 116-122. [Indonesia] DOI: 10.33230/JLSO.1.2.2012.15
- Syamsudin TS, Faizal A, Kirana, R. 2019. Dataset on antixenosis and antibiosis of chili fruit by fruit fly (*Bactrocera dorsalis*) infestation Tati. *Data Br.* 23: 1-4. DOI: 10.1016/j.dib.2019.103758
- Thaochan N, Sausa-Ard W. 2017. Occurrence and effectiveness of indigenous *Metarhizium anisopliae* against adults *Zeugodacus cucurbitae* (Coquillett) (Diptera: Tephritidae) in Southern Thailand. *Songklanakarin J. Sci. Technol.* 39: 325-334. DOI: 10.14456/sjstpsu.2017.35
- Tompe AA, Hole UB, Kulkarni SR, Chaudhari CS, Chavan SK. 2020. Studies on seasonal incidence of leaf eating caterpillar, *Spodoptera litura* (Fab.) infesting capsicum under polyhouse condition. *J. Entomol. Zool. Stud.* 8: 761-764.
- Vijayalakshmi P, Vijayalakshmi T, Naidu LN. 2016. Evaluation of certain insecticide molecules against chillipod borer, *Spodoptera litura* in Andhra Pradesh. *J. Res. ANGRAU* 44: 26-30.
- Visagie CM, Jacobs K. 2012. Three new additions to the genus *Talaromyces* isolated from Atlantis sandveld fynbos soils. *Persoonia* 28: 14-24. DOI: 10.3767/003158512X632455
- Visagie CM, Roets F, Jacobs K. 2009. A new species of *Penicillium*, *P. ramulosum* sp. nov., from the natural environment. *Mycologia* 101: 888–895. DOI: 10.3852/08-149
- White T, Bruns T, Lee S, Taylor J. 1990. Amplification and direct sequencing of fungal ribosomal RNA genes for phylogenetics, in: Innis MA, Gelfand DH, Sninsky JJ, White TJ (eds) *PCR Protocols: A Guide to Methods and Applications*. Academic Press Inc., New York.
- Yang H, Qin CS, Chen YM, Zhang GY, Dong LH, Wan SQ. 2019. Persistence of *Metarhizium* (Hypocreales: Clavicipitaceae) and *Beauveria bassiana* (Hypocreales: Clavicipitaceae) in tobacco soils and potential as biocontrol agents of *Spodoptera litura* (Lepidoptera: Noctuidae). *Environ. Entomol.* 48: 147-155. DOI: 10.1093/ee/nvy161
- Youssef NA. 2014. Effect of certain entomopathogenic fungi and nematode on the desert locust *Schistocerca gregaria* (Forsk.). *Ann. Agric. Sci.* 59: 125131. DOI: 10.1016/j.aos.2014.06.017.

#### 4. Bukti tagihan untuk penerbitan artikel





**Biodiversitas & Nusantara Biosci.**

**Society for Indonesian Biodiversity**  
 Jl. Ir. Sutami 36 A Surakarta 57126  
 Tel./Fax. 0271-663375, email: unsjournals@gmail.com  
 http://biodiversitas.mipa.uns.ac.id/  
 http://biosains.mipa.uns.ac.id/nusbioscience.htm

**INVOICE**  
for AUTHOR

DATE	10/10/2020
CUSTOMER ID	2681
DUE DATE	17/10/2020

**BILL TO**

**SITI HERLINDA**  
 Department of Plant Pests and Diseases, Faculty of Agriculture, Universitas Sriwijaya  
 Indralaya 30662, Indonesia  
 Email: sitihertinda@unsri.ac.id

**Title:**

New emerging entomopathogenic fungi isolated from soil in South Sumatra (Indonesia) and their filtrate and conidial insecticidal activity against Spodoptera litura

DESCRIPTION	TAXED	AMOUNT (IDR)
Payment for manuscript publication		3.500.000,00
Payment for English improvement		-
Cost reduction for manuscript presented in the SIB Meeting		-

**OTHER COMMENTS**

1. Currency exchange: USD 1 = IDR 14,000
2. Total bill: IDR 3.500.000,-
3. Transfer to BNI (BNINIDJA), Acc. no. 0356986994 (Dewi Nur Pratiwi)
4. Send the proof of payment to unsjournals@gmail.com and a carbon copy (CC) to dewinp11@gmail.com

Subtotal	3.500.000,00
Taxable	-
Tax rate	0,000%
Tax due	-
Other	-
<b>TOTAL IDR</b>	<b>3.500.000,00</b>

If you have any questions about this invoice, please contact  
 Dewi NP. HP +62-812-9165-0588, email: dewinp11@gmail.com  
**Terimakasih atas partisipasi anda**

