

THE EFFECT OF CHITOSAN IN THE RATION ON TEGAL DUCK PERFORMANCE

by Eli Sahara

Submission date: 17-May-2023 01:37PM (UTC+0700)

Submission ID: 2095243563

File name: CEK_THE_EFFECT_OF_CHITOSAN_IN_THE_RATION_2_1.docx (406.15K)

Word count: 2012

Character count: 13366

THE EFFECT OF CHITOSAN IN THE RATION ON TEGAL DUCK PERFORMANCE

Eli SAHARA¹, Tuti WIDJASTUTI², Roostita L. BALIA², ABUN²

¹Faculty of Agriculture, University Sriwijaya, Palembang, Indonesia

²Faculty of Animal Husbandry, University Padjadjaran, Bandung, Indonesia

Corresponding author email: elisahara.unsri@gmail.com

Abstract

The aim of the study is to examine the effects of chitosan in the ration on Tegal duck performance. The dose of chitosan used ranging from 0.0%, 0.5%, 2% and 2.5%, mixed into the basal ration. The basal diet used iso-protein and iso-energy, with protein content of 15.34% and Metabolic Energy 2809 kcal / kg (NRC, 1994). Parameters measured were feed intake, duck day production (DDP), total egg weight and feed conversion. This study uses a completely randomized design (CRD) consisting of 4 treatments and 5 replicates and each replicate consisted of two ducks. The basal diet (R0) = without chitosan as a control, R1 = 0.5% chitosan, R2 = 2% chitosan, and R3 = 2.5% chitosan. Data were analyzed using SPSS 16 statistical program (Statistical Package for Social Science). Results indicated that chitosan showed no significant effect ($P > 0.05$) on feed consumption, duck day production, total egg weight and feed conversion. In this study, treatment of chitosan 2.5% (R2) gives the best results on day duck production, total egg weight and feed conversion. From the daily egg production (DDP), treatment R2 has a result of 59%, larger than R0 (57.76%), R1 (44.1%) and R3 (46.9%). Total egg weight for R2 = 3597.7 (73.42 g / grain) also show a greater number than the treatment R0 (2769.42), R1 (2662.46), and R3 (3403.14). On feed conversion, R2 showed the smallest (1.93) compared to R0 (3.06), R1 (7.4) and R3 (2.31). This means duck in treatment R2 more efficient of feed consumption,

1.93 kilograms to produce one kilogram of eggs.

Key words: chitosan, rations, performance, Tegal ducks.

INTRODUCTION

Performance of duck production is highly dependent on the farm management such as seed, feed and disease prevention. Some of the advantages of duck eggs by reference are as follows:

1. Duck egg was spot used as an option to meet community nutrition. This is due to the nutrient content of duck eggs are very complete and easy to digest. Total protein content of duck eggs is 13.10% (Winamo and Koswara, 2002);
2. Economically, the selling price of duck eggs is more expensive than chicken eggs, so it is an alternative for farmers' additional income;
3. The advantages of duck are more resistant to disease and more tolerant of crude fiber, making it easier to choose the raw material feed.

Productivity of duck is determined by the farm management, especially feeding factors. Feed should contain nutrition according to the

needs of duck, especially for basic living needs and production. Feed can also be added with a feed supplement or feed additives to improve livestock productivity.

Chitosan is poly-glicosamin, an animal fiber origin of crustaceans which are very abundant in nature. Chitosan has the characteristics of an anti-germ, antioxidants, enzymes mobilization and fat binder. If fed to livestock as feed additive, it is predicted to be able to launch the body's metabolism.

Anti-bacterial characteristic of chitosan when mixed in the ration will protect feed from pathogenic bacteria contamination; inhibit the growth of pathogenic bacteria in the gastrointestinal tract of duck. Therefore, it would increase a large number of good bacteria to optimize digestive metabolism. Optimizing metabolism of digestive enzymes would optimize the absorption of nutrient in the small intestine.

MATERIALS AND METHODS

converted,

The material used in this study was 40 Tegal ducks of production period (aged 10 months) were kept in a cage colony (2 ducks per unit). Size per unit is 1 x 1 meter², with a total of 20 units equipped with feed and drink. Cages are than Alabio ducks also equipped with a lamp as lighting at night.

Rations are prepared with 15.34% and protein 2809 kcal/kg metabolizable energy (NRP, 2004). 900-1.100

Feed materials yellow corn, rice are bran, soybean fish meal, coconut meal, flour shells and premix. Chitosan is given as treatment with doses of 0%, 0.5%, 2% and 2.5%. Rations are given in the form of pasta, two times daily (morning and afternoon), while drinking water is given *ad libitum*.

The observation was carried out for 7 weeks. Ducks lay eggs per unit enclosure is recorded and weighed each day. The leftovers ration to the ducks also collected and weighed once a week. The parameters measured in this study were: feed intake, daily egg production, egg weight and total feed conversion. Feed consumption during the study (7 weeks) is calculated based on the amount of rations given minus the leftovers ration for a week.

Daily egg production (Duck Day Productions) is calculated based on the number that indicates the average number of eggs entirely on ducks produce at a certain time and stated in percentage (Scott, 1992, cited by Manin, 2003). Total weight of the eggs is obtained from multiplying the number of eggs with egg weights.

Feed conversion is the number which indicates the duck ability to change the sum

ranging from 6.759-7.222 g/duck. If

the amount of feed consumed by ducks every week was 965.6 to 1,031.7 grams (Figure 1). This means that the experiment ducks spend ration amount from 137.9 to 147.39 grams/head/day.

This figure is much lower feed consumption which is 215-248 g/head/day. The local ducks feed intake over 20 weeks of age, in one week between grams (Hardjosworo, 2001).

Chitosan which acts as a feed additive in duck ration is thought to contribute to protect the quality of rat on nutrition from extreme environmental influences, so the predicted balance of protein energy available in the an feed is sufficient for life and production of duck.




Figure 1. Diagram of Tegal Ducks Feed Consumption

Effect of Treatment of Egg Production (Duck Day Production)

The average of egg production (duck day production) for 7 weeks between 44.1 to 59%. This figure is slightly lower than the egg of rations

production¹ of Alabio duck, which is into the¹ production of one kilogram (kg) of eggs within a certain time unit. The weight of³ the egg is the number which indicates the average weight of the eggs produced in a given period expressed in grams per¹ egg. Data were statistically analyzed using SPSS 16, following the pattern completely randomized design (4 treatments, 5 replicates), if the treatment showed significant differences then it is continued with Duncan's multiple range test.

RESULTS AND DISCUSSIONS

Effect of Treatment of Consumption Rations

The average feed consumption¹ of ducks trial during³ the study (7 weeks) in all treatments

was 58.92 to 64.63% (Setioko, 2001). This is thought because of the difference¹ of age and the ducks¹ species used in the research.

Beside genetic factors and the duck age, the amount³ and quality of feed will affect the number of eggs¹ produced. If seen from the production percentage of trial duck eggs produced per treatment (Figure 2), the treatment with 2.5% chitosan in the ration, provided a better yield (59%) and a small ration consumption figure (984.9 g/head/week), although statistically did not show significant differences with the other treatments ($P > 0.05$). Giving of 2.5% chitosan is thought to provide a good¹ influence in protecting the nutritional quality of feed because of its characteristics as

anti-oxidant, in addition to chitosan role as that the treatment R2 has the highest efficiency dietary fiber and prebiotic (gives a positive with a reduced ration feed conversion by 6.93% effect on microflora proliferation which is beneficial in the duck gastrointestinal tract, so that it will facilitate the digestive system.

(from 3.06 to 1.93) compared with the control diet (R0 = ration without chitosan).




Figure 2. 7 Weeks Tegal Duck Day Production

Effect of Treatment on Mean of Eggs Total Weight

Total weight of the eggs is representing the egg production (egg weights) obtained and multiplied by the number of eggs with egg weights (Figure 3). The average total weight of egg during 7- weeks is 2662.46 to 3597.7 gram. This means that the weight of the egg treatment R0 = 56.52, R1 = 54.34, R2 = 73.42 and R3 = 45 grams/egg. Based on the observation, total weight of the eggs in treatment R2 with 2.5% chitosan shows heavier than treatments (R0, and other R1, R3). The total egg weight of 3597.7 (R2 = 2.5% chitosan) is presumably this higher, because dose chitosan is able to activated the digestive enzymes in the duck gastrointestinal tract; so the nutrient absorption is better, especially




Figure 3. Total Egg Weight of Tegal Duck




Figure 4. Tegal Ducks Feed Conversion

This result is more efficient than the feed conversion for the Indonesian ducks is 3.2 to 5.0 (Ketaren, 2007). It is further explained that the poor feed conversion in laying ducks in Indonesia is allegedly caused by three factors, namely genetic quality, the number of scattered

pellet, and nutritional value of the ration supplied which does not match the needs of the ducks.

In this study it is predicted that, with a dose of the protein and fat; as the egg primary material The egg composition consisted of the albumin with 11% protein content and 0%

fat, while the yolk

contains 17% protein and 35%.

Effect of Treatment on Feed Conversion

The average of feed conversion was between

1.93 to 7.4. (R0 = 3.06 ± 1.98 , R1 = 7.4 ± 11.23 , R2 = 1.93 ± 0.16 , and R3 = 1.02 ± 2.31).

The lowest of the feed average conversion (1.93) achieved from treatment R2 is (2.5% chitosan in the ration). These results indicated

2.5% chitosan, it is able to protect the quality of the nutrient rations from extreme contamination of germs or other environmental influences, so the health of duck is well maintained and able to efficiently utilize the ration.

CONCLUSIONS

The 2.5% chitosan in the ration (R2) is able to provide better production performance (feed consumption 984.9 grams/week, duck day production 59%, egg total of 3597.7 grams or 73.42 per egg, and feed conversion of 1.93), when compared to other treatments (R0, R1 and R3).

REFERENCES

- Winarno F.G., S. Koswara, 2002. *Telur: Komposisi, Penanganan dan Pengolahannya*. M-Brio Press. Bogor, p.6.
- Nutrient Requirements of Poultry (NRP), 1994. National Academy Press. Washington DC, p.42.
- Manin F., 2003. Efektivitas Kultur *Bacillus* sp dan *Sacharomyces cerevisiae* sebagai Sumber Probiotik dan Implikasinya terhadap Produktivitas Ternak Itik Lokal Kerinci. Disertasi. Program Pasca Sarjana Universitas Padjadjaran, Bandung, p.70.
- Steel R.G.D., Torrie J.H., 1991. *Prinsip dan Prosedur Statistika. Suatu Pendekatan Biometrik*. Alih Bahasa Bambang Sumantri. Jakarta: PT. Gramedia, p. 168- 205.
- Setioko R., 2001. Keragaan Produksi Telur Pada Sentra Pengembangan Agribisnis Komoditas Unggulan (SPAKU) Itik Alabio Di Kabupaten Hulu Sungai Utara, Kalimantan Selatan. Prosiding Pengembangan Agribisnis Unggas Air Sebagai Peluang Usaha Baru. Kerjasama Fakultas Peternakan IPB, Balai Penelitian Ternak Pusat Penelitian dan Pengembangan Peternakan, dan Yayasan Kehati. Ciawi Bogor. Indonesia, p.139 – 145.
- Hardjosworo P.S., A. Setioko, P.P. Ketaren, L.H. Prasetyo, A.P. Sinurat and Rukmiasih, 2001. Perkembangan Teknologi Peternakan Unggas Air di Indonesia. Prosiding Pengembangan Agribisnis Unggas Air sebagai Peluang Usaha Baru. Kerjasama Fakultas Peternakan IPB, Balai Penelitian Ternak, Pusat Penelitian dan Pengembangan Peternakan, dan Yayasan Kehati. Ciawi, Bogor, p.22-41.
- Ngo D.H., Kim S.K., 2014. Antioxidant Effects of Chitin, Chitosan, and Their Derivatives. *Advance in Food and Nutrition Research*. Vol. 73: 15-31.
- Ketaren P.P., 2007. Peran Itik Sebagai Penghasil Telur dan Daging Nasional. *Wartazoa*. Vol. 17 (3), 117 – 127.

THE EFFECT OF CHITOSAN IN THE RATION ON TEGAL DUCK PERFORMANCE

ORIGINALITY REPORT

22%

SIMILARITY INDEX

22%

INTERNET SOURCES

1%

PUBLICATIONS

4%

STUDENT PAPERS

PRIMARY SOURCES

1	cdpp.irap.omp.eu Internet Source	16%
2	mafiadoc.com Internet Source	2%
3	docplayer.hu Internet Source	1%
4	hl.udogs.net Internet Source	1%
5	Submitted to 8398 Student Paper	1%
6	peda.net Internet Source	1%

Exclude quotes On

Exclude bibliography On

Exclude matches < 15 words