

# Analysis of Factors Affecting Exposure...

*by* Muhammad Irsan Saleh

---

**Submission date:** 17-May-2023 08:49AM (UTC+0700)

**Submission ID:** 2095062383

**File name:** of\_the\_Public\_Health\_Center\_in\_Agung,\_Musi\_Banyuasin\_Regency.pdf (139.59K)

**Word count:** 5055

**Character count:** 26901

# Analysis of Factors Affecting Exposure to Nicotine in Breastfeeding Mothers (ASI) in the Work Area of the Public Health Center in Agung, Musi Banyuasin Regency

M Marlini<sup>1\*</sup>, Amin Rejo<sup>2</sup>, S Suheryanto<sup>2</sup>, Irsan Saleh<sup>2</sup>  
{marliniyalin05@gmail.com}

<sup>1</sup>Department of Environmental Health  
Environmental Science Masters Program, Sriwijaya University  
Jl. Padang Tuesday No. 524 Bukit Besar Ilir Barat I Palembang, Indonesia, 30139  
<sup>2</sup>Lecturer at the Department of Environmental Health  
Sriwijaya University Environmental Science Postgraduate Program  
Jl. Padang Tuesday No. 524 Bukit Besar Ilir Barat I Palembang, Indonesia, 30139

**Abstract.** Cigarette smoke is a common air pollutant that affects the health of both active and passive smokers. Passive smoking refers to individuals who are exposed to the smoke exhaled by smokers or volatile compounds from burning tobacco products. The percentage of smoking behavior in homes with babies and toddlers continues to rise, with 85% of family members smoking in their homes as of May 2022. This study utilized a quantitative research design with a case-control study to investigate the factors affecting nicotine exposure in breastfeeding mothers in the Balai Agung Public Health Center's working area. The study utilized a point-time approach, with each object observed only once. The results of the study showed that age and education level had no significant impact on nicotine exposure, while knowledge, length of exposure to cigarette smoke per day, and the ventilation conditions of the house were significant factors affecting exposure to nicotine in nursing mothers. To address this issue, the Musi Banyuasin District Health Office has issued regulations prohibiting smoking in homes. Furthermore, Regular health checks focused on the condition of breastfeeding mothers and their babies are also essential to ensure their well-being. The Balai Agung Health Center must be more proactive in taking measures to protect their patients from the hazards of smoking.

**Keywords:** Passive Smokers, Breastfeeding Mothers, Exposure to Cigarette Smoke, Nicotine

## 1. Introduction

Air is a determining factor that plays a role in assessing the quality of sustainable public health up to a certain regional health level[3]. Human survival is largely determined by the quality of the surrounding air, either directly or indirectly. Since the beginning of the industrial revolution, technological developments have certainly been very close and linear with the increasing amount of pollutants in the air[4]. Until now, open public spaces with healthy air quality have become a rarity in big cities. Air quality is assessed by reviewing existing pollutant level parameters, such as measurements of PM 2.5, PM 10, COx, NOx., organic gas components and other volatile airborne dust particles.[5], [6]. Currently, air that has pollutant levels that exceed the threshold value is an interesting topic to study. In particular, intensive discussion of its impact on health. Air pollution can originate from activities that still use conventional energy sources. Not only that, pollutants can also come from vehicle exhaust emissions, a number of deep-rooted activities such as burning household waste, clearing land by burning, even from burning cigarettes.[6]–[8].

Based on data obtained from the Musi Banyuasin District Health Office in 2019, 76% of family members who have babies and toddlers smoke indoors. In 2020 the percentage will increase to 81% of family members who have babies and toddlers who smoke in the house. By 2021, smoking in the home will increase by 83%. The percentage of smoking behavior in the home continues to increase so that in May 2022 it was recorded that 85% of family members with babies and toddlers smoked in the house. In 15 sub-districts in Musi Banyuasin Regency, one of the sub-districts with

the highest number of smokers is Sekayu District, especially at the Balai Agung Health Center with the percentage of family members who smoke inside the house as much as 82% in 2021.

The results of the analysis by the Musi Banyuasin Regency Health Office regarding this increase were the impact of the Covid-19 pandemic. Family members spend a lot of time at home, as well as family members who are active smokers [32]. Based on these data, the authors are interested in conducting research with the title "Analysis of Factors Affecting Nicotine Exposure in Breastfeeding Mothers (ASI) in the Work Area of the Balai Agung Health Center, Musi Banyuasin Regency".

## 2. Research methods

### 2.1 Research design

The quantitative research used in this study was a case-control study design. The method used in this study was a one-time approach, meaning that each research subject was only observed once and measurements were taken about the state of the nature or variable of the object at the time of the study. research. test time by quantitative analysis.

#### 2.1.1 Time and Place of Research

This research will be conducted in the working areas of the Balai Agung Health Center, namely Balai Agung Village, Serasan Jaya Village, Soak Baru Village & Sukarami Village. The research was conducted in June-July 2022.

#### 2.1.2 Techniques for collecting data

The sampling technique in this study was purposive sampling, namely sampling based on a specific consideration made by the researcher himself based on previously known characteristics or traits. Breastfeeding mothers included in the inclusion and exclusion criteria were 70 people using 35 problem respondents and 35 control respondents. So the number of respondents in this study amounted to 70 people in one month.

### 2.2 Data collection

Data collection techniques used are:

1. Determine the sample to be observed/data taken first.
2. Primary data collection with questionnaires and interviews
3. Analyzing the relationship between age, mother's education level, mother's knowledge, length of exposure to husband's cigarette smoke, ventilation conditions at home on the quality of breast milk in the working area of the Balai Agung Health Center, Musi Banyuasin Regency.
4. By analyzing the relationship between the dependent variable and several independent variables that are suspected to have a relationship based on multivariate analysis, it can be seen which variable is most relevant to the dividend variable. This study uses multiple logistic regression analysis.

### 2.3 Data analysis

#### 2.3.1 Univariate

Analysis of each variable is done from the search results. This analysis describes each variable presented in the frequency distribution table. In this study, univariate analysis was carried out on the variables of age of breastfeeding mothers, education level of breastfeeding mothers,

knowledge of breastfeeding mothers, length of daily exposure to cigarette smoke, and house ventilation.

### 2.3.2 Bivariate

Two variables, namely the independent variable and the dependent variable which will be seen the relationship then used the Chi-Square statistical test.

## 3. Results and Discussion

### 3.1 Bivariate

The relationship between the two variables, namely dependent and independent, will be seen.

#### 3.1.1 Correlation between Maternal Age and Nicotine Exposure in Breastfeeding Mothers

Based on the trial of the relationship between maternal age and nicotine exposure in breastfeeding mothers in the working area of the Balai Agung Treatment Center, Musi Banyuasin Regency, the following results were obtained:

**Table 1.** Maternal Age With Exposure to Nicotine in Breastfeeding Mothers

Age	Nicotine Exposure				P-value	OR	95% CI
	Active Smoker Husband		Passive Smoker's Husband				
	N	%	N	%			
25-35 Years	25	71.4	21	60.0	0.450	1.667	0.615-4.519
<25 and >35 Years	10	28.6	14	40.0			
Amount	115	50	35	50			

Source: Research Primary Data, 2022

Based on table 2.1 it is known that the results of bivariate analysis with the chi-square test obtained a P-value of 0.450. The value of  $p > \alpha$  (0.05) means that  $H_0$  is accepted and  $H_a$  is rejected, which means that there is no relationship between maternal age and exposure to nicotine in breastfeeding mothers at the Balai Agung Health Center, Musi Banyuasin Regency. The Odd Ratio value was 1.667 (95% CI = 0.615-4.519), which means that mothers aged <25 and >35 years had a 1.667 times higher risk of exposure to nicotine in nursing mothers.

#### 3.1.2 Correlation between Mother's Education Level and Nicotine Exposure in Breastfeeding Mothers

Based on testing the relationship between the education level of the mother and nicotine exposure in breastfeeding mothers in the Working Area of the Balai Agung Health Center, Musi Banyuasin Regency, the following results were obtained:

**Table 2.** Mother's Education Level With Exposure to Nicotine in Breastfeeding Mothers

Mother's Education Level	Nicotine Exposure				P-value	OR	95% CI
	Active Smoker Husband		Passive Smoker's Husband				
	N	%	N	%			
Low (Elementary-junior high school graduates)	3	8.6	5	14.3	0.707	0.563	0.124-2.560

Mother's Education Level	Nicotine Exposure				P-value	OR	95% CI
	Active Smoker Husband		Passive Smoker's Husband				
	N	%	N	%			
High (High School Graduates-Completed D3/S1/S2/S3 College)	32	91.4	30	85.7			
Amount	11	35	50	35	50		

Source: Research Primary Data, 2022

Based on table 2.2 it is known that the results of bivariate analysis with the chi-square test obtained a P-value of 0.707. The value of  $p > \alpha$  (0.05) means that  $H_0$  is accepted and  $H_a$  is rejected, which means that there is no relationship between the education level of the mother and exposure to nicotine in breastfeeding mothers at the Balai Agung Health Center, Musi Banyuasin Regency.

### 3.1.3 Relationship between Mother's Knowledge and Nicotine Exposure in Breastfeeding Mothers

Based on testing the relationship between mother's knowledge and nicotine exposure in breastfeeding mothers in the Working Area of the Balai Agung Health Center, Musi Banyuasin Regency, the following results were obtained:

Table 3. Mother's Knowledge With Nicotine Exposure in Breastfeeding Mothers

Mother Knowledge	Nicotine Exposure				P-value	OR	95% CI
	Active Smoker Husband		Passive Smoker's Husband				
	N	%	N	%			
Adequate (<56%-75%)	14	40.0	25	71.4	0.016	0.267	0.098-0.723
Good (76%-100%)	21	60.0	10	28.6			
Amount	11	35	50	35			

Source: Research Primary Data, 2022

Based on table 2.3 it is known that the results of bivariate analysis with the chi-square test obtained a P-value of 0.016. The value of  $p < \alpha$  (0.05) means that  $H_0$  is rejected and  $H_a$  is accepted, which means that there is a relationship between mother's knowledge and exposure to nicotine in breastfeeding mothers at the Balai Agung Health Center, Musi Banyuasin Regency. The Odd Ratio value was 0.267 (95% CI = 0.098-0.723), which means that the level of knowledge of mothers is low/sufficient, 3.273 times more at risk of exposure to nicotine in breast milk of mothers who are breastfeeding.

### 3.1.4 Relationship between duration of exposure to cigarette smoke smoked by mothers per day and exposure to nicotine in breastfeeding mothers

Based on the correlation test between long exposure to cigarette smoke smoked by mothers per day and exposure to nicotine in breastfeeding mothers in the Working Area of the Balai Agung Health Center, Musi Banyuasin Regency, the following results were obtained:

Table 4. Duration of Exposure to Cigarette Smoke Per Day With Exposure to Nicotine in Breastfeeding Mothers

Length of exposure to cigarette smoke per day	Nicotine Exposure				P-value	OR	95% CI
	Active Smoker Husband		Passive Smoker's Husband				
	N	%	N	%			
>6-12 hours	21	60.0	11	31.4	0.031	3,273	1224-8748
0-6 hours	14	40.0	24	68.6			
Amount	35	50	35	50			

Source: Research Primary Data, 2022

Based on table 2.4 it is known that the results of bivariate analysis with the chi-square test obtained a P-value of 0.031. The p value  $< \alpha$  (0.05) so that  $H_a$  is accepted, which means that there is a relationship between exposure to cigarette smoke per day and exposure to nicotine in breastfeeding mothers at the Balai Agung Health Center, Musi Banyuasin Regency. The Odd Ratio value was 3.273 (95% CI = 1.224-8.748), meaning that the duration of exposure to cigarette smoke per day was 3.273 times the risk of exposure to nicotine in nursing mothers who were breastfeeding.

### 3.1.5 Correlation between Home Ventilation Conditions and Nicotine Exposure in Breastfeeding Mothers

Based on testing the relationship between home ventilation conditions and nicotine exposure in breastfeeding mothers in the Working Area of the Balai Agung Health Center, Musi Banyuasin Regency, the following results were obtained:

Table 5. Conditions of House Ventilation with Nicotine Exposure in Breastfeeding Mothers

Home Ventilation Conditions	Nicotine Exposure				P-value	OR	95% CI
	Active Smoker Husband		Passive Smoker's Husband				
	N	%	N	%			
Not Good (> 10% x Room Floor Area)	23	65.7	10	28.6	0.004	4,792	1741-13188
Good (<10% x Room Floor Area)	12	34.3	25	71.4			
Amount	35	50	35	50			

Source: Research Primary Data, 2022

Table 2.5 shows that the results of bivariate analysis with the chi-square test obtained a P-value of 0.004. The value of  $p < \alpha$  (0.05) means that  $H_0$  is rejected, which means that there is a relationship between the ventilation conditions of the house and exposure to nicotine in breastfeeding mothers at the Balai Agung Health Center, Musi Banyuasin Regency. The Odd Ratio value was 4.792 (95% CI = 1.741-13.188), meaning that indoor ventilation conditions were 4.792 times faster at risk of exposure to nicotine in breast milk of mothers who are breastfeeding.

## 3.2 Discussion

### 3.2.1 Maternal Age Relationship with Exposure to Nicotine in Breastfeeding Mothers

The results of this study were in line with research (Masni, 2012) which showed that there was no significant relationship between maternal age and nicotine exposure in nursing mothers. Masni's study found that the characteristics of the respondents in the case group were that there were more mothers aged between 20 to 35 years compared to mothers aged 35 years. In addition, the sample size in Masni's study (2012) between cases and controls had a ratio of 1:2 due to the limited sample of cases, in contrast to this study, namely the sample size of cases and controls (1:1) with a wider coverage. research area.

### 3.2.2 Correlation between Mother's Education Level and Nicotine Exposure in Breastfeeding Mothers

In research [12], in line with the results of this study, it was explained that the results of the analysis showed a significance level = 0.131 while there was no significant relationship between education and the practice of preventing exposure to cigarette smoke. Education affects the way a person learns because the higher the education, the easier it is to receive information from the formal education they have received as well as from other people and the media. The more information a person has, the more his knowledge develops. This is also supported by research findings [13] which confirms that there is no relationship between education and childbirth practices in an effort to protect families from exposure to cigarette smoke.

According to Notoamodjo (2012) explains the concept of education as a learning process, meaning that in education there is a process of growth, development or change in individual groups or society. The higher a person's education, the easier it is to accept new things and adapt to them. Low educational attainment made it difficult to receive instruction, including information about the dangers of nicotine exposure in nursing mothers. A good level of education will make it easier to obtain information, especially related to meeting the nutritional needs of children to ensure their nutritional adequacy.

According to [14], the low level of maternal education causes a lack of knowledge of mothers to solve problems, especially to receive information about the dangers of nicotine exposure to mothers. This knowledge is obtained both formally and informally. Meanwhile, mothers with more education are often more open to accepting changes or things to maintain their health. Educational attainment is thought to be related to knowledge of breastfeeding mothers, this is related to the level of knowledge of mothers that someone with higher education will have broader knowledge than those with low education.

### 3.2.3 Mother Knowledge Relationship with Exposure to Nicotine in Breastfeeding Mothers

According to Notoatmodjo (2007), the result of human perception or the result of someone experiencing an object through their senses is an understanding of knowledge. Knowledge is an important area in shaping behavior. Behavior based on knowledge will last longer and be more accepted in society than behavior that is not based on knowledge. So someone who already has good knowledge will behave well because of knowledge-based behavior [36].

The results of this study also have the same results as [12] showing that knowledge and actions to avoid exposure to cigarette smoke actually have a relationship. The results of this study also lead to [15] which confirms that there is a relationship between subject knowledge and practice in prevention practices. The more someone knows or understands something, the better their practice will be. On the other hand, bad knowledge will lead to bad practice.

### 3.2.4 Correlation between duration of exposure to cigarette smoke per day and exposure to nicotine in breastfeeding mothers

Based on (Wardaningtyas., AK Prakoso DA 2012) a family with 1 or more smokers in the house has a major effect on the weight gain of a growing baby. More than 5,300 compounds have been identified in tobacco smoke from tobacco or cigarette smoke, namely exposure to a mixture of compounds from tobacco smoke and the smoke exhaled by the smoker. This group of compounds includes carbon and amides, imides, lactams, metals, lactones, N-nitrosamines, aldehydes, nitrogen oxides of ketones, carboxylic acids, alcohols, phenols, esters, amines, N-heterocyclics, nitriles, nitro compounds, anhydrides, carbohydrates, ethers and hydrocarbons (IARC, 2012).

Passive smoking is as dangerous as active smoking because passive smokers indirectly or forcibly inhale active smokers' cigarette smoke and unknowingly contaminate the people around them. Or in general passive smokers are people who do not smoke but inhale environmental tobacco

smoke (ETS), especially primary tobacco smoke and second hand cigarette smoke exhaled by smokers.[16]

The amount of exposure to tobacco smoke that a person receives depends on the environment in which that person lives. If someone lives with a family member who smokes, that person will be exposed to secondhand smoke for a longer time. The longer a person is exposed to secondhand smoke, the higher the risk of adverse health effects. Where the health problems that occur in passive smokers depend on the type and length of exposure to tobacco smoke in the environment they live in [17] [18].

Breastfeeding mothers who are constantly exposed to cigarette smoke will experience a decrease in their supply of breast milk (ASI) and a shorter duration of breastfeeding because their body stops producing breast milk.[19] [20][21][22] Secondhand smoke or nicotine alters the mother's hormone levels, causing decreased lactation. The results of research and experiments clearly show that the levels of the hormone prolactin will be reduced by the presence of nicotine in the blood. Even though the role of the hormone prolactin is what encourages lactation and stimulates the development of the mammary glands. This is what causes milk production to stop.[23] [24][25][26].

The results of [9]'s study [27] also provide evidence that mothers with a lot of household exposure to smokers have a much higher risk of stopping breastfeeding than vice versa. The practitioner can assess the current regimen of smoking family members to provide smoking cessation education and support for this high-risk group. Further research is needed to analyze the mechanism of smoking in breastfeeding practice.

### 3.2.5 Correlation between Home Ventilation Conditions and Nicotine Exposure in Breastfeeding Mothers

Smoking behavior does not only affect smokers but also non-smokers around or commonly known as passive smokers. Passive smokers receive 75% of the light from passive smokers due to direct exposure [33]. Most smoking is the habit of family members to smoke in the house, which is bad for the health of other family members who do not smoke. Children and women are the group most at risk for smoke-related disorders, especially at home because of the time they spend in the bedroom. again [28]. The increasing problem of passive smoking in closed living or work environments allows the effects of secondhand smoke to occur. This shows the double danger of smoking not only for the smoker himself but also for the people around him. Therefore, controlling smokers who cause cigarette smoke is very dangerous for the health of active smokers and passive smokers is a solution to breathing fresh air without exposure to cigarette smoke or often called building a smoke-free living space. [29].

Exposure to cigarette smoke can reduce milk production because it interferes with the hormones prolactin and oxytocin in milk production. Smoking stimulates the release of adrenaline where adrenaline inhibits the release of oxytocin (Haryono & Setianingsih, 2014). According to research [30], it shows that there is an effect of passive smoking on regular milk production in Bendan Village, Banyudono Regency, Boyolali Province (p value = 0.010 < 0.05). This study is also in line with research (Susanna, 2003) which used descriptive research which showed that nicotine was 4-6 times higher in passive smokers compared to regular cigarette smoke, so that smokers have a higher risk of exposure. . passive smokers compared to active smokers. for health[31].

Many family members, especially men, are active smokers but do not understand and are not aware of the conditions around them when they smoke. Researchers have seen husbands and grandfathers smoking indoors even when they are around their wives, children and grandchildren. Awareness among breastfeeding mothers that smoking should be avoided is also lacking in allowing mothers to breastfeed in places where there is a risk of exposure to secondhand smoke.[30]. Exposure to secondhand smoke increases a child's risk of developing lung infections, asthma, and sudden infant death syndrome [31].



#### 4. Conclusion

1. Mother's age with nicotine exposure in breastfeeding mothers in the Working Area of the Balai Agung Health Center, Musi Banyuasin Regency has no significant relationship.
2. The education level of mothers with nicotine exposure in breastfeeding mothers in the working area of the Balai Agung Health Center, Musi Banyuasin Regency also does not have a significant relationship which can be interpreted that nicotine exposure in breast milk is not due to mother's education.
3. Mother's knowledge of nicotine exposure in breastfeeding mothers has a significant relationship.
4. Length of exposure to cigarette smoke per day with exposure to nicotine in nursing mothers has a significant relationship.
5. The condition of home ventilation with nicotine exposure in nursing mothers has a significant relationship, which means that good ventilation in the home will reduce the impact of exposure to nicotine if the family member is an active smoker, but you should not smoke in the house.

#### 5. Suggestion

1. To the Musi Banyuasin District Health Office to issue regulations regarding the prohibition of smoking in the house.
2. To the Balai Agung Health Center to be more active in preventive and promotive activities related to the dangers of active and passive smoking and to carry out regular health checks focused on the condition of breastfeeding mothers and their babies.
3. Examination in the laboratory regarding the content of breast milk (ASI) in nursing mothers who are exposed to cigarette smoke which contains nicotine and others as well as the long-term danger to the baby's growth.

#### References

- [1] W. Schober, L. Fembacher, A. Frenzen, and H. Fromme, "Passive exposure to pollutants from conventional cigarettes and new electronic smoking devices (IQOS, e-cigarettes) in passenger cars," *int. J. Hyg. environment. Health*, vol. 222, no. 3, p. 486–493, 2019, doi: <https://doi.org/10.1016/j.ijheh.2019.01.003>.
- [2] International Agency for Research on Cancer, "Monography on the Evaluation of Carcinogenic Risk in Humans - Volume 89," *Smokeless Tobacco. Some Tobacco-specific N-Nitrosamines*, vol. 89, 2007.
- [3] ML Melamed, J. Schmale, and E. von Schneidmesser, "Policy sustainability—key considerations for air quality and climate change," *Kur. Opinion. environment. maintain.*, vol. 23, p. 85–91, 2016, doi: <https://doi.org/10.1016/j.cosust.2016.12.003>.
- [4] D. Vallero, "Chapter 29 - Air Pollutant Emissions," in *Air Pollution Fundamentals (Fifth Edition)*, DBT-F. of the AP (Fifth E. Vallero, Boston Ed.: Academic Press, 2014, pp. 787–827. doi: <https://doi.org/10.1016/B978-0-12-401733-7.00029-3>.
- [5] H. Bravo Alvarez, R. Sosa Echeverria, P. Sanchez Alvarez, and S. Krupa, "Air Quality Standards for Particulate Matter (PM) in Upland Cities," *environment. Pollution.*, vol. 173, p. 255–256, 2013, doi: <https://doi.org/10.1016/j.envpol.2012.09.025>.
- [6] Z. Salameh, "Chapter 1 - Factors Promoting Renewable Energy Applications," in *Renewable Energy System Design*, ZBT-RES D Salameh, Ed. Boston: Academic Press, 2014, p. 1–32. doi: <https://doi.org/10.1016/B978-0-12-374991-8.00001-5>.
- [7] E. von Schneidmesser, K. Steinmar, EC Weatherhead, B. Bonn, H. Gerwig, and J. Quedenau, "Air pollution at the human scale in urban environments: Impact of local and vehicle environments on particle count concentrations," *science. Total Environment.*, vol. 688, p. 691–700, 2019, doi: <https://doi.org/10.1016/j.scitotenv.2019.06.309>.
- [8] SC van der Zee, PH Fischer, and G. Hoek, "Air pollution in perspective: The health risks of air pollution are expressed in terms of the equivalent of passively smoking a cigarette," *environment. res.*, vol. 148, p. 475–483, 2016, doi: <https://doi.org/10.1016/j.envres.2016.04.001>.
- [9] NL Benowitz, "Smoking and Nicotine Addiction," *Clinical Medicine. Am North.*, vol. 76, no. 2, p.

- 415–437, 1992, doi: [https://doi.org/10.1016/S0025-7125\(16\)30360-1](https://doi.org/10.1016/S0025-7125(16)30360-1).
- [10] J. Hukkanen, P. 3rd Jacob, and NL Benowitz, "Metabolism and kinetics of nicotine disposition." *Pharmacol. round.*, vol. 57, no. 1, p. 79–115, March 2005, doi:10.1124/pr.57.1.3.
- [11] International Agency for Research on Cancer, "i," *Tob. Involuntary Smoke Smoke.*, vol. 83, p. Volumes 83, 2004.
- [12] TB Indrajati, T. Istiarti, and A. Kusumawati, "Factors Associated with Mother's Actions in Preventing Exposure to Cigarette Smoke in Toddlers of Passive Smokers," *J. Health. Mass.*, vol. 5, no. 5, p. 1123–1131, 2017.
- [13] S. Rahayu, B. Widjanarko, and A. Kusumawati, "Factors related to the practice of protecting families from exposure to secondhand smoke," *J. Health. Mass.*, vol. 7, no. 4, p. 219–225, 2019.
- [14] F. Bancin, "Factors Influencing Working Mothers Not Giving Exclusive Breastfeeding in the Work Area of the Singkohor Health Center, Kec. Singkohor Regency. Aceh Singkil, Aceh Province in 2019," p. 20, 2019.
- [15] VF Kasenda, JAM Rattu, FRR Maramis, F. Health, M. University, and S. Ratulangi, "The Relationship between Knowledge and Attitudes About the Dangers of Smoking and Smoking in Students at SMP N VIII Manado City," 2015.
- [16] L. Nurwidayanti and CU Wahyuni, "Analysis of the effect of exposure to cigarette smoke at home in women on the incidence of hypertension," *J. Berk. Epidemioles.*, vol. 1, no. 2, p. 244–253, 2013.
- [17] MST Dayu, "The relationship between a long history of smoking and the incidence of coronary heart disease (PKK) at the cardiac polyclinic at dr. H. Abdul Moeloek, Lampung Province in 2015," 2015.
- [18] I. Mawardi, Nurdin, and Zulkarnain, "Postharvest Coffee Appropriate Technology Program: Business Units for Production, Marketing, and Coffee Processing Machines" *Indonesia. J. Community Engagement.*, vol. 5, no. 2, p. 267–284, 2019.
- [19] MS Clements, EA Mitchell, SP Wright, A. Esmail, DR Jones, and RPK Ford, "Effects on breastfeeding in southeast England," *Acta Paediatr. int. J. Paediatr.*, vol. 86, no. 1, p. 51–56, 1997, doi:10.1111/j.1651-2227.1997.tb08831.x.
- [20] BL Horta, MS Kramer, and RW Platt, "Maternal smoking and risk of early weaning: A meta-analysis," *I. J. Public Health.*, vol. 91, no. 2, p. 304–307, 2001, doi:10.2105/AJPH.91.2.304.
- [21] A. Knudsen, H. Pedersen, and J. Klebe, "Effect of smoking on duration of breastfeeding in mothers with insulin-dependent diabetes mellitus," *Acta Paediatr. int. J. Paediatr.*, vol. 90, no. 8, p. 926–930, 2001, doi:10.1111/j.1651-2227.2001.tb02459.x.
- [22] GW Letson, KD Rosenberg, and L. Wu, "Relationship between Smoking During Pregnancy and Breastfeeding at Around 2 Weeks of Age," *J. Hum. lacquer.*, vol. 18, no. 4, p. 368–372, 2002, doi:10.1177/089033402237910.
- [23] CA Blake and CH Sawyer, "Nicotine blocks lactation-induced increase in prolactin in lactating mice," *Science (80- )*, vol. 177, no. 4049, p. 619–621, 1972, doi:10.1126/science.177.4049.619.
- [24] F. Vio, G. Salazar, and C. Infante, "Smoking during pregnancy and breastfeeding and its effect on breast milk volume," *I. J. Clin. nutrition.*, vol. 54, no. 6, p. 1011–1016, 1991, doi:10.1093/ajcn/54.6.1011.
- [25] BL Horta, CG Victora, AM Menezes, and FC Barros, "Environmental tobacco smoke and duration of breastfeeding," *I. J. Epidemioles.*, vol. 146, no. 2, p. 128–133, 1997, doi:10.1093/oxfordjournals.aje.a009243.
- [26] B. Bahadori, ND Riediger, SM Farrell, E. Uitz, and MF Moghadasian, "Hypothesis: Smoking reduces the duration of breastfeeding by suppressing prolactin secretion," *Hypothesis Medicine.* vol. 81, no. 4, p. 582–586, 2013, doi:10.1016/j.mehy.2013.07.007.
- [27] KY. Lok, MP Wang, VH. Chan, and M. Tarrant, "The Effect of Secondary Cigarette Smoke from Household Members on Breastfeeding Duration," *Sch. nurse. Univ. Hong Kong*, vol. XX, no. xx, p. 1–6, 2018, doi:10.1089/bfm.2018.0024.
- [28] N. Maulina and H. Sawitri, "VILLAGE HEALTH EMPOWERMENT WITH THE CONSTRUCTION OF CIGARETTE FRIENDLY HOUSE IN UTEUNKOT VILLAGE, MUARA DUA DISTRICT, LHOKSEUMAWE CITY," vol. 6, no. 2, p. 152–157, 2022.
- [29] RD Alnur and M. Veruswati, "Information and Community Commitment Initiation in Creating a Smoke Free Area as an Effort to Protect Passive Smokers," *aksologiya, J. Pembdi. to Mass.*, vol. 6, no. 1, p. 8–17, 2022.
- [30] I. Safitri, "FACTORS AFFECTING THE EQUITY OF BREAST MILK PRODUCTION IN BREASTFEEDING MOTHERS IN BENDAN VILLAGE, BANYUDONO DISTRICT, BOYOLALI DISTRICT," 2016.
- [31] D. Susanna, B. Hartono, and H. Fauzan, "Determination of nicotine levels in cigarette smoke," *So Health.*, vol. 7, no. 2, p. 2–5, 2003.
- [32] Weiser TM., Lin M., Garikapaty V., Feyerharm RW., Bensyl DM., Zhu BP. 2009. Association of Maternal Smoking Status With Breastfeeding Practices: Missouri. *Pediatrics*, 124(6):1603–1610.

- [33] Profile of the Banyuasin District Health Office. Profile of the Musi Banyuasin District Health Office in 2022. MusiBanyuasin: Health Office.
- [34] P2PTM Ministry of Health of the Republic of Indonesia. (2018). Dangers of Cigarette Smoke. Directorate of Prevention and Control of Non-Communicable Diseases, Ministry of Health of the Republic of Indonesia. Accessed on 16 August 2022 from <http://www.p2ptm.kemkes.go.id/>
- [35] Indonesian Health Ministry. 2018. Indonesia Health Profile 2017. Jakarta: Indonesian Ministry of Health. Accessed August 20, 2022 from <http://www.depkes.go.id/resources/download/pusdatin/profil-kesehatanindonesia/Profil-Kesehatan-Indonesia-tahun-2017.pdf>
- [36] Notoatmodjo, S. 2007. The Science and Art of Public Health. Jakarta: Rineka Cipta.

# Analysis of Factors Affecting Exposure...

## ORIGINALITY REPORT

16%

SIMILARITY INDEX

11%

INTERNET SOURCES

16%

PUBLICATIONS

3%

STUDENT PAPERS

## PRIMARY SOURCES

1	<a href="http://icoehp.poltekkesdepkes-sby.ac.id">icoehp.poltekkesdepkes-sby.ac.id</a> Internet Source	2%
2	"1st Annual Conference of Midwifery", Walter de Gruyter GmbH, 2020 Publication	2%
3	<a href="http://journal.unnes.ac.id">journal.unnes.ac.id</a> Internet Source	2%
4	Indra Setia W, Nur Isnaini. "The correlation between self-efficacy and motivation on elderly's daily independence", Proceedings Series on Health & Medical Sciences, 2020 Publication	1%
5	<a href="http://repository.stikim.ac.id">repository.stikim.ac.id</a> Internet Source	1%
6	<a href="http://repository.ung.ac.id">repository.ung.ac.id</a> Internet Source	1%
7	Marta Napierala, Jan Mazela, T. Allen Merritt, Ewa Florek. "Tobacco smoking and breastfeeding: Effect on the lactation process, breast milk composition and infant	1%

development. A critical review",  
Environmental Research, 2016

Publication

---

8

Paweł Gać, Przemysław Jaźwiec, Małgorzata Poręba, Grzegorz Mazur et al. "The risk of coronary artery disease estimated non-invasively in patients with essential hypertension environmentally exposed to cigarette smoke", Environmental Toxicology and Pharmacology, 2017

Publication

---

1 %

9

Kris Y.W. Lok, Man Ping Wang, Vincci H. S. Chan, Marie Tarrant. "Effect of Secondary Cigarette Smoke from Household Members on Breastfeeding Duration: A Prospective Cohort Study", Breastfeeding Medicine, 2018

Publication

---

1 %

10

Submitted to Eden Prairie High School

Student Paper

---

1 %

11

eudl.eu

Internet Source

---

1 %

12

Sri Yuniarti. "Relation between Knowledge and Mother's Occupation with Growth and Development of Children between Ages 5-6 in Tk At-Taqwa Cimahi", KnE Life Sciences, 2021

Publication

---

1 %

13

Dyah Dwi Astuti, Tri Widyastuti Handayani,  
Duwi Pudji Astuti. "CIGARETTE SMOKE  
EXPOSURE AND INCREASED RISKS OF  
STUNTING AMONG UNDER-FIVE CHILDREN",  
Clinical Epidemiology and Global Health, 2020

Publication

1 %

14

nusantarahasanajournal.com

Internet Source

1 %

Exclude quotes On

Exclude matches < 1%

Exclude bibliography On