

# The Effect of Consumption Nata De Cocolawak on Lipid Serum Levels on Healthy Women

*by* Miksusanti Salbi

---

**Submission date:** 20-May-2023 07:09AM (UTC+0700)

**Submission ID:** 2097485728

**File name:** ion\_Nata\_De\_Cocolawak\_on\_Lipid\_Serum\_Levels\_on\_Healthy\_Women.pdf (162.3K)

**Word count:** 4631

**Character count:** 25956

# The Effect of Consumption Nata De Cocolawak on Lipid Serum Levels on Healthy Women

Indah Solihah<sup>1</sup>, Rennie Puspa Novita<sup>1</sup>, Ina Suci Pratiwi<sup>1</sup>, Miksusanti<sup>2\*</sup>

<sup>1</sup>Department of Pharmacy, Faculty of Mathematic and Natural Sciences, Sriwijaya University, Ogan Ilir 30662, Indonesia

<sup>2</sup>Department of Chemistry, Faculty of Mathematic and Natural Sciences, Sriwijaya University, Ogan Ilir 30662, Indonesia

\*Corresponding author: [miksusanti@gmail.com](mailto:miksusanti@gmail.com)

Received September 18, 2019; Revised October 21, 2019; Accepted November 05, 2019

**Abstract** Temulawak (*Curcuma xanthorrhiza* Roxb.), traditionally known as a medicinal plant which is effective in reducing blood lipid levels. In this study we made a nata de coco product fortified with temulawak juice called nata de cocolawak. The purpose of this study was to determine the effect of supplementation of nata de cocolawak on total serum cholesterol, LDL-cholesterol, HDL-cholesterol, triglyceride-cholesterol, and serum pancreatic lipase levels. Respondents in this study were healthy adult women who were divided into 2 groups, there are group 1 consuming nata de coco products and group 2 consuming nata de cocolawak products. Each product was given as much as 100 grams consumed 3 times a day for 30 days. Nata de cocolawak products contain ascorbic acid, catechins, rutin, quercetin, alkaloids, and curcumin. Supplementation of nata de cocolawak products can significantly reduce serum total cholesterol, LDL-cholesterol, and serum pancreatic lipase levels ( $p < 0.05$ ) and significantly increase HDL-cholesterol levels ( $p < 0.05$ ) compared to baseline data. While it is not significantly ( $p > 0.05$ ) increase triglyceride-cholesterol levels. In this study, we can prove that regular consumption of nata de cocolawak products can affect blood lipid profile.

**Keywords:** Nata de coco, *Curcuma xanthorrhiza* Roxb., Lipid

**Cite This Article:** Indah Solihah, Rennie Puspa Novita, Ina Suci Pratiwi, and Miksusanti, "The Effect of Consumption Nata De Cocolawak on Lipid Serum Levels on Healthy Women." *Journal of Food and Nutrition Research*, vol. 7, no. 11 (2019): 766-771. doi: 10.12691/jfnr-7-11-2.

## 1. Introduction

Cardiovascular disease due to atherosclerosis and thrombosis is the leading cause of death in the world. The main clinical entities of the disease are chronic heart disease, ischemic stroke, and peripheral arterial disease. The cause of the disease is multifactorial, where some of it can be modified. One risk factor that can be modified is dyslipidemia [1,2,3]. Dyslipidemia is a condition of increased levels of lipids in the serum which include increased triglyceride levels, total serum cholesterol levels, LDL, and decreased HDL.

Management of dyslipidemia can be done through pharmacological and non-pharmacological therapy. Changes in a healthy lifestyle are one of types of non-pharmacological therapy. Non-pharmacological efforts to improve dyslipidemia conditions include reducing saturated fat intake, increasing fiber intake, reducing carbohydrate and alcohol intake, increasing daily physical activity, reducing excess weight, and stopping smoking.

Diets of water-soluble fiber such as peas, vegetables, fruits, and cereals have a hypolipidemic effect [4]. Diets of 5-10 grams of fiber per day can reduce LDL cholesterol by 5% [5,6]. The recommended diet for water-soluble fiber to reduce LDL cholesterol is 5-15 grams/day [7].

Nata de coco is one of the foods that are rich in fiber with maximum crude fiber of 450mg/g product [8]. Fortification of temulawak (*Curcuma xanthorrhiza* Roxb.) rhizome extract on nata de coco products can synergize the effect of nata de coco in reducing blood lipid profile. Temulawak (*Curcuma xanthorrhiza* Roxb.) contains curcumin compounds which are known to have hypolipidemic effects. The hypolipidemic effect of curcumin is thought to be influenced by its antioxidant activity [9].

Based on the results of previous studies, fiber content in the nata de cocolawak product was measured at 211mg/g and contained total curcumin of 129.355 ± 0.032mg/100g products [10]. One of the activities to decrease blood lipid was influenced by the activity of antioxidant compounds. Some sources of natural antioxidants are ascorbic acid, phenolic compounds, and flavonoid compounds. Based on its feasibility as a food, nata de cocolawak products also meet Indonesian National Standards for nata de coco products [10]. This study aims to look at the effect of consumption of nata de cocolawak on decreasing blood lipid profiles.

## 2. Experimental Section

The materials used in this study, nata de coco and nata de cocolawak, was made by home industry in Palembang

city, South Sumatera, Indonesia. All chemicals used in this study was pro analytical standard.

## 2.1. Determination of Phytochemical Content

Nata de cocolawak and nata de coco was subjected to phytochemical estimation of parameters such as ascorbic acid [11], catechins content [12], rutin content [13], quercetin [14], and total alkaloids [15].

### 2.1.1. Determination of Ascorbic Acid Content

The quantitative ascorbic acid level of the samples was measured using HPLC-PDA at quality testing service laboratory, Indonesian University. Mobile phase composition was tested with methanol/phosphate buffer with tetrabutylammonium (95:5 and 70:30, v/v) in C8 column ( $\lambda=254\text{nm}$ ); and 0.2% metaphosphoric acid in water solution, 0.2% metaphosphoric acid/methanol (95:5 and 90:10, v/v), 0.2% metaphosphoric acid/acetonitrile (95:5 and 90:10, v/v) and 0.2% metaphosphoric acid/methanol/acetonitrile (90:5:5, v/v/v) in C8 column ( $\lambda=254\text{nm}$ ).

The column used was Superspher RP-18 (250x4.6mm) while the mobile phase (pH 2.6) consisted of 1,5g ascorbic acid dissolved in 500 mL of acetic acid (99.8%) and mixed well. Routine degassing of the mobile phase was carried out by passing it through a 0,45 $\mu\text{m}$  membrane filter. The mobile phase was pumped isocratically at a flow rate of 0.7mL/min at 20°C. The injection volume was 50 $\mu\text{L}$ .

Approximately 250 mg of samples were weighed precisely and dissolved separately in 50mL mobile phase. The mixtures were centrifuged at 3000 rpm for 5 min at room temperature (20°C). The supernatants were collected and aliquots of the samples were diluted to 0.2mg/mL.

### 2.1.2. Determination of Catechin Content

The samples were extracted using ethyl acetate. The extract was read spectrophotometrically at maximum absorbance 279nm. The standard calibration curve (10-50  $\mu\text{L/mL}$ ) was plotted using catechins standard.

### 2.1.3. Determination of Rutin Content

The quantitative rutin level of the samples were measured using  $\text{AlCl}_3$  colorimetric method. The samples were extracted using 70% ethanol. Briefly, 200  $\mu\text{L}$  of sample, 1,5mL of 70% of ethanol, 100  $\mu\text{L}$  of 10%  $\text{AlCl}_3$  and 100  $\mu\text{L}$  of 1M  $\text{CH}_3\text{COONa}$  were homogenized with ethanol until volume reaches 5mL in a volumetric flask. The solution was incubated for 30 minutes. Each solution was read spectrophotometrically at maximum absorbance 420nm. The standard calibration curve of rutin (5, 10, 15, 20, and 25  $\mu\text{L/mL}$ ) was plotted.

### 2.1.4. Determination of Quercetin Content

The quercetin level of the samples were evaluated using  $\text{AlCl}_3$  colorimetric method. The samples were extracted using 96% ethanol. Briefly, 200 $\mu\text{L}$  of sample, 1,5mL of 95% of ethanol, 100 $\mu\text{L}$  of 10%  $\text{AlCl}_3$  and 100 $\mu\text{L}$  of 1M  $\text{CH}_3\text{COONa}$  were homogenized with ethanol until volume reaches 5mL in a volumetric flask. The solution was incubated for half an hour. Each solution was

read spectrophotometrically at maximum absorbance 434nm. The standard calibration curve of quercetin (10-50  $\mu\text{L/mL}$ ) was plotted.

### 2.1.5. Determination of Total Alkaloids Content

Preparation of solutions. Bromocresol green (BCG) solution was prepared by heating 69.8mg BCG with 3mL of 2N NaOH and 5 mL distilled water until completely dissolved and the solution was diluted to 1000 mL with distilled water. Phosphate buffer solution (pH 4.7) was prepared by adjusting the pH of 2M sodium phosphate (71.6 g  $\text{Na}_2\text{HPO}_4$  in 1 L distilled water) to 4.7 with 0.2M citric acid (42.02g citric acid in 1 L distilled water). Atropine standard solution was made by dissolving 1 mg pure atropine in 10 mL distilled water.

The samples were extracted with methanol then heated until dry. A part of this residue was dissolved in 2 N HCl and then filtered. One mL of this solution was transferred to a separatory funnel and wash with 10 mL chloroform (3 times). The pH of this solution was adjusted to neutral with 0,1 N NaOH. Then 5 mL of BCG solution and 5 mL of phosphate buffer were added to this solution. The mixture was shaken and the complex formed was extracted with 4 mL chloroform by vigorous shaking. The chloroform phase was collected and diluted with chloroform until 10 mL in a volumetric flask. The solutions were read at maximum absorbance 416 nm spectrophotometrically. The standard calibration curve (10, 15, 20, 25, and 30 %v/v) was plotted using atropine standard.

## 2.2. Experimental Study

Design of this experimental study was randomized controlled trial. Respondents were randomly selected with 36 healthy women, divided into placebo and nata de cocolawak groups. Respondents were woman, adults (17-40 years), and healthy. Respondents consumed 100g of nata de coco for placebo and 100g of nata de cocolawak for treatment groups three times a day for 30 days. All groups respondents were given same nutritional intake during the experiment. This clinical experimental study was registered in Health Research Review Committee Mohammad Hoesin Central General Hospital and Faculty of Medicine Sriwijaya University with number 062/kepkrsmhfkunsri/2019. Examination of blood samples was carried out before and after the treatment. Blood sampling was carried out through a 3 mL median cubital vein. Before taking blood samples, all equipment is cleaned first with 70% alcohol. The blood taken was put in a clean and dry venoject<sup>®</sup> tube, then centrifuged at 4000 rpm for 20 minutes. The separated serum was sent immediately for biochemical analysis.

## 2.3. Lipid and Serum Pancreatic Level Examination

Total serum cholesterol, triglyceride-cholesterol, HDL-cholesterol, LDL-cholesterol, and serum pancreatic lipase levels were measured by commercially available kits in spectrophotometer at Sriwijaya University Clinic.

## 2.4. Sensory Evaluation

The sensory evaluation of nata de cocolawak and nata de coco was performed following the method described by reference [16] with slight modification. The sensory analysis of nata de cocolawak and nata de coco performed using 36 respondents consisted of the women adult population. The panelists were asked to evaluate each sample for taste, aroma, texture, color, and overall acceptability. A 5-point hedonic scale was used where 1-dislike very much, 2-dislike, 3-neither like nor dislike, 4-like, and 5-like very much. The panelists were instructed to rate the attributes indicating their degree of liking or disliking by putting a number as provided in the hedonic scale according to their preference [17].

## 2.5. Statistical Analysis

All data were shown as mean  $\pm$  SEM for at least three replications for each prepared sample. The IBM SPSS® ver.25 for windows was used to determine the significant differences of comparisons data. Results with p-value less than 0.05 were considered as statistically significant.

## 3. Result and Discussion

Prevention of hyperlipidemia condition can be through a healthy lifestyle, such as by consume of functional foods. Functional foods are foods that are specifically designed and utilize certain bioactive compounds that have a role in preventing certain diseases. Temulawak has been studied as having anti hyperlipidemia activity [9,18]. Temulawak product development in the form of dessert nata de cocolawak is intended to improve acceptability.

Several bioactive compounds have been investigated to have antihyperlipidemic effects. The majority of these bioactive compounds have antioxidant effects. Some compounds that have antioxidant activity are ascorbic acid, flavonoids, alkaloids, and phenolic derivatives such as

catechins and curcumin. Table 1 shows the levels of several phytochemical compounds in nata de coco and nata de cocolawak products.

Table 1 shows that the fortification of temulawak extract in nata de coco products has higher levels of phytochemical compounds. Ascorbic acid, as an antioxidant, can influence lipid profile. In cholesterol metabolism, ascorbic acid plays a role in increasing the rate of cholesterol removed in the form of bile acids and also increasing HDL-cholesterol levels [19]. Catechin reported reducing circulating total cholesterol and LDL-cholesterol concentration on serum lipid postmenopausal women [20]. Rutin was identified to be antioxidant compound that reduces serum total cholesterol and LDL-cholesterol on *in vitro* study [21,22]. Quercetin reported can decrease in cholesterol, triglycerides and LDL value with parallel increase in HDL on clinical study [23]. *In vitro* studies investigated the anti-hyperlipidemia effect of the alkaloid rich extract some medicinal plants [24,25]. Curcumin reported had properties to lowered oxidative damage by neutralizing free radical species when they attack the lipid membrane [26]. This antioxidative activity of curcumin provides protection against lifestyle-related disorders, such as hyperlipidemia [27].

Table 2 shows the effect of nata de cocolawak product on serum lipid profiles of respondents. The lipid serum in nata de cocolawak supplementation decreased significantly ( $p < 0.05$ ) both total serum cholesterol and LDL-cholesterol levels when compared with baseline levels. Nata de cocolawak supplementation had no significant ( $p > 0.05$ ) increase of triglyceride serum level. Supplementation of nata de cocolawak product also increased significantly ( $p < 0.05$ ) HDL-cholesterol levels when compared with baseline levels. The respondents who were received nata de coco supplementation have significantly ( $p < 0.05$ ) lower triglyceride level, and higher HDL-cholesterol level compares with baseline. Though, they have no significant ( $p > 0.05$ ) effect on other lipid serum parameters.

Table 1. Phytochemical contents in Nata de coco and Nata de cocolawak product

Phytochemical contents	Nata de coco (Mean $\pm$ SEM)	Nata de cocolawak (Mean $\pm$ SEM)
Ascorbic acid (mg/100g)	2.575 $\pm$ 0.159	2.625 $\pm$ 0.055
Catechin (mg/g)	419.487 $\pm$ 3.589	536.923 $\pm$ 3.203
Rutin (mg/g)	38.969 $\pm$ 0.061	44.242 $\pm$ 0.061
Quercetin (mg/g)	17.494 $\pm$ 0.115	27.494 $\pm$ 0.115
Total Alkaloid (mg/g)	41.296 $\pm$ 0.185	47.963 $\pm$ 0.185
Curcumin (mg/100g)	0	129.355 $\pm$ 0.032*

Source: \*Reference [10].

Table 2. Effect of nata de coco and nata de cocolawak product on the Serum Lipid Levels

Parameters (mg/dL)	Groups					
	Nata de coco			Nata de cocolawak		
	Baseline (Mean $\pm$ SEM)	End (Mean $\pm$ SEM)	p-value	Baseline (Mean $\pm$ SEM)	End (Mean $\pm$ SEM)	p-value
Total serum Chol	140.333 $\pm$ 6.495	131.778 $\pm$ 4.393	0.248	149.222 $\pm$ 4.144	128.444 $\pm$ 3.744	0.001*
Triglyceride-Chol	122.788 $\pm$ 5.273	107.667 $\pm$ 2.762	0.023*	112.278 $\pm$ 3.622	116.611 $\pm$ 3.333	0.318
LDL-Chol	73.056 $\pm$ 4.528	76.833 $\pm$ 3.879	0.485	82.667 $\pm$ 3.071	70.889 $\pm$ 3.169	0.018*
HDL-Chol	33.500 $\pm$ 0.764	42.778 $\pm$ 2.015	0.001*	33.611 $\pm$ 0.647	44.389 $\pm$ 1.319	0.001*

\*The significant difference ( $p < 0.05$ ) is examined using *t*-test.

**Table 3. Comparison Lipid Serum Levels after treatment**

Parameters	Nata de coco (Mean±SEM)	Nata de cocolawak (Mean±SEM)	p-value
Total serum-Chol	131.778±4.393	128.444±3.744	0.454
Triglyceride-Chol	107.667±2.762	116.611±3.333	0.098
LDL-Chol	76.833±3.879	70.889±3.169	0.186
HDL-Chol	42.778±2.015	44.389±1.319	0.873

Statistical analysis of lipid serum levels after treatment in both nata de coco and nata de cocolawak groups are exhibit in Table 3. Independent t-test was use to examine the mean of lipid serum levels. The p-value of lipid serum levels shows no significant ( $p>0.05$ ) difference between groups.

The phytochemical compound that is contained in the nata de coco dan nata de cocolawak product affect the serum lipid profile. Besides that, the raw material, coconut water in products also has activity to affect serum lipid profile. Nata de coco and nata de cocolawak was made from coconut water. Coconut water has been reported rich in vitamins, amino acids, enzymes and minerals. Coconut water contains ascorbic acid 2.2-3.7mg/100mL has been reported scavenges DPPH, ABTS, and superoxide radicals [28]. Ascorbic acid as antioxidant compound in coconut water can reduces lipid peroxidation [29] and inhibit the formation of hydroxyl radical species [30].


Fortification temulawak (*Curcuma xanthorrhiza* Roxb.) extract in nata de coco intended to increase bioactivity of the product. Traditionally, temulawak (*Curcuma xanthorrhiza* Roxb.) had been used as blood purifier [31]. Many studies had been prove that temulawak (*Curcuma xanthorrhiza* Roxb.) have antihyper-cholesterolemic effect in vivo or in vitro studies [32,33,34,35,36].

Curcumin as active compound was reported to modulate 3-hydroxy-3-methylglutaryl-coenzyme A (HMG-CoA) reductase activity [35]. HMG-CoA reductase is enzyme that lead to decrease in the level of serum cholesterol, triglycerid, and free fatty acid. Curcumin also reported stimulate hepatic cholesterol-7 $\alpha$ -hydroxylase enzyme, an enzyme that regulates cholesterol catabolism [36].

In previous study, nata de cocolawak contains 58.25mg/Kg of calcium and 25.73 mg/Kg of zinc [10]. The reference [37] reported that a low calcium intake was related to increase total serum cholesterol and LDL-cholesterol concentration significantly. Zinc supplementation also reduced total cholesterol, LDL-cholesterol, and triglycerides significantly [38].

Pancreatic lipase enzyme is a key enzyme related to dietary fats absorption [11]. Among various lipase, pancreatic lipase performs the hydrolysis of 50-70% of total dietary fats to fatty acids and monoglycerides [40]. The reduction of fat absorption through pancreatic lipase inhibition is known to benefit the regulation of dyslipidemia [41].

In our current study, the obtained results in Figure 1 show that nata de cocolawak supplementation reduced the level of serum pancreatic lipase significantly ( $p<0.05$ ) compared to its baseline. Whereas, nata de coco supplementation not significantly ( $p>0.05$ ) reduced the level of pancreatic lipase compared to its baseline. Many in vitro studies reported that curcumin [42] and *Curcuma xanthorrhiza* Roxb. [43], tannins, flavonoids, and alkaloids are active inhibitors of pancreatic lipase [44,45,46].



**Figure 1.** Serum pancreatic lipase levels (Mean±SEM) (\*The significant difference between baseline ( $p$ -value = 0.040) is examined using *t*-test.)

**Table 4.** Sensory Score of nata de coco and nata de cocolawak product

Parameters	Nata de coco (Mean±SEM)	Nata de cocolawak (Mean±SEM)	p-value
Color	4.22±0.101	4.38±0.118	0.044*
Texture	3.88±0.196	4.16±0.167	0.838
Aroma	3.83±0.159	4.11±0.145	0.827
Taste	4.22±0.129	4.11±0.111	0.240
Overall acceptability	4.04±0.132	4.19±0.118	0.945

\*The significant difference ( $p<0.05$ ) is examined using *t*-test.

Fortification temulawak juice on nata de coco product may affect the organoleptic of product. The sensory scores of nata de coco and nata de cocolawak products are shown in Table 4. The result shows that there was significantly ( $p<0.05$ ) difference color in nata de coco and nata de cocolawak products. Nata de coco products were white, while nata de cocolawak products were yellow. This yellow color comes from the content of curcumin in nata de cocolawak products. The sensory score of texture, aroma, taste, and overall acceptability are no significantly difference ( $p>0.05$ ). This sensory score results might indicate that fortification temulawak extract on nata de coco product might not play any sensory role.

## 4. Conclusions

Nata de cocolawak product contains ascorbic acid, catechin, rutin, quercetin, alkaloids, and curcumin. Supplementation of nata de cocolawak to healthy women significantly reduced total serum-cholesterol, LDL-cholesterol, and serum pancreatic lipase levels compared to its baseline. It is also significantly increased HDL-cholesterol levels compared to its baseline. Based on these results, nata de cocolawak supplementation potential to be used as adjunct therapy for dyslipidemia condition.

15

## Conflict of Interest

The authors declare no conflict of interest.

## Acknowledgements

The authors would like to express their gratitude toward Sriwijaya University Clinic for facilitating clinical biochemical analysis during this study.

## References

- [1] Hokanson J.E., Austin M.A., "Plasma triglyceride level is a risk factor for cardiovascular disease independent of high-density lipoprotein cholesterol level: a metaanalysis of population-based prospective studies", *J Cardiovasc Risk*, 3, 213-219. 1996.
- [2] Karthikeyan G., Teo K.K., Islam S., McQueen M.J., Pais P., Wang X., Sato H., Lang C.C., Sithi-Amorn C., Pandey M.R., Kazmi K., Sanderson J.E., Yusuf S., "Lipid Profile, Plasma Apolipoproteins, and Risk of a First Myocardial Infarction Among Asians: An Analysis From the INTERHEART Study", *J Am Coll Cardiol*, 53, 244-253. 2009.
- [3] Nugroho, A.E., Malik, A., Pramono, S., "Total Phenolic and Flavonoid Contents, and In Vitro Antihypertension Activity of Purified Extract of Indonesian Cashew Leaves (*Anacardium occidentale* L.)", *International Food Research Journal*, 20(1), 299-305. 2013.
- [4] Brown L., Rosner B., Willet W., Sacks S.M., "Cholesterol-lowering effects of dietary fiber: a meta-analysis", *Am J Clin Nutr*, 69, 30-42. 1999.
- [5] U.S. Department of Health and Human Services. Food and Drug Administration, *Food labeling: health claims; soluble fiber from certain foods and coronary heart disease: proposed rule*, Federal Register, 1997, 62.28234-45.
- [6] U.S. Department of Health and Human Services. Food and Drug Administration, *Food labeling: health claims; soluble fiber from certain foods and coronary heart disease: final rule*, Federal Register, 1998, 63.8103-8121.
- [7] Rideout T.C., Harding S.V., Jones P.J., Fan M.Z., "Guar gum and similar soluble fibers in the regulation of cholesterol metabolism: current understandings and future research priorities", *Vasc Health Risk Manag*, 4.1023-1033. 2008.
- [8] National Standardization Department, *Indonesian National Standard for nata in packaging*. SNI 01-4317-1996.
- [9] Zingg, J.M., Hasan, S.T., Meydani, M., "Review Article: Molecular Mechanism of Hypolipidemic Effect of Curcumin", *BioFactors*, 39(1), 101-121. 2013.
- [10] Solihah, I., Miksusanti, Novita, R.P., Ramadhan, S.Y., "Determination of quality parameters in nata de cocolawak as hepatoprotector functional food", *J Phys: Conf Ser*, 1282, 012067. 2019.
- [11] Mitic S.S., Kostic D.A., Naskovic-Dokic D.C., Mitic, M.N., "Rapid and Reliable HPLC method for the determination of Vitamin C in Pharmaceutical samples", *Trop. J. Pharmaceutical Res.*, 10(1), 105-111. 2011.
- [12] Ibrahim, Y.A., Musa, A., Yakasai, I.A., "Spectrophotometric method for determination of catechins in green tea and herbal formulations", *Nigerian Journal of Pharmaceutical Sciences*, 6 (1), 25-30. 2017.
- [13] Xu, H., Li, Y., Tang, H.W., Liu, C.M., Wu, Q.S., "Determination of rutin with UV-Vis spectrophotometric and laser-induced fluorimetric detection using a non-scanning spectrometer", *Analytical Letters*, 43, 893-904. 2010.
- [14] Shiyani, S., Hertiani, T., Martini, R., Nugroho, A.K. "Optimization of a novel kinetic-assisted infundation for rich-egg and polyphenols of white tea (*Camellia sinensis*) using central composite design", *Int J of Applied Pharmaceutics*. 10(6), 259-268. 2018.
- [15] Shamsa, F., Monsef, H., Ghamooshi, R. & Verdian-rizi, M., "Spectrophotometric determinator of total alkaloid in Peganum harmala L. using bromocresol green", *Res J Phytochem*, 1(2), 79 – 82. 2007.
- [16] Larmond, E. *Laboratory methods for sensory evaluation of food* (Publication No. 1637), Ottawa: Research Branch, Canada Department of Agriculture. 19-63. 1977.
- [17] Rahim A., Kadir S., Jusman J., Zulkipli Z., Hambali T.N.A., "Physical, chemical and sensory characteristics of bread with different concentrations of acetylated arenga starches", *Int. Food Res. Journal*. 26(3), 841-848. 2019.
- [18] Wientarsih, I., Chakeredza, S., Meulen, U.t., "Influence of curcuma (*Curcuma xanthorrhiza* Roxb.) on lipid metabolism in rabbits", *J.Sci Food Agric*, 82, 1875-1880. 2002.
- [19] McRae MP. "Vitamin C supplementation lowers serum low-density lipoprotein cholesterol and triglycerides: a meta-analysis of 13 randomized controlled trials", *Journal of Chiropractic Medicine*; 7, 48-58. 2008.
- [20] Samavat, H., Newman, A.R., Wang, R., Yuan, J.M., Wu, A.H., Kurzer, M.S., "Effect of green tea catechin extract on serum lipids in postmenopausal women: a randomized, placebo-controlled clinical trial", *Am J Clin Nutr*, 104, 1671-82. 2016.
- [21] Katsube T, Imawaka N, Kawano Y, Yamazaki Y, Shiwaku K, Yamane Y., "Antioxidant flavonol glycosides in mulberry (*Morus alba* L.) leaves isolated based on LDL antioxidant activity", *Food Chem*, 97, 25-31. 2006.
- [22] Ziaee, A., Zamansoltani, F., Nassiri-Asl, M., Abbasi, E., "Effect of Rutin on Lipid Profile in Hypercholesterolaemic Rats", *Basic & Clinical Pharmacology & Toxicology*, 104, 253-258. 2008.
- [23] Talirevic, E. and Sehovic, J., "Quercetin in the treatment of dyslipidemia", *MED ARH*, 66(2), 87-88. 2012.
- [24] He, K., Kou, S., Zou, Z., Hu, Y., Feng, M., Han, B., Li, X., Ye, X., "Hypolipidemic effects of alkaloid from rhizome *Coptidis* in diet-induced hyperlipidemic Hamsters", *Planta Med*, 82, 690-697. 2015.
- [25] Zhang, X., Jin, Y., Wu, Y., Zhang, C., Jin, D., Zheng, Q., Li, Y., "Anti-hyperglycemic and anti-hyperlipidemia effect of the alkaloid-rich extract from barks of *Litsea glutinosa* in ob/ob mice", *Scientific Reports*, 8, 12646. 2018.
- [26] Chattopadhyay I., Biswas K., Bandyopadhyay U., Banerjee R.K., "Turmeric and curcumin: biological actions and medicinal applications", *Current Science*, 87, 44-53. 2004.
- [27] Manikandan P., Sumitra M., Aishwarya S., Manohar B.M., Lokanadam B., Puvanakrishnan R. "Curcumin modulates free radical quenching in myocardial ischaemia in rats", *International Journal of Biochemistry and Cell Biology*, 6, 1967-1980. 2004.
- [28] Mantena, S.K., Jagadish, Badduri S.R., Siripurapu K.B., Unnikrishnan, M.K., "In vitro evaluation of antioxidant properties of *Cocos nucifera* Linn.water", *Nahrung*, 47(2).126-131. 2003
- [29] Halliwell, B., Gutteridge, J.M.C., *Biochem. J.* 1-14. 1984
- [30] Krishnankutty, S., in: Thampan, P.K., Markose, V.T., Pankajakshan, A.S., Edachal, M., Nandakumar, T.B., (Eds), *Processing and marketing of coconuts in India*, *Coconut Development Board*, Ministry of Agriculture, Government of India, Kochi, 70-76. 1987
- [31] Elliot, S., Brimacombe, J., "The Medicinal Plants of Gunung Leuser National Park, Indonesia", *Ethnopharmacol.* 19(3), 285-317. 1987.
- [32] Mauren, FM., Yanti, Lay, BW., "Efficacy of oral curcuminoid fraction from *Curcuma xanthorrhiza* and curcuminoid cider in high-cholesterol fed rats", *Pharmacognosy Res.* 8(3), 153-159. 2016.
- [33] Bertges LC., Souza JB., Cardoso VA, "Hyperlipidemia induced by triton WR1339 (Tyloxapol) in wistar rats", *Braz J Med Sci Health.* 1, 32-34. 2011.
- [34] Ramirez-Tortosa MC., Mesa MD., Aguilera MC., Quiles JL., Baro L., Ramirez-Tortosa CL, et al, "Oral administration of a turmeric extract inhibits LDL oxidation and has hypocholesterolemic effects in rabbits with experimental atherosclerosis", 147, 371-378. 1999.
- [35] Murugan P., Pari L, "Effect of tetrahydrocurcumin on lipid peroxidation and lipids in streptozotocin-nicotinamide-induced diabetic rats", *Basic Clin Pharmacol Toxicol.* 99, 122-127. 2006.
- [36] Babu PS., Srinivasan K, "Hypolipidemic action of curcumin, the active principle of turmeric (*Curcuma longa*) in streptozotocin induced diabetic rats", *Mol Cell Biochem.* 166, 169-175. 1997.
- [37] Jacqmain M, Doucet E, Despres JP, Bouchard C, Tremblay A., "Calcium intake, body composition, and lipoprotein-lipid concentrations in adults", *Am J Clin Nutr*, 77, 1448-52. 2003.

- [38] Ranasinghe, P., Wathurapatha W.S., Ishara, M.H., Jayawardana, R., Galappaththy P., Katulanda, P., Constantine, G.R., "Effect of Zinc supplementation on serum lipids: a systematic review and -meta-analysis", *Nutrition & Metabolism*, 12(26). 1-16 2015.
- [39] Veeramachaneni, G.K., Raj, K. K., Chalasani, L. M., Annamraju, S. K., B. Js, and Talluri, V. "Shape based virtual screening and molecular docking towards designing novel pancreatic lipase inhibitors," *Bioinformation*, 11(12). 535-542. 2015.
- [40] Birari, R. B. and Bhutani, K. K. "Pancreatic lipase inhibitors from natural sources: unexplored potential," *Drug Discovery Today*, 12 (19-20). 879-889. 2007.
- [41] Ahn, J.H., Liu, Q., Lee, C. et al., "A new pancreatic lipase inhibitor from *Broussonetia kanzinoki*," *Bioorganic and Medicinal Chemistry Letters*, 22(8). 2760-2763. 2012.
- [42] Kim, T.H., Kim, J.K., Ito, H., Jo, C., "Enhancement of pancreatic lipase inhibitory activity of curcumin by radiolytic transformation", *Bioorganic and Medicinal Chemistry Letters*, 21. 1512-1514. 2011.
- [43] Ado, M.A., Abas, F., Mohammed A.S., Ghazali, M., "Anti-and pro-Lipase activity of selected medicinal, herbal, and aquatic plants, and structure elucidation of an anti-lipase compound", *Molecules*, 18. 14651-69. 2013.
- [44] Thomas, S.; Patil, D.A.; Patil, A.G.; Chandra, N. "Pharmacognostic evaluation and physicochemical analysis of *Averrhoa carambola* L. fruit", *J. Herbal Med. Toxicol*, 2. 51-54. 2008.
- [45] Liu, H., Di, Y., Yang, J., Teng, F., Lu, Y., Ni, W., Chen, C., Hao, X., "Three novel 3,4-seco-podocarpene trinorditerpenoids from *Aleurites moluccana*", *Tetrahedron Lett*, 49. 5150-5151. 2008.
- [46] Mackeen, M.M., Ali, A.M., Abdullah, M.A., Nasir, R.M., Mat, N.B., Razak, A.R., Kawazu, K., "Antinematodal activity of some Malaysian plant extracts against the pine wood nematode, *Bursaphelenchus xylophilus*", *Pestic. Sci.* 51. 165-170. 1997.



© The Author(s) 2019. This article is an open access article distributed under the terms and conditions of the Creative Commons Attribution (CC BY) license (<http://creativecommons.org/licenses/by/4.0/>).

# The Effect of Consumption Nata De Cocolawak on Lipid Serum Levels on Healthy Women

## ORIGINALITY REPORT

12%

SIMILARITY INDEX

7%

INTERNET SOURCES

6%

PUBLICATIONS

6%

STUDENT PAPERS

## PRIMARY SOURCES

1	<a href="http://bu.edu.eg">bu.edu.eg</a> Internet Source	1%
2	Submitted to Institute of Technology, Nirma University Student Paper	1%
3	Submitted to October University for Modern Sciences and Arts (MSA) Student Paper	1%
4	<a href="http://www.phcogres.com">www.phcogres.com</a> Internet Source	1%
5	RJ Mongi, BK Ndabikunze, BE Chove, P Mamiro, C. C Ruhembe, J.G Ntwenya. "Proximate composition, bread characteristics and sensory evaluation of cocoyam-wheat composite breads", African Journal of Food, Agriculture, Nutrition and Development, 2011 Publication	1%
6	Mohamed S Karawya, Mohamed G Ghourab, Faten H Elrakhawy. "Estimation of Glycoalkaloids as Solasodine in Solanum	1%



laciniatum", Journal of AOAC INTERNATIONAL,  
1975

Publication

---

7	Submitted to Jaypee University of Information Technology Student Paper	1 %
8	fieldresearch.msf.org Internet Source	1 %
9	Submitted to Sriwijaya University Student Paper	1 %
10	downloads.hindawi.com Internet Source	1 %
11	vdoc.pub Internet Source	1 %
12	Aliyu Muhammad, David Ebuka Arthur, Sanusi Babangida, Ochuko L. Erukainure et al. "Modulatory role of rutin on 2,5-hexanedione-induced chromosomal and DNA damage in rats: validation of computational predictions", Drug and Chemical Toxicology, 2018 Publication	1 %
13	www.docstoc.com Internet Source	1 %
14	Submitted to Indian School of Mines Student Paper	1 %

---

15

Internet Source

1 %

16

Titilope Ruth Manuwa, Afolabi Clement Akinmoladun, Olamide Olajusi Crown, Kayode Komolafe, Mary Tolulope Olaleye.

1 %

"Toxicological Assessment and Ameliorative Effects of Parinari curatellifolia Alkaloids on Triton-Induced Hyperlipidemia and Atherogenicity in Rats", Proceedings of the National Academy of Sciences, India Section B: Biological Sciences, 2015

Publication

Exclude quotes On

Exclude matches < 1%

Exclude bibliography On