

# Antioxidant Compound of Endophytic Fungi *Penecillium oxalicum* Isolated from *Swietenia mahagoni* L. Jacq

*by* Hary Widjajanti

---

**Submission date:** 21-Sep-2022 12:07PM (UTC+0700)

**Submission ID:** 1905173523

**File name:** PHYTIC\_FUNGI\_\_TROPICAL\_JOURNAL\_OF\_NATURAL\_PRODUCT\_MARET\_2021.pdf (519.9K)

**Word count:** 3819

**Character count:** 21147

Antioxidant Compound of Endophytic Fungi *Penicillium oxalicum* Isolated from *Swietenia mahagoni* L. JacqFadhillah Fadhillah<sup>1</sup>, Elfita Elfita<sup>2\*</sup>, Muharni Muharni<sup>2</sup>, Heni Yohandini<sup>2</sup>, Hary Widjajanti<sup>3</sup><sup>1</sup>Magister Program of Chemistry, Department of Chemistry, Faculty of Mathematics and Natural Sciences, University of Sriwijaya, Palembang, South Sumatra, 30662 Indonesia.<sup>2</sup>Department of Chemistry, Faculty of Mathematics and Natural Sciences, University of Sriwijaya, Inderalaya, Ogan Ilir, South Sumatra, 30662 Indonesia<sup>3</sup>Department of Biology, Faculty of Mathematics and Natural Sciences, University of Sriwijaya, Inderalaya, Ogan Ilir, South Sumatra, 30662 Indonesia

## ARTICLE INFO

43  
ABSTRACT

## Article history:

Received 22 January 2021

Revised 09 March 2021

Accepted 18 March 2021

Published online 01 April 2021

Copyright: © 2021 Fadhillah et al. This is an open-access article distributed under the terms of the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.

Endophytic fungi are microorganisms that are symbionts in plant tissues. Endophytic fungi isolated from medicinal plants such as *Swietenia mahagoni* may produce a plethora of bioactive compounds. In this study, endophytic fungi was isolated from the leaves of *S. mahagoni* and cultivated in potato-dextrose broth (PDB) for six weeks at static condition. The antioxidant compound was isolated from the ethyl acetate extract of endophytic fungi by chromatographic method, and the chemical structure was determined by spectroscopic analysis. The antioxidant activity was tested by the 2,2-diphenyl-1-picrylhydrazyl (DPPH) method, and the endophytic fungi was analyzed to identify its species by molecular method. Based on the spectroscopic analysis (<sup>1</sup>H-NMR, <sup>13</sup>C-NMR, HMQC, and HMBC spectra), the antioxidant compound was identified as mono-*o*-methylcurvulinic acid with IC<sub>50</sub> of 38.01 µg/mL. Phylogenetic analysis showed the endophytic fungi as *Penicillium oxalicum*.

**Keywords:** Antioxidant compound, Endophytic fungi, *Penicillium oxalicum*, *Swietenia mahagoni*

## Introduction

Lifestyle factors, stress, and environmental influences can cause excessive free radical formation and trigger oxidative stress. Oxidative stress is an important risk factor in the pathogenesis of chronic diseases. Reactive oxygen species (ROS), such as superoxide anions (O<sub>2</sub><sup>-</sup>) and hydrogen peroxide (H<sub>2</sub>O<sub>2</sub>), can cause oxidative damage to macromolecules such as DNA, proteins, lipids, and small molecules. ROS is recognized as an agent involved in the pathogenesis of diseases such as cardiovascular, asthma, arthropathy, inflammation, diabetes, cancer, anemia, ischemia, stroke, parkinson, alzheimer, atherosclerosis, neurodegenerative and premature aging.<sup>1-4</sup> Therefore, the balance between oxidation and antioxidants is vital to maintain the health of the biological system. Antioxidants are substances that inhibit oxidative damage to target molecules.<sup>3</sup> The antioxidant activity will stop the chain reaction by eliminating free radical intermediates, inhibiting other oxidation reactions.<sup>2</sup> Antioxidant compounds have the ability to trap free radicals such as peroxides, hydroperoxides, or peroxy lipids by inhibiting oxidative mechanisms that cause degenerative diseases.<sup>3</sup> Antioxidant compounds are needed to overcome cell damage. However, synthetic antioxidant compounds such as butylated hydroxytoluene (BHT) and butylated hydroxyanisole (BHA) can cause various side effects.<sup>5</sup> Another alternative used to minimize synthetic antioxidants' side effects are to take advantage of medicinal plant extracts' benefits because medicinal plants have low toxicity and good therapeutic performance.<sup>4</sup> Antioxidant compounds of medicinal

plants displays good protection against free radicals and an inhibitor of lipid peroxidation.<sup>1</sup> Medicinal plants also contain polyphenol and flavonoid compounds, useful as antioxidants.<sup>2</sup> Studies on plant activity are increasing rapidly due to interest in the potential used as a source of natural antioxidants.<sup>4</sup> Endophytic fungi are type of fungi that can live in plant tissue at certain periods of their life without causing damage to the host.<sup>5,6</sup> Bioactive compounds from endophytic fungi are a potential source of new natural antioxidants for medicinal, agricultural, and industrial purposes.<sup>7</sup> The secondary metabolites of endophytic fungi are useful for plant defence, and some have potential use in modern medicine such as antibiotic, antidiabetic, anticancer and antioxidant.<sup>8</sup> Endophytic fungi isolated from different hosts showed significant differences in biological activity.<sup>9</sup> The same isolates inhabit different hosts exhibit different and varied biological activities.<sup>10</sup> In any ecosystem, the production of secondary metabolites is based on the mutualistic relationship of endophytes and their hosts.<sup>9</sup> Antioxidant compounds such as flavonoid and phenol derivatives have been obtained from *S. mahagoni*.<sup>11-14</sup> Endophytic fungi isolated from medicinal plants are a good source of bioactive compounds because they produce important and novel bioactive compounds in a short time and do not require large areas to grow.<sup>15,16</sup>

71  
Materials and Methods

## Plant material

Fresh leaves of mahoni (*Swietenia mahagoni*) were collected during last week of December 2018 from Ogan Ilir in the province of South Sumatra, Indonesia with identification number 24/UN9.1.7/4/PP/2018.

## Isolation of endophytic fungi

The PDA medium was inserted in a petri dish using physical sterilization by heating. The medium was left to solidify. Leaves of mahoni were cleaned by running water, sterilized with 70% alcohol then rinsed with sterile distilled water. This sterilization was done three times. Then the samples were re-sterilized by 1% NaOCl (1

\*Corresponding author. E mail: [elfita.elfita.69@gmail.com](mailto:elfita.elfita.69@gmail.com)

Tel: +62 852-7974-5106

Citation: Fadhillah F, Elfita E, Muharni M, Yohandini H, Widjajanti H. Antioxidant Compound of Endophytic Fungi *Penicillium oxalicum* Isolated from *Swietenia Mahagoni* L. Jacq. Trop J Nat Prod Res. 2020; 5(3):543-548. doi.org/10.26538/tjnpr/v5i3.20

Official Journal of Natural Product Research Group, Faculty of Pharmacy, University of Benin, Benin City, Nigeria.

min). Then, samples were cleaned again by sterile aquades. Leaf samples were cut to 0.5 cm in width and 2 cm in length. Then the leaf pieces were plated into a petri dish containing a solid field PDA medium containing chloramphenicol (150 mg/L).<sup>15</sup> The sample was incubated for 1–2 weeks at room temperature. Colonies were grown in a medium and observed after 3 to 14 days. Colonies that show differences characteristic was considered as different isolates, then separated and re-cultured in the PDA medium until a similar culture were obtained.<sup>17</sup>

#### Cultivation and extraction

Each fungal strain was cultivated into potato dextrose broth (PDB) medium (20 g dextrose monohydrates, 200 g potatoes, and 1,000 mL aquades) under static conditions at room temperature. The mycelia agar plugs of all strain (5 x 5 mm<sup>2</sup>, 6 pieces) were cultivated into PDB medium (200 mL) for six weeks. After cultivation, The broth culture was extracted by partition technique with ethyl acetate (1 : 1) and concentrated under vacuum at 45°C.<sup>17,18</sup>

#### Antioxidant activity test

The extract was weighed 0.002 g and dissolved with 2 mL of methanol. Then the mixture was diluted to 1000, 500, 250, 125, 62.5, 31.2 ppm. Each concentration was added by 3.8 mL DPPH 0.05 mM and left in a dark room for ± 30 minutes. Each sample was determined for its absorbance value using a UV spectrometer at a wavelength of 517 nm. Each sample was determined by % inhibition, using the following formula.<sup>19</sup>

$$\% \text{ Inhibition} = \frac{\text{Standard Absorbance} - \text{Absorbance of sample}}{\text{Standard Absorbance}} \times 100\%$$

#### Identification of endophytic fungus

Endophytic fungus was identified using the internal transcribed spacer (ITS) method conducted at LIPI Cibinong biological research center. Purified DNA extracts were analyzed by PCR method using ITS-4 and ITS-5 primers. Furthermore, nucleotide sequences were compared for homology with BLAST search (<http://www.ncbi.nlm.nih.gov>). That file was converted to txt, MSF, and PHYLIP files using the ClustalW program. The run file was renamed and entered into the SEQBOOT program then calculated by replication 1000 times. The resulting file was run with the DNAdist and Neighbor program with a lot of data processing 1000 times. The file was entered into the CONSENSE program. The phylogenetic tree was visualized by the Treeview and Mega4 program.<sup>20</sup>

#### Isolation and identification of pure compound

Secondary metabolite of the active extract was separated using column chromatography with silica gel as the stationary phase and a gradient solvent as the mobile phase. Eluate was collected in vials and grouped into sub-fraction by thin layer chromatographic (TLC) analysis. The secondary metabolite of sub-fraction was separated and purified by column chromatography and recrystallization techniques to yield pure compound. The chemical structures were determined by NMR spectra, including NMR 1 and 2 Dimension (Agilent DD2 <sup>1</sup>H-500 MHz and <sup>13</sup>C-125 MHz).

## Results and Discussion

A total of three fungal isolates (labeled: DMP1, DMQ2, and DMR3) were isolated from leaves *S. mahagoni* collected from Ogan Ilir in the province of South Sumatra, Indonesia in December 2018. Each fungi was harvested and the medium extracted with ethyl acetate until concentrated. Each extract was tested antioxidant activity with a variety of sample concentrations. Antioxidant activity tests was carried out using the DPPH (2,2-diphenyl-1-picrylhydrazyl) method. The method was used because a sample's antioxidant reactivity can be easily detected when the DPPH solution changes colour (from violet to yellow) upon addition of an antioxidant compound.<sup>21-24</sup> DPPH is an active free radical that has very reactive free electrons. The presence of antioxidants can reduce the reactivity of free radicals by donating electrons. The 2,2-diphenyl-1-picrylhydrazine (violet) when reacted

with antioxidant compounds produces 2,2-diphenyl-1-picrylhydrazine (yellow) which is non-reactive. The IC<sub>50</sub> concentration was calculated based on the results of the percentage inhibition of each sample concentration. Based on these data, we find that the DMP1 extract had a higher activity compared other extracts (Table 1).

The DNA molecule of DMP1 was identified using the PCR method. The obtained sequences were subjected to BLAST program at the National Center for Biotechnology Information (NCBI). The ITS sequences of isolate DMP1 showed homology with 20 strain of fungus as *penicillium family*. The following is DNA of DMP1 counting:

```
TATGCTTAAGTTCAGCGGGTATCCCTACCTGATCCGAGGTCACCTGGTTAAGATTGATGGTGTTCGCGCGCGGGCGCCGGCCGGCCTACAGAGCGGGTGACGAAGCCCCATACGCTCGAGGACCGGACGCGGTGCCCGCGCTGCCTTTCGGGCCCGCCCGCCGGAGCGGGGGCGAGAGCCCAACACACAAGCCGTGTTGAGGGCAGCAATGACGCTCGGACAGGCATGCCCCCGGAATACCAGGGGGCGCAATGTGCGTTCAAAGACTCGATGATTACATGAATTCTGCAATTCACATTACTATCGCATTTTCGCTGCGTTCTTCATCGATCCCGGAACCAAGAGATCCGTTGTGAAAGTTTAACTGATTTAGTCAAGTACTCAGACTGCAATCTTCAGACAAGTTCGTTGTGTGCTTCGCGCGCGGTGAGGGCGGGCCCGCAAGCAACAAGGTACGATAAACACGGGTGGG
```

The closest relationship of each sequence was further analyzed using the phylogenetic method by using the neighbor-joining method with 1000 matches. It was found that the DMP1 fungus had DNA similarity with the fungus *Penicillium oxalicum* isolate HC6 (MH58553)<sup>25</sup> with 100% similarity (figure 1).

*Penicillium oxalicum* is one of the fungi whose extracts are widely used as an anti-fungal, anti-microbial and antioxidant.<sup>26,27,28</sup> The ethyl acetate extract of the fungus *Penicillium oxalicum* inhibited the peroxidation of linoleic acid by 78%.<sup>29</sup> An antioxidant compound of *Penicillium oxalicum* chloroform extract had an IC<sub>50</sub> value of 17.47 µg/ml for antioxidant test with the DPPH method.<sup>30</sup>

The crude extract DMP1 (3 g) was separated using column chromatography (60 g silica gel 70-230 mesh) and eluted with a gradient solvent system, n-Hexane-ethyl acetate (10: 0 → 0: 10) and ethyl acetate-methanol (10: 0 → 8: 2) to give six fractions (F1-F6). Fraction 3 (F3, 402 mg) was further separated by column chromatography and eluted with n-Hexane-ethyl acetate (7:3 → 0:10) to give four subfractions. Fraction F3.3 (96 mg) was subjected to column chromatography and eluted with n-Hexane-ethyl acetate (4:6) to yield compound 1 (42 mg). The chemical structure of compound 1 was identified by NMR spectra. The <sup>1</sup>H-NMR spectra (figure 2) identified the presence of five proton signals at δ<sub>H</sub> 2.5 - 6.5 ppm, which is typical for aromatic oxymethyl, methyl, and methylene protons. The spectrum showed the presence of two signals in the aromatic region at δ<sub>H</sub> 6.36 (1H, d, J = 2.5 Hz) and 6.73 (1H, d, J = 2.5 Hz) ppm as doublet meta-coupled protons. The proton signal at δ<sub>H</sub> 2.53 (3H, s) and 3.78 (3H, s) indicated the presence of one methyl and one methoxyl group. Other signals appeared at δ<sub>H</sub> 3.70 (2H, s) as methylene protons of the acetate group. The <sup>13</sup>C-NMR spectrum (figure 2) showed the presence of eleven signals carbon at δ<sub>C</sub> 30-210 ppm which consists of oxymethyl (δ<sub>C</sub> 55.8), ketone (δ<sub>C</sub> 206.4), methyl ketone (δ<sub>C</sub> 32.4), carboxylate (δ<sub>C</sub> 137.8), methylene acetate (δ<sub>C</sub> 40.8) carbon, and six aromatic carbons. The presence of two oxyaryl (= C-O) of aromatic carbons exhibited at δ<sub>C</sub> 163.7 and 160.5 ppm.

**Table 1: Antioxidant activity using DPPH method**

Samples	IC <sub>50</sub> (µg/mL)
DMP1	50.28
DMQ2	1963.02
DMR3	1036.15
Compound 1	38.01
Ascorbic acid	10.54

The HMQC spectrum (Figure 2) showed the presence of five  $^1\text{H}$ - $^{13}\text{C}$  correlations through one bond consisting of two correlations in the aromatic ring, one correlation to the methoxyl, methyl ketone, and methylene acetate groups, respectively. The HMBC spectrum (Figure 2) showed a correlation between the methylene acetate proton signal at  $\delta_{\text{H}}$  3.70 ppm with the carboxylate carbon ( $\delta_{\text{C}}$  175.4), and aromatic carbons ( $\delta_{\text{C}}$  110.7; 121.9; and 137.8 ppm). The correlation indicates that the methylene acetate group is attached directly to the aromatic ring. The methyl ketone proton at  $\delta_{\text{H}}$  2.53 showed correlation with the carbonyl carbon of the ketone group at  $\delta_{\text{C}}$  206.4 ppm. The proton  $\delta_{\text{H}}$  3.78 is far from the carbon in the aromatic ring with a chemical shift value of  $\delta_{\text{C}}$  163.7. The relationship between HMQC and HMBC can be seen in table 2. Best on the spectroscopy data, compound 1 was

identified as 2-(2-acetyl-3-hydroxy-5-methoxyphenyl)acetic acid or mono-O-methylcurvilinic acid (Figure 3) with good antioxidant activity  $\text{IC}_{50}$  38.01  $\mu\text{g}/\text{mL}$  (Table 1).

Mono-*o*-methylcurvilinic acid is not produced by the host *S. mahagoni*. However, the compound was also found in another endophytic fungi, namely *Drechslera indica* purchased from the Commonwealth Mycological Institute (CMI)<sup>31</sup> and *Penicillium mariae-crucis* was isolated from *Paeonia lactiflora* Pall.<sup>32</sup> The production of the same compound in some endophytic fungi is common. An example is (5-hydroxy-2-oxo-2H-pyran-4-yl) methyl acetate produced by the endophytic fungus *Cladosporium tenuissimum* from *S. mahagoni*,<sup>20</sup> *Trichoderma* sp. from *Tinospora crispa*,<sup>33</sup> and *Aspergillus* sp from *Andrographis paniculata*.<sup>34</sup>




Figure 1: Phylogenetic tree of DMP1

Table 2: The data of NMR spectrum of compound 1 ( $^1\text{H}$ -500 MHz,  $^{13}\text{C}$ -125 MHz, in  $\text{CD}_3\text{OD}$ ) and mono-*o*-methylcurvilinic acid 1\* ( $^1\text{H}$ -400 MHz in acetone)<sup>31</sup>

No. C	$\delta_{\text{C}}$ (ppm)	$\delta_{\text{H}}$ (ppm), $\Sigma\text{H}$ , mult, $J$ (Hz)	HMBC	$\delta_{\text{H}}$ (ppm), $\Sigma\text{H}$ , mult, $J$ (Hz)
	1	1	1	1*
1	137.8			
2	121.9			
3	160.5			
4	101.2	6.36 (1H, d, $J = 2.5$ Hz)	110.7; 160.5; 163.7	6.45 (1H, d, $J = 2.4$ Hz)
5	163.7			
6	110.7	6.34 (1H, d, $J = 2.5$ Hz)	40.8; 101.2; 121.9	6.55 (1H, d, $J = 2.4$ Hz)
1'	40.8	3.70 (2H, s)	110.7; 121.9; 137.8; 175.4	3.81 (2H, s)
2'	175.4			
3'	206.4			
4'	32.4	2.53 (3H, s)	206.4	2.56 (3H, s)
5-OCH <sub>3</sub>	55.8	3.78 (3H, s)	163.7	3.82 (3H, s)





Figure 2: The NMR spectra of compound 1

Figure 3: The  $\delta$ -assignment and HMBC correlation of compound 1 (mono-*O*-methylcurvilinic acid)

**Conclusion**

The endophytic fungus *Penicillium oxalicum* isolated from the leaves of *S. mahagoni* is able to produce mono-*o*-methylcurvulinic acid as an antioxidant compound. The compound apart from those produced by the host plant has been found in endophytic fungi.

**Conflict of interest**

The authors declare no conflict of interest.

**Authors' Declaration**

The authors hereby declare that the work presented in this article is original and that any liability for claims relating to the content of this article will be borne by them.

**Acknowledgements**

The research of this article was funded by DIPA of Universitas Sriwijaya, through "Penelitian Unggul Profesi" 2018-2020.

**References**

- Jahan N, Parvin MS, Das N, Islam MS, Islam ME. Studies on the antioxidant activity of ethanol extract and its fractions from *Pterygota alata* leaves. JACME 2024; (4/3):103-108.
- Dey TK, Emran TB, Saha D, Rahman MA, Hosen SZ, Chowdhury N. Antioxidant activity of ethanol extract of *Cassia hirsuta* (L.) Leaves. Bull Pharm Res. 2012; 2(2):78-81.
- Mahdi-Pour B, Jothy SL, Latha LY, Chen Y, Sasidharan S. Antioxidant activity of methanol extracts of different parts of *Lantana camara*. Asian Pac J Trop Biomed. 2013; 2(12):960-965.
- Ekin S, Bayramoglu M, Goktasoglu A, Ogoke F, Kiziltas H. Antioxidant Activity of Aqueous and Etanol Extracts of *Crataegus meyeri pojark* Leaves and Contents of Vitamin, Trace Element. J Chil Chem Soc. 2017; 62(4):3661-3667.
- Kumara PM, Shweta S, Vasanthakumari MM, Sachin N, Manjunatha BL, Jadhav SS, Ravikanth G, Ganeshiah KN, Shaanker RU. Endophytes and plant secondary metabolite synthesis: molecular and evolutionary perspective. In: V.C. Verma and A.C. Gange (eds.), *Advances in Endophytic Research*. New Delhi: Springer; 2014; 117-190p.
- Mawati N, Isfandito AR, Astuti DI, Aditiawati P. Endophytic Fungi from Surian (*Toona sinensis roem*) and Antioxidant potency from Its Culture. Asian J Plant Sci. 2011; 5(15):8-15.
- Huang W-Y, Cai Y-Z, Xing J, Corke H, Sun M. A Potential Antioxidant Resource: Endophytic Fungi from Medicinal Plants. Econ Bot. 2007; 61(1):14-30.
- Prihantini AI and Tachibana S. Antioxidant Compounds Produced By *Pseudocercospora* sp. ESL 02, an Endophytic fungus Isolated from *Eleocharis sylvestris*. Asian Pac J Trop Biomed. 2017; 7(2):110-115.
- Selim KA, Elkhateeb WA, Tawila AM, El-Beih AA, Abdel-Rahman TM, El-Diwanly AI, Ahmed EF. Antiviral and Antioxidant Potential of Fungal Endophytes of Egyptian Medicinal Plants. Fermentation. 2018; 4(49):2-11.
- Caicedo NH, Davalos AF, Puente PA, Rodriguez AY, Caicedo PA. Antioxidant Activity of Exo-Metabolites Produced By *Fusarium oxysporum*: An Endophytic Fungus from Leaves of *Otoba gracilipes*. Microbiologyopen. 2019; 8(1):1-7.
- Hajra S, Mehta A, Pandey P. Phenolic Compound and Antioxidant Activity of *Swietenia mahagoni* seed. Int J Pharm Sci. 2011; 3:431-434.
- Kahman SMA, Akter M, Hira TE, Mian MdY, Ahmed I, Rahman MM. Antioxidant and Antimicrobial Activities of Flower and Bark Extract of *Swietenia mahagoni* (L.) Jacq. J Pharmacogn Phytochem. 2104; 2(6):185-188.
- Bera TK, Chatterjee K, Ghosh D. In-vitro Antioxidant Properties of The Hydro-methanol Extract of The Seeds of *Swietenia mahagoni* (L.) Jacq. Biomark. Genom Med. 2015; 7(1):18-24.
- Sukardiman and Ervina M. The Recent Use of *Swietenia mahagoni* (L.) Jacq. As Antidiabetes Type 2 Phytomedicine: A Systematic Review. Heliyon. 2020; 6(e03536):1-8.
- Elfita, Muharni, Munawar, Aryani S. Secondary metabolite from Endophytic Fungi *Aspergillus niger* of The Stem Bark of Kandi Gajah (*Garcinia griffithii*). Indon J Chem. 2012; 12(2):195-200.
- Budiono EM, Yohandini H, Widjanti H. Antioxidant activity of *Syzygium samarangense* L. and their endophytic fungi. Molecules. 2019; 14(1):48-55.
- Elfita, Mardiyanto, Fitriya, Larasati JE, Julinar, Widjanti H, Muharni. Antibacterial activity of *Cordyline fruticosa* leaf extract and its endophytic fungi extract. Biodiv. 2019; 15(12):3804-3812.
- Gustianingtyas M, Herlinda S, Suwandi S, Hamidson H, Hasbi, Setiawan A, Verawaty M, Elfita A. Toxicity of entomopathogenic fungal culture filtrate of lowland and highland soil of South Sumatra (Indonesia) against *Spodoptera litura* larvae. Biodiversitas. 2020; 21(5):1839-1849.
- Metasari S, Elfita, Muharni, Yohandini H. Antioxidant compounds from the stem bark of *Syzygium samarangense*. Molecules. 2020; 15(3):175-183.
- Fadhillah, Elfita, Muharni, Yohandini H, Widjanti H. Chemical Compound Isolated from Antioxidant Active Extract of Endophytic Fungi *Cladosporium tenuissimum* in *Swietenia mahagoni* Leaf Stalk. Biodiv. 2019; 20(9):2645-2650.
- Gangwar M, Gautam MK, Sharma AK, Tripathi YB, Goel RK, Nath G. Antioxidant Capacity and Radical Scavenging Effect of Polyphenol Rich *Mallotus philippensis* Fruit Extract on Human Erythrocytes: An In Vitro Study. Sci World J. 2014; 2014:1-12.
- Akar Z, Kucuk M, Dogan H. A New Colorimetric DPPH Scavenging Activity Method with No Need for A Spectrophotometer Applied On Synthetic and Natural Antioxidant and Medicinal Herbs. J Enzym Inhib Med Chem. 2017; 32(10):60-647.
- Ivbulkovit K, Nouanthavong S, Sameenoi Y. Paper-based DPPH Assay for Antioxidant Activity Analysis. Anal Biochem. 2018; 34:795-800.
- Sadeer NB, Montesano D, Albrizio S, Zengin G, Mahmoodaly MF. Review The Versality of Antioxidant Assays Food Science and Safety-Chemistry, Applications, Strengths, and Limitations. MDPI. 2020; 9(709):2-39.
- Sun Y-X, Shen B-B, Han H-Y, Lu Y, Zhang B-X, Gao Y-F, Hu B-Z, Hu X-M. Screening of Potential Cellulases and Their Efficient Saccharification of IL-Pre-treated Cellulose. RSC Adv. 2018; 8(54):30957-30965.
- Kaur R, Kaur J, Kaur M, Kalotra V, Chandha P, Kaur A, Kaur A. An Endophytic *Penicillium oxalicum* Isolated from *Citrus limon* Possesses Antioxidant and Genoprotective Activity. J Appl Microbiol. 2019; 128(5):1400-1413.
- Bisht R, Sharma D, Agarwal PK. Antimicrobial and Antidiabetic Activity of An *Penicillium oxalicum* Isolated from *Cupressus torulosa*. Int J Biotechnol Biomed Sci. 2018; 2(2):119-122.
- Mamat S, Shah UKMd, Remli NAM, Shaari K, Laboh R, Rahman NAA. Characterization of Antifungal Activity of Endophytic *Penicillium oxalicum* T 3.3 for Anthracnose Biocontrol in Dragon Fruit (*Hylocereus* sp). Int J Environ.

- 428; 4(1):65-76.
29. Yen G-C and Lee C-A. Antioxidant Activity Of Extract  
21 in Molds. *J Food Prot.* 1996; 59(12): 1327-1330.
30. Bisht R, Bhatt A, Agarwal PK. Antimicrobial, Antioxidant  
Activity and Phytochemical Analysis Of An Endophytic  
Fungi *Penicillium oxalicum* Isolated From A *Gymnosperm*  
*Tree Cupressus Torulosa* D.Don. of Garhwal Region.  
24 *EM.* 2016; 1(2):90-96.
31. Kenfield D, Hallock Y, Clardy J, Strobel G. Curvulin and  
O-Methylcurvulinic Acid: Phytotoxic Metabolites of  
*Drechlera indica* which Caused Necroses on *Purslane* and  
*Spiny Amaranth*. *Plant Sci.* 1989; 60(1989):123-127.
32. Jutao W, Zonhui 1, 39 Luokai W, Fengqing X, Yang Y, Gang  
W, Jinsong L. Secondary Metabolites of Endophytic  
Fungus *Penicillium mariae-crucis* from *Paeonia lactiflora*  
31. *Chin J Org Chem.* 2019; 39:1136-1141.
33. Elfta E, Munawar M, Muharni M, and Sudrajat, M A.  
Identification of new lactone derivatives isolated from  
*Trichoderma sp.*, an endophytic fungus of brotowali  
(*Tinasporacrispa*). *Hayati J* 32 *sci.* 2014; 21(1):15-20.
34. Elfta, Munawar, Muharni, Ivantri I. Chemical constituents  
from an endophytic fungus *Aspergillus sp.* (Sbd5) isolated  
from sambilot (*Andrographis paniculata* Nees). *Microbiol*  
*Indon.* 2015; 9(2):82-88.

# Antioxidant Compound of Endophytic Fungi *Penicillium oxalicum* Isolated from *Swietenia mahagoni* L. Jacq

## ORIGINALITY REPORT

**41** %  
SIMILARITY INDEX

**34** %  
INTERNET SOURCES

**36** %  
PUBLICATIONS

**20** %  
STUDENT PAPERS

## PRIMARY SOURCES

1	<a href="http://seer.sct.embrapa.br">seer.sct.embrapa.br</a> Internet Source	1 %
2	<a href="http://biointerfaceresearch.com">biointerfaceresearch.com</a> Internet Source	1 %
3	Submitted to CVC Nigeria Consortium Student Paper	1 %
4	<a href="http://mdpi.com">mdpi.com</a> Internet Source	1 %
5	Submitted to Mizoram University Student Paper	1 %
6	<a href="http://old.biomedcentral.com">old.biomedcentral.com</a> Internet Source	1 %
7	<a href="http://cmuir.cmu.ac.th">cmuir.cmu.ac.th</a> Internet Source	1 %
8	<a href="http://ejournal2.litbang.kemkes.go.id">ejournal2.litbang.kemkes.go.id</a> Internet Source	1 %
9	Hui Zhang, Junhui Wu. "Statistical optimization of aqueous ammonia pretreatment and	1 %



enzymatic saccharification of corn stover for enhancing sugars production", Environmental Technology & Innovation, 2022

Publication

---

10

Q W Sari, P A Utari, D Setiabudidaya, I Yustian, E Siswanto, I Iskandar. "Variability of surface chlorophyll-a distributions in the northwestern coast of Sumatra revealed by MODIS", Journal of Physics: Conference Series, 2018

Publication

---

1 %

11

[www.thaiscience.info](http://www.thaiscience.info)

Internet Source

---

1 %

12

Khaled Selim, Waill Elkhateeb, Ahmed Tawila, Ahmed El-Beih, Tahany Abdel-Rahman, Ahmed El-Diwany, Eman Ahmed. "Antiviral and Antioxidant Potential of Fungal Endophytes of Egyptian Medicinal Plants", Fermentation, 2018

Publication

---

1 %

13

Selorm Torgbo, Prapassorn Rugthaworn, Udomlak Sukatta, Prakrit Sukyai. " Biological Characterization and Quantification of Rambutan ( L.) Peel Extract as a Potential Source of Valuable Minerals and Ellagitannins for Industrial Applications ", ACS Omega, 2022

Publication

---

1 %

14	Internet Source	1 %
15	Submitted to Sriwijaya University Student Paper	1 %
16	nveo.org Internet Source	1 %
17	Indah Solihah, Herlina Herlina, Inayatul Munawwaroh, Riana Sari Puspita Rasyid. "In vivo study of the antioxidant test of ethanolic extract of Chromolaena odorata Linn. leaves", MEDISAINS, 2020 Publication	1 %
18	M.A. Balendres, J.C. Bengoa. "Diseases of dragon fruit (Hylocereus species): Etiology and current management options", Crop Protection, 2019 Publication	1 %
19	wjgnet.com Internet Source	1 %
20	Nelson H. Caicedo, Andres F. Davalos, Paula A. Puente, Ana Y. Rodríguez, Paola A. Caicedo. " Antioxidant activity of exo - metabolites produced by An endophytic fungus isolated from leaves of ", MicrobiologyOpen, 2019 Publication	1 %

21	R. Kaur, J. Kaur, M. Kaur, V. Kalotra, P. Chadha, A. Kaur, A. Kaur. " An endophytic isolated from possesses antioxidant and genoprotective potential ", Journal of Applied Microbiology, 2020 Publication	1 %
22	<a href="http://www.appconnect.in">www.appconnect.in</a> Internet Source	1 %
23	<a href="http://nopr.niscair.res.in">nopr.niscair.res.in</a> Internet Source	1 %
24	Antonio Evidente, Anna Andolfi, Alessio Cimmino, Maurizio Vurro, Mariano Fracchiolla, Raghavan Charudattan. " Herbicidal Potential of Ophiobolins Produced by ", Journal of Agricultural and Food Chemistry, 2006 Publication	1 %
25	<a href="http://ijpsr.com">ijpsr.com</a> Internet Source	1 %
26	<a href="http://repository.maranatha.edu">repository.maranatha.edu</a> Internet Source	1 %
27	<a href="http://soar-ir.repo.nii.ac.jp">soar-ir.repo.nii.ac.jp</a> Internet Source	1 %
28	Maria Michela Salvatore, Anna Andolfi, Rosario Nicoletti. "The Genus Cladosporium: A Rich Source of Diverse and Bioactive Natural Compounds", Molecules, 2021	1 %

- 
- 29 [journal.umpr.ac.id](http://journal.umpr.ac.id) 1 %  
Internet Source
- 
- 30 [www.ijpsonline.com](http://www.ijpsonline.com) 1 %  
Internet Source
- 
- 31 Submitted to Pukyong National University 1 %  
Student Paper
- 
- 32 Suryelita Suryelita, Riga Riga, Sri Benti Etika,  
Mariam Ulfah, Muh Ade Artasasta. 1 %  
"Antibacterial Screening of Endophytic Fungus  
Xylaria sp. derived from Andrographis  
paniculata (Sambiloto)", Open Access  
Macedonian Journal of Medical Sciences, 2021  
Publication
- 
- 33 [nbn-resolving.org](http://nbn-resolving.org) 1 %  
Internet Source
- 
- 34 [www.ascidatabase.com](http://www.ascidatabase.com) 1 %  
Internet Source
- 
- 35 ELFITA, MUNAWAR, MUHARNI, MASTUR ADHY  
SUDRAJAT. "Identification of New Lactone  
Derivatives Isolated from Trichoderma sp., An  
Endophytic Fungus of Brotowali (Tinaspora  
crispa)", HAYATI Journal of Biosciences, 2014  
Publication
- 
- 36 [repository.unsri.ac.id](http://repository.unsri.ac.id) 1 %  
Internet Source
-

37	<a href="http://www.researchgate.net">www.researchgate.net</a> Internet Source	1 %
38	<a href="http://new.esp.org">new.esp.org</a> Internet Source	<1 %
39	Chun - Mei Chen, Wei - Hao Chen, Hua - Ming Tao, Bin Yang, Xue - Feng Zhou, Xiao - Wei Luo, Yong - Hong Liu. " Diversified Polyketides and Nitrogenous Compounds from the Mangrove Endophytic Fungus 41025 ", Chinese Journal of Chemistry, 2021 Publication	<1 %
40	<a href="http://pubag.nal.usda.gov">pubag.nal.usda.gov</a> Internet Source	<1 %
41	Submitted to Nguyen Tat Thanh University Student Paper	<1 %
42	Yen, Gow-Chin, Yung-Chi Chang, Fuu Sheu, and Hsin-Chang Chiang. "Isolation and Characterization of Antioxidant Compounds from <i>Aspergillus candidus</i> Broth Filtrate", Journal of Agricultural and Food Chemistry, 2001. Publication	<1 %
43	<a href="http://www.lenus.ie">www.lenus.ie</a> Internet Source	<1 %
44	"Steroids and isoquercetin from <i>Lygodium microphyllum</i> ", Journal of Applied	<1 %

# Pharmaceutical Science, 2017

Publication

---

45 M.J. Fang, H. Fang, W.J. Li, D.M. Huang, Z. Wu, Y.F. Zhao. " A new diphenyl ether from sp. strain, SHZK-2 ", Natural Product Research, 2012

Publication

---

46 Submitted to Universidade Estadual de Campinas

Student Paper

---

47 etd.aau.edu.et

Internet Source

---

48 Galvan, I.J.. "Antifungal and antioxidant activities of the phytomedicine pipsisewa, Chimaphila umbellata", Phytochemistry, 200802

Publication

---

49 Submitted to University of Massachusetts

Student Paper

---

50 ir.cnu.edu.tw

Internet Source

---

51 Miyata, O.. "Development of radical addition-cyclization-elimination reaction of oxime ether and its application to formal synthesis of (+/-)-martinelline", Tetrahedron, 20071001

Publication

52

Internet Source

&lt;1 %

53

[etd.repository.ugm.ac.id](http://etd.repository.ugm.ac.id)

Internet Source

&lt;1 %

54

[garuda.kemdikbud.go.id](http://garuda.kemdikbud.go.id)

Internet Source

&lt;1 %

55

[repositorio.unicamp.br](http://repositorio.unicamp.br)

Internet Source

&lt;1 %

56

[www.hindawi.com](http://www.hindawi.com)

Internet Source

&lt;1 %

57

[www.thieme-connect.de](http://www.thieme-connect.de)

Internet Source

&lt;1 %

58

[www.ukm.my](http://www.ukm.my)

Internet Source

&lt;1 %

59

[getjson.sid.ir](http://getjson.sid.ir)

Internet Source

&lt;1 %

60

A. I. Khodair, H. I. El-subbagh, A. M. Al-obaid. " SYNTHESIS, CONFORMATIONAL ANALYSIS AND ANTITUMOR TESTING OF 5-( )-ARYLIDENE-4-IMIDAZOLIDINONE DERIVATIVES ", Phosphorus, Sulfur, and Silicon and the Related Elements, 1998

Publication

&lt;1 %

61

Tao Yang, Changhong Wang, Hongjuan Liu, Guixin Chou, Xuemei Cheng, Zhengtao Wang.

&lt;1 %

" A new antioxidant compound from ",  
Pharmaceutical Biology, 2010

Publication

62

[docksci.com](http://docksci.com)

Internet Source

<1 %

63

[gsmis.gs.kku.ac.th](http://gsmis.gs.kku.ac.th)

Internet Source

<1 %

64

[sciencetechindonesia.com](http://sciencetechindonesia.com)

Internet Source

<1 %

65

[www.jcps.ac.cn](http://www.jcps.ac.cn)

Internet Source

<1 %

66

Khozirah Shaari, Suryati Safri, Faridah Abas,  
Nordin HJ Lajis, D. A. Israf. " A  
geranylacetophenone from the leaves of ",  
Natural Product Research, 2006

Publication

<1 %

67

Risyandi Anwar, Susi Sukmasari, Lilis Siti Aisya,  
Faridia Puspita Le, Delia Ilfani, Yenny Febriani  
Y, Pebrian Diki Prest. "Antimicrobial Activity of  
 $\beta$ -Sitosterol Isolated from Kalanchoe  
tomentosa Leaves Against Staphylococcus  
aureus and Klebsiella pneumonia", Pakistan  
Journal of Biological Sciences, 2022

Publication

<1 %

68

Sugandha Pant, Dhruv Mishra, Shubhpriya  
Gupta, Preeti Chaturvedi. "Fungal endophytes

<1 %



as a potential source of therapeutically important metabolites", Elsevier BV, 2021

Publication

69

[archives.gjasr.com](https://archives.gjasr.com)

Internet Source

<1 %

70

[pubs.rsc.org](https://pubs.rsc.org)

Internet Source

<1 %

71

[www.dovepress.com](http://www.dovepress.com)

Internet Source

<1 %

72

Íris Carolina dos Santos Lacerda, Júlio Cesar Polonio, Halison Correia Golias. "ENDOPHYTIC FUNGI AS SOURCES OF ANTIVIRAL COMPOUNDS - A REVIEW", *Chemistry & Biodiversity*, 2022

Publication

<1 %

73

Andri Frediansyah, Fajar Sofyantoro, Saad Alhumaid, Abbas Al Mutair et al. "Microbial Natural Products with Antiviral Activities, Including Anti-SARS-CoV-2: A Review", *Molecules*, 2022

Publication

<1 %

74

Kannan Baskar, Ragavendran Chinnasamy, Karthika Pandey, Manigandan Venkatesan et al. "Larvicidal and histopathology effect of endophytic fungal extracts of *Aspergillus tamarii* against *Aedes aegypti* and *Culex quinquefasciatus*", *Heliyon*, 2020

Publication

<1 %

75	Submitted to Mahidol University Student Paper	<1 %
76	Microbial Diversity and Biotechnology in Food Security, 2014. Publication	<1 %
77	dergipark.org.tr Internet Source	<1 %
78	patents.justia.com Internet Source	<1 %
79	Priyamvada Gupta, Ashish Verma, Nilesh Rai, Anurag Kumar Singh, Santosh Kumar Singh, Brijesh Kumar, Rajiv Kumar, Vibhav Gautam. "Mass Spectrometry-Based Technology and Workflows for Studying the Chemistry of Fungal Endophyte Derived Bioactive Compounds", ACS Chemical Biology, 2021 Publication	<1 %
80	"Fungi and their Role in Sustainable Development: Current Perspectives", Springer Science and Business Media LLC, 2018 Publication	<1 %

Exclude quotes      On  
Exclude bibliography      Off

Exclude matches      Off