

Diversity of endophytic fungi in *Syzygium aqueum*

by Hary Widjajanti

Submission date: 21-Sep-2022 12:08PM (UTC+0700)

Submission ID: 1905173907

File name: 22_DIVERSITY_OF_ENDOPHYTIC_FUNGI_BIODIVERSITAS_MARET_2021.pdf (437.96K)

Word count: 5231

Character count: 29874

Diversity of endophytic fungi in *Syzygium aqueum*

UMMI H. HABISUKAN^{1,2,*}, ELFITA^{3,}, HARY WIDJAJANTI⁴, ARUM SETIAWAN⁴,
ALFIA R. KURNIAWATI²**

¹Graduate School of Sciences, Faculty of Mathematics and Natural Sciences, Universitas Sriwijaya. Jl. Padang Selasa No. 524, Palembang 30139, South Sumatra, Indonesia. *email: ummhirash@gmail.com

²Department of Biology Education, Faculty of Tarbiyah and Teacher Training, Universitas Islam Negeri Raden Fatah. Jl. Prof. K.H. Zainal Fikri, Palembang 30126, South Sumatra, Indonesia

³Department of Chemistry, Faculty of Mathematics and Natural Sciences, Universitas Sriwijaya. Jl. Raya Palembang-Prabumulih Km 32, Indralaya, Ogan Ilir 30662, South Sumatra, Indonesia. Tel.: +62-711-580056, **email: elfita69@gmail.com

⁴Department of Biology, Faculty of Mathematics and Natural Sciences, Universitas Sriwijaya. Jl. Raya Palembang-Prabumulih Km 32, Indralaya, Ogan Ilir 30662, South Sumatra, Indonesia

Manuscript received: 15 October 2020. Revision accepted: 2 February 2021.

1
Abstract. Habisukan UH, Elfita, Widjajanti H, Setiawan A, Kurniawati AR. 2021. Diversity of endophytic fungi in *Syzygium aqueum*. *Biodiversitas* 22: 1129-1137. Exploring endophytic fungi may provide alternative, plant-based ethnomedicines. The discovery of endophytic fungi can produce many plant-derived drugs that give new horizons to the pharmaceutical industry for the availability and production of such medicines on a large scale. Extracts from *Syzygium aqueum* Alston (jambu air), a fruit-bearing plant native to Indonesia, have long been used in traditional medicine. Therefore, the purpose of this study was to identify, investigate, the antimicrobial activity of the endophytic fungi found in *S. aqueum*. Results revealed that total 16 fungi were isolated from different tissues (stem bark, root bark, and leaves) of the plant. Fungal endophytes were identified through morphological characterization and subsequently compared with key fungal identification books. The ethyl acetate extracts of the isolated endophytic fungi were screened for their antimicrobial activity through paper disc diffusion assay. Fungi isolated from *S. aqueum* were identified as: *Aspergillus niger* (Isolate code R21, R41, T72), *Cylindrocarpus* sp. (Isolate code R11, R31), *Trichoderma aureoviridae* (Isolate code T11), *Trichoderma harzianum* (Isolate code T21), *Trichoderma* sp. (Isolate code T31), *Pestalotia* sp. (Isolate code T41 and T62), *Beltrania* sp. (Isolate code T52), *Chaetomium* sp. (Isolate code L11), *Cochliobolus* sp. (Isolate code L22), *Penicillium* sp. (Isolate code L32), *Cylindrocladium* sp. (Isolate code L42), and an unidentified Ascomycota (T51). The values of Shannon-Weiner and Simpson diversity indexes for the overall fungal community were $H' = 2.133$ and $1-D = 0.859$ respectively. All isolates showed potential antimicrobial activity against *Escherichia coli*, *Salmonella Typhi*, and *Staphylococcus aureus*, but the lowest activity against *Candida albicans*. The results indicate that the bioactive compounds and secondary metabolites of these isolates should be further investigated for pharmaceutical applications.

Keywords: Antimicrobial activity, diversity, endophytic fungi, fungal identification, *Syzygium aqueum*

INTRODUCTION

Syzygium aqueum Alston, which belongs to the Myrtaceae family, is a fruit-bearing plant native to Indonesia and Malaysia. In Indonesia, it is well known as jambu air. The leaves, bark, roots, and other tissues of *S. aqueum* have long been used in traditional medicine (Cock and Chessman 2018). It has antibiotic, anti-inflammatory, antioxidant, and anti-tyrosinase properties (Palanisamy et al. 2011; Cock and Chessman 2018). Of these properties, the current researchers have chosen to analyze the antimicrobial activity of *S. aqueum*, which has also been highlighted by Mapatac and Mapaoag (2014) and Chua et al. (2019). Microbial infections are serious challenges to human health because of microbial resistance to antibiotics (Dos Santos et al. 2015). Plant-based ethnomedicines are derived from compounds occurring in plant extracts. Investigating the properties of endophytic fungi could be an alternative prospect for producing plant-derived compounds, which will benefit the pharmaceutical industry (Nicoletti and Florentino 2015).

Endophytic fungi are polyphyletic microorganisms, which inhabit plant tissues without inciting disease

symptoms and eventually establish mutualistic associations with their host plants. Numerous studies have been conducted on the diversity and isolation of endophytic fungi, which have revealed that fungi in plants are important and quantitative components of biodiversity, which have an impact on the diversity and composition of the plant community (Porras-Alfaro and Fayman 2011; Potshangbam et al. 2017). Exploration of the biosynthetic potential of endophytes has gained attention due to the ongoing discovery of strains that can synthesize plant compounds—a property that may reflect the adaptive functional role of endophytes in biocenosis (Nicoletti and Florentino 2015). Secondary metabolites serve multiple physiological functions, and, in a way, it is intuitive that similar compounds may be produced by ecologically associated entities (Jumpathong et al. 2010; Elfita et al. 2013; Budiono et al. 2019). Medicinal plants usually contain endophytic fungi, which produce natural compounds, similar to those found in the host plant such as saponins, alkaloids, flavonoids, tannins, phenolics, glycosides, and terpenoids (Budiono et al. 2019; Madhavi Ram 2015).

This is an initial step in exploring the potential of endophytic fungi as alternative strategies for the production of natural antimicrobial agents, and since there is no literature available for fungal communities associated with *S. aqueum*, the objectives of this study were: (i) to identify the endophytic fungi present in *S. aqueum*, (ii) to investigate the diversity of fungi associated with the stem bark, root bark, and leaves of *S. aqueum*, and (iii) to examine the antimicrobial activity of endophytic fungal extract.

MATERIALS AND METHODS

Sample collection

Syzygium aqueum plant samples were obtained from Ogan Ilir Regency, South Sumatra, Indonesia. Plant identification was confirmed at the Biosystematics Laboratory, Biology Department, Sriwijaya University, Ogan Ilir, Indonesia with certificate no. 329//UN9.1.7/4/EP/2020. Fresh and healthy samples of *S. aqueum* stem bark, root bark, and leaves were immediately transported to the laboratory.

Isolation of endophytic fungi

The *S. aqueum* stem bark, root bark, and leaf samples were washed with running tap water for 10 min and then air-dried. Sample fragments were successively surface sterilized by immersing each sample in 70% alcohol for ± 1 min, followed by immersion in 3% (w/v) sodium hypochlorite (NaOCl) for 1 min. After being rinsed in sterile distilled water for ± 1 min, the outer tissue of sample was removed with a sterile scalpel. Small pieces of stem bark, root bark, and leaves were placed in Petri dishes containing potato dextrose agar (PDA) media (200 g potato, 20 g dextrose, and 15 g agar in 1 L of H₂O) supplemented with chloramphenicol (0.2 g/L). Petri dishes were then incubated at $30 \pm 2^\circ\text{C}$ for seven days in the dark. All experiments were performed in triplicates. For the fungal growth from the leaf segments, the plates were monitored every day. Individual hyphal tips were transferred into fresh PDA and incubated at 30°C for seven days. The pure cultures were numbered, maintained on PDA slants, and kept at 4°C (Muharni et al. 2014; Elfita et al. 2019).

Identification of endophytic fungi

Macroscopic and microscopic features were used to characterize the morphology of the endophytic fungal isolates. Macroscopic characterization was based on the colony growth pattern, texture, margins, color, and other features. The microscopic characterization employed the slide culture method. Endophytic fungal cultures were placed on slides and mixed with one drop of lactophenol blue reagent. These slides were examined under a light microscope. The morphological characteristics data were then compared with key fungi identification books (Barnett 1969; Gandjar 1999; Watanabe 2002; Pitt and Hocking 2009).

Fungal diversity and ecological associations

Fungal diversity was estimated for each tissue sample and for the total sample population using the Shannon-Weiner (Shannon and Weiner 1949) and the Simpson (Simpson 1949) diversity indexes. Principal Component Analysis (PCA) was used to study the ecological interrelationships between the fungal species and different plant tissue types (PAST software, Hammer et al. 2001).

Cultivation and extraction

For the cultivation inoculum of fungi ($\pm 10^6$ spores/mL) was inoculated as much as 5% (v/v) in 300 ml of a potato dextrose broth (PDB) medium (20 g dextrose, 80 g potato, and 1,000 mL aquadest) in 1 L bottles. They were then incubated at room temperature for four to eight weeks. Color change in the samples indicated that secondary metabolite compounds had been formed. As a comparison, 300 ml of the PDB medium was placed into a separate 1 L bottle along with secondary metabolites, which had been partitioned in ethyl acetate and extracted via evaporation (Elfita et al. 2019; Gustianingtyas et al. 2020). The concentrated extracts were used for antimicrobial assay.

Antimicrobial activity

Antimicrobial activity was carried out via paper disc diffusion assay. Each sample was dissolved in dimethyl sulfoxide 10% (DMSO; Merck, Germany), and the antibacterial activity was evaluated against one gram-positive bacterium (*S. aureus*), two gram-negative bacteria (*S. typhi* and *E. coli*), and one yeast (*C. albicans*). These organisms were maintained on Muller-Hinton agar (MHA). The resulting suspension turbidity was diluted with sterile aquadest to match a 0.5 McFarland turbidity standard (1 McFarland is equivalent to approximately 3.0×10^8 CFU/mL) (Singh et al. 2015). The extract concentration used for this assay was 1,000 $\mu\text{g/mL}$. Absorbent disc (Whatman, 6.0 mm in diameter) was impregnated with 10 μL of the solution and then placed on the surface of inoculated plates (90 mm). Positive control discs of ciprofloxacin 10 $\mu\text{g/mL}$ were excluded from the assay. Diameters of the growth inhibition zones were measured after incubation at 37°C for 24 h. All experiments were performed in triplicates (Talukdar et al. 2020).

RESULTS AND DISCUSSION

A total of 16 endophytic fungi were isolated from different tissues (stem bark, root bark, and leaves) of *S. aqueum*. The growth of endophytic fungi isolates was indicated by the appearance of hyphae around the plant tissue segments (Figure 1). The colonies of fungal isolates exhibited various physical appearances. Massive hyphae growth appeared in the stem bark samples, with white colony dominance and yellow pigmentation. White filament colonization appeared around the root bark segments, and small, dark green and white colonies appeared on the leaf segments.

Identification of endophytic fungi

The 16 fungal colonies exhibited diversity in colors (white, black, green, and yellow) and texture (velvety, powdery, cottony, and granularly). Only isolates T11 and T21 evidenced exudates droplets and emitted yellow pigment. The macroscopic features of the isolated fungi are presented in Table 1.

Microscopic analysis showed that 16 isolates belong to ten different genera. Nine identified genera were (*Aspergillus*, *Cylindrocarpon*, *Trichoderma*, *Pestalotia*, *Beltrania*, *Chaetomium*, *Penicillium*, *Cochliobolus*, and *Cylindrocladium*) and one unidentified genus from the Ascomycota phylum

(Table 2). The endophytic fungi isolated from *S. aqueum* showed diversity in their spore shapes (cylindrical, globose, subglobose, spindle, and ellipsoidal). Most of the isolates had septate hyphae, except for isolates R11, R31, T51, and T11, which had coenocytic hyphae.

Four endophytic fungi were isolated from the root bark of *S. aqueum* (Figure 2), with the codes R11, R21, R31, and R41. Isolates R21 and R41 were spreading black colonies with white margins (Figure 2), and the microscopic characteristics showed that they had hyaline straight conidiophores, round vesicles, filial circling vesicles, and round, black conidia.




Figure 1. Mycelium appearance of endophytic fungi around *Syzygium aqueum* segments; A. Root bark; B. Stem bark; C. Leaves




Figure 2. Morphology of endophytic fungi isolated from *Syzygium aqueum* root bark. R11. *Cylindrocarpon* sp., R21. *A. niger*, R31. *Cylindrocarpon* sp., R41. *A. niger*; A. Front view colony; B. Reverse colony; C. Conidia without staining; D. Conidia with lactophenol blue dye

According to the key identification books, these isolates were identified as *A. niger*, which has specific features including black mass spores with globose vesicles. Isolate T72 had similar morphological characteristics to isolates R21 and R41, therefore isolate T72 was also identified as *A. niger*.

Isolates R11 and R31 were yellowish-white colonies with umbonate topography. Their microscopic features showed simple conidiospores (phialides). They were hyaline, erect, perceptic, and mostly 4-5-septate. They also evidenced ellipsoidal conidia and were apiculate at one end, slightly straight on one side, and curved on one side. According to their morphology, the closest species resembling R11 and R31 was *Cylindrocarpon* sp.

Isolate T41 was a white flowery colony with a cottony surface (Figure 3, T41 a-b). Microscopically, it showed short conidiospores, in spindle or ellipsoidal shapes, with four to five cells, two to three centrally pigmented cells, two to four appendages (setulae) in the apical cells, and one short appendage (pedicel) at the basal cell. Based on these characteristics, isolate T41 was identified as *Pestalotia* sp.

Isolate T52 (Figure 3, T52 a-b) was showing greenish brown with white margins and grew in a flowery pattern. Isolate T52 had pale brown conidiophores, had simple and occasionally proliferated forming nodes, and was inflated slightly at the apex. *Beltrania* sp. was the closest match to the fungus.

Four endophytic isolates i.e. L11, L22, L32, and L42 were isolated from the leaves of *S. aqueum* (Figure 4). Isolate L11 showed round, white colony with a cottony surface. Microscopically, it had thick perithecia and subglobose or ovate spores. It was covered with terminal hairs on the upper surface and was rhizoidal at its base. Thus, isolate L11 was thought to be closest to *Chaetomium* sp. Isolate L22 had a white cottony surface with a dark reverse color. Microscopically, it had septate hyphae, and its conidiophores were upright, brown, branched, straight or curved, porous, and subellipsoidal. It was mostly four-celled and was darker brown at the two-cell center, with a larger middle at the peripheral cells and indistinct hilar at the base. Thus, the characteristics of isolate L22 were considered to be closest to those of *Cochliobolus* sp.

Table 1. Macroscopic characteristics of endophytic fungi isolated from *Syzygium aqueum*

| Isolate | Colony color | Reverse colony color | Texture | Topography | Pattern | Exudate drops | Radial line | Concentric circle |
|---------|--------------------------|---------------------------|----------|------------|---------|---------------|-------------|-------------------|
| R11 | Yellowish white | Dark brown | Velvety | Umbonate | Zonate | - | √ | - |
| R21 | Black | Broken white | Powdery | Rugose | Spread | - | - | - |
| R31 | Yellowish white | Dark brown | Velvety | Umbonate | Zonate | - | √ | - |
| R41 | Black | Broken white | Powdery | Rugose | Spread | - | - | - |
| T11 | White with green spots | Light yellow | Granular | Flat | Radiate | √ | - | √ |
| T21 | White with green spots | Light yellow | Granular | Flat | Radiate | √ | - | √ |
| T31 | Dark green and white | Greenish white | Cottony | Umbonate | Zonate | - | √ | √ |
| T41 | White | Yellowish white | Cottony | Flat | Flowery | - | √ | - |
| T51 | Light green and white | Greenish white | Granular | Umbonate | Zonate | - | √ | √ |
| T52 | Brown and greenish-white | Army green | Cottony | Flat | Flowery | - | √ | √ |
| T62 | Dark green and white | Greenish white | Cottony | Umbonate | Zonate | - | √ | √ |
| T72 | Black | Broken white | Powdery | Rugose | Spread | - | - | - |
| L11 | White | Yellowish white | Cottony | Convex | Zonate | - | - | - |
| L22 | Grey with marginal white | Black with marginal white | Cottony | Verrucose | Zonate | - | - | √ |
| L32 | Dark green and white | Broken white | Powdery | Rugose | Spread | - | - | - |
| L41 | White | White | Cottony | Verrucose | Flowery | - | √ | - |

Note: - : characteristic does not appear; √: characteristic appear

Table 2. Microscopic characteristics and list of endophytic fungi isolated from *Syzygium aqueum*

| Isolate | Type of spore | Shape of spore | Hyphae | Specific characteristic | Genus / species |
|---------|---------------|----------------|------------|--|---------------------------------|
| R11 | Conidia | Cylindrical | Coenocytic | Macroconidia cylindrical, 4-5 septate | <i>Cylindrocarpon</i> sp. |
| R21 | Conidia | Globose | Septate | Globose vesicle, conidia black in mass | <i>Aspergillus niger</i> |
| R31 | Conidia | Cylindrical | Coenocytic | Macroconidia cylindrical, 4-5 septate | <i>Cylindrocarpon</i> sp. |
| R41 | Conidia | Globose | Septate | Globose vesicle, conidia black in mass | <i>Aspergillus niger</i> |
| T11 | Conidia | Subglobose | Septate | Phialide verticillate, chlamydospore subglobose | <i>Trichoderma aureoviridae</i> |
| T21 | Conidia | Subglobose | Septate | Spore masses apically at verticillate phialides | <i>Trichoderma harzianum</i> |
| T31 | Conidia | Globose | Septate | Phialide verticillate, chlamydospore subglobose | <i>Trichoderma</i> sp. |
| T41 | Conidia | Spindle | Septate | Conidia spindle, pigmented cell, apparent appendages | <i>Pestalotia</i> spp. |
| T51 | Sporangia | Globose | Coenocytic | Branched hyphae, apparent asci with granular spore | Unidentified |
| T52 | Conidia | Globose | Septate | Apex bearing 3-4 conidia sympodially | <i>Beltrania</i> sp. |
| T62 | Conidia | Ellipsoidal | Septate | Phialide verticillate, chlamydospore subglobose | <i>Trichoderma</i> sp. |
| T72 | Conidia | Globose | Septate | Globose vesicle, conidia black in mass | <i>Aspergillus niger</i> |
| L11 | Ascospore | Subglobose | Coenocytic | Ascocarps with well-developed hair, perithecia black | <i>Chaetomium</i> sp. |
| L22 | Conidia | Ellipsoidal | Septate | Subellipsoidal, mostly four-celled, central dark brown | <i>Cochliobolus</i> sp. |
| L32 | Conidia | Globose | Septate | Branch apex, 2-3 metulae, verticillate phialides | <i>Penicillium</i> sp. |
| L42 | Conidia | Cylindrical | Septate | Terminal vesicle subglobose, clavate, lanceolate | <i>Cylindrocladium</i> sp. |

Macroscopically, isolate L32 (Figure 4) was a spreading, dark green color colony with white edges. The reverse colony was yellowish white. Microscopically (Figure 1E), it had upright hyaline conidiophores, which branched into two to three metulae; each metula contained verticillate phialides, and each phialide contained dark, round, green conidia. After comparison, isolate L22 was identified as *Penicillium* sp.

Isolate L42 identified as *Cylindrocladium* sp. The colony was white with a more yellowish reverse colony. Its surface was cottony, and it grew in a flowery pattern (Figure 4, L42 a-b). Its conidiophores were erect and branched, with elongated primary and secondary branches. The phialide branches carried spore masses with cylindrical stalk and terminal vesicles. The conidia were cylindrical and phialosporous (Figure 4, L42 c-d).

Endophytic fungal diversity and ecological association

Eight isolates were obtained from the stem bark of *S. aqueum*, and four were obtained from the root bark and leaves. The diversity index values (Shannon-Wiener and Simpson) of *S. aqueum* tissues are listed in Table 3. The α -diversity values were obtained via the PAST software (Shadegi et al. 2019). The Shannon-Weiner and Simpson diversity index values were highest in leaves ($H' = 1.386$ and $1-D = 0.750$, respectively) and lowest in root bark ($H' = 0.693$ and $1-D = 0.500$, respectively). In addition, the overall diversity values of the endophytic fungal populations were $H' = 1.133$ and $1-D = 0.859$ (Table 3).

The two principal components or axes (1-2) obtained via PCA explained 68% of the total fungal variation (Figure 5). The analysis showed that some isolates had an affinity for a specific tissue. Host affinity could reflect functional differences in phenology, chemistry, or other traits, which vary among fungi and host plant taxa. Differences in diversity and species composition between stems and leaves may also reflect their functional similarities.

Antimicrobial activity of endophytic fungi

Antimicrobial activity of the endophytic fungi isolated from *S. aqueum* is presented in Table 4. Ten out of 16 isolates were used in the antimicrobial analysis. Overall, the endophytic fungal extracts inhibited the growth of the three bacterial species, but they had little effect on the yeast (*C. albicans*). Isolate R42 (*A. niger*) showed strong antimicrobial activity against *E. coli*, and isolate T31 (*T. harzianum*) showed strong antimicrobial activity against *S. aureus* and *S. typhi*. On average, the lowest level of antimicrobial activity was presented by isolate L22 (*Cochliobolus* sp.).

Discussion

The sampling of *S. aqueum* endophytes indicated that a diverse community of fungi live on the plant's various tissues. The purpose of surface sterilization during the isolation process was to remove microorganisms from the plant surface so that the fungi growing in the media are only endophytic fungi (Strobel and Daisy 2003). 3%

NaOCl and 70% alcohol were used in chemical solutions for surface sterilization as both are widely employed as disinfectants. Hypochlorite ions regenerated from hypochlorous acid (HOCl) and dissolved in water are capable of damaging bacterial plasma membranes (Rahayu et al. 2019).

Results showed that all endophytic fungi isolated from the different tissues of *S. aqueum* were categorized into three classes of Ascomycota: Eurotiomycetes, Sordariomycetes, and Ascomycetes. To date, most fungi reported as endophytes have been identified as Ascomycetes and their anamorphs. Basidiomycetous endophytes have only been reported in a limited number of studies (Rivera-Orduña et al. 2011; Sridhar and Raviraja 1995; Wang et al. 2005). The most evenly distributed genus *Trichoderma* isolated only from *S. aqueum* stem bark. *Trichoderma* species are typically soilborne organisms commonly associated with plant roots and stems. Some strains of *Trichoderma*, such as *T. harzianum*, *T. stromaticum*, and *T. asperellum*, have antagonistic effects on some fungal diseases (Bailey et al. 2008; Gazis and Chaverri 2010).

Some isolated endophytic taxa prefer specific tissue types. Some sapwood endophytes may invade tree hosts through their root systems, moving from the soil to their new niche inside the plant (Gazis and Chaverri 2010). In this research, PCA showed an increase in occurrence and frequency of tissue specificity (Rodrigues 1994; Photita et al. 2001; Kumar and Hyde 2004). Certain endophytes were found only in one *S. aqueum* tissue (stem bark, root bark, or leaves), indicating that they may have an affinity for a particular tissue type (Rivera-Orduña et al. 2011). The fungal endophytes from *S. aqueum* had a greater affinity for the stem bark than for any other tissue.

Antimicrobial analysis results showed that endophytic fungal extracts have the ability to inhibit the growth of bacterial species. Harjoto et al. (2015) reported that *S. aqueum* leaf extract inhibited the growth of *Shigella dysenteriae*, *Vibrio cholerae*, *S. aureus*, *E. Coli*, and *S. typhi*. He also reported that *S. aqueum* leaf extract showed strong activity against *E. Coli*, *S. typhi*, and *S. aureus* at 50% concentration. Sobeh et al. (2016) have also reported that some bioactive compounds are found in the essential oils of *S. aqueum* leaves, which exhibit antimicrobial activity. Phenolic and flavonoid compounds are dominant in the *S. aqueum* plant, and both have strong antioxidant and antimicrobial properties (Tehrani et al. 2011; Palanisamy et al. 2011; Marinova 2005). The in vitro antibacterial activity of several classes of polyphenols can be due to direct action against bacteria, viruses, and fungi, as well as the suppression of microbial virulence factors (Daglia 2012). The most common constituents in *S. aqueum* with antimicrobial properties are α -selinene, β -caryophyllene, and δ -selinene (Sobeh et al. 2016). *Trichoderma* species showed strong antimicrobial activity against *S. aureus* and *S. typhi*. *Trichoderma* species can protect their hosts from direct parasitism, antibiosis, and nutrient competition via enhanced plant growth or induced resistance (Gazis and Chaverri 2010).




Figure 3. Morphology of endophytic fungi isolated from *Syzygium aqueum* stem bark; T11. *T. aureoviridae*, T21. *T. harzianum*, T31. *Trichoderma* sp., T41. *Pestalotia* sp., T51. Unidentified, T52. *Beltrania* sp., T62. *Trichoderma* sp., T72. *A. niger*; A. Front view of colony; B. Reverse colony; C. Conidia without staining; D. Conidia with lactophenol blue dye




Figure 4. Morphology of endophytic fungi isolated from *Syzygium aqueum* leaves; L11. *Chaetomium* sp., L22. *Cochliobolus* sp., L32. *Penicillium* sp., L42. *Cylindrocladium* sp.; A. Front view of colony; B. Reverse colony; C. Conidia without staining; D. Conidia with lactophenol blue dye

Table 3. Fungal diversity and fungal endophytes isolated from different *Syzygium aqueum* tissues

| Taxon (Genera) | Tissues of <i>Syzygium aqueum</i> | | | |
|------------------------------------|-----------------------------------|-----------|-----------|----------------|
| | Leaves | Stem bark | Root bark | Total isolated |
| <i>Cylindrocarpon</i> | 0 | 0 | 2 | 2 |
| <i>Aspergillus</i> | 0 | 1 | 2 | 3 |
| <i>Trichoderma</i> | 0 | 4 | 0 | 4 |
| <i>Pestalotia</i> | 0 | 1 | 0 | 1 |
| Unidentified | 0 | 1 | 0 | 1 |
| <i>Beltrania</i> | 0 | 1 | 0 | 1 |
| <i>Chaetomium</i> | 1 | 0 | 0 | 1 |
| <i>Cochliobolus</i> | 1 | 0 | 0 | 1 |
| <i>Penicillium</i> | 1 | 0 | 0 | 1 |
| <i>Cylindrocladium</i> | 1 | 0 | 0 | 1 |
| 76 of total fungal isolates | 4 | 8 | 4 | 16 |
| Simpson's index (D) | 0.250 | 0.313 | 0.500 | 0.141 |
| Simpson's index of diversity (1-D) | 0.750 | 0.688 | 0.500 | 0.859 |
| Shannon index of diversity (H') | 1.386 | 1.386 | 0.693 | 2.133 |




Figure 5. PCA of fungal endophytes isolated from different tissues of *Syzygium aqueum*. Components 1 and 2 explained 68% of the total fungal variation

Table 4. Antimicrobial activity of endophytic fungi

| Code isolate | Isolate name | ⁷⁹ <i>Escherichia coli</i> | <i>Staphylococcus aureus</i> | <i>Salmonella typhi</i> | <i>Candida albicans</i> |
|-------------------------------|--------------------------------|---------------------------------------|------------------------------|-------------------------|-------------------------|
| R31 | <i>Cylindrocarpon</i> sp. | ++ | ++ | ++ | + |
| R41 | <i>Aspergillus niger</i> | +++ | ++ | + | + |
| T11 | <i>Trichoderma aureoviride</i> | ++ | ++ | ++ | - |
| T31 | <i>Trichoderma</i> sp. | ++ | +++ | +++ | + |
| T41 | <i>Pestalotia</i> sp. | ++ | + | + | + |
| T52 | <i>Beltrania</i> sp. | + | ++ | ++ | + |
| L11 | <i>Chaetomium</i> sp. | ++ | ++ | + | + |
| L22 | <i>Cochliobolus</i> sp. | + | + | + | + |
| L32 | <i>Penicillium</i> sp. | ++ | + | ++ | + |
| L42 | <i>Cylindrocladium</i> sp. | ++ | ++ | + | + |
| Positive control ^a | - | +++ | ⁵⁹ +++ | +++ | + |

Note: ^a: Ciprofloxacin 30 µg/m ⁶³ ++: strong activity with an inhibition zone >20 mm; +: moderate activity with an inhibition zone 10-20 mm; +: low activity with an inhibition zone 1-9 mm; -: negative activity with an inhibition zone <1 mm (Islamia et al. 2019).

In conclusion, 16 endophytic fungi ³⁹ were isolated from *S. aqueum* stem bark, root bark, and leaves. These endophytic fungi belonged to nine genera such as *Aspergillus*, *Cylindrocarpon*, *Trichoderma*, *Pestalotia*, *Beltrania*, *Chaetomium*, *Penicillium*, *Cochliobolus*, and *Cylindrocarpon*. The diversity values of the endophytic fungal populations were $H' = 2.133$ according to the Shannon-Weiner index and $1-D = 0.859$ according to the Simpson index. Therefore, it shows that *S. aqueum* can supply a comfortable habitat and nutrients for fungal endophyte survival. Ten endophytic fungal extracts exhibited antimicrobial activity. The high antimicrobial potential was shown to be exclusively by *Trichoderma* sp. (T31), which acted strongly against *S. aureus* and *S. typhi* but had low antimicrobial activity against *C. albicans*. Therefore, *S. aqueum* is worth further investigating its bioactive secondary metabolites and isolating their bioactive compounds to produce antioxidant and antimicrobial agents.

ACKNOWLEDGEMENTS

This research ¹⁴ was funded by *Daftar Isian Pelaksanaan Anggaran (DIPA)* of the Public Service Agency ¹⁴ at Universitas Sriwijaya, Indonesia 2020. SP DIPA-023.17.2.677515/2020, revision IV, July 2, 2020, in accordance with Rector's Decree number 0009/UN9/SK.LP2M.PT/2020, September 28, 2020.

⁴⁹ REFERENCES

- Bailey BA, Bae H, Strem MD, Crozier J, Thomas SE, Samuels GJ, Vinyard BT, Holmes KA. 2008. Antibiosis, mycoparasitism, and colonization success for endophytic *Trichoderma* isolate with biological control potential in *Theobroma cacao*. *Biol Control* 46 (1): 24-35.
- ⁶⁰ Barnett HL. 1969. *Illustrated Genera of Imperfect Fungi*. Burgess Publishing Company, Virginia.
- ¹³ Budiono B, Elfita E, Muhami M, Yohandini H and Widjajanti H. 2019. Antioxidant activity of *Syzygium marangense* L. and their endophytic fungi. *Molekul* 14 (1): 48-55. DOI: <http://dx.doi.org/10.20884/1.jm.2019.14.1.503>.
- ¹² Chua LK, Lim CL, Ling AP, Chye SM, Koh RY. 2019. Anticancer potential of *Syzygium* species: a review. *Plant Foods Human Nutr* 74 (1): 18-27.
- Cock IE, Cheesman M. 2018. Plants of the genus *Syzygium* (Myrtaceae): A review on ethnobotany, medicinal properties and phytochemistry. In: Goyal MR, Ayeleso AO (eds) *Bioactive Compounds of Medicinal Plants: Properties and Potential for Human Health*. Apple Academic Press, Inc., Palm Bay, FL.
- ⁶ Daglia M. 2012. Polyphenols as antimicrobial agents. *Curr Opin Biotechnol* 23 (2): 174-181.
- ⁵² Santos Ferreira da Silva J, López Malo D, Anceschi Bataglion G, Nogueira Eberlin M, Machado Ronconi C, Alves Júnior S, de Sá GF. 2015. Adsorption in a fixed-bed column and stability of the antibiotic oxytetracycline supported on Zn (II)-[2-methylimidazole] frameworks in aqueous media. *PLoS One* 10 (6): e0128436. DOI: <https://doi.org/10.1371/journal.pone.0128436>.
- ¹³ Elfita E, Mardiyanto M, Pitrya F, Larasati J, Julinar J, Widjajanti H, and Muharni M. 2019. Antibacterial activity of *Cordyline fruticosa* leaf extracts and its endophytic fungi extracts. *Biodiversitas* 20 (12): 3804-3812.
- ¹⁷ Elfita E, Munawar M, Muhami M, Suprayetno S. 2013. New pyran of an endophytic fungus *Fusarium* sp. isolated from the leaves of brotowali (*Tinospora crispa*). *J Chem* 13 (3): 209-215.
- Gandjar I, Rifai MA. 1999. *Pengenalan kapang tropik umum*. Yayasan Obor Indonesia. [Indonesian]
- Gaziz R, Chaverri P. 2010. Diversity of fungal endophytes in leaves and stems of wild rubber trees (*Hevea brasiliensis*) in Peru. *Fungal Ecol* 3 (3): 240-254.
- ⁶⁹ Kristianingtyas M, Herlinda S, Suwandi, Suparman, Hamidson H, Hasbi, Setiawan A, Verawaty M, Elfita, Arsi. 2020. Toxicity of entomopathogenic fungal culture filtrate of lowland and highland soil of South Sumatra (Indonesia) against *Spodoptera litura* larvae. *Biodiversitas* 21: 1839-1849.
- Hammer, Harper DAT, Ryan PD. 2001. PAST: Paleontological statistics software package for education and data analysis. *Paleontol Electron* 4: 4-9.
- ²³ Hariyati T, Soelistya D, Jekti DSD, Andayani Y. 2015. Pengaruh ekstrak etanol daun jambu air (*Syzygium aqueum*) terhadap bakteri isolat klinis. *Jurnal Penelitian Pendidikan IPA* 1 (2): 31-38. [Indonesian]
- ⁶⁴ Jumpathong J, Abdalla MA, Lumyong S, Laatsch H. 2010. Stempholgalactoside, a new stemphol derivative isolated from the tropical endophytic fungus *Gaeumannomyces cesamomi*. *Nat Prod Commun* 5 (4): 1934578X1000500415.
- ³² Kumar DSS, Hyde KD. 2004. Biodiversity and tissue-recurrence of endophytic fungi in *Tripterium wilfordii*. *Fungal Divers* 17: 69-90.
- Madhavi M, Ram MR. 2015. Phytochemical screening and evaluation of biological activity of root extracts of *Syzygium samarangense*. *Int J Res Pharm Chem* 5 (4): 753-763.
- ²⁶ Mapatae LC, Mamaog NR. 2014. Efficacy of three varieties of *Syzygium aqueum* (Tambis) as antimicrobial agent and its bioactive component. *Int J Sci Clin Lab* 9: 1.

- 10 Mannova D, Ribarova F, Atanassova M. 2005. Total phenolics and total flavonoids in Bulgarian fruits and vegetables. *J Univ Chem Technol Metallurgy* 40 (3): 233-260.
- Muharni M, Fitriya F, Ruliza MO, Susanti DA, Elfita E. 2014. Di-(2-ethylhexyl) phthalate and pyranone derived from endophytic fungi *Penicillium sp.* the leave of Kunyit Putih (*Curcuma zedoaria*). *Indon J Chem* 14 (3): 290-296.
- 31 Nicoletti R, Fiorentino A. 2015. Plant bioactive metabolites and drugs produced by endophytic fungi of Spermatophyta. *Agriculture* 5 (4): 918-970.
- 18 Palanisamy UD, Ling LT, Manaharan T, Appleton D. 2011. Rapid isolation of geraniin from *Nephelium lappaceum* rind waste and its anti-hyperglycemic activity. *Food Chem* 127: 21-27.
- Palanisamy U D, Ling L T, Manaharan T, Sivapalan V, Subramaniam T, Helme M H and Masilamani T. 2011. Standardized extract of *Syzygium aqueum*: a safe cosmetic ingredient. *Int J Cosmet Sci* 33 (3): 269-275.
- 6 Photita W, Lumyong S, Lumyong P, Hyde KD. 2001. Endophytic fungi of wild banana (*Musa acuminata*) at Doi Suthep Pui National Park, Thailand. *Mycol Resour* 105: 1508-1513.
- 46 Pitt JI, Hocking AD. 2009. *Fungi and food spoilage* (Vol. 519, p. 388). Springer, New York.
- 7 Potshangbam M, Devi SI, Sahoo D, Strobel GA. 2017. Functional characterization of endophytic fungal community associated with *Oryza sativa* L. and *Zea mays* L. *Front Microbiol* 8: 325.
- 20 Porras-Alfaro A, Bayman P. 2011. Hidden fungi, emergent properties: endophytes and microbiomes. *Annu Rev Phytopathol* 49: 291-315.
- 86 Rahayu ES, Widiatningrum T, Herlina L, Hermayani N, Amalia AR. 2019. The optimal sterilizing compound and culture medium in *Elaeocarpus grandiflorus* L. in vitro shoot induction. *J Phys: Conf Ser* 1321 (3): 032040.
- Rivera-Orduñ FN, Suarez-Sanchez RA, Flores-Bustamante ZR, Gracida-Rodriguez JN, Flores-Cotera LB. 2011. Diversity of endophytic fungi of *Taxus globosa* (Mexican yew). *Fungal Divers* 47 (1): 65-74.
- Rodrigues KF. 1994. The foliar fungal endophytes of the Amazonian palm *Euterpe oleracea*. *Mycologia* 86: 376-385.
- 53 Sadeghi F, Samsampour D, Seyahooei MA, Bagheri A, Soltani J. 2019. Diversity and spatiotemporal distribution of fungal endophytes associated with *Citrus reticulata* cv. Siyadoo. *Curr Microbiol* 76 (3): 279-289.
- 30 Singh R, Shushni MA, Belkheir A. 2015. Antibacterial and antioxidant activities of *Mentha piperita* L. *Arabian J Chem* 8 (3): 322-328.
- 50 Sobeh M, Braun, MS, Krstin S, Youssef FS, Ashour ML, Wink M. 2016. Chemical profiling of the essential oils of *Syzygium aqueum*, *Syzygium samarangense*, and *Eugenia uniflora* and their discrimination using chemometric analysis. *Chem Biodivers* 13 (11): 1537-1550.
- 45 Sridhar KR, Raviraja NS. 1995. Endophytes—a crucial issue. *Curr Sci* 69: 570-571.
- 33 Strobel G, Daisy B. 2003. Bioprospecting for microbial endophytes and their natural products. *Microbiol Mol Biol Rev* 67 (4): 491-502.
- 51 Talukdar R, Wary S, Mili C, Roy S, Tayung K. 2020. Antimicrobial secondary metabolites obtained from endophytic fungi inhabiting healthy leaf tissues of *Houttuynia cordata* Thunb., an ethnomedicinal plant of Northeast India. *J Appl Pharmaceutical Sci* 10 (09): 099-106.
- 22 Tehrani M, Sharif Hossain ABM, Nasrulhaq-Boyce A. 2011. Post-harvest physicochemical and mechanical changes in Jambu Air (*Syzygium aqueum* Alston) fruits. *Aust J Crop Sci* 5 (1): 32.
- 10 Wang Y, Guo LD, Hyde KD. 2005. Taxonomic placement of sterile morphotypes of endophytic fungi from *Pinus tabulaeformis* (Pinaceae) in northeast China based on rDNA sequences. *Fungal Divers* 20: 235-260.
- 15 Watanabe T. 2002. *Pictorial Atlas of Soil and Seed Fungi: Morphologies of Cultured Fungi and Key to Species* 2nd ed. CRC Press LLC, Florida.

Diversity of endophytic fungi in *Syzygium aqueum*

ORIGINALITY REPORT

38%

SIMILARITY INDEX

30%

INTERNET SOURCES

32%

PUBLICATIONS

15%

STUDENT PAPERS

PRIMARY SOURCES

1

www.sciencegate.app

Internet Source

5%

2

Submitted to Institut Pertanian Bogor

Student Paper

2%

3

doaj.org

Internet Source

2%

4

www.mdpi.com

Internet Source

1%

5

Nurhaida Nurhaida, Syarifah A. Rashid, Darah Ibrahim. " The Potency of endophytic fungi from the bark of (*L.*) Skeels as antifungal agents ", E3S Web of Conferences, 2020

Publication

1%

6

ebin.pub

Internet Source

1%

7

www.frontiersin.org

Internet Source

1%

8

Robert Jankowiak, Piotr Bilański, Jarosław Paluch, Zbigniew Kołodziej. "Fungi associated

1%

with dieback of *Abies alba* seedlings in naturally regenerating forest ecosystems", *Fungal Ecology*, 2016

Publication

9

Hui Jin, Zhiqiang Yan, Quan Liu, Xiaoyan Yang, Jixiang Chen, Bo Qin. "Diversity and dynamics of fungal endophytes in leaves, stems and roots of *Stellera chamaejasme* L. in northwestern China", *Antonie van Leeuwenhoek*, 2013

Publication

1 %

10

academicjournals.org

Internet Source

1 %

11

H. Marzouki, A. Piras, K. Bel Haj Salah, H. Medini, T. Pivetta, S. Bouzid, B. Marongiu, D. Falconieri. " Essential oil composition and variability of *L.* growing in Tunisia, comparison and chemometric investigation of different plant organs ", *Natural Product Research*, 2009

Publication

1 %

12

Submitted to University of Aberdeen

Student Paper

1 %

13

iopscience.iop.org

Internet Source

1 %

14

www.atlantis-press.com

Internet Source

1 %

| | | |
|----|---|------|
| 15 | hal.univ-lorraine.fr Internet Source | 1 % |
| 16 | docplayer.net Internet Source | 1 % |
| 17 | eprints.unsri.ac.id Internet Source | 1 % |
| 18 | Hong Sheng Cheng, So Ha Ton, Khalid Abdul Kadir. "Ellagitannin geraniin: a review of the natural sources, biosynthesis, pharmacokinetics and biological effects", <i>Phytochemistry Reviews</i> , 2016 Publication | 1 % |
| 19 | drum.lib.umd.edu Internet Source | 1 % |
| 20 | mafiadoc.com Internet Source | 1 % |
| 21 | ouci.dntb.gov.ua Internet Source | 1 % |
| 22 | epdf.tips Internet Source | <1 % |
| 23 | media.neliti.com Internet Source | <1 % |
| 24 | biosains.mipa.uns.ac.id Internet Source | <1 % |

| | | |
|----|---|------|
| 25 | pilotscholars.up.edu Internet Source | <1 % |
| 26 | www.phcogres.com Internet Source | <1 % |
| 27 | Mohamed Sheashea, Seham El-Hawary, Fatma A. Ahmed, Mohamed A. Salem, Marwa I. Ezzat. "UPLC-MS-based metabolomics profiling and chemometric analysis for <i>Hypericum sinaicum</i> Boiss and the endophytic <i>Aspergillus foetidus</i> in comparison to <i>Hypericum perforatum</i> L", Research Square Platform LLC, 2022 Publication | <1 % |
| 28 | www.degruyter.com Internet Source | <1 % |
| 29 | Romina Gazis, Priscila Chaverri. "Diversity of fungal endophytes in leaves and stems of wild rubber trees (<i>Hevea brasiliensis</i>) in Peru", <i>Fungal Ecology</i> , 2010 Publication | <1 % |
| 30 | dergipark.org.tr Internet Source | <1 % |
| 31 | journal.unnes.ac.id Internet Source | <1 % |
| 32 | Satish K. Verma, Surendra K. Gond, Ashish Mishra, Vijay K. Sharma et al. "Impact of | <1 % |

environmental variables on the isolation, diversity and antibacterial activity of endophytic fungal communities from *Madhuca indica* Gmel. at different locations in India", *Annals of Microbiology*, 2013

Publication

33

d-nb.info

Internet Source

<1 %

34

Hao Wu, Hongyan Yang, Xiangling You, Yuhua Li. "Isolation and Characterization of Saponin-Producing Fungal Endophytes from *Aralia elata* in Northeast China", *International Journal of Molecular Sciences*, 2012

Publication

<1 %

35

Watanabe, . "'Anamorphic Fungi : Deuteromycetes, Fungi Imperfecti, Mitosporic Fungi'", *Pictorial Atlas of Soil and Seed Fungi Morphologies of Cultured Fungi and Key to Species Third Edition*, 2010.

Publication

<1 %

36

www.geneticsmr.com

Internet Source

<1 %

37

A. B. M. Neshar Uddin, Farhad Hossain, A. S. M. Ali Reza, Mst. Samima Nasrin, A. H. M. Khurshid Alam. " Traditional uses, pharmacological activities, and phytochemical constituents of the genus : A review ", *Food Science & Nutrition*, 2022

<1 %

38

[bmccomplementmedtherapies.biomedcentral.com](https://www.bmccomplementmedtherapies.biomedcentral.com) <1 %
Internet Source

39

Pascale Clivio, Bernard Richard, Jean-Marc Nuzillard, Monique Zèches-Hanrot. "16-epi-silicine, an alkaloid of the ervatamine-type from *Pandaca caducifolia*", *Phytochemistry*, 1995
Publication

40

Sumit Bansal, Shivani Choudhary, Manu Sharma, Suthar Sharad Kumar, Sandeep Lohan, Varun Bhardwaj, Navneet Syan, Saras Jyoti. "Tea: A native source of antimicrobial agents", *Food Research International*, 2013
Publication

41

Nur Arfa Yanti, Jamili, Ardiansyah, Syaiful Anwarrudin. "Antibacterial activity of fungi endophytic isolated from leaves the mangrove *acanthus ilicifolius* I", *Journal of Physics: Conference Series*, 2021
Publication

42

E S Rahayu, T Widiatningrum, L Herlina, N Hermayani, A R Amalia. "The optimal sterilizing compound and culture medium in *Elaeocarpus grandiflorus* L. in vitro shoot induction", *Journal of Physics: Conference Series*, 2019
Publication

43

Submitted to King Mongkut's Institute of
Technology Ladkrabang

Student Paper

<1 %

44

Naidu, Yuvarani, Abu Seman Idris, Ahmad
Zairun Madihah, and Norman Kamarudin. "In
vitro Antagonistic Interactions Between
Endophytic Basidiomycetes of Oil Palm (*Elaeis
guineensis*) and *Ganoderma boninense*",
Journal of Phytopathology, 2016.

Publication

<1 %

45

[epdf.pub](#)
Internet Source

<1 %

46

[www.springerprofessional.de](#)
Internet Source

<1 %

47

Jae-Hak Moon, Seung-Jin Ma, Hyang-Hee Lee,
Naoharu Watanabe, Akihito Yagi, Kanzo
Sakata, Keun-Hyung Park. " Isolation and
Structural Determination of a Novel
Antimicrobial Compound from the Root of ",
Natural Product Letters, 2000

Publication

<1 %

48

V. V. Mulwad, B. P. Langi, J. Chaskar, A.
Chaskar. "Synthesis and antimicrobial activity
of [6'-methyl-4'-methoxy-2-oxo-2H-[1]-
benzopyran)-2",4"-dihydro-[1", 2", 4"]-triazol-
3"-one and 3"-phenyl-thiazolidin-4"-one-
phenoxymethyl derivatives of

<1 %

49

docksci.com

Internet Source

<1 %

50

ftp.academicjournals.org

Internet Source

<1 %

51

Submitted to Management & Science
University

Student Paper

<1 %

52

ehp.niehs.nih.gov

Internet Source

<1 %

53

jtrolis.ub.ac.id

Internet Source

<1 %

54

B.A. Bailey, H. Bae, M.D. Strem, J. Crozier, S.E. Thomas, G.J. Samuels, B.T. Vinyard, K.A. Holmes. "Antibiosis, mycoparasitism, and colonization success for endophytic Trichoderma isolates with biological control potential in Theobroma cacao", Biological Control, 2008

Publication

<1 %

55

Elfita Elfita, Muharni Muharni, Heni Yohandini, Fadhillah Fadhillah. " Antioxidant activity of endophytic fungi isolated from the stem bark of (L.) Jacq ", IOP Conference Series: Materials Science and Engineering, 2021

<1 %

56

Gurpreet Singh, Archana Katoch, Mod Razak, Surinder Kitchlu, Anindya Goswami, Meenu Katoch. "Bioactive and biocontrol potential of endophytic fungi associated with *Brugmansia aurea* Lagerh", *FEMS Microbiology Letters*, 2017

Publication

<1 %

57

JianQiu Sun, LiangDong Guo, Wei Zang, WenXiang Ping, DeFu Chi. "Diversity and ecological distribution of endophytic fungi associated with medicinal plants", *Science in China Series C: Life Sciences*, 2008

Publication

<1 %

58

Kandasamy Premalatha, Sivaraman Gokul, Amit Kumar, Priyanka Mishra, Pooja Mishra, Kaliamoorthy Ravikumar, Alok Kalra. "Molecular profiling of fungal assemblages in the healthy and infected roots of *Decalepis arayalpathra* (J. Joseph & V. Chandras) Venter, an endemic and endangered ethnomedicinal plant from Western Ghats, India", *Annals of Microbiology*, 2014

Publication

<1 %

59

cmm.mazums.ac.ir

Internet Source

<1 %

60

issuu.com

Internet Source

<1 %

61

repository.unair.ac.id

Internet Source

<1 %

62

www.scielo.br

Internet Source

<1 %

63

Fares Zeidán-Chuliá, José L. Rybarczyk-Filho, Mervi Gursoy, Eija Könönen et al. "

Bioinformatical and approaches to essential oil-induced matrix metalloproteinase inhibition ", *Pharmaceutical Biology*, 2012

Publication

<1 %

64

Juangjun Jumpathong, John Peberdy, Isao Fujii, Saisamorn Lumyong. "Chemical investigation of novel ascomycetes using PCR based screening approaches", *World Journal of Microbiology and Biotechnology*, 2011

Publication

<1 %

65

Lee Kee Chua, Chooi Ling Lim, Anna Pick Kiong Ling, Soi Moi Chye, Rhun Yian Koh. "Anticancer Potential of Syzygium Species: a Review", *Plant Foods for Human Nutrition*, 2018

Publication

<1 %

66

Tiago de Souza Leite, Andréia Cnossen-Fassoni, Olinto Liparini Pereira, Eduardo Seiti Gomide Mizubuti et al. "Novel and highly diverse fungal endophytes in soybean

<1 %

revealed by the consortium of two different techniques", Journal of Microbiology, 2013

Publication

67

W. Tian, Y.H. Bi, W. Zeng, W. Jiang, Y.H. Xue, G.X. Wang, S.P. Liu. "Diversity of endophytic fungi of *Myricaria laxiflora* grown under pre- and post-flooding conditions", Genetics and Molecular Research, 2015

Publication

68

Yu Li, Pachaiyappan Saravana Kumar, Quansheng Tan, Xuhui Tan, Mingyuan Yuan, Jianqiong Luo, Meijun He. "Diversity and chemical fingerprinting of endo-metabolomes from endophytes associated with *Ampelopsis grossedentata* (Hand.-Mazz.) W. T. Wang possessing antibacterial activity against multidrug resistant bacterial pathogens", Journal of Infection and Public Health, 2021

Publication

69

sciencetechindonesia.com

Internet Source

70

www.pjps.pk

Internet Source

71

Asiri Perera, So Ha Ton, Mohanambal Moorthy, Uma Devi Palanisamy. " The insulin-sensitising properties of the ellagitannin geraniin and its metabolites from rind in 3T3-

<1 %

<1 %

<1 %

<1 %

<1 %

L1 cells ", International Journal of Food Sciences and Nutrition, 2020

Publication

72

ELFITA, MUNAWAR, MUHARNI, MASTUR ADHY SUDRAJAT. "Identification of New Lactone Derivatives Isolated from Trichoderma sp., An Endophytic Fungus of Brotowali (*Tinaspora crispa*)", HAYATI Journal of Biosciences, 2014

Publication

73

Fábio de Azevedo Silva, Rhavena Graziela Liotti, Ana Paula de Araújo Boleti, Érica de Melo Reis et al. "Diversity of cultivable fungal endophytes in *Paullinia cupana* (Mart.) Ducke and bioactivity of their secondary metabolites", PLOS ONE, 2018

Publication

74

J. D. P. Bezerra, M. G. S. Santos, V. M. Svedese, D. M. M. Lima, M. J. S. Fernandes, L. M. Paiva, C. M. Souza-Motta. "Richness of endophytic fungi isolated from *Opuntia ficus-indica* Mill. (Cactaceae) and preliminary screening for enzyme production", World Journal of Microbiology and Biotechnology, 2012

Publication

75

Kanika Chowdhary, Nutan Kaushik. "Fungal Endophyte Diversity and Bioactivity in the

<1 %

<1 %

<1 %

<1 %

Indian Medicinal Plant *Ocimum sanctum*
Linn", PLOS ONE, 2015

Publication

76

Qinghua Zhang, Jing Zhang, Long Yang, Lei Zhang, Daohong Jiang, Weidong Chen, Guoqing Li. "Diversity and biocontrol potential of endophytic fungi in *Brassica napus*", *Biological Control*, 2014

Publication

<1 %

77

Sana Sheik, K. R. Chandrashekar, Sukesh Shirlal. "Endophytic fungi associated with *Psychotria flavida* Talbot., an endemic plant of Western Ghats (India) and their bioactive potential", *Oriental Pharmacy and Experimental Medicine*, 2018

Publication

<1 %

78

You-Kun Zheng, Xin-Guo Qiao, Cui-Ping Miao, Kai Liu, You-Wei Chen, Li-Hua Xu, Li-Xing Zhao. "Diversity, distribution and biotechnological potential of endophytic fungi", *Annals of Microbiology*, 2015

Publication

<1 %

79

journal.abu.edu.ng

Internet Source

<1 %

80

repositorio.uam.es

Internet Source

<1 %

81

scitepress.org

Internet Source

<1 %

82

www.mycosphere.org

Internet Source

<1 %

83

Elaissi, Ameer, Zyed Rouis, Samia Mabrouk, Karima Bel Haj Salah, Mahjoub Aouni, Mohamed Larbi Khouja, Farhat Farhat, Rachid Chemli, and Fethia Harzallah-Skhiri.

"Correlation Between Chemical Composition and Antibacterial Activity of Essential Oils from Fifteen Eucalyptus Species Growing in the Korbous and Jbel Abderrahman Arboreta (North East Tunisia)", *Molecules*, 2012.

Publication

<1 %

84

Hui Jin, Xiaoyan Yang, Dengxue Lu, Chunjie Li, Zhiqiang Yan, Xiuzhuang Li, Liming Zeng, Bo Qin. "Phylogenic diversity and tissue specificity of fungal endophytes associated with the pharmaceutical plant, *Stellera chamaejasme* L. revealed by a cultivation-independent approach", *Antonie van Leeuwenhoek*, 2015

Publication

<1 %

85

Thamilvaani Manaharan, David Appleton, Hwee Ming Cheng, Uma D. Palanisamy. "Flavonoids isolated from *Syzygium aqueum* leaf extract as potential antihyperglycaemic agents", *Food Chemistry*, 2012

<1 %

86 Kasahun Amare, Geleta Dugassa. "Plant tissue culture challenges in Ethiopia and alternative options for low-cost media", F1000Research, 2022 $<1\%$
Publication

87 Soil Biology, 2006. $<1\%$
Publication

88 T. K. Lim. "Edible Medicinal And Non Medicinal Plants", Springer Science and Business Media LLC, 2012 $<1\%$
Publication

89 ijpsr.com $<1\%$
Internet Source

Exclude quotes On

Exclude matches Off

Exclude bibliography Off