

A Phenolic Compound of Endophytic Fungi Isolated from Stem of *Syzygium aqueum* and its Diuretic Activity

by Hary Widjajanti

Submission date: 21-Sep-2022 12:10PM (UTC+0700)

Submission ID: 1905175138







File name: 26_A_PHENOLIC_COMPOUND_BIOINTERFACE_21_NOPEMBER_2021.pdf (637.85K)

Word count: 5835

Character count: 30374

47

A Phenolic Compound of Endophytic Fungi Isolated from Stem of *Syzygium aqueum* and its Diuretic Activity

Ummi H. Habisukan^{1,2}, Elfita^{3,*}, Hary Widjajanti⁴, Arum Setiawan⁴, Salni⁴, Rian Oktiansyah²

42

¹ Graduate School of Sciences, Faculty of Mathematics and Natural Sciences, University of Sriwijaya, Jl. Padang Selasa 524, Palembang 30139, South Sumatra, Indonesia; ummihirashabisukan@radenfatah.ac.id (U.H.H.);

² Universitas Islam Negeri Raden Fatah Palembang, South Sumatra, Indonesia; ummihirashabisukan@radenfatah.ac.id (H.H.); rianoktiansyah@radenfatah.ac.id (R.O.);

³ Department of Chemistry, Faculty of Mathematics and Natural Sciences, University of Sriwijaya, Jl. Palembang 4 abumulih, Indralaya, South Sumatra, 30662 Indonesia; elfita.elfita.69@gmail.com (E.)

⁴ Department of Biology, Faculty of Mathematics and Natural Sciences, University of Sriwijaya, Jl. Palembang Prabumulih, Indralaya, South Sumatra, 30662 Indonesia; hary_widjajanti@unsri.ac.id (H.W.); arum.setiawan@unsri.ac.id (A.S.); salnibasir@unsri.ac.id (S.);

* Correspondence: elfita.elfita.69@gmail.com (E.);

3

Scopus Author ID 55532411800

Received: 20.09.2021; Revised: 25.10.2021; Accepted: 28.10.2021; Published: 21.11.2021

Abstract: Guava (*Syzygium aqueum*), a typical Indonesian crop plant, has diuretic properties. Plants used as diuretic medicines can also be valuable sources for endophytic fungi containing diuretic drugs. The aims of this study were to isolate and identify endophytic fungi from *S. aqueum* stems, to isolate their secondary metabolites, and to conduct in vivo tests of the diuretic activity of an endophytic fungus extract on white male rats of the Wistar strain. Endophytic fungi were isolated by a direct plating method, and fungal isolates were identified molecularly. A phenolic compound was isolated by chromatography, and the chemical structure was identified spectroscopically. Doses of 100, 200, and 300 mg/kg BW were administered to white male rats divided into six groups (normal control, negative control, positive control, and the three treatment groups). Diuretic tests included urine volume; sodium, chloride, and calcium electrolyte levels; and pH. Phylogenetic analysis identified the endophytic fungal isolate as *Trichoderma ghanense* (isolation code SA1). The secondary metabolite isolated from the *T. ghanense* extract was a phenolic compound. Diuretic tests using the endophytic fungus extract showed the highest urine volume with a dose of 300 mg/kg BW. The diuretic test results showed an increase in urine volume and levels of sodium, potassium, and chloride ions in the urine. The extract of *T. ghanense* isolated from the stem of *S. aqueum* has the potential as initial therapy for hypertension because it contains phenolic secondary metabolites that show diuretic activity.

Keywords: diuretic; endophytic fungi; secondary metabolite; *Syzygium aqueum*.

© 2021 by the authors. This article is an open-access article distributed under the terms and conditions of the Creative Commons Attribution (CC BY) license (<https://creativecommons.org/licenses/by/4.0/>).

1. Introduction

Treating ailments such as hypertension, nephrotic syndrome, cirrhosis, and heart conditions includes diuretics to flush out excessive fluids within the body [1]. Diuretics cause an increase in the production and excretion of urine by the kidneys to promote the removal of toxins, excess water, and accumulated metabolic products, such as urea, from the body [2]. However, their use is sometimes accompanied by a loss of electrolytes, such as sodium, chloride, and potassium. Diuretics function to rid the body of excess fluid (edema) accumulated in tissues due to various disease states [3,4].

One plant with traditionally reported diuretic properties is guava (*Syzygium aqueum*), a typical crop plant from Indonesia. This plant has also been used in various parts of the world to treat various diseases, including hypertension, diabetes, diarrhea, asthma, inflammation, skin diseases, fever, digestive complaints, and high cholesterol [5], [6]. One study stated that most of the 87 secondary metabolites isolated from parts of the *Syzygium aqueum* plant were phenolics or phenolic derivatives [7]. People in South Sumatra, Indonesia, have used a decoction of the leaves and stems of *S. aqueum* as a diuretic medicine and to treat hypertension and diabetes [8].

The implantation of *S. aqueum* and other medicinal plants in Indonesia still faces many obstacles in terms of production, including the unprofessional implementation of medicinal plant implantation activities, the inability of farmers to maintain the quality and quality of medicinal plants, and the lack of attention from the medicinal plant industry regarding the results of scientific research in product development. One potential strategy for working around these obstacles is to use endophytic fungi, which are fast becoming important biotechnological tools [9,10]. Endophytic fungi are microscopic living organisms that live in plant tissues for a certain period by forming colonies without harming their hosts [11]. These endophytic fungi can produce metabolites similar to those produced by their host or different new compounds; therefore, this microbial group has huge biotechnological potential for new drug discovery [12–14]. Compounds produced by endophytic fungi from plants are often used as antioxidants, antidiabetics, cancer, and diseases caused by oxidative damage [15–17]. *Syzygium aqueum* contains compounds such as flavonoids, phenolics, tannins, terpenoids which are often used as antioxidants, anticancer, antiviral, anti-inflammatory [7]. The endophytic fungi are also believed to contain the same compounds and activities. From two of *Syzygium aqueum*, an antioxidant compound identified as 4-hydroxy-3-(4-hydroxyphenyl)-5-oxotetrahydrofuran-2-yl has been isolated [11]. The endophytic fungi associated with the plant *Syzygium aqueum* could represent ideal sources of diuretic drugs.

2. Materials and Methods

2.1. Chemical and drug.

Alcohol 70%, distilled water, sodium hypochlorite (NaOCl), chloramphenicol (Chloromex[®] A, Pabris, Indonesia), Potato Dextrose Agar (PDA) Merck, Potato Dextrose Broth (PDB) Merck, Silica gel 60 (70–230 mesh) Merck, thin layer chromatography (TLC), Silica gel 60 F254 Merck (Art.5554), methanol, acetone, ethyl acetate, chloroform, n-hexane (Merck), furosemide (Lasix[®], Aventis Indonesia Pharma, Indonesia), tween 80 1%.

75

2.2. Plant material.

The fresh stem of *Syzygium aqueum* was collected from Martapura, South Sumatra, Indonesia (4.17248 ° S, 104.20472° E). Plant identification was carried out at the Indonesian Institute of Sciences, Plant Conservation Center, Purwodadi Botanical Gardens, Certificate of Identification No.: B-301/III /KS.01.03/1/2021.

2.3. Isolation and identification of endophytic fungi.

Syzygium aqueum stems were washed under running water to remove dirt. The stem surface was sterilized by immersion in 70% ethanol for 2 min, followed by 4% NaOCl for 1

min and two rinses in sterile distilled water. The plant stems were dried on sterile filter paper and then cut aseptically into 1 × 1 cm pieces which were then planted directly in a Petri dish containing PDA plus chloramphenicol and incubated at 25 °C for 2–5 days. The tip of the hyphae growing out of each segment was transferred to a Petri dish containing new PDA and then re-incubated for approximately 14 days. Pure cultures were obtained after several subcultures and grown on PDA plates without antibiotics [18].

Identification of endophytic fungi was carried out based on partial genetic analysis on the Internal Transcribed Spacer (ITS) of fungal ribosomal DNA. PCR amplification using ITS Primer 4:5' - TCC TCC GCT TAT TGA TAT GC – 3' and ITS Primer 5:5' -GGA AGT AAA AGT CGT AAC AAG G –3'. The purified polymerase chain reaction (PCR) product was then precipitated, followed by cyclic sequencing. The result was purified again by the ethanol purification method. Sequencing was performed using an automated DNA sequencer. Sequence data were trimmed and assembled, followed by alignment BLAST using genomic data at the NCBI (National Center for Biotechnology Information) for analysis of homology/similarity and creation of a phylogenetic tree [19].

2.4. Cultivation and extraction.

Endophytic fungal suspensions were inoculated in 300 mL potato dextrose broth (PDB), placed in 1 L bottles (9 L in 30 bottles), and incubated for 4 weeks at room temperature under static conditions. The endophytic fungal mycelium in each bottle was removed using tweezers, and the medium was filtered through filter paper. The filtrate was extracted with ethyl acetate at a 1:1 ratio. The filtrate and solvent were separated in a separating funnel to obtain an ethyl acetate extract, which was concentrated using a rotary evaporator [11].

2.5. Isolation and identification of secondary metabolites.

The endophytic fungal extracts from *Syzygium aqueum* were separated by column chromatography (stationary phase: silica gel) and eluted with a graded eluent (n-hexane: ethyl acetate: methanol). The eluate droplets were collected in a 10 mL bottle and analyzed by TLC to group the bottles into subfractions. Subfractions identified as having secondary metabolites were further purified by column chromatography until pure compounds were obtained. The structures were identified by 1D and 2D NMR [20].

2.6. Design of the test animal experiments animals.

The test animals were male Wistar rats (weight 150–250 g) obtained from Abduh Tikus Palembang, South Sumatra, Indonesia. All animals were housed at 25 ± 2°C and a 12 h light/dark cycle in polypropylene cages with clean rice husks as bedding. The test animals were acclimatized in the laboratory for 7 days. All animals were provided standard rodent food and drinking water ad libitum. The animal experiment protocol was approved by the Research Ethics Committee of Ahmad Dahlan University, Yogyakarta, Indonesia, with approval number: 022107033 dated August 27, 2021.

Table 1. Test animal treatment group.

| No | Category | Treatment |
|----|------------------|---|
| 1 | Normal control | Aquadest |
| 2 | Negative control | Aquadest + NaCl 4,5% + tween 80 suspension 1% |
| 3 | Positive control | Aquadest + NaCl 4,5% + furosemide suspension |

| No | Category | Treatment |
|----|-------------|---|
| 4 | Treatment 1 | Aquadest + NaCl 4,5% + mushroom extract suspension dose 100 mg/kgBB |
| 5 | Treatment 2 | Aquadest + NaCl 4,5% + mushroom extract suspension dose 200 mg/kgBB |
| 6 | Treatment 3 | Aquadest + NaCl 4,5% + mushroom extract suspension dose 300 mg/kgBB |

2.7. Preparation for assays.

Suspensions of endophytic fungal extract at doses of 100 mg/kg BW, 200 mg/kg BW, and 300 mg/kg BW were prepared by grinding the endophytic fungal extract from *Syzygium aqueum* in a mortar according to the dose calculation (1.5 g in 50 mL for the highest dose of 300 mg/kg BW), then dissolving in phosphate-buffered saline containing tween 80 1%. The next dose was prepared by diluting the mother liquor (the highest dose).

2.8. Test of diuretic effect.

The test animal was divided into 6 treatment groups, consisting of 4 rats (Table 1). Before testing, the animals fasted for 18 h. Samples were given using a gastric probe. After treatment, the test animals were put into metabolic cages to collect their urine. Urine volume was measured after administration of the extract at 1, 2, 3, 4, 5, and 6 h, and cumulative urine volume was determined after 24 h. Drinking water at 1 to 6 h was controlled by giving 2.5 mL of 4.5 ml NaCl every 3 h orally to each test animal, except for the normal group, which was given 2.5 mL distilled water. During the treatment, the test animals were not fed. The percentage potency of the diuretics was determined using the following equation [21]:

$$\% \text{ Diuretic Potency} = \frac{\text{extract treatment volume} - \text{negative treatment volume}}{\text{Positive treatment volume} - \text{negative treatment volume}} \times 100\%$$

Urinary Na⁺, K⁺, and Cl⁻ levels were measured by atomic absorption spectroscopy (AAS) at wavelengths of 766.5 nm and 589.0 nm [22]. The pH was determined with a pH meter. The urine was fresh or preserved so that ammonia was not produced by urea breakdown.

Data Analysis

The results of the study were expressed as mean ± standard error of the mean (SEM). The data were statistically analyzed by one-way analysis of variance (ANOVA) followed by a Tukey post hoc multiple comparison test. P-values less than 0.05 were considered statistically significant.

3. Results and Discussion

3.1. Isolation and identification of endophytic fungi.

The endophytic fungi isolated from the stems of *Syzygium aqueum* were given the isolation code SA1 (Figure 1). The results of the molecular identification of the 649 bp sequence of the SA1 endophytic fungus were as follows: GATCCTTCCGTAGGTGAACCTGCGGAGGGATCATTACCGAGTTTACAACCTCCCAAACCCAATGTGAACGTTACCAATCTGTTGCCTCGGCGGGATTCTCTGCCCCGGGC GCGTCGCAGCCCCGATCCCATGGCGCCCCGCCGAGGACCAACCAAACCTCTTTTT TCTCTCCGTCGCGGCTTCCGTCGCGGCTCTGTTTTAACCTTTGCTCTGAGCCTTTCT CGGCGACCCTAGCGGGCGTCTCGAAAATGAATCAAACTTTCAACAACGGATCT CTTGGTTCTGGCATCGATGAAGAACGCAGCGAAATGCGATAAGTAATGTGAATT

GCAGAATTCAGTGAATCATCGAATCTTTGAACGCACATTGCGCCCGCCAGTATTC
TGCGGGCATGCTGTCCGAGCGTCATTTCAACCCTCGAACCCCTCCGGGGTCC
GGCGTTGGGGATCGGCACGCCCTCACACGGGTGCCGCCCCGAAATCCAGTGGC
GGTCTCGCCGACGCTCTCCTGCGCAGTAGTTTGCACACTCGCACCCGGGAGCGCG
GCGCGGCCACAGCCGTAAACACCCCAAACCTTCTGAAAATGTTGACCTCGGATCA
GGTAGGAATACCCGCTGAACCTTAAGCATATCAATAAGCCGGAGGAA.

The SA1 strain was given the accession number SUB10223461 SA1 MZ855361. The SA1 endophytic fungus had the highest homology with *Trichoderma ghanense* MF078652. The phylogenetic analysis followed the described method shown in Figure 2.




Figure 1. Endophytic fungi isolation results from *Syzygium aqueum* stem. (a) SA1 isolate in front; (b) SA1 isolate in back.




Figure 2. Phylogenetic tree from endophytic fungi SA1.

3.2. Isolation of secondary metabolic compounds from the endophytic fungus.

The separation of 2 g of the concentrated SA1 endophytic fungal extract (silica gel: 70–230 mesh; 20 g) generated subfraction SA1.7 (128.7 mg), giving a purple stain on TLC analysis. The compound was purified using column chromatography (silica gel 70–230 mesh;

10 g) and eluted with hexane: ethyl acetate (7:3–4:6) to yield four subfractions (SA1.7.1–SA1.7.4). The SA1.7.3 fraction (55 mg) was further purified by column chromatography until compound 1 (435 µg) was obtained.

The $^1\text{H-NMR}$ spectrum of compound 1 (Figure 3 (A)) revealed the presence of seven proton signals; four for the sp^3 proton, including a methyl signal at δ_{H} 1.25 ppm (3H, d, $J=1.5$ Hz) and a methine proton signal at δ_{H} 4.17 ppm (1H, m), and three oxygenated methine signals at δ_{H} 4.02 (2H, m), 5.34 (1H, d, $J=2.5$ Hz), and 5.79 ppm (1H, s). The spectrum is in the aromatic chemical shift region, at δ_{H} 7.58 and 8.22 ppm. The spectrum of each signal shows a doublet with the same plot constant $J = 8.0$ Hz, indicating the integration of two protons. Therefore, compound 1 is a para-substituted aromatic compound, so it has two pairs of equivalent protons. Analysis of the $^1\text{H-NMR}$ spectrum revealed compound 1 as a para-substituted aromatic compound with 12 protons attached to a carbon atom. The solvent used in this measurement is CDCl_3 , so the protons bound to the heteroatoms do not appear in the spectrum.




Figure 3. (a) $^1\text{H-NMR}$ spectrum; (b) $^{13}\text{C-NMR}$ of compound 1.

The $^{13}\text{C-NMR}$ spectrum of compound 1 (Figure 3(B)) showed the presence of nine signals: five sp^3 carbon signals, including three signals in the oxygenated carbon area at δ_{C} 64.1, 66.3, and 73.6 ppm. Four other signals appear in the aromatic region as two high-intensity signals, indicating that compound 1 has two pairs of equivalent aromatic carbons (δ_{C} 126.8 and 123.8 ppm), and as two low-intensity signals, indicating the presence of quaternary carbon, one of which is an oxyaryl carbon in the low field (δ_{C} 164.6 ppm). Analyses of the proton and carbon NMR spectra were confirmed by the data on the heteronuclear multiple quantum coherence (HMQC) spectrum shown in Figure 4 and Table 2, namely the $^1\text{H-}^{13}\text{C}$ correlation through one bond. The HMQC spectrum showed seven correlations consisting of two correlations on the aromatic ring, three correlations on oxygenated $^1\text{H-}^{13}\text{C}$, one proton correlation methine, and one correlation of proton methyl.

The heteronuclear multiple bond correlation (HMBC) spectrum (Figure 5) shows the $^1\text{H-}^{13}\text{C}$ correlation through two or three bonds. The aromatic proton signal at δ_{H} 8.22 ppm indicates the existence of two correlations, one to carbon δ_{C} 123.9, which is the carbon equivalent, and the other to δ_{C} 147.7, which is the aromatic quaternary carbon. Another aromatic proton at δ_{H} 7.58 ppm has three correlations, each with carbon at δ_{C} 126.9 ppm, which is the carbon

equivalent, δ_C 147.7 and δ_C 73.6 ppm, which is oxygenated carbon directly bonded to the aromatic ring. Oxygenated methine protons at δ_H 5.34 ppm also have three correlations to carbons at δ_C 66.3, 147.7, and 126.9 ppm.




Figure 4. The HMQC spectrum and placement of the ^1H and ^{13}C shift in compound 1.

55

Table 2. The NMR data of Compound 1.

| No. C | δ_C ppm | δ_H ppm (Σ H, multiplicity, J (Hz)) | HMBC | COSY |
|-------|----------------|--|--------------------|------|
| 1 | 73.6 | 5.34 (1H, d, J=2.5 Hz) | 66.3; 147.7; 126.9 | 4.17 |
| 2 | 55.7 | 4.17 (1H, m) | | 5.34 |
| 3 | 66.3 | 5.0 (1H, s) | | |
| 4 | 29.9 | 1.25 (3H, d, J=1.5 Hz) | 55.7 | |
| 5 | 64.1 | 4.02 (2H, m) | 73.6 | |
| 1' | 147.7 | 1.5 | | |
| 2' | 126.9 | 7.58 (1.5 d, J=8.0 Hz) | 126.9; 147.7; 73.6 | 8.22 |
| 3' | 123.9 | 8.22 (1H, d, J=8.0 Hz) | 123.9; 147.7 | 7.58 |
| 4' | 164.6 | 1.5 | | |
| 5' | 123.9 | 8.22 (1.5 d, J=8.0 Hz) | 123.9; 147.7 | 7.58 |
| 6' | 126.9 | 7.58 (1H, d, J=8.0 Hz) | 126.9; 147.7; 73.6 | 8.22 |

The correlation indicates that the oxygenated methine group at δ_H 5.34 ppm is directly bound to the aromatic ring and is para-substituted with a hydroxyl group. The proton hydroxyl signal does not appear on the spectrum because the pure compound is measured with CDCl_3 as the solvent. The 1D and 2D NMR spectral data for compound 1 are shown in Table 2.

The COSY spectrum in Figure 6 shows the ^1H - ^1H correlation through three bonds. Two types of correlation are evident: the correlation of the aromatic proton at δ_H 8.22 ppm with the aromatic proton at δ_H 7.57 ppm and the proton sp^3 correlation of the substituent group, namely the oxygenated proton of methine at δ_H 5.34 ppm with the proton of methine at 4.17 ppm.




Figure 5. The HMBC spectrum and the correlation ^1H - ^{13}C through two or three bonds in compound 1.

This indicates that the two aromatic equivalent protons are in the ortho position. This strengthens the proposed structure of compound 1 as a benzene ring directly bonded to the oxygenated methine carbon at the para position with a hydroxyl group.




Figure 6. COSY spectrum and the correlation ^1H - ^1H in compound 1.

The spectrum analyses by ^1H -NMR, ^{13}C -NMR, HMQC, HMBC, and COSY can explain that compound 1 has a benzene ring that is substituted at the para position with a hydroxyl group and an oxygenated methine group that is part of the substituent group, namely 2- (hydroxymethyl)butane-1,3-diol. The proton at H 5.78 ppm appears as a singlet, probably

because the neighboring proton, which is three bonds apart, is in the farthest geometric position; therefore, cleavage and/or other factors, such as inappropriate resolution, do not occur. Thus, the proposed chemical structure of compound 1 is 2-(hydroxymethyl)-1-(4-hydroxyphenyl)butane-1,3-diol, as shown in Figure 7.




Figure 7. Structure compound of 3-(hydroxymethyl)-1-(4-hydroxyphenyl)butane-1,3-diol.

3.3. Test of diuretic effect.

The total urine volume produced was larger in the treatment group than in the normal and negative groups. This indicates a diuretic effect of the extract on urine excretion. The highest urine volume was produced by the positive control group; the data for the treatment group can be seen in Table 3.

Table 3. Total Urine Volume and Diuretic Potency.

| Group | 1 hour | 2 hour | 3 hour | 4 hour | 5 hour | 6 hour | 24 hour | % Diuretic Potency |
|-------------|------------|------------|------------|------------|------------|-------------------------|-------------|--------------------|
| Normal | 1.42±0.572 | 2.42±0.453 | 2.85±0.433 | 3.30±0.248 | 3.42±0.319 | 3.57±0.278 ^a | 6.37±0.327 | 0 |
| Negative | 1.17±0.411 | 2.02±0.652 | 2.32±0.850 | 3.77±1.048 | 4.37±0.851 | 4.57±0.797 ^a | 7.50±0.659 | 0 |
| Treatment 1 | 0.62±0.252 | 1.52±0.564 | 2.20±0.696 | 3.87±1.295 | 4.75±1.420 | 5.17±1.546 ^a | 8.27±1.981 | 20 |
| Treatment 2 | 0.37±0.217 | 1.20±0.535 | 1.75±0.590 | 3.90±0.994 | 4.77±1.220 | 5.67±0.900 ^b | 8.67±1.697 | 21.27 |
| Treatment 3 | 1.45±0.132 | 2.67±0.406 | 2.92±0.314 | 6.70±0.817 | 7.05±0.790 | 7.70±0.994 ^c | 11.00±1.511 | 93.95 |
| Positive | 1.46±0.221 | 2.65±0.457 | 4.35±0.800 | 7.32±1.145 | 8.07±1.416 | 8.42±1.353 ^d | 11.22±1.915 | 100 |

Each value represents the mean ± S.E.M (Standard Error of Mean); n=4. Positive: furosemide 20 mg/kgBW, Treatment 1 (100 mg/kgBW), Treatment 2 (200mg/kgBW), Treatment 3 (300 mg/kgBW)

Treatment with the extract increased urine volume or diuresis as a ratio between the dose and diuretic potency (R=0.78). The increase in urine volume occurs due to the inhibition of water reabsorption in the kidneys [23]. The increase in urine volume in the treatment group compared to the negative and normal groups occurred at 4–6 hours, due to the working time of furosemide at 4–6 hours, while the treatment groups 1–3 also showed a similar response, indicating that the working time of the extract was the same as the working time of furosemide [24] The treatment dose that had the highest diuretic potential was treatment 3, with a diuretic potency of 93.95% (P<0.003), whereas it was considered 0% (P<0.00) for the control and negative groups because those were the baseline references for the diuretic effect of the extract.

Previous studies have revealed that phenolics can inhibit the carbonic enzyme anhydrase in the renal tubules. Phenolics increase diuresis because they inhibit the angiotensin-converting enzyme (ACE) [25,26]. Cumulatively, the results for the urine volumes were greater for each treatment group than for the negative and normal control groups. The results of one-way ANOVA for the total urine volume (p = 0.003) indicate a significant difference between the test groups. The homogeneity test resulted in a Levene statistic of 2.257 with p = 0.523, showing that the test data have the same variance or are homogeneous.

The pH value of urine (Table 4) is determined by acid-base regulation in the kidneys. If a certain amount of HCO_3^- is continuously filtered into the renal tubules and excreted in the urine, the urine will be alkaline, as was observed in this study, where the pH value was pH 9–10 [27]. By contrast, if H^+ ions were continuously filtered into the kidney tubules and excreted into the urine, the urine would be acidic [28]. The kidneys have several mechanisms for regulating the concentration of H^+ ions, including secretion of H^+ ions into the tubules, reabsorption of HCO_3^- ions, and production of new HCO_3^- ions. All these mechanisms reduce and neutralize excess H^+ ions in the body [29,30].

Table 4. pH measurement.

| Treatment category | Urine pH |
|--------------------|---------------|
| Normal | 10,00 ± 0,000 |
| Negative | 10,00 ± 0,000 |
| Treatment 1 | 09,50 ± 0,288 |
| Treatment 2 | 09,00 ± 0,000 |
| Treatment 3 | 09,50 ± 0,288 |
| Positive | 10,00 ± 0,000 |

Each value represents the mean ± S.E.M (Standard Error of Mean); n=4. Positive: furosemide 20 mg/kgBW, Treatment 1 (100 mg/kgBW), Treatment 2 (200mg/kgBW), Treatment 3 (300 mg/kgBW)

increase in diuresis caused by the endophytic fungal from *S. aqueum* was also evident in the excretion of electrolytes. The extract significantly increased the excretion of urinary electrolytes, such as Na^+ , K^+ , and Cl^- (Table 5). This increase in electrolyte excretion is thought to be due to the endophytic fungi extract acting through a loop diuretic mechanism, thereby increasing the urine flow rate and in turn causing a decrease in the absorption of Na^+ , K^+ , and Cl^- in the loop of Henle in the kidney [31,32]. This will increase the amount of water and salt excreted in the urine [33]. However, in the present study, the amount of K^+ excreted was not as large as the amounts of Na^+ and Cl^- . This condition results from the antioxidant activity of the endophytic fungal extract, which reduces the side effects of loop diuretics so that the body can still absorb K^+ [34,35]. The saluretic activity of the three test groups was above 100 percent, compared to the positive control using furosemide. These results indicate that the extract has saluretic potential.

Table 5. Measurement of Na^+ , K^+ , and Cl^- levels.

| Treatment Category | Na^+ level (ppm) | K^+ level (ppm) | Cl^- level (ppm) | Saturetic level of Na (%) | Saturetic level of Cl (%) | Saturetic level of K (%) |
|--------------------|---------------------------|--------------------------|---------------------------|---------------------------|---------------------------|--------------------------|
| Normal | 200±1,154 | 141,5±3,175 | 141,5±3,175 | 0 | 0 | 0 |
| Negative | 247,5±8,948 | 330±16,743 | 386±73,323 | 0 | 0 | 0 |
| Treatment 1 | 363±16,454 | 360,5±47,342 | 360,5±47,342 | 417,9 | 161,6 | 86,9 |
| Treatment 2 | 272,5±7,794 | 338±31,754 | 338±31,754 | 184,6 | 145 | 77,97 |
| Treatment 3 | 363±15,011 | 360,5±4,330 | 360,5±4,330 | 417,9 | 161,2 | 86,9 |
| Positive | 239±3,464 | 393,5±14,145 | 277±81,406 | 100 | 100 | 100 |

Each value represents the mean ± S.E.M (Standard Error of Mean); n=4. Positive: furosemide 20 mg/kgBW, Treatment 1 (100 mg/kgBW), Treatment 2 (200mg/kgBW), Treatment 3 (300 mg/kgBW)

4. Conclusions

The endophytic fungi extract from *Syzygium aqueum* contains a phenolic compound 2-(hydroxymethyl)-1-(4-hydroxyphenyl) butane-1,3-diol that has diuretic activity and increases the volume of excreted urine and its ion concentration.

Funding

The research/publication of this article was funded by the DIPA of the Public Service Agency of Universitas Sriwijaya 2021. SP DIPA -023.17.2.677515/2021, on November 23, 2020. In accordance with the Rector's Decree Number: 0014/ UN9/ SK.LP2M.PT/2021, on May 25, 2021.

Acknowledgments

The authors wish to respect the facility's support of the Organic Chemistry and Microbiology Laboratory, Faculty of Mathematics and Natural Sciences, University of Sriwijaya also Palembang Health Laboratory Center, Ministry of Health of the Republic of Indonesia.

Conflicts of Interest

The authors declare no conflict of interest.

References

- Jayanthi, M.K.; Aswathi, K.; Krishna, K.L.; Ramu, R. Evaluation of antioxidant and diuretic activities of *Portia ternatea* leaf extracts in *Wistar albino* rats. *J Appl Pharm Sci* **2021**, *11*, 152-157. <https://dx.doi.org/10.7324/JAPS.2021.110118>.
- Trujillo, H.; Fontán, F.C.; Caro, J.; Morales, E.; Praga, M. The forgotten antiproteinuric properties of diuretic. *Am J Nephrol* **2021**, *2021*, 435-449. <http://dx.doi.org/10.1159/000517020>.
- Divya B.; Fakrudin, B.; Devappa, V. Expression profiling of resistance gene analogs from french bean (*Phaseolus vulgaris* L.) for the manifestation of rust (*Uromyces phaseoli* L.). *Int J Curr Microbiol Appl Sci* **2021**, *9*, 1760-1773. <http://dx.doi.org/10.20546/ijcmas.2019.803.206>.
- Shah, N.; Madanieh, R.; Alkan, M.; Dogar, M.U.; Kosmas, C.E.; Vittorio, T.J. A perspective on diuretic resistance in chronic congestive heart failure. *Ther Adv Cardiovasc Dis* **2017**, *11*, 271-278. <https://dx.doi.org/10.1177/1753944717718717>.
- Cock, I.E.; Cheesman, M. Chapter 23 - The Potential of Plants of the Genus *Syzygium* (Myrtaceae) for the Prevention and Treatment of Arthritic and Autoimmune Diseases. In: *Bioactive food as dietary interventions for arthritis and related inflammatory diseases*. 2nd ed.; Academic Press: Elsevier Inc., **2019**; pp. 401-424. <https://dx.doi.org/10.1016/B978-0-12-813820-5.00023-4>.
- Itam, A.; Wati, M.S.; Agustini, V.; Sabri, N.; Jumanah, R.A.; Efdi, M. Comparative study of phytochemical, antioxidant, and cytotoxic activities and phenolic content of *Syzygium aqueum* (burm. f. *alston* f.) extracts growing in Sumatra Indonesia. *Sci World J* **2021**, *2021*, 1-9. <https://doi.org/10.1155/2021/5537597>.
- Sobeh, M. *Syzygium aqueum*: a polyphenol-rich leaf extract exhibits antioxidant, hepatoprotective, pain-killing and anti-inflammatory activities in animal models. *Front Pharmacol* **2018**, *9*, 1-14. <https://dx.doi.org/10.3389/fphar.2018.00566>.
- Habisukan, U.H.; Elfita; Widjajanti, H.; Setiawan, A.; Kurniawati, A.R. Antioxidant and antimicrobial activity of endophytic fungi isolated from *Syzygium aqueum* leaves. *J Phys Conf Ser* **2021**, *1751*, 1-8.
- Park, C.H. In vitro antioxidant and antimicrobial properties of flower, leaf, and stem extracts of Korean mint. *Antioxidants* **2019**, *8*, 1-12. <http://dx.doi.org/10.3390/antiox8030075>.
- Wei, L.; Yang, M.; Huang, L.; Li, J.L. Antibacterial and antioxidant flavonoid derivatives from the fruits of *Staplexis japonica*. *Food Chem* **2019**, *289*, 308-312. <http://dx.doi.org/10.1016/j.foodchem.2019.03.070>.
- Habisukan, U.H.; Elfita; Widjajanti, H.; Setiawan, A. Chemical characterization of secondary metabolite from the endophytic fungus *Trichoderma reesei* isolated from the twig of *Syzygium aqueum*. *Sci Technol Indones* **2021**, *6*, 137-143. <http://dx.doi.org/10.26554/sti.2021.6.3.137-143>.
- Cruz, J.S.; da Silva, C.A.; Samerski, L. Natural products from endophytic fungi associated with Rubiaceae species. *J. Fungi* **2020**, *6*, 1-26. <http://dx.doi.org/10.3390/jof6030128>.
- El-Hawary, S.S.; Moawad, A.S.; Bahr, H.S.; Abdelmohsen, U.R.; Mohammed, R. Natural product diversity from the endophytic fungi of the genus *Aspergillus*. *RSC Adv* **2020**, *10*, 22058-22079. <https://dx.doi.org/10.1039/d0ra04290k>.
- Ababutain, I.M.; Aldosary, S.K.; Aljuraifani, A.A.; Alghamdi, A.I.; Alabdall, A.H.; Al-Khaldi, E.M.; Aldakeel, S.A.; Almandil, N.B.; Azeez, S.A.; Borgio, J.F. Identification and antibacterial characterization of endophytic fungi from *Artemisia sieberi*. *Int J Microbiol* **2021**, *2021*, 1-11. <https://doi.org/10.1155/2021/6651020>.

15. Rai, N. Plant associated fungal endophytes as a source of natural bioactive compounds. *Mycology* **2021**, *12*, 111–159, <http://dx.doi.org/10.1080/21501203.2020.1870579>.
16. Singh, A.; Singh, D.K.; Kharwar, R.N.; White, J.F.; Gond, S.K. Fungal endophytes as efficient sources of plant-derived bioactive compounds and their prospective applications in natural product drug discovery: Insights, avenues, and challenges. *Microorganisms* **2021**, *9*, 1–42, <https://doi.org/10.3390/microorganisms9010197>.
17. Adeleke, B.S.; Babalola, O.O. Pharmacological potential of fungal endophytes associated with medicinal plants: A review. *J. Fungi* **2021**, *7*, 1–16. <http://doi.org/10.3390/jof7020147>.
18. Habiskan, U.H.; Elfita; Widjajanti, H.; Setiawan, A.; Kurniawati, A.R. Diversity of endophytic fungi in *Pygium aqueum*. *Biodiversitas* **2021**, *22*, 1129–1137, <http://dx.doi.org/10.13057/biodiv/d220307>.
19. Yohandini, H.; Julinar; Muharni. Isolation and phylogenetic analysis of thermophile community within tanjung sakti hot spring, South Sumatera, Indonesia. *HAYATI J Biosci* **2015**, *22*, 143–148, <http://dx.doi.org/10.1016/j.hjb.2015.10.006>.
20. Muharni; Fitriya; Ruliza, M.O.; Susanti, D.A.; Elfita. Di-(2-ethylhexyl)phthalate and pyranon derived from endophytic fungi *Penicillium* sp the leave of kunyit putih (*Curcuma zedoaria*). *Indones J Chem* **2014**, *14*, 292–296.
21. Jadhav, N.; Patil, C.R.; Chaudhari, K.B.; Wagh, J.P.; Surana, S.J.; Jadhav, R.B. Diuretic and natriuretic activity of two mistletoe species in rats. *Phcog Res* **2010**, *2*, 50–57, <http://dx.doi.org/10.4103/0974-2700.60576>.
22. Korespondesi, P.; Maulidar, N.P.; Haro, G. Penetapan kadar kalium, kalsium, natrium dan magnesium dalam bunga nangka (*Artocarpus heterophyllus* lam.) jantan secara spektrofotometri serapan atom. *Media Farm. J. Ilmu Farm* **2018**, *15*, 81–87, <http://dx.doi.org/10.12928/mf.v15i2.12659>.
23. Tufer, S.; Engidawork, E.; Ayele, A.G.; Bashea, C. Evaluation of the diuretic activity of aqueous and 80% methanol extracts of *Conyza macrostachyus* (euphorbiaceae) leaves in saline-loaded rats. *J Exp Pharmacol* **2021**, *2021*, 213–221, <http://dx.doi.org/10.2147/JEP.S294062>.
24. Mahmoodpoor, A.; Zahedi, S.; Pourakbar, A.; Hamishehkar, H.; Shadvar, K.; Asgharian, P.; Shahabi, F.; Hamishehkar, H. Efficacy of furosemide-albumin compared with furosemide in critically ill hypoalbuminemia patients admitted to intensive care unit: a prospective randomized clinical trial. *J Pharm* **2020**, *1*, 1–7, <http://dx.doi.org/10.1007/s40199-020-00339-8>.
25. Huyut, Z.; Beydemir, Ş.; Gülçin, I. Antioxidant and antiradical properties of selected flavonoids and phenolic compounds. *Biochem Res Int* **2017**, *2017*, 1–10, <http://dx.doi.org/10.1155/2017/7616791>.
26. Madyastuti, R.; Wientarsih, I.; Widodo, S.; Purwaningsih, E.H.; Harlina, E. The diuretic activity and urin mineral analysis of cat whisker's extract (*Orthosiphon stamuneus* benth) effect in male rats. *Acta Vet Indones* **2020**, *8*, 16–23, <https://doi.org/10.33263/BRIAC125.70407051>.
27. Ramadhian, M.R.; Pahmi, K. Aktivitas diuresis *Leucaena leucocephala* l. pada mencit jantan (*Mus musculus*). *Journal Syifa Sciences and Clinical Research* **2021**, *3*, 19–28, <https://doi.org/10.37311/jsscr.v3i1.9894>.
28. Oster, D.C. Diuretic therapy in congestive heart failure. *Congest Hear Fail* **2000**, *6*, 197–201, <https://doi.org/10.1080/00015385.2021.1878423>.
29. Nalwaya, N.; Jarald, E.; Asghar, S.; Ahmad, S. Diuretic activity of a herbal product UNEX. *Int J Green Pharm* **2009**, *3*, 224–226.
30. Ding, D. Ototoxic effects and mechanisms of loop diuretics. *J Otol* **2016**, *11*, 145–156, <http://dx.doi.org/10.1080/10801203.2016.10.001>.
31. Bolda Mariano, L.N. Prolonged diuretic and renoprotective effects of a xanthone obtained from *Garcinia achachairu* rusby in normotensive and hypertensive rats. *Evidence-based Complement Altern Med* **2021**, *2021*, 1–9, <https://doi.org/10.1155/2021/5510053>.
32. Schlickmann, F. Gallic acid, a phenolic compound isolated from *Mimosa bimucronata* (DC.) Kuntze leaves, induces diuresis and saluresis in rats. *Naunyn Schmiedeberg's Arch Pharmacol* **2018**, *391*, 649–655, <http://dx.doi.org/10.1007/s00210-018-1502-8>.
33. Sarker, M. Diuretic and laxative activities of *Kandelia candel* and *Brownlowia tera* in experimental mice. *J Med Plants Stud* **2021**, *9*, 59–61, <http://dx.doi.org/10.22271/plants.2021.v9.i2a.1267>.
34. Meharie, B.G.; Tunta, T.A. Evaluation of diuretic activity and phytochemical contents of aqueous extract of the shoot apex of *Podocarpus falcatus*. *J Exp Pharmacol* **2020**, *12*, 629–641, <https://doi.org/10.2147/JEP.S287277>.
35. Jackson, A.M.; Dewan, P.; Anand, I.S.; Bělohávek, J.; Bengtsson, O.; de Boer, R.A.; Böhm, M.; Boulton, D.W.; Chopra, V.K.; DeMets, D.L.; Docherty, K.F.; Dukát, A.; Greasley, P.J.; Howlett, J.G.; Inzucchi, S.E.; Katova, T.; Køber, L.; Kosiborod, M.N.; Langkilde, A.M.; Lindholm, D.; Ljungman, C.E.A.; Martinez, F.A.; O'Meara, E.; Sabatine, M.S.; Sjöstrand, M.; Solomon, S.D.; Tereshchenko, S.; Verma, S.; Jhund, P.S.; McMurray, J.J.V. Dapa-gln and diuretic use in patients with heart failure and reduced ejection fraction in dapa-hf. *J Am Heart Assoc* **2020**, *2020*, 1040–1054, <http://doi.org/10.1161/CIRCULATIONAHA.120.047077>.

A Phenolic Compound of Endophytic Fungi Isolated from Stem of Syzygium aqueum and its Diuretic Activity

ORIGINALITY REPORT

35%
SIMILARITY INDEX

30%
INTERNET SOURCES

27%
PUBLICATIONS

16%
STUDENT PAPERS

PRIMARY SOURCES

1 Submitted to University of Strathclyde
Student Paper 1%

2 media.neliti.com
Internet Source 1%

3 www.researchgate.net
Internet Source 1%

4 www.jcps.ac.cn
Internet Source 1%

5 ouci.dntb.gov.ua
Internet Source 1%

6 www.japsonline.com
Internet Source 1%

7 www.sciencetechindonesia.com
Internet Source 1%

8 scholar.unand.ac.id
Internet Source 1%

9 Dalia H. Abdelkader, Walaa A. Negm, Engy Elekhawy, Duaa Eliwa, Basmah N. Aldosari, 1%

Alanood S. Almurshedi. "Zinc Oxide Nanoparticles as Potential Delivery Carrier: Green Synthesis by Aspergillus niger Endophytic Fungus, Characterization, and In Vitro/In Vivo Antibacterial Activity", Pharmaceuticals, 2022

Publication

10

Soghra Bagheri, Ali A. Saboury. "Hypothesis of using albumin to improve drug efficacy in cancers accompanied by hypoalbuminemia", Xenobiotica, 2021

Publication

1 %

11

encyclopedia.pub

Internet Source

1 %

12

A. Sureda, M. Monserrat-Mesquida, S. Pinya, P. Ferriol, S. Tejada. "Hypotensive effects of the triterpene oleanolic acid for cardiovascular prevention", Current Molecular Pharmacology, 2020

Publication

1 %

13

www.ijcmas.com

Internet Source

1 %

14

research-repository.griffith.edu.au

Internet Source

1 %

15

Vanessa Chapla, Maria Zeraik, Ioanis Leptokarydis, Geraldo Silva et al. "Antifungal Compounds Produced by Colletotrichum

1 %

gloeosporioides, an Endophytic Fungus from
Michelia champaca", Molecules, 2014

Publication

16

techscience.com

Internet Source

1 %

17

www.dovepress.com

Internet Source

1 %

18

Xiaoyu Li, Xianglei Wang, Menglu Zhao, He Zhang, Chao Liu. "The Chemical and Pharmacological Research Progress on a Kind of Chinese Herbal Medicine, Fructus Malvae", Molecules, 2022

Publication

1 %

19

Zhenyou Tan, Jun Deng, Qiongqian Ye, Zhenfeng Zhang. "The Antibacterial Activity of Natural-derived Flavonoids", Current Topics in Medicinal Chemistry, 2022

Publication

1 %

20

Submitted to University of St La Salle

Student Paper

1 %

21

Chenchen Peng, Xuehua Wan, Jingjing Zhang, Bingling Zhang, Shuo Wang, Ting Ma, Ya Bian, Wei Wang. "Bacterial diversity and competitors for degradation of hazardous oil refining waste under selective pressures of temperature and oxygen", Journal of Hazardous Materials, 2022

Publication

1 %

| | | |
|----|---|------|
| 22 | Igor Ukrainets, Lyudmila Sidorenko, Mykola Golik, Igor Chernenok, Lina Grinevich, Alexandra Davidenko. "N-Aryl-7-hydroxy-5-oxo-2,3-dihydro-1H,5H-pyrido-[3,2,1-ij]quinoline-6-carboxamides. The Synthesis and Effects on Urinary Output", Scientia Pharmaceutica, 2018 Publication | 1 % |
| 23 | dergi.fabad.org.tr Internet Source | 1 % |
| 24 | Submitted to School of Business and Management ITB Student Paper | 1 % |
| 25 | www.tandfonline.com Internet Source | 1 % |
| 26 | pt.scribd.com Internet Source | 1 % |
| 27 | repository.akfarsurabaya.ac.id Internet Source | <1 % |
| 28 | www.plantsjournal.com Internet Source | <1 % |
| 29 | Bartholomew Adeleke, Olubukola Babalola. "Pharmacological Potential of Fungal Endophytes Associated with Medicinal Plants: A Review", Journal of Fungi, 2021 Publication | <1 % |

30

Luísa Nathália Bolda Mariano, Daiane Aline Pontoli, Ana Aline da Silva, Rivaldo Niero et al. "Diuretic and antiurolithic effect of *Garcinia humilis* (Vahl) C.D.Adams leaves, a medicinal plant native to South American countries", *Chemistry & Biodiversity*, 2022

Publication

<1 %

31

[Www.dovepress.com](http://www.dovepress.com)

Internet Source

<1 %

32

ejurnal.ung.ac.id

Internet Source

<1 %

33

Hatice Kiziltas. "Determination of LC-HRMS profiling, antioxidant activity, cytotoxic effect and enzyme inhibitory properties of *Satureja avromanica* using in vitro and in silico methods", *Process Biochemistry*, 2022

Publication

<1 %

34

journals.plos.org

Internet Source

<1 %

35

Elfita Elfita, Munawar Munawar, Muharni Muharni, Gusti Pratiwi, Rahmadania Rahmadania. "A New Benzoyl Compound Isolated from the Endophytic Fungi of *Kandis Gajah* (*Garcinia griffithii*) and *Asam Kandis* (*Garcinia cowa*)", *Makara Journal of Science*, 2016

Publication

<1 %

- | | | |
|----|---|------|
| 36 | Kaoru Okazaki, Noriyasu Fukuoka, Toru Kuboi, Jun Unemoto, Masatoshi Kondo, Takashi Kusaka, Susumu Itoh. "Furosemide clearance in very preterm neonates early in life: A pilot study using scavenged samples", Pediatrics International, 2021 Publication | <1 % |
| 37 | Submitted to Universitas Muhammadiyah Sumatera Utara Student Paper | <1 % |
| 38 | Www.mdpi.com Internet Source | <1 % |
| 39 | academic.oup.com Internet Source | <1 % |
| 40 | austinpublishinggroup.com Internet Source | <1 % |
| 41 | www.medworld.gr Internet Source | <1 % |
| 42 | doaj.org Internet Source | <1 % |
| 43 | journal.ummat.ac.id Internet Source | <1 % |
| 44 | Tanmay Sarkar, Molla Salauddin, Arpita Roy, Nikita Sharma et al. "Minor tropical fruits as a potential source of bioactive and functional | <1 % |

foods", Critical Reviews in Food Science and Nutrition, 2022

Publication

-
- | | | |
|----|--|------|
| 45 | Salni Salni, Herlina Herlina, Aprila Purnamasari. "A subchronic toxicity test of combination n-hexane and ethyl acetate fraction of rose myrtle leaves (<i>Rhodomyrtus tomentosa</i> [Ait.] Hassk) on male white rats", <i>Journal of Applied Biology & Biotechnology</i> , 2022 Publication | <1 % |
| 46 | repository.unmuhjember.ac.id Internet Source | <1 % |
| 47 | Submitted to Higher Education Commission Pakistan Student Paper | <1 % |
| 48 | Submitted to Myongji University Graduate School Student Paper | <1 % |
| 49 | lume.ufrgs.br Internet Source | <1 % |
| 50 | patentscope.wipo.int Internet Source | <1 % |
| 51 | ejbpc.springeropen.com Internet Source | <1 % |
| 52 | f1000research.com Internet Source | <1 % |

53 Submitted to Mount Kenya University <1 %
Student Paper

54 journalsaintek.uinsby.ac.id <1 %
Internet Source

55 ojs.jmolekul.com <1 %
Internet Source

56 "Acute Heart Failure", Springer Science and <1 %
Business Media LLC, 2008
Publication

57 coek.info <1 %
Internet Source

58 www.cellmolbiol.org <1 %
Internet Source

59 www.imedpub.com <1 %
Internet Source

60 www.thaiscience.info <1 %
Internet Source

61 Ibtisam Mohammed Ababutain, Sahar <1 %
Khamees Aldosary, Amal Abdulaziz Aljuraifani,
Azzah Ibrahim Alghamdi et al. "Identification
and Antibacterial Characterization of
Endophytic Fungi from Artemisia sieberi",
International Journal of Microbiology, 2021
Publication

62 Submitted to Pukyong National University

<1 %

63

Ata Mahmoodpoor, Sahra Zahedi, Arezou Pourakbar, Hamed Hamishehkar et al. "Efficacy of furosemide-albumin compared with furosemide in critically ill hypoalbuminemia patients admitted to intensive care unit: a prospective randomized clinical trial", DARU Journal of Pharmaceutical Sciences, 2020

Publication

<1 %

64

Submitted to Universidade do Porto

Student Paper

<1 %

65

text-id.123dok.com

Internet Source

<1 %

66

www.ijpba.info

Internet Source

<1 %

67

Pernilla Videhult Pierre, Jesper Edvardsson Rasmussen, Sahar Nikkhou Aski, Peter Damberg, Göran Laurell. "High-Dose Furosemide Enhances the Magnetic Resonance Signal of Systemic Gadolinium in the Mammalian Cochlea", Otology & Neurotology, 2020

Publication

<1 %

68

Seham S. El-hawary, Abeer S. Moawad, Hebatallah S. Bahr, Usama Ramadan

<1 %

Abdelmohsen, Rabab Mohammed. " Natural product diversity from the endophytic fungi of the genus ", RSC Advances, 2020

Publication

69

Toga Pangihotan Napitupulu, Indriati Ramadhani, Atit Kanti, I Made Sudiana.

"Diversity, phosphate solubilizing, and IAA production of culturable fungi associated with healthy and wilt banana", Archives of Phytopathology and Plant Protection, 2021

Publication

<1 %

70

biota.ac.id

Internet Source

<1 %

71

journal.uad.ac.id

Internet Source

<1 %

72

www.inabj.org

Internet Source

<1 %

73

www.phcogj.com

Internet Source

<1 %

74

Chia Yook Chin, Ooi Pei Boon, Hwang Jung Shan, Teow Sin Yeang, Badariah Ahmad, Peh Suat Cheng (Eds). "Diabetes: Disarming the Silent Killer", Diabetes, 2022

Publication

<1 %

75

Didjour Albert Kambiré, Jean Brice Boti, Zana Adama Ouattara, Thierry Acafou Yapi, Ange Bighelli, Félix Tomi, Joseph Casanova. " Leaf

<1 %

essential oil from Ivorian (Annonaceae):
Chemical composition and structure
elucidation of four new natural
sesquiterpenes ", Flavour and Fragrance
Journal, 2020

Publication

76

Esraa Nasser Mohammed, Amel M. Soliman,
Ayman S. Mohamed. " Modulatory effect of
on myocardial infarction induced by
epinephrine in rats ", Journal of Food
Biochemistry, 2022

Publication

<1 %

77

Zhang-Gui Ding. "¹H and ¹³C NMR
assignments of eight nitrogen containing
compounds from *Nocardia alba* sp.nov (YIM
30243^T)", Magnetic Resonance in Chemistry,
04/2009

Publication

<1 %

78

bmccomplementmedtherapies.biomedcentral.com

Internet Source

<1 %

79

downloads.hindawi.com

Internet Source

<1 %

80

ebin.pub

Internet Source

<1 %

81

eprints.utm.my

Internet Source

<1 %

| | | |
|----|--|------|
| 82 | revistagastroenterologiamexico.org Internet Source | <1 % |
| 83 | rsdjournal.org Internet Source | <1 % |
| 84 | www.diva-portal.org Internet Source | <1 % |
| 85 | www.hindawi.com Internet Source | <1 % |
| 86 | www.pubfacts.com Internet Source | <1 % |
| 87 | www.science.gov Internet Source | <1 % |
| 88 | www.vliz.be Internet Source | <1 % |
| 89 | Sara Tufer, Ephrem Engidawork, Akeberegn Gorems Ayele, Chala Bashea. "Evaluation of the Diuretic Activity of Aqueous and 80% Methanol Extracts of Croton macrostachyus(Euphorbiaceae) Leaves in Saline-Loaded Rats", Journal of Experimental Pharmacology, 2021 Publication | <1 % |
| 90 | www.ncbi.nlm.nih.gov Internet Source | <1 % |

91

Ji-Hae Park, Ye-Jin Jung, Jae-Woo Jung, Sabina Shrestha, Dong Wook Lim, Daeseok Han, Nam-In Baek. " A new flavonoid glycoside from the root bark of L. ", Natural Product Research, 2014

Publication

<1 %

92

Priscila de Souza, Luísa Nathália Bolda Mariano, Camile Cecconi Cechinel-Zanchett, Valdir Cechinel-Filho. "Promising Medicinal Plants with Diuretic Potential Used in Brazil: State of the Art, Challenges, and Prospects", Planta Medica, 2020

Publication

<1 %

93

www.wjgnet.com

Internet Source

<1 %

Exclude quotes On

Exclude matches Off

Exclude bibliography Off