

Secondary Metabolite and Antioxidant Activity of Endophytic Fungi Isolated from Syzygium aqueum Leaves Stalk

by Hary Widjajanti

Submission date: 21-Sep-2022 12:15PM (UTC+0700)

Submission ID: 1905177017

File name: 28_SECONDARY_METABOLITE_JURNAL_BRIAC_25_NOPEMBER_2021.pdf (743.26K)

Word count: 5575

Character count: 30513

Secondary Metabolite and Antioxidant Activity of Endophytic Fungi Isolated from *Syzygium aqueum* Leaves Stalk

Ummi H. Habisukan^{1,2}, Elfita^{3,*}, Hary Widjajanti⁴, Arum Setiawan⁴

¹ Graduate School of Sciences, Faculty of Mathematics and Natural Sciences, University of Sriwijaya, Jl. Padang Selasa 524, Palembang 30139, South Sumatra, Indonesia; ummihirashabisukan@radenfatah.ac.id (U.H.H.);

² Universitas Islam Negeri Raden Fatah Palembang, South Sumatra, Indonesia; ummihirashabisukan@radenfatah.ac.id (U.H.H.);

³ Department of Chemistry, Faculty of Mathematics and Natural Sciences, University of Sriwijaya, Jl. Palembang Prumulih, Indralaya, South Sumatra, 30662 Indonesia; elfita.elfita.69@gmail.com (E.);

⁴ Department of Biology, Faculty of Mathematics and Natural Sciences, University of Sriwijaya, Jl. Palembang Prumulih, Indralaya, South Sumatra, 30662 Indonesia; hary_widjajanti@unsri.ac.id (H.W.); arum.setiawan@unsri.ac.id (A.S.);

* Correspondence: elfita.elfita.69@gmail.com (E.);

Scopus Author ID 55532411800

Received: 23.10.2021; Revised: 19.11.2021; Accepted: 21.11.2021; Published: 25.11.2021

Abstract: Medicinal plants are a promising host for endophytic fungi to produce secondary metabolites relevant for food and health. In this study, we evaluate antioxidant activity to determine the species of endophytic fungi isolated from *Syzygium aqueum*. Endophytic fungi were isolated from leaf stalks through surface sterilization. The fungi's isolate was identified with morphology and molecular analysis (ITS-rDNA). The pure fungi in was cultivated on PDB media for 4 weeks, and metal were extracted using ethyl acetate. The crude extract of endophytic fungi was examined for its antioxidant activity using 2,2-diphenyl-1-picrylhydrazyl (DPPH). The pure compound was isolated using the chromatography method, and its structure was determined using spectroscopy analysis involving NMR 1D and 2D. In total, four obtained endophytic fungi were isolated from leaf stalks. The fungi with good antioxidant activity (IC_{50} 59.2 μ g/mL) were identified as *Beltrania rhombica*. The characteristics of the pure compound are white-yellowish powder with IC_{50} 44.2 μ g/mL. Based on spectroscopy analysis, the pure compound was identified as 3-(hydroxyl(3,4,5-trihydroxyphenyl)methyl)-3,4-dihydro-2H-pyran-4,5,6-triol.

Keywords: secondary metabolite; endophytic fungi; *Beltrania rhombica*; antioxidant; *Syzygium aqueum*.

© 2021 by the authors. This article is an open-access article distributed under the terms and conditions of the Creative Commons Attribution (CC BY) license (<https://creativecommons.org/licenses/by/4.0/>).

1. Introduction

Antioxidants play an important role in hindering or preventing degenerative diseases caused by free radicals [1,2]. At the normal limit, the body has natural defenses to help it overcome the rise of free radicals [3,4]. Superoxide compounds can be pressed by antioxidants [5]. Antioxidants have become an interesting topic nowadays because of their ability to protect the human body from various diseases caused by free radical reactions and reduce oxidative stress [6]. Antioxidants are compounds that are widely used as industrial resources in the food and health fields. The use of synthetic antioxidants to protect against free radical damage has been reported to cause harmful effects [7,8]. Therefore, it is necessary to find a safer source of

antioxidants from natural ingredients, such as jambu air (*Syzygium aqueum*), which has been reported to contain active antioxidant compounds, especially in their leaves [9].

In various parts of the world, including Indonesia, *Syzygium aqueum* has been used as medicine for its antioxidant activity [10,11]. The most abundant chemical compounds in *Syzygium aqueum* are flavonoids and phenols, which are found in almost all parts, such as leaves, flowers, fruit, seeds, roots, and bark. They have been reported to have antioxidant, antibacterial, antiviral, anti-inflammatory, anti-allergic, and antidiabetic activities and play a role in cancer prevention [12,13]. Currently, the search for natural bioactive is not only from plants but also from their endophytic fungi.

Endophytic fungi are microscopic living organisms that live in plant tissues (leaves, fruit, seeds, stems, and roots) at certain periods and form colonies without harming their hosts, even maintaining mutually beneficial relationships [14,7]. Endophytic fungi have been identified as a source of secondary metabolites, such as antioxidants, antibiotics, antivirals, antiprotozoal, antidiabetic, and anticancer agents. These compounds include alkaloids, terpenoids, steroids, isocoumarin derivatives, lactones, quinones, flavonoids, phenols, indole derivatives, lignins, tannins, anthraquinones, xanthenes, phenylpropanoids, phenolic acids, and peptide peptides [15-17]. Natural products derived from endophytic fungi are considered as one of the most relevant sources of discovery and molecular diversity for new drugs. This is due to the large diversity of endophytic fungi species that can produce secondary metabolites with biological activity, plus the ease of cultivating endophytic fungi in large quantities and in a short time. [18-20]. Therefore, research on endophytic fungi that produce antioxidant compounds needs to be carried out on the host plant *S. aqueum*.

66

2. Materials and Methods

2.1. Plant material.

The plant material is *Syzygium aqueum* leaf stalk taken in the Sriwijaya University area, Indralaya, Ogan Ilir, South Sumatera, and identified at the Biosystematics Laboratory, Biology, FMIPA, Sriwijaya University, with certificate number: 329//UN9.1.7/4/EP/2020.

48

2.2. Isolation and identification of endophytic fungi.

The leaf stalks of *S. aqueum* were washed with running tap water and then dried. Fragments of it were surface-sterilized by immersing each sample in 70% alcohol for 1 min, then in 3% sodium hypochlorite (NaOCl) for 1 min. After rinsing with sterile distilled water for 1 min, the outer tissue was removed with a sterile scalpel. Small pieces of leaf stalks were placed in a Petri dish containing PDA media supplemented with chloramphenicol (0.2 g/L) then incubated for 7 d at $30 \pm 2^\circ\text{C}$. All experiments were carried out in triplicate. Fungal growth from the leaf stalk segments was monitored every day. The individual hyphae tips were transferred to fresh PDA and incubated at 30°C for 7 d. Pure cultures were numbered and maintained in a PDA slope and stored at 4°C .

Identification of endophytic fungi through molecular analysis followed the procedure described [21], which was based on partial genetic analysis at the Internal Transcribed Spacer (ITS) ribosomal DNA of fungi. The phylogenetic tree construction was carried out using several applications: Clustal W, SEQBOOT, DNA dist, Neighbor, CONSENSE, and FigTree.

2.3. Cultivation and extraction.

Endophytic fungal cultures were prepared on PDB media by placing 6 cm diameter agar blocks of pure culture into five 1 L Erlenmeyer flasks containing 300 mL of medium each. The flasks were incubated under static conditions at room temperature for 4 weeks. The culture was filtered through filter paper to separate the mycelium. The liquid broth was collected and extracted with ethyl acetate (1:1) in a separating funnel with vigorous shaking for 1 hour, then filtered. Extraction was carried out with two repetitions. The ethyl acetate extract was evaporated with a rotary evaporator to produce a concentrated extract [3,22,23].

2.4. Antioxidant activity assay.

The antioxidant activity assay used the DPPH method [24]. A 0.05 mM solution of DPPH in methanol was prepared, and 3.8 mL of this solution was mixed with 0.2 ml of the test sample in methanol at a series concentration. 200, 100, 50, 25, 12.5, 6.75, and 0 µg/mL were prepared. The reaction mixture was thoroughly vortexed and left in the dark for 30 min. The absorbance of the mixture was measured spectrophotometrically (Shimadzu, Uv-1900) at a wavelength of 517 nm. Gallic acid is used as a standard antioxidant. The percentage of DPPH radical scavenging activity was calculated using the formula:

$$\% \text{ Inhibition} = \frac{\text{Control Absorbance} - \text{Sample Absorbance}}{\text{Control Absorbance}} \times 100\%$$

2.5. Isolation and identification of secondary metabolites.

The active extract (selected by IC₅₀ highest value) was analyzed by thin-layer chromatography (TLC) of silica gel G-60 F 254 using solvents with various eluents to see the staining pattern. Column chromatography extract used a stationary phase such as silica gel. Samples that have been prepared by pre-absorption are fed into the chromatographic column evenly and eluted with a gradient system. The eluate was collected in vials every 10 ml, and each had a chromatographed thin layer to be grouped into column fractions based on the staining pattern. The visible stains were UV lamp at 254 nm and cerium sulfate spray reagent. Fractions containing potential compounds in column chromatography were used to obtain pure compounds. Identification of the chemical structure was carried out by spectroscopic analysis, which included ¹H-NMR, ¹³C-NMR, DEPT 135, HMQC, HMBC, and COSY (Agilent DD2 500 MHz (¹H) dan 125 MHz (¹³C)).

3. Results and Discussion

The isolation of endophytic fungi from leaf stalks of *S. aqueum* obtained four isolates, three of which were the same as endophytic fungi obtained from leaves: *Cochliobolus* sp., *Penicillium* sp., and *Fusarium* sp. The difference was one fungus labeled B52. The results of the molecular identification of Sequence Assembly 572 bp are

```
CGTAGGTGAACCTGCGGAGGGATCATTACAGAGTTTTCTAAACTCCCAAACCCAT
GTGAACTTACCATTGTTGCCTCGGCGGAGCCTACCCTGTAGCTACCCTATAAGGT
GGTACCCTGTAGCGCCCGCCGGTGGATTTTCAAACCTCTTGTTATTTATAGTAATC
TGAGAGTCTTATTTTAAATAAGTCAAACTTTCAACAACGGATCTCTTGGCTCTG
GCATCGATGAAGAACGCAGCGAAATGCGATAAGTAATGTGAATTGCAGAATTCA
```

GTGAATCATCGAATCTTTGAACGCACATTGCGCCCATTAGTATTCTAGTGGGCAT
GCCTGTTTCGAGGTCATTTCAACCCTTAAGCCTAGCTTAGTGTGGGAGTCTACGA
GTGGGCCACGTTACCCTGTAGCGTGGTTACCCTGTAGTTCCTGAAAATCAACGGC
GGATTTACAGTATCCTCTGAGCGTAGTAATTCTTTATCTCGCTTCTGTTTAGGTGC
TGTGACTTCGGCCGCTAAACCCACAATTTTTTTGTGGTTGACCTCGGATCAGGT
AGGAATACCCGCTGAACTTAAGCAT

The strain B52 has accession number OK376219. The results of the B52 endophytic fungal phylogeny tree construction are shown in Figure 1. The B52 phylogenetic tree showed that the isolate sequence was in the same branch as *Beltrania rhombica*. Phylogenetic analysis indicated that isolate B52 was *Beltrania rhombica*. It was carried out using Clustal W, Seqboot, DNA dist, Neighbor, Consense, and FigTree.

The endophytic fungus isolate culture B52 in 5 Erlenmeyer flasks containing 300 mL PDB medium was incubated for 4 weeks at room temperature and dark conditions. Extraction of endophytic fungi using ethyl acetate solvent, after evaporation, produces a concentrated extract of ethyl acetate 6.2 g. The antioxidant activity test gave IC₅₀ a value of 59.2 µg/mL, while the IC₅₀ values for the other three endophytic fungi were 98.26 g/mL (*Cochliobolus* sp.), 92.3 g/mL (*Penicillium* sp.), and 64.33 g/mL (*Fusarium* sp.).




Figure 1. Phylogenetic tree of the endophytic fungal B52 was carryout using applications: Clustal W, Seqboot, DNA dist, Neighbor, Consense, and Fig Tree.

The results of the TLC analysis of the concentrated ethyl acetate extract showed that there was a major purple stain at R_f 0.55, with n-hexane:ethyl acetate as the eluent (2:8). Therefore, the eluent to be used in column chromatography can use a stepwise system. Ethyl acetate extract 2 g was pre-absorbed with silica gel as stationary phase 70–230 mesh in a ratio

1). Separation of pure compounds was carried out by column chromatography, which was eluted with a mixture of n-hexane and ethyl acetate (10:0–0:10), and a mixture of ethyl acetate and methanol (9:1). The eluate was collected every 10 mL in the vial until 65 vials were obtained. Based on TLC analysis, four subfractions (F1–F4) were obtained. Subfraction F3 showed major stains and continued purification by column chromatography until pure compound (compound 1) was obtained in the form of a yellowish-white solid (44 mg). The antioxidant activity test gave an IC_{50} value of 44.2 $\mu\text{g/mL}$.

3.1. Determination of chemical structure.

The $^1\text{H-NMR}$ spectrum of compound 1 (Fig. 2) shows the presence of 11 proton signals, including two doublet signals, in the aromatic chemical shift region, namely, at δ_{H} 7.70 and 8.19 ppm, each of which has a double integration and coupling constant of $J=8.5$ Hz. This indicates that compound 1 has a para-substituted aromatic structure, with two pairs of equivalent protons. Five other signals appear in the sp^3 proton region, namely, two oxygenated methine signals, namely, at δ_{H} 5.33 (1H, m); 6.37 (1H, s); and two oxygenated methylene proton signals that appear at different chemical shifts, namely, at δ_{H} 3.70 (1H, m) and 3.79 ppm (1H, m), while the other sp^3 signal is proton methine at δ_{H} 4.16 ppm (1H, m). In addition, there are four proton signals that, after being confirmed with the HMQC spectrum (Fig. 4), turn out not to be bound to the carbon atom. This indicates that these signals are proton signals bound to heteroatoms, such as hydroxyl protons.

The $^{13}\text{C-NMR}$ spectrum of compound 1 (Fig. 3) showed 10 signals: four sp^3 carbon signals and six other signals that appeared in the sp^2 carbon region. After confirmation with DEPT 135 spectrum, it was found that the four sp^3 carbon signals were two oxygenated methine carbon signals (δ_{C} 71.2 and 67.5 ppm), one oxygenated methylene carbon signal (δ_{C} 62.2 ppm), and one tertiary carbon signal (δ_{C} 58.0 ppm). Two high-intensity signals in the sp^2 carbon region indicate that compound 1 has two pairs of equivalent aromatic carbons (δ_{C} 123.9 and 128.2 ppm). Four other sp^2 carbon signals were at δ_{C} 148.1, 151.5, 164.4, and 164.5 ppm, respectively, after being confirmed with DEPT 135 spectrum. It is known that the four carbons are quaternary.




Figure 2. The $^1\text{H-NMR}$ spectra of compound 1.




Figure 3. The ^{13}C -NMR (A) DEPT 135 (B) spectra of compound 1.

80

The analysis of the proton and carbon NMR spectra is confirmed by the data on the HMQC spectrum (Fig. 4) the ^1H - ^{13}C correlation through one bond. The HMQC spectrum showed six correlations consisting of two ^1H - ^{13}C correlations on the aromatic ring, three correlations on oxygenated ^1H - ^{13}C , and one ^1H - ^{13}C correlation for methine. Thus, the 11 signals that appear on proton NMR consist of seven signals bound to the carbon atom (two non-equivalent methylene proton signals) and four protons bound to the heteroatom.




Figure 4. The HMQC spectra of compound 1.

The HMBC spectrum (Fig. 5) showed a ^1H - ^{13}C correlation through two or three bonds. The aromatic proton signal at H 8.19 ppm showed three correlations: a correlation each with ortho aromatic carbon (δ_{C} 128.2 ppm), its equivalent aromatic carbon (δ_{C} 123.9 ppm), and quaternary aromatic carbon (δ_{C} 148.1 ppm). Another aromatic proton at H 7.70 ppm has four correlations: a correlation each with ortho aromatic carbon (δ_{C} 123.9 ppm), its equivalent aromatic carbon (δ_{C} 128.2 ppm), quaternary aromatic carbon (δ_{C} 148.1 ppm), and oxygenated side-chain carbon (δ_{C}) 71.2 ppm.

Furthermore, oxygenated methine protons at δ_{H} 5.33 ppm have three ^1H - ^{13}C correlations via three bonds: aromatic carbon (δ_{C} 128.2 ppm) and oxygenated carbon (δ_{C} 58.0

and 62.2 ppm). The correlation indicates that the oxygenated methine group at δ_H 5.33 ppm is directly attached to the aromatic ring and is para-substituted with a hydroxyl group. The oxygenated methine proton (δ_H A= 3.70; B= 3.79 ppm) had two 1H - ^{13}C correlations with two bonds with oxygenated methine carbon at δ_C 58.0 ppm and a triple correlation with oxygenated methine carbon at δ_C 71.2 ppm. The 1D and 2D NMR spectral data for compound 1 are shown in Table 1.




Figure 5. The HMBC spectra of compound 1.

24 Table 1. The NMR data of Compound 1.

No. C	δ_C ppm	DEPT 135	δ_H ppm (ΣH , multiplicity, J (Hz))	HMBC	COSY
2	62.2	CH ₂	A= 3.70 (1H, m) B= 3.79 (1H, m)	58.0; 71.2 58.0; 71.2	4.16 4.16
3	58.0	CH	4.16 (1H, m)	62.2	5.33; 3.70; 3.79
4	67.5	CH	6.37 (1H, s)	164.4	-
5	151.5	C	-	-	-
6	164.4	C	-	-	-
1'	148.1	53	-	-	-
2'	128.2	CH	7.70 (1H, d, J=8.5)	71.2; 128.2; 148.1; 123.9	8.19
3'	123.9	CH	8.19 (1H, d, J=8.5)	123.9; 148.1; 128.2	7.70
4'	164.5	C	-	-	-
5'	123.9	53	8.19 (1H, d, J=8.5)	123.9; 148.1; 128.2	7.70
6'	128.2	CH	7.70 (1H, d, J=8.5)	71.2; 128.2; 148.1; 123.9	8.19
7'	71.2	CH	5.33 (1H, m)	128.2; 58.0; 62.2	4.16; 7.70
OH	-	-	3.82	67.5	-
OH	-	-	4.33	-	-
OH	-	-	5.29	58.0; 151.5	-
OH	-	-	7.56	164.5	-

The COSY spectrum in Figure 6 appears to show that the aromatic proton at δ_H 7.70 ppm has a 1H - 1H correlation through three bonds with an aromatic proton δ_H 8.19 ppm and a 1H - 1H correlation through more than three bonds with the oxygenated sp³ methine proton at δ_H 5.33 ppm bound to an aromatic ring. The methine proton at δ_H 4.16 ppm, part of the side chain, has a 1H - 1H correlation through three bonds with the oxygenated sp³ methine proton (δ_H

5.33 ppm) and oxygenated methylene proton (δ_H 3.70 and 3.79 ppm). This 1H - 1H correlation of the COSY spectrum indicates that the two aromatic equivalent protons are in the ortho position. This strengthens the proposed structure: compound 1 is a benzene ring directly bonded to the oxygenated methine carbon at the para position with a hydroxyl group.




Figure 6. The COSY spectra of compound 1.

Based on the spectrum analysis of 1H -NMR, ^{13}C -NMR, DEPT 135, HMQC, HMBC, and COSY, it can be explained that compound 1 has a para-substituted frame on the benzene ring. It was identified from the presence of two aromatic proton signals with the integration of two protons. The multiplicity of the two signals is doublet with the same coupling constant ($J = 8.5$ Hz). This indicates that the two protons bound to the aromatics are in the ortho position, and the two substituents are bound to para (Fig. 2). One of the substituents is a hydroxyl group, which is characterized by a chemical shift value of oxyaryl carbon at δ_C 164.5 ppm, while the second substituent is carbon methine sp^3 (δ_C 71.2 ppm), which has a hydroxyl group and a pyran-4,5,6-triol group (Fig. 3). Furthermore, the three hydroxyl groups attached to the pyran ring can be validated from the HMBC spectrum in the presence of a 1H - ^{13}C correlation through two or three bonds. In addition, the binding of the pyran group to C methine (C-7') can be identified from the COSY spectrum (Fig. 6) through the 1H - 1H three-bond correlation.

The proton at δ_H 6.37 ppm appears singlet, possibly due to the transposition with the proton at δ_H 4.16 ppm, as shown in Figure 7, so the distant position does not cause cleavage. Thus, the proposed chemical structure of compound 1 is 3-(hydroxy(4-hydroxyphenyl)methyl)-3,4-dihydro-2H-pyran-4,5,6-triol, as shown in Figure 7.

Compound 1 has good antioxidant activity due to the easy abstraction of hydroxyl protons by DPPH free radicals and generates new free radicals that can be stabilized by radical delocalization in the compound. Compounds with 1,2-dihydroxy units in the $C=C$ bond are more susceptible to proton abstraction by free radicals to form more stable free radicals. This stability is due to the free radicals produced, forming intramolecular hydrogen bonds with the adjacent hydroxyl group and a diketone. Likewise, compounds with a hydroxyl group at the para position are prone to proton abstraction by free radicals to form new, more stable free radicals. This stability is due to the distribution and delocalization of electrons [25-27].




Figure 7. Compound 1 as 3-(hydroxy(4-hydroxyphenyl)methyl)-3,4-dihydro-2H-pyran-4,5,6-triol.

The antioxidant activity of compound 1 can be increased by carrying out a semisynthetic reaction similar to the position of the hydroxyl group on gallic acid (antioxidant standard). The OH group on the aromatic ring of compound 1 is ortho directive so that with the electrophilic substitution reaction, Nitration (2 mol $\text{HNO}_3/\text{H}_2\text{SO}_4$), followed by a reduction of the nitro group (Fe/HCl), will produce two amine groups at C3 and C5. The next step is transforming the amine functional group into aryldiazonium chloride ($\text{ArN}_2^+ \text{Cl}^-$) and its final replacement by $-\text{OH}$ through the reaction between the diazonium salt and hot aqueous acid [28]. Through this series of reactions, it is expected to produce 3-(hydroxyl (3,4,5-trihydroxyphenyl)methyl)-3,4-dihydro-2H-pyran-4,5,6-triol compound with antioxidant activity close to gallic acid, where gallic acid has a strong antioxidant with an IC_{50} value of $11.4 \mu\text{g}/\text{mL}$. The production of compound 1 can be carried out within 4 weeks of incubation, and the enrichment technique needs to be investigated further so that it can be produced as needed.

Endophytic fungi from the Beltraniaceae family can produce a variety of bioactive compounds [29]. *Beltrania* species is one of the rare endophytic fungi that are rarely found [30]. From the ethyl acetate extract of the culture broth of *Beltrania rhombica*, two new eudesmane sesquiterpenes, named rhombidiol and rhombitriol, were isolated. These secondary metabolites have antibacterial and antifungal activity [31]. *Beltrania queiroza* isolated from Amazonian medicinal plants contains active antiviral compounds [32]. *Beltrania* species showed their importance for the production of antitumor agents. Two new sesquiterpenes were isolated from the culture broth of *Beltrania* species [33], and sesquiterpenes are well known to have a potential application as an anticancer agent [34].

The results of the literature study [12,35] produced a report that the phenyl compounds from *S. aqueum* and other *Syzygium* spp. (Fig. 8) were not the same as compound 1. This indicates that compound 1 is typically produced by the endophytic fungus *Beltrania rhombica* of *S. aqueum*. There is no genetic evolution between fungal and host genes in producing compound 1.




Figure 8. Phenyls of *S. aqueum* and other *Syzygium* spp.

Some endophytic fungi produce certain phytochemical compounds that are also produced by their host plants. This is thought to be the result of genetic transfer from the host plant to endophytic fungi. Studies have shown that about 18% of plant-derived metabolites can also be derived from related fungi. That is the case with anticancer compounds, such as taxol and podophyllotoxin. Taxol is produced from the medicinal plant *Taxus* spp. and the endophytic fungus *Taxomyces andreanae*. Podophyllotoxin compounds are produced from the *Sinopodophyllum hexadrum* plant and its endophytic fungus *Fusarium solani* [36-38]. In recent years, endophytic fungi have become a source of secondary metabolite production. A total of 449 new secondary metabolites were produced from different tissues and plants with different chemical structures and biological activities. Among these new compounds, terpenoids constituted the largest proportion (26%), followed by ketones (22%), lactones (7%), anthraquinones (5%), steroids, penylpropanoids, alkaloids each 3% and other compounds (31%) [39].

4. Conclusions

Endophytic fungi B52 were isolated from *Syzygium aqueum* leaves stalk identified as *Beltrania rhombica* produced compound 1 as 3-(hydroxy(4-hydroxyphenyl)methyl)-3,4-dihydro-2H-pyran-4,5,6-triol, which is a phenolic compound that has antioxidant activity that can be developed as a source of new antioxidants. The compound produced by *Beltrania rhombica* is a promising potential source as a drug raw material in the future through several stages of simple semisynthetic reactions.

Funding

The research was funded by Daftar Isian Pelaksanaan Anggaran (DIPA) Universitas Sriwijaya, Indonesia, through Unggulan Profesi grants 2020, number 0009/UN9/SK.LP2M.PT/2020, September 28, 2020.

Acknowledgments

The authors wish to respect the facilities' support of the Organic Chemistry and Microbiology Laboratory, Faculty of Mathematics and Natural Sciences, University of Sriwijaya.

Conflicts of Interest

The authors declare no conflict of interest.

References

1. Etiana, E.; Bustanussalam; Yadi; Rachman, F; Hapsari, Y; Izzati, F.N.; Rahmawati, S.I.; Simanjuntak, P. Antioxidant activity of endophytic fungi from young and old leaves of cinnamon plants from Berau, Indonesia. *IOP Conf. Series: Earth and Environmental Science* **2021**, *762*, 1-10. <https://doi.org/10.1088/1755-1315/762/1/012038>.
2. Stagos, D. Antioxidant Activity of Polyphenolic Plant Extracts. *Antioxidants* **2020**, *9*, 1-7, <https://doi.org/10.3390/antiox9010019>.
3. Elfita; Mardiyanto, M.; Fitrya, F.; Larasati; Julinar, J.; Widjajanti, H.; Muharni, M. Antibacterial activity of *Cordyline fruticosa* leaf extracts and its endophytic fungi extracts. *Biodiversitas* **2019**, *20*, 3804-3812, <https://doi.org/10.13057/biodiv/d201245>.
4. Mehdi, S.R.; Nanjangud, V.; Anil, K.; Paolo, Z.; Elena, M.V.; Luciana, D.; Elisa, P.; Jovana, R.; Patrick, V.; Tsouh, F.; Elena, A.; Ilaria, P.; Abhay, P.M.; Manisha, N.; Youssef, E.I.R.; Marc, E.I.B.; Letizia, P.; Marcello, I.; Natália, M.; Miquel, M.; Anca, O.D.; William, N.; Setzer; Daniela, C.; William, C.C.; Javad,

- 65 R. Lifestyle, oxidative stress, and antioxidants: back and forth in the pathophysiology of chronic diseases. *Front Physiol* **2021**, *11*, <https://doi.org/10.3389/fphys.2020.00694>.
5. Mucha, P.; Skoczynska, A.; Malecka, M.; Hikiş, P.; Budzisz, E. Overview of the oxidant and anti-inflammatory activities of selected plant compounds and their metal ions complexes. *Molecules* **2021**, *26*, 4886, <https://doi.org/10.3390/molecules26164886>.
 6. Aragon, M.O.H.; Martinez, M.F.; Isiordia, E.M.A.; Contreras, D.L.; Juan, M.J.R.Z.; Gonzalez, R.M.L. Bacterial resistance and failure of clinical cure could be produced by oxidative stress in patients with diabetes or cardiovascular diseases during fluoroquinolone therapy. *Medical Hypotheses* **2017**, *103*, 32-34, <https://doi.org/10.1016/j.mehy.2017.04.004>.
 7. Ibrahim, M.; Oyebanji, E.; Fowora, M.; Aiyeolemi, A.; Orabuchi, C.; Akinnawo, B.; Adekunle, A. Extracts of endophytic fungi from leaves of selected Nigerian ethnomedicinal plants exhibited antioxidant activity. *JAC Complementary Medicine and Therapies* **2021**, *21*, 1-13, <https://doi.org/10.1186/s12906-021-03269-3>.
 8. Habiskan, U.H.; Elfita, W.H.; Setiawan, A.; Kurniawati, A.R. Diversity of endophytic fungi in *Syzygium Aqueum*. *Biodiversitas* **2021**, *22*, 1129-1137, <https://doi.org/10.13057/biodiv/d220307>.
 9. Sobeh, M. *Syzygium aqueum*: a polyphenol-rich leaf extract exhibits antioxidant, hepatoprotective, pain-killing and anti-inflammatory activities in animal models. *Front Pharmacol* **2018**, *9*, 1-14, <https://doi.org/10.3389/fphar.2018.00566>.
 10. Habiskan, U.H.; Elfita, W.H.; Setiawan, A.; Kurniawati, A.R. Antioxidant and antimicrobial activity of endophytic fungi isolated from *Syzygium aqueum* Leaves. *Journal of Physics: Conference Series* **2021**, *1751*, 03, <https://doi.org/10.1088/1742-6596/1751/1/012059>.
 11. Sobeh, M.; Braun, M.S.; Krstin, S.; Youssef, F.S.; Ashour, M.L.; Wink, M. Chemical profiling of the essential oils of *Syzygium aqueum*, *Syzygium samarangense*, and *Eugenia uniflora* and their discrimination using chemometric analysis. *Chem Biodivers* **2016**, *13*, 1537-1550, <https://doi.org/10.1002/cbdv.201600089>.
 12. Aung, E.E.; Kristanti, A.N.; Aminah, N.S.; Takaya, Y.; Ramadhan, R. Plant description, phytochemical constituents and bioactivities of *Syzygium* genus: A review. *A Review De Gruyter Open Chemistry* **2020**, *18*, 1436-1281, <https://doi.org/10.1515/chem-2020-0175>.
 13. Tandi, J. Pengaruh ekstrak etanol daun jambu air (*Syzygium aqueum*(Burm f.)Alston) terhadap glukosa darah, ureum dan kreatinin tikus putih (*Rattus norvegicus*) *J. Trop. Pharm. Chem* **2017**, *4*, 43-51, <https://doi.org/10.25026/jtpc.v4i2.137>.
 14. Khan, R.; Naqvi, S.T.; Ousain; Fatima, N.; Muhammad, S.A. Study of antidiabetic activities of endophytic fungi isolated from plants. *Pure and Applied Biology* **2019**, *8*, 1287-1295, <https://dx.doi.org/10.19045/bspab.2019.80071>.
 15. Manganyi, M.C.; Ateba, C.N. Untapped potentials of endophytic fungi: A review of novel bioactive compounds with biological applications. *Microorganisms* **2020**, *8*, 1-25, <https://doi.org/10.3390/microorganisms8121934>.
 16. Gakuubi, M.M.; Munusamy, M.; Liang, Z.X.; Ng, S.B. Fungal endophytes: a promising frontier for discovery of novel bioactive compounds. *J. Fungi* **2021**, *7*, <https://doi.org/10.3390/jof7100786>.
 17. Cruz, J.S.; da Silva, C.A.; Hamerski, L. Natural products from endophytic fungi associated with rubiaceae species. *Journal of Fungi* **2020**, *6*, 1-26, <https://doi.org/10.3390/jof6030128>.
 18. El-Hawary, S.S.; Moawad, A.S.; Bahr, H.S.; Abdelmohsen, U.R.; Mohammed, R. Natural product diversity from the endophytic fungi of the genus *Aspergillus*. *RSC Advances* **2020**, *10*, 22058-22079, <https://doi.org/10.1039/D0RA04290K>.
 19. Co, I.E.; Cheesman, M. Plants of the genus *Syzygium* (Myrtaceae): A review on ethnobotany, medicinal properties and phytochemistry. In: *Bioactive Compounds of Medicinal Plants: Properties and Potential for Human Health*. Academic Press: Elsevier Inc **2018**; pp. 35-84.
 20. Rai, N.; Keshri, P.K.; Verma, A.; Kamble, S.C.; Mishra, P.; Barik, S.; Singha, S.K.; Gautam, V.G. Plant associated fungal endophytes as a source of natural bioactive compounds. *Mycology* **2021**, *12*, 139-159, <https://doi.org/10.1080/21501203.2020.1870579>.
 21. Yohandini, H.; Julinar; Muharni. Isolation and phylogenetic analysis of thermophile community within Gunung Sakti hot spring, South Sumatera, Indonesia. *Hayati Journal of Biosciences* **2015**, *22*, 143-148, <https://doi.org/10.1016/j.hjb.2015.10.006>.
 22. Pansanit, A.; Pripdeevech, P. Antibacterial secondary metabolites from an endophytic fungus, *Arthrinium* sp. MFLUCC16-1053 isolated from *Zingiber cassumunar*. *Mycology* **2018**, *9*, 264-272, <https://doi.org/10.1080/21501203.2018.1481154>.
 23. Marcellano, J.P.; Collanto, A.S.; Fuentes, R.G. Antibacterial activity of endophytic fungi isolated from the bark of *Cinnamomum mercadoi*. *Pharmacognosy Journal* **2017**, *9*, 405-409, <https://doi.org/10.5530/pj.2017.3.69>.
 24. Muharni, M.; Elfita, E.; Yohandini, H.; Julinar, J.; Yasrina, Y.; Miranti, M. Chemical constituents from stem bark of *Lacourtia rukam* Zoll. & Mor. and their antioxidant activities. *Sains Malaysiana* **2019**, *48*, 1899-1906, <http://dx.doi.org/10.17576/jsm-2019-4809-10>.
 25. Sawai, Y.; Sakata, K. NMR Analytical approach to clarify the antioxidative molecular mechanism of catechins using 1,1-Diphenyl-2-picrylhydrazyl. *Journal of Agricultural and Food Chemistry* **1998**, *46*, 111-114, <https://doi.org/10.1021/jf970342x>.

26. Procházková, D.; Boušová, I.; Wilhelmová, N. Antioxidant and prooxidant properties of flavonoids. *Fitoterapia* **2011**, *82*, 513–519, <https://doi.org/10.1016/j.fitote.2011.01.018>.
27. Brunetti, C.; Martina, D.; Ferdinando, M.D.; Fini, A.; Pollastri, S.; Tattini, M. Flavonoids as antioxidants and developmental regulators: relative significance in plants and humans. *International Journal of Molecular Sciences* **2013**, *14*, 3540–3555, <https://doi.org/10.3390/ijms14023540>.
28. Fessenden, R.J.; Fessenden, J. *Organic Chemistry*. Jakarta: Erlangga. **1992**.
29. Helaly, S.E.; Thongbai, B.; Stadler, M. Diversity of biologically active secondary metabolites from endophytic and saprotrophic fungi of the ascomycete order xylariales. *Natural Product Report* **2018**, *35*, 1014–1014, <https://doi.org/10.1039/c8np00010g>.
30. Elkhateeb, W.A.; Daba, G.M.; Elnahas, M.O.; Thomas, P.W. The rarely isolated fungi: *Arthrinium sacchari*, *Beltrania querna*, and *Papulaspora immersa*, Potentials and Expectations. *ARC Journal of Pharmaceutical Sciences (AJPS)* **2019**, *5*, 1–6.
31. Rukachaisirikul, V.; Kaewbumrung, C.; Phongpaichit, S.; Hajiwangoh, Z. Eudesmane sesquiterpenes from aquatic fungus *Beltrania rhombica*. *Chem. Pharm. Bull* **2005**, *53*, 238–240, <https://doi.org/10.1248/cpb.53.238>.
32. Ferreira, M.C.; Vieira, M.S.L.A.; Zani, C.L.; Alves, T.M.D.A.; Junior, P.A.S.; Murta, S.M.F.; Romanha, A.J.; Gil, L.H.V.G.; Carvalho, A.G.D.O.; Zilli, J.E.; Vital, M.J.S.; Rosa, C.A.; Rosa, L.H. Molecular phylogeny, diversity, symbiosis and discover of bioactive compounds of endophytic fungi associated with the medicinal Amazonian plant *Carapa guianensis Aublet* (Meliaceae). *Biochemical Systematics and Ecology* **2015**, *59*, 36–44, <https://doi.org/10.1016/j.bse.2014.12.017>.
33. Ga, B.M. Natural sesquiterpenoid. *Natural product report* **2003**, *20*, 392–413, <https://doi.org/10.1039/B208084M>.
34. Modzelewska, A.; Sur, S.; Kumar, S.K.; Khan, S.R. Sesquiterpenes: natural product that decrease cancer growth. *Current Medical Chemistry-Anti-Cancer Agent* **2005**, *5*, 477–499, <https://doi.org/10.2174/1568011054866973>.
35. Habisukan, U.H.; Elfita, W.H.; Setiawan, A. Chemical characterization of secondary metabolite from the endophytic fungus *Trichordema reecei* isolated from The Twig of *Syzygium aqueum*. *Science and Technology Indonesia* **2021**, *6*, 137–143, <https://doi.org/10.26554/sti.2021.6.3.137-143>.
36. Singh, A.; Singh, D.K.; Kharwar, R.N.; White, J.F.; Gond, S.K. Fungal Endophytes as Efficient Sources of Plant-Derived Bioactive Compounds and Their Prospective Applications in Natural Product Drug Discovery: Insights, Avenues, and Challenges. *Microorganisms* **2021**, *9*, 1019, <https://doi.org/10.3390/microorganisms9010197>.
37. Tan, X.; Zhou, Y.; Zhou, X.; Xia, X.; Wei, Y.; He, L.; Tang, H.; Yu, L. Diversity and bioactive potential of culturable fungal endophytes of *Dyosma versipellis*: A rare medicinal plant endemic to China. *Scientific Reports* **2018**, *8*, 1–9, <https://doi.org/10.1038/s41598-018-24313-2>.
38. Adeleke, B. S.; Babalola, O. O. Pharmacological potential of fungal endophytes associated with medicinal plants: A review. *Journal of Fungi* **2021**, *7*, 1–16, <https://doi.org/10.3390/jof7020147>.
39. Zheng, R.; Li, S.; Zhang, X.; Zhao, C. Biological activities of some new secondary metabolites isolated from endophytic fungi: A review study. *Int. J. Mol. Sci* **2021**, *22*, <https://doi.org/10.3390/ijms22020959>.

Secondary Metabolite and Antioxidant Activity of Endophytic Fungi Isolated from Syzygium aqueum Leaves Stalk

ORIGINALITY REPORT

37%
SIMILARITY INDEX

31%
INTERNET SOURCES

31%
PUBLICATIONS

19%
STUDENT PAPERS

PRIMARY SOURCES

1 periodicos.ifap.edu.br 1%
Internet Source

2 www.sciencetechindonesia.com 1%
Internet Source

3 ojs3.unpatti.ac.id 1%
Internet Source

4 Souvik Kusari, Marc Lamshöft, Sebastian Zühlke, Michael Spiteller. " An Endophytic Fungus from that Produces Hypericin ", *Journal of Natural Products*, 2008 1%
Publication

5 www.arcjournals.org 1%
Internet Source

6 nanobioletters.com 1%
Internet Source

7 Submitted to clsu 1%
Student Paper

8	<p>Heba A. S. El-Nashar, Wagdy M. Eldehna, Sara T. Al-Rashood, Amal Alharbi, Razan O. Eskandrani, Shaza H. Aly. "GC/MS Analysis of Essential Oil and Enzyme Inhibitory Activities of Syzygium cumini (Pamposia) Grown in Egypt: Chemical Characterization and Molecular Docking Studies", <i>Molecules</i>, 2021</p> <p>Publication</p>	1 %
9	<p>repository.futminna.edu.ng:8080</p> <p>Internet Source</p>	1 %
10	<p>Submitted to Salem State University</p> <p>Student Paper</p>	1 %
11	<p>Submitted to University of South Alabama</p> <p>Student Paper</p>	1 %
12	<p>docslib.org</p> <p>Internet Source</p>	1 %
13	<p>dspace.udel.edu</p> <p>Internet Source</p>	1 %
14	<p>scholar.unand.ac.id</p> <p>Internet Source</p>	1 %
15	<p>Bartholomew Adeleke, Olubukola Babalola. "Pharmacological Potential of Fungal Endophytes Associated with Medicinal Plants: A Review", <i>Journal of Fungi</i>, 2021</p> <p>Publication</p>	1 %

16	Ramar Mohankumar, Soosai Eugene Leo Prakash, Navabshan Irfan, Subramanian Mohanraj, Chidambaram Kumarappan. "Evaluation of analgesic, anti-inflammatory, and antipyretic activities of Ziziphus mauritania Lam leaves in animal models", Pharmacological Research - Modern Chinese Medicine, 2022 Publication	1 %
17	Submitted to University of St La Salle Student Paper	1 %
18	pubs.acs.org Internet Source	1 %
19	Ji Liu, Guangming Ma, Yan Wang, Yonggen Zhang. " leaf flavonoids protect bovine mammary epithelial cells from hydrogen peroxide-induced oxidative stress in vitro ", Reproduction in Domestic Animals, 2020 Publication	1 %
20	www.mapress.com Internet Source	1 %
21	Submitted to American University in Cairo Student Paper	1 %
22	Bing-Chao Yan, Wei-Guang Wang, Ling-Mei Kong, Jian-Wei Tang, Xue Du, Yan Li, Pema-Tenzin Puno. "Cytochalasans from the	1 %

Endophytic Fungus *Phomopsis* sp. shj2 and Their Antimigratory Activities", Journal of Fungi, 2022

Publication

23	Submitted to University of the Philippines - Main Library Student Paper	1 %
24	eprints.unsri.ac.id Internet Source	1 %
25	faculty.uobasrah.edu.iq Internet Source	1 %
26	peerj.com Internet Source	1 %
27	res.mdpi.com Internet Source	1 %
28	www.j3.jstage.jst.go.jp Internet Source	1 %
29	www.tjnpr.org Internet Source	1 %
30	Submitted to Universidade do Porto Student Paper	1 %
31	www.intechopen.com Internet Source	1 %
32	Elfita Elfita, Muharni Muharni, Mardiyanto Mardiyanto, Fitriya Fitriya, Feti Fera, Hary	<1 %

Widjajanti. "Antibacterial Activity of Traditional Medicine *Scurrula atropurpurea* (BL) DANS and their Endophytic Fungi", Key Engineering Materials, 2020

Publication

33

www.jeeng.net

Internet Source

<1 %

34

etd.repository.ugm.ac.id

Internet Source

<1 %

35

www.hindawi.com

Internet Source

<1 %

36

www.thepab.org

Internet Source

<1 %

37

Sushma Mishra, Priyanka, Shilpi Sharma.
"Metabolomic Insights Into Endophyte-Derived Bioactive Compounds", Frontiers in Microbiology, 2022

Publication

<1 %

38

jcreview.com

Internet Source

<1 %

39

revistabionatura.com

Internet Source

<1 %

40

www.ikprpress.org

Internet Source

<1 %

41

K. Tayung, D. K. Jha. "Antimicrobial endophytic fungal assemblages inhabiting bark of Taxus

<1 %

baccata L. of Indo-Burma mega biodiversity hotspot", Indian Journal of Microbiology, 2010

Publication

42

science.yru.ac.th

Internet Source

<1 %

43

Li Feng, Jia Wang, Shuai Liu, Xue-Jia Zhang, Qi-Rui Bi, Yan-Yun Hu, Zhe Wang, Ning-Hua Tan.

" Colletopeptides A–D, Anti-inflammatory Cyclic Tridepsipeptides from the Plant

Endophytic Fungus sp. S8 ", Journal of Natural Products, 2019

Publication

<1 %

44

Muharni Muharni, Julinar Julinar, Heni Yohandini, Fitrya Fitrya, Rima Melati.

"Antihyperlipidemia activity of ethanol extract the stem bark of Flacourtia rukam on propylthiouracil-induced albino rats, Rattus noverticus (Wistar strain)", IOP Conference Series: Materials Science and Engineering, 2021

Publication

<1 %

45

Submitted to Universitas Diponegoro

Student Paper

<1 %

46

irep.iiium.edu.my

Internet Source

<1 %

47

Li Pan, Ulyana Muñoz Acuña, Jie Li, Nivedita Jena et al. " Bioactive Flavaglines and Other

<1 %

Constituents Isolated from ", Journal of Natural Products, 2013

Publication

48

Srichandan Padhi, Kumananda Tayung. "In vitro antimicrobial potentials of endolichenic fungi isolated from thalli of Parmelia lichen against some human pathogens", Beni-Suef University Journal of Basic and Applied Sciences, 2015

Publication

<1 %

49

www.finechem-mirea.ru

Internet Source

<1 %

50

H Widjajanti, Muharni, E Nurnawati, V Triuspita. "The potency of endophytic fungi isolated from Hippobroma longiflora (L) G. Don as an antioxidant sources", IOP Conference Series: Earth and Environmental Science, 2022

Publication

<1 %

51

H. J. Ridgway. "Detection of botryosphaeriaceous species in environmental samples using a multi-species primer pair : Multi-species primers to amplify botryosphaeriaceous species", Plant Pathology, 12/2011

Publication

<1 %

52

Xiao-Shan Shi, Ling-Hong Meng, Xiao-Ming Li, Xin Li, Dun-Jia Wang, Hong-Lei Li, Xing-Wang

<1 %

Zhou, Bin-Gui Wang. " Trichocadinins B–G: Antimicrobial Cadinane Sesquiterpenes from QA-8, an Endophytic Fungus Obtained from the Medicinal Plant ", Journal of Natural Products, 2019

Publication

53

www.freepatentsonline.com

Internet Source

<1 %

54

Submitted to Institute of Graduate Studies,
UiTM

Student Paper

<1 %

55

Submitted to Turun yliopisto

Student Paper

<1 %

56

T. T. L. Tra, N. M. Chi, D. T. K. Anh, P. Q. Thu,
N. P. Nhung, B. Dell. "Bacterial endophytes
from Chukrasia tabularis can antagonize
Hypsipyla robusta larvae", Phytoparasitica,
2022

Publication

<1 %

57

media.neliti.com

Internet Source

<1 %

58

Maria L. Gonzalez-Rivera, Flavio Martinez-
Morales, Angel J. Alonso-Castro, Juan F.
Lopez-Rodriguez et al. "Validated and rapid
measurement of the ferric reducing
antioxidant power in plasma samples",
Chemical Papers, 2018

Publication

<1 %

59

www.banglajol.info

Internet Source

<1 %

60

Arakkaveettil Kabeer Farha, Abdulla Mohamed Hatha. " Bioprospecting potential and secondary metabolite profile of a novel sediment-derived fungus sp. ArCSPf from continental slope of Eastern Arabian Sea ", *Mycology*, 2019

Publication

<1 %

61

I Hairunisa, I A Mentari, T Julianti, E R Wikantyasning, Z Choliso, S C Ningsih, M R F Muslim. "Antioxidant Activities in Different Parts of Pulasan (*Nephelium mutabile* Blume) from East Borneo", *IOP Conference Series: Earth and Environmental Science*, 2021

Publication

<1 %

62

chemrj.org

Internet Source

<1 %

63

pustaka.unpad.ac.id

Internet Source

<1 %

64

rebus.us.edu.pl

Internet Source

<1 %

65

Luca Elli, Donatella Barisani, Valentina Vaira, Maria Teresa Bardella et al. "How to manage celiac disease and gluten-free diet during the COVID-19 era: proposals from a tertiary

<1 %

referral center in a high-incidence scenario",
BMC Gastroenterology, 2020

Publication

66

academica-e.unavarra.es

Internet Source

<1 %

67

www.ejournal.radenintan.ac.id

Internet Source

<1 %

68

Submitted to De La Salle University

Student Paper

<1 %

69

Souvik Kusari, Sebastian Zühlke, Ján Košuth, Eva Čellárová, Michael Spiteller. " Light-Independent Metabolomics of Endophytic Provides Insight into Microbial Hypericin Biosynthesis ", Journal of Natural Products, 2009

Publication

<1 %

70

www.acgpubs.org

Internet Source

<1 %

71

www.atlantis-press.com

Internet Source

<1 %

72

www.ukm.my

Internet Source

<1 %

73

H. J. Park, H. K. Jeong, M. I. Chang, M. H. Im et al. "Structure determination of new analogues of vardenafil and sildenafil in dietary

<1 %

supplements", Food Additives and Contaminants, 2007

Publication

74

Submitted to KYUNG HEE UNIVERSITY

Student Paper

<1 %

75

Ranjani S, Janani Prabha G, Karunya J R, Hemalatha S. "Differential actions of nanoparticles and nanoemulsion synthesized from Colletotrichum siamense on food borne pathogen", LWT, 2022

Publication

<1 %

76

Si-Xian WANG, Rui-Lin ZHAO, Cui GUO, Bao-Song CHEN, Huan-Qin DAI, Gao-Qiang LIU, Hong-Wei LIU. "New meroterpenoid compounds from the culture of mushroom Panus lecomtei", Chinese Journal of Natural Medicines, 2020

Publication

<1 %

77

bmccomplementmedtherapies.biomedcentral.com

Internet Source

<1 %

78

ejbpc.springeropen.com

Internet Source

<1 %

79

www.igi-global.com

Internet Source

<1 %

80

Fábio de Azevedo Silva, Rhavena Graziela Liotti, Ana Paula de Araújo Boleti, Érica de Melo Reis et al. "Diversity of cultivable fungal

<1 %

endophytes in *Paullinia cupana* (Mart.) Ducke and bioactivity of their secondary metabolites", PLOS ONE, 2018

Publication

81

George R. Pettit, Sylvie Ducki, Rui Tan, Roberta S. Gardella et al. " Isolation and Structure of Pedilstatin from a Republic of Maldives sp. ", *Journal of Natural Products*, 2002

Publication

82

Hong-Tao Li, Hao Zhou, Rong-Ting Duan, Hong-Yu Li, Lin-Huan Tang, Xue-Qiong Yang, Ya-Bin Yang, Zhong-Tao Ding. " Inducing Secondary Metabolite Production by Co-culture of the Endophytic Fungus sp. and the Symbiotic Fungus sp. ", *Journal of Natural Products*, 2019

Publication

83

Jin-Gyeong Cho, Na-Young Song, Tae-Gyu Nam, Sabina Shrestha et al. " Flavonoids from the Grains of C1/R-S Transgenic Rice, the Transgenic *Oryza sativa* spp. , and Their Radical Scavenging Activities ", *Journal of Agricultural and Food Chemistry*, 2013

Publication

84

Kushan M. Ukwatta, Jennifer L. Lawrence, C. D. Wijayarathna. " The study of antimicrobial, anti-cancer, anti-inflammatory and α -glucosidase inhibitory activities of

<1 %

<1 %

<1 %

<1 %

Nigronaphthaphenyl, isolated from an extract of ", Mycology, 2019

Publication

85

Martin Muthee Gakuubi, Kuan Chieh Ching, Madhaiyan Munusamy, Mario Wibowo et al. "Enhancing the Discovery of Bioactive Secondary Metabolites From Fungal Endophytes Using Chemical Elicitation and Variation of Fermentation Media", *Frontiers in Microbiology*, 2022

Publication

<1 %

86

Mónica Flores-Reséndiz, Patricia Lappe-Oliveras, Martha Lydia Macías-Rubalcava. "Mitochondrial damage produced by phytotoxic chromenone and chromanone derivatives from endophytic fungus *Daldinia eschscholtzii* strain GsE13", *Applied Microbiology and Biotechnology*, 2021

Publication

<1 %

87

berkalahayati.org

Internet Source

<1 %

88

repository.unair.ac.id

Internet Source

<1 %

89

rsdjournal.org

Internet Source

<1 %

90

www.science.gov

Internet Source

<1 %

91

Joo, Se-Jin, Hee-Jung Park, Ji-Hae Park, Jin-Gyeong Cho, Ji-Hyun Kang, Tae-Sook Jeong, Hee Kang, Dae-Young Lee, Hack-Soo Kim, Sang-Yo Byun, and Nam-In Baek. "Flavonoids from *Machilus japonica* Stems and Their Inhibitory Effects on LDL Oxidation", *International Journal of Molecular Sciences*, 2014.

Publication

<1 %

92

"Medically Important Plant Biomes: Source of Secondary Metabolites", Springer Science and Business Media LLC, 2019

Publication

<1 %

93

Alfinda Novi Kristanti, Ei Ei Aung, Nanik Siti Aminah, Yoshiaki Takaya, Hnin Thanda Aung, Rico Ramadhan. " Bioactive triterpenoids from Indonesian medicinal plant ", *Open Chemistry*, 2022

Publication

<1 %

94

Kazuyoshi Kawazoe, Naoko Shimogai, Yoshihisa Takaishi, K.Sundar Rao, Yasuhiro Imakura. "Four stilbenes from *Salacia lehmbachii*", *Phytochemistry*, 1997

Publication

<1 %

95

SooWhan Choi, Jae Wook Lee, Young Ho Ko, Kimoon Kim. "Pseudopolyrotaxanes Made to Order: Cucurbituril Threaded on Polyviologen", *Macromolecules*, 2002

<1 %

96

Tao Xu, Mohammed Gherib, Chahrazed Bekhechi, Fewzia Atik-Bekkara et al. " Thymyl esters derivatives and a new natural product modhephanone from Coss (Asteraceae) root oil ", Flavour and Fragrance Journal, 2015

<1 %

Publication

Exclude quotes On

Exclude matches Off

Exclude bibliography Off