

Preparation and FTIR-ATR combined with chemometrics analysis of selfemulsifying loaded sungkai extract from *Peronema canecens*

by Shaum Shiyan

Submission date: 12-May-2023 03:48PM (UTC+0700)

Submission ID: 2091194092

File name: iyan-Q2_RJPT_15_1_79-85_Preparation_and_FTIR-ATR_combination.pdf (329.65K)

Word count: 5007

Character count: 26071

RESEARCH ARTICLE

Preparation and FTIR-ATR combined with chemometrics analysis of self-emulsifying loaded sungkai extract from *Peronema canecens*

Shaum Shiyani^{1,2,*}, Nafisah Ramadona¹, Widea Fitri Utami^{1,2}, Natascha Depriyanti^{1,2}, Anazir Mukafi¹, Wanda Novianndhani^{1,2}

¹Department of Pharmacy, Faculty of Mathematics and Natural Sciences, Universitas Sriwijaya, Indralaya (OI) Sumatera Selatan 30662, Indonesia.

²Phytopharmaceutical Reseach Center (PRC) at Laboratory of Pharmaceutical Biology, Department of Pharmacy, Faculty of Mathematics and Natural Sciences, Universitas Sriwijaya, Indralaya (OI) Sumatera Selatan 30662, Indonesia.

*Corresponding Author E-mail: shaumshiyani@unsri.ac.id

ABSTRACT:

The use of immunomodulators is one strategy in maintaining the immune system during the Covid-19 pandemic. Sungkai leaf extract from *Peronema canecens* keeps the immune system in good shape. Therefore, in this study, we formulated a self-emulsifying loaded sungkai leaves extract (SE-SLE) with oleic acid and virgin coconut oil (VCO) oil phases, span 80 and tween 80 as surfactants and co-surfactants in the form of PEG-400 and PG. Chemometric analysis was conducted by observing the typical pattern in each FTIR-ATR spectra. The pattern is divided into several groups based on the wavenumber and analyzed using principal component analysis (PCA) to identify the compounds contained therein. Grouping based on chemical properties via IR spectra on SE-SLE resulted in two large groups. The results obtained are beneficial as initial information in developing and optimizing the self-nano emulsifying drug delivery system formula.

KEYWORDS: Chemometrics, FTIR-ATR, *Peronema canecens*, Self-emulsifying, Emulsion, Sungkai, Principal component analysis, cluster analysis.

INTRODUCTION:

The immune system has an essential role in the Covid-19 pandemic situation. The immune system is intended to protect the body from infection by producing protein molecules (antibodies) that bind to antigens¹. One of the plants believed to act as an immunomodulator is sungkai (*Peronema canecens* Linn.). An immunomodulatory is any substance that can increase immunity². Sungkai leaf extract contains phenolic, tannin, steroid, saponin, and flavonoid activities as antioxidant, antipyretic, antiplasmodial and can increase the immune system^{2,3,4}.

Phytochemical compounds such as alkaloids and flavonoids act as immunomodulators by increasing the activity of interleukin 2 (IL-2) and lymphocyte proliferation⁵. Local people usually use the sungkai leaves directly by boiling them and making them fresh vegetables. This direct use method has several drawbacks: it cannot be stored for more prolonged time consumption and less stable. In addition, the phytochemical compounds are still not specific but using the medicinal plants as herbal immunomodulators have positive effects such as reducing the side effects of using conventional chemical compound⁶. Therefore, the self-emulsifying formulation of sungkai leaf extract is an exciting development innovation.

Self-emulsifying is a delivery system design in an isotropic mixture of oil, surfactant, co-surfactant, and extract. The formulation forms an emulsion spontaneously after being introduced into the aqueous phase by peristaltic motion, producing mild agitation in the gastrointestinal tract⁷. Self-emulsifying can increase

oral absorption and dissolution speed so that bioavailability and stability can be increased⁸. In addition, it can reduce the dose and frequency of drug administration when it is formulated with self-emulsifying⁹. This carrier system can be formulated with various oils, surfactants, and co-surfactants. Therefore, preliminary studies related to the composition of the constituents are needed to determine the grouping of formulas based on their constituent components. Self-emulsifying drug delivery systems are classified as superior methods for oral drug delivery¹⁰.

This research formulates self-emulsifying with several different formula compositions. The types of oil used were oleic acid and virgin coconut oil (VCO), surfactants in the form of tween 80 and span 80, co-surfactants selected PEG 400, and propylene glycol. Combining these different constituent components provides a different pattern of interactions with the phytochemical compounds present in the extract. Therefore, it is exciting to evaluate using the FTIR-ATR spectra pattern with chemometric analysis. The chemometric analysis approach aims to classify and study the correlation of the responses generated. The chemometric approach is carried out by multivariate statistical modeling in data processing and evaluation and interpretation with much data¹¹⁻¹³.

MATERIAL AND METHODS:

Materials and Chemicals:

Sungkai leaves (*Peronema canescens*) were obtained from Sekayu, Musi Banyuas¹¹ Sumatera Selatan. Ingredients such as oleic acid oil, virgin coconut oil (VCO), span 80, tween 80, PEG-400, and PG were purchased from Bratachem (Jakarta, Indonesia). Solvents such as alcohol, aqua pro injection, and distilled water were obtained from Embacang (Palembang, Indonesia).

Prepara¹⁶ of sungkai leaf extract (SLE):

Sungkai leaves were dried in the sun covered with black cloth for 3 x 24 hours. Dried leaves are converted into powder by a gr²²ng process. Sungkai leaf powder was extracted using the ultrasound-assisted extraction (UAE) method with 70% ethanol solvent in a ratio of 1:10.¹⁴

Preparation of self-emulsifying loaded sungkai leaf extract (SE-SLE):

A total of 16 formulas were designed using the fractional factorial design method¹⁵. Self-emulsifying (SE) preparation was started by dissolving the extract with a carrier oil, then vortexed and ultrasonicated for 5 minutes at room temperature. Then added surfactant and co-surfactant in the oil-extract solution and vortexed until homogeneously mixed.

Evaluation of SE-SLE:

The SE-SLE viscosity was determined at 25-30 °C using Ostwald viscometer instrumentation¹³. The emulsification time was measured by adding SE-SLE in 500 mL of distilled water at a temperature of 37 °C using a magnetic stirrer with a speed of 120 rpm¹⁶. Density was measured with a pycnometer by weighing an empty pycnometer using an analytical balance, filling the pycnometer with microemulsion preparations to the brim, and weighing with an analytical balance¹⁷. SE-SLE pH¹³ termination uses a universal pH indicator. A total of 10 µL of the sample was dropped evenly on the surface of the universal pH indicator. The color pattern formed is matched with the standard pH color that has been provided in the pH indicator box. The uniform color pattern indicates the pH of SE-SLE¹⁸.

Clarity using spectrophotometer:

Self-emulsifying nanoemulsion of Sungkai leaf extract was taken 1 mL and diluted 100 times using distilled water. The transmission percentage was measured with a UV-Vis spectrophotometer Biobase BK-UV1000 (Shandong, China) at 638 nm with distilled water as blank¹⁹.

FTIR-ATR fingerprinting:

The analysis was carried out by observing the interaction b²⁰een the constituent materials of SE-SLE qualitatively using Fourier transform infrared-attenuated total reflectance (FTIR-ATR) fingerprinting. The FTIR-ATR uses the Nicolet iS10 series instrumentation (Thermo Scientific, USA) equipped with Omnic software. The FTIR-ATR spectrum was analyzed based on the vibrations of the functional groups of each component of the material. The analysis was carried out at 4000 cm⁻¹ to 500 cm⁻¹ with three times replication²⁰. ATR crystal has to be cleaned with ethanol p.a before used to analyze the sample for minimize the noise²¹.

Chemometrics analysis:

¹⁵a analysis used a chemometric approach with principal component analysis (PCA) and cluster analysis (CA) methods. The PCA-CA method was processed using Minitab^{12,13}.

RESULT:

SE-SLE was designed into 16 formulas using various carrier oils in the form of oleic acid and VCO, variations of surfactants tween 80 and span 80, and co-surfactants with types of PEG-400 and PG. Each formula is composed of different components and different concentrations. SE-SLE preparation was performed by mixing the oil and extract until homogeneous using a vortex at a constant speed. The next process is sonication with an ultrasonicator which aims to reduce the particle size and homogenize the self-emulsifying

mixture so that its solubility can increase²². Visualization of SE-SLE can be seen in Figure 1. Each formula has a different color starting from light yellow to brownish-yellow. These different colour influenced by the type of oil, the amount of extract and the concentration of each component. Self-emulsifying has light yellow color indicated by F1, F4, F6, F8, F13, and F15, a formula with the dark yellow color indicated by F2, F5, F7, F10, F12, and F14, while F3, F11, and F16 are yellow-brown.




Figure 1. SE-SLE visualization from formulas 1 to 16

The self-emulsifying characterization of sungkai leaves included viscosity tests, specific gravity, pH, and transmittance (Table 1). The viscosity in this study is related to the ability of the SE-SLE formula to flow. The viscosity is influenced by several factors such as molecular size, solution concentration, attractive intermolecular forces, and temperature. Viscosity measurements on self-emulsifying sungkai leaves showed that F12, F7, and F8 were the three formulas with the largest viscosity values. These three formulas were surfactants span 80 (F7 and F12) and tween 80 (F8). The co-surfactant used is propylene glycol (PG).

Table 1. Evaluation results of SE-SLE and the emulsion formed

Formula	Oil	Surfactant	Co-surfactant	Color	Viscosity (N/m ²)	Density (g/cm ³)	pH	Clarity (%T)
1	Oleic acid	Span 80	PEG-400	Light yellow	1.76	0.62	5	66.40
2	VCO	Tween 80	PG	Dark yellow	1.23	0.43	6	46.10
3	VCO	Tween 80	PG	Brownish yellow	1.38	0.29	6	82.30
4	Oleic acid	Tween 80	PEG-400	Light yellow	0.53	0.56	5	73.30
5	Oleic acid	Span 80	PG	Dark yellow	0.38	0.20	6	87.20
6	Oleic acid	Span 80	PG	Light yellow	0.49	0.52	6	47.90
7	VCO	Span 80	PG	Dark yellow	1.95	0.69	6	86.10
8	Oleic acid	Tween 80	PG	Light yellow	1.93	0.68	6	41.90
9	VCO	Tween 80	PEG-400	Light yellow	0.76	0.80	6	92.20
10	VCO	Span 80	PEG-400	Dark yellow	0.41	0.43	6	90.80
11	VCO	Span 80	PEG-400	Brownish yellow	0.35	0.37	5	78.30
12	VCO	Span 80	PG	Dark yellow	2.07	0.43	6	19.30
13	Oleic acid	Span 80	PEG-400	Light yellow	0.54	0.57	6	73.70
14	Oleic acid	Tween 80	PEG-400	Dark yellow	0.94	0.25	6	80.30
15	Oleic acid	Tween 80	PG	Light yellow	1.73	0.60	6	88.70
16	VCO	Tween 80	PEG-400	Brownish yellow	2.84	0.49	6	92.00

21

The type of surfactant and co-surfactant used to affect the viscosity of a self-emulsifying agent.

Observation of transmittance of SE-SLE at a wavelength of 638 nm shows a formula with high transmittance, namely F9, F16, F10. Emulsions with a transmittance value close to distilled water, namely 100, indicate that the emulsion is getting clearer and has smaller droplets with an estimated size of 10-200 nm²³. The pH value ranges from 5-6, which does not follow the intestinal pH requirements of 6.8, and the stomach has a pH of 2.0²⁴.

Spectra of FTIR-ATR:

The FTIR data is processed based on the resulting pattern to obtain data in numbers. Absorption regions with peaks will be grouped by wavenumber and analyzed using chemometrics. The obtained vibrations are then read to determine the functional groups present at the peak of the spectra. Fundamental vibrations are generally in the 4000-2500 cm⁻¹ area with O-H, C-H, and N-H stretching clusters. Absorption in the area with a wavenumber of 2500-2000 cm⁻¹ indicates a triplicate group. The bands in the 2000-1500 cm⁻¹ area are caused by C=C and C=O stretching.

Spectra data obtained from 16 formulas that have been analyzed based on certain wavenumbers are shown in Table 2. Infrared spectra in 16 formulas show medium-intensity absorption in the 3580-3207 region, indicating O-H functional groups (free alcohol). The second absorption is at wavenumber 3001-2953 cm⁻¹ with a strong intensity indicating the C-H group of stretched alkane. C-H absorption is also found in wave numbers 2890-2811 cm⁻¹, with weak intensity indicating the presence of aldehydes. Absorption at wavenumbers 858-816 cm⁻¹ with strong intensity indicates aromatic groups (bend, out of plane).

Table 2. Wavenumbers and functional groups based on FTIR-ATR spectra from SE-SLE

Functional groups	Wavenumbers (cm ⁻¹)	Type of vibration	Intensity
O-H	3580 – 3207	Alcohol (free)	Medium
C-H	3001-2953	Alkane (stretch)	Strong
	2890-2811	Aldehyde	Weak
	858-816	Aromatic (bend, out of plane)	Strong
C ≡ N	2379-2314	Nitrile	Medium
C=O	1750-1733	Ester	Strong
	1724-1702	Ketones	Strong
N=O	1475-1428	Nitro	Strong
C-O	1269-1222	Alcohols, ethers, esters, carboxylic acids, anhydrides	Strong
	1133-1001		
C-X	739-696	Chloride	Strong

The nitrile group C N was obtained at wavenumber 2379-2314 cm⁻¹ with medium intensity. Wavenumbers 1750-1733 cm⁻¹ indicate an ester group with a strong intensity, and wavenumbers 1724-1702 cm⁻¹ indicate a ketone group with strong intensity. The band in wavenumbers 1475-1428 cm⁻¹ indicates a strong intensity nitro group. The absorption in the 1269-1001 cm⁻¹ region indicated the presence of alcohol, ether, ester, carboxylic acid, anhydride groups of strong intensity, and absorption at 739-696 cm⁻¹ indicated the presence of a strong chloride group.

Chemometrics analysis:

Figure 3 shows the results of PCA and CA from the overall absorbance data of the FTIR-ATR spectra on the SE-SLE sample. Chemometric analysis using PCA resulted in a score plot (Figure 3a), scree plot (Figure 3b), loading plot (Figure 3c), and dendrogram (Figure 3d) of CA. The score plot shows the classification of formulas grouped at different distances from each other. The score plot results show that 15 of the 16 existing formulas have similarities. The score plot analysis on

PCA showed that there were two different groups. Group A has many groups, and group B consists of F15 only. The close distance between group A evidences this, while group B has a considerable distance from the other formula points. The further apart a sample is in the score plot analysis, the less similarity of the sample ²⁵.




Figure 2. FTIR-ATR Spectra of SE-SLE(1) F1, (2) F2, (3) F3, (4) F4, (5) F5, (6) F6, (7) F7, (8) F8, (9) F9, (10) F10, (11) F11, (12) F12, (13) F13, (14) F14, (15) F15, (16) F16

Table 3. Absorbance value at selected wavenumber of SE-SLE

Formula	Absorbance at wavenumber										
	1	2	3	4	5	6	7	8	9	10	11
1	0.021	0.035	0.163	0.068	0.044	0.108	0.093	0.055	0.054	0.045	0.047
2	0.027	0.160	0.127	0.160	0.070	0.180	0.064	0.055	0.060	0.053	0.055
3	0.078	0.118	0.097	0.073	0.075	0.302	0.110	0.116	0.065	0.103	0.101
4	0.030	0.138	0.125	0.083	0.066	0.253	0.078	0.071	0.081	0.069	0.074
5	0.059	0.124	0.090	0.065	0.054	0.148	0.066	0.069	0.091	0.078	0.063
6	0.040	0.118	0.087	0.054	0.050	0.131	0.064	0.065	0.071	0.053	0.050
7	0.043	0.169	0.120	0.151	0.077	0.180	0.071	0.070	0.087	0.073	0.061
8	0.023	0.130	0.119	0.083	0.058	0.237	0.084	0.072	0.069	0.060	0.047
9	0.020	0.149	0.133	0.068	0.057	0.134	0.126	0.059	0.054	0.050	0.032
10	0.021	0.135	0.156	0.040	0.145	0.077	0.167	0.109	0.055	0.047	0.044
11	0.043	0.157	0.129	0.045	0.098	0.091	0.089	0.075	0.083	0.087	0.087
12	0.021	0.035	0.163	0.068	0.044	0.108	0.093	0.055	0.054	0.045	0.047
13	0.027	0.160	0.127	0.160	0.070	0.180	0.064	0.055	0.060	0.053	0.055
14	0.078	0.118	0.097	0.073	0.075	0.302	0.110	0.116	0.065	0.103	0.101
15	0.030	0.138	0.125	0.083	0.066	0.253	0.078	0.071	0.081	0.069	0.074
16	0.059	0.124	0.090	0.065	0.054	0.148	0.066	0.069	0.091	0.078	0.063




Figure 3. Result of chemometric analysis using PCA method *self-emulsifying sungkai leaves extract*, (A) Score plot, (B) Scree Plot, (C) Loading Plot, (D) Dendrogram

DISCUSSION:

The analysis of SE-SLE spectra with FTIR-ATR aims to determine the vibration of functional groups and interactions between spectra in each formula. FTIR spectra were read at wavenumbers of 4000 cm^{-1} to 400 cm^{-1} ¹⁷. A total of 16 self-emulsifying formulas of sungkai leaf were tested using FTIR-ATR, with each formula being run for three replications. This replication aims to obtain more accurate data based on similarities or differences in peaks, wavenumbers, and absorbance of each run. The peak at a certain wave number will be shown by FTIR spectra based on the vibration of the functional group of a material in the formula.

Chemometric analysis was carried out by observing the typical pattern in each FTIR spectra to identify a compound conveniently. A distinctive pattern is what distinguishes one compound from another²⁶. The resulting FTIR spectra pattern will be divided into certain groups based on the wavenumber for further analysis using PCA chemometric method. Principal

component analysis (PCA) is used to process multivariate data with unknown samples to simplify the uncertain elements by reducing their dimensions²⁷. PCA analysis was conducted to determine the characteristics and characteristics of each formula and evaluate the degree of similarity^{21,22}. The chemometric analysis method was carried out by grouping all the formulas into several small groups based on the similarity of characteristics and the closeness of the responses possessed by each formula. Chemometrics becomes an important method for analysing active compounds in pharmaceutical formulations²⁸. Chemometric analysis can work as a new analytical tool for many studies, combining formulas with the desired parameters²⁹.

The dendrogram in PCA analysis shows the formula group with the same variable and has a bond in a group based on the similarity of the value³⁰. The results of the calculation of the dendrogram similarity are shown in Figure 3d. SE-SLE formulas are grouped based on their similarity. F2 and F7 have a similarity of 89.95%, F4

and F8 91.27%, F1 and F11 71.86%, F13 and F14 84.79%. The loading plot determines the formula that has the most role in forming the principal component³¹. The loading plot analysis describes the angle that shows the correlation between the responses in the 16 formulas.

17 SE-SLE is an isotropic mixture of oil, surfactant, and cosurfactant that spontaneously forms nanoemulsions when mixed with water. Surfactants and cosurfactants that can mix well with the oil phase will increase the formation of the nanoemulsion system. Surfactants and cosurfactants in the nanoemulsion system work together to form a good and flexible interface system and reduce the surface tension value to near zero to support the formation of stable nano-sized globules³¹. SE-SLE is designed in several 16 formulas with eight variations of the constituent components, namely from oil, surfactant, and co-surfactant. The differences in the composition of the SE-SLE make each formula have differences both visually and the results of characteristic testing. The carrier oils used in preparing SE-SLE were oleic acid and VCO. The surfactants used were tween 80 and span 80 and co-surfactants PG and PEG-400.

VCO belongs to the group of triglycerides with medium-chain fatty acids and is an oil suitable for manufacturing nanoemulsions. SE-SLE made with VCO carrier oil showed a light yellow color with surfactant components such as Tween 80 and co-surfactant PEG-400. PG and PEG are organic solvents suitable for oral delivery, which in some range concentrations can be hydrophilic surfactant^{32,33}. This phenomenon indicates good results visually and is evidenced by the highest transmittance value of 92.20. Using VCO as a carrier oil with tween 80 as a surfactant and PEG-400 as a co-surfactant can produce nanoemulsions with nanometer particle size²⁵. Solubility of the drug in the oil phase is influential in the formulation of self-emulsifying and affects the process of absorption in the gastrointestinal area. Each oil in the formula has different solubility and physicochemical properties^{32,34}.

CONCLUSION:

FTIR-ATR spectra and chemometric analysis were successfully applied in the initial evaluation of the self-emulsifying formula. There are two major groups based on the similarity of the FTIR-ATR spectra pattern. Formula 15 into its separate group.

ACKNOWLEDGEMENT:

This study was supported by Program Kreativitas Mahasiswa – Riset Eksakta (PKM-RE) 2021, Ministry of Education, Culture, Research, and Technology, Directorate General of Higher Education, Research, and Technology of Republic Indonesia No. 99/E2/KM/2021

SPP Number 0043/BELMAWA/690437/2021. The author is grateful, and this research is facilitated by the Laboratory of Pharmaceutical Biology, Faculty of Mathematics and Natural Sciences, Universitas Sriwijaya. Thanks to the Phytopharmaceutical Research Center (PRC) and LP2M Universitas Sriwijaya.

24 CONFLICT OF INTEREST:

The authors declare no conflict of interest.

REFERENCES:

1. Nilashi, M. Saminah S. Leila S. Hossein A. Elnaz A. Tarik A. The Covid-19 infection and the immune system: The Role of complementary and alternative medicines. *Biomedical Research*. 2020; 30 (2).
2. Khokra SL, Parashar B, Dhamija HK, Bala M. Immunomodulators: Immune system modifiers. *Research Journal of Pharmacy and Technology*. 2012; 5(2): 169-174.
3. Okfrianti Y, Irmameria D, Bertalina. Antioxidant activity of sungkai leaf (*Peronema canescens* Jack) ethanol extract. 2022; 13(2): 333-339.
4. Yamin R, Sartinah A, Ihsan S, Kasmawati H, Suryani, Andriyani R, Asma, Adjeng ANT, Arba M. Radical scavenging assay and determination Flavonoid and Phenolic total of extract and Fractions of Raghu bark (*Dracontomelon dao* (Blanco) Merr). *Research Journal of Pharmacy and Technology*. 2020; 13(5): 2335-2339. <https://doi.org/10.5958/0974-360X.2020.00420.5>
5. Dillasamola D, Aldi Y, Wahyuni FS, Rita RS, Dachriyanus, Salman Umar S, Rivai H. Study of sungkai (*Peronema canescens* Jack) leaf extract activity as an immunostimulators with in vivo and in vitro methods. *Pharmacognocny Journal*. 2021; 13(6): 1397-1407. <https://doi.org/10.5530/pj.2021.13.177>
6. Debnath S, Chakravorty R, Devi D. A review on role of medicinal plants in immune system. *Asian Journal of Pharmacy and Technology*. 2020; 10(4): 273-277. <https://doi.org/10.5958/2231-5713.2020.00045.8>
7. Abbas A, Lichtman AH, Pillai S. Cellular and Molecular Immunology. WB Saunders Company Philadelphia.2007; 6thEd.
8. Abd-Elhakeem E, Teaima MHM, Abdelbary GA, El Mahrouk GM. Bioavailability enhanced clopidogrel-loaded solid SNEDDS: Development and in-vitro/in-vivo characterization. *Journal of Drug Delivery Science and Technology*, 2019; 49: 603–614. <https://doi.org/10.1016/j.jddst.2018.12.027>
9. Makadia HA, Bhatt AY, Parmar RB, Paun JS, Tank HM. Self-nano emulsifying drug delivery system (SNEDDS): Future aspects. *Asian Journal of Pharmaceutical Research*. 2013; 3(1): 21–27.
10. Gurudutta P, Jeetesh P, Sajid AM, Tahir AM. Self-emulsifying drug delivery systems: And attempt to improve oral absorption of poorly Soluble Drugs. *Research Journal of Pharmaceutical Dosage Forms and Technology*. 2010; 2(3): 206-214.
11. Chaudhary S, Aqil M, Sultana Y, Kalam MA. Self-nanoemulsifying drug delivery system of nabumetone improved its oral bioavailability and anti-inflammatory effects in rat model. *Journal of Drug Delivery Science and Technology*. 2019; 51: 736–745. <https://doi.org/10.1016/j.jddst.2018.04.009>.
12. Shiyani S, Arifin A, Amriani A, Pratiwi G. Immunostimulatory activity of ethanol extract from *Calotropis gigantea* L. flower in rats against *Salmonella typhimurium* infection. *Research Journal of Pharmacy and Technology*. 2020; 13(11): 5244-5250. <https://doi.org/10.5958/0974-360X.2020.00917.8>
13. Pratiwi G, Ramadhiani AR, Shiyani S. Understanding the combination of fractional factorial design and chemometrics analysis for screening super-saturable quercetin-self nano emulsifying components. *Pharmacia*. 2022; 69(2): 273–284. <https://doi.org/10.3897/pharmacia.69.e80594>.
14. Shafii NZ, Saudi ASM, Pang JC, Abu IF, Sapawe N, Kamarudin

- MK. Saudi SFM. Application of chemometrics techniques to solve environmental issues in Malaysia. *Heliyon*. 2019; 5(10): e02534. <https://doi.org/10.1016/j.heliyon.2019.e02534>
15. Shiyani S. Hertiani T. Martien R. Nugroho AK. Optimization of a novel kinetic-assisted infundation for rich-EGCG and polyphenols of white tea (*Camellia Sinensis*) using central composite design. *International Journal of Applied Pharmaceutics*. 2018; 10(6): 259-267. <https://doi.org/10.22159/ijap.2018v10i6.29654>
 16. 7. tiwi G. Susanti S. Shiyani S. Application of factorial design for optimization of PVC-HPMC polymers in matrix film ibuprofen patch-transdermal drug delivery system. *Indonesian Journal of Chemometrics and Pharmaceutical Analysis*. 2020; 1(1): 11-22. <https://doi.org/10.2255/146/jcpa.486>.
 17. 8. Maryatno P. Chabib L. Hayati F. Awaluddin R. Stability study of Ipomoea reptans extract self-nanoemulsifying drug delivery system (SNEDDS) as anti-diabetic therapy. *Journal of Applied Pharmaceutical Science*. 2018; 8(9): 11-14. <https://doi.org/10.7324/japs.2018.8903>.
 18. Zhao T, Maniglio D, Chen J, Chen B, Motta A, Migliaresi C. Self-nanoemulsifying drug delivery systems (SNEDDS) for the oral delivery of lipophilic drugs. *Nanotechnology*. 2015; 26(12). <http://dx.doi.org/10.1088/0957-4484/26/12/125102>
 19. Patmayuni D. Sulaiman TNS. Zulkarnain AK. Shiyani S. Method validation of simvastatin in PCL-PEG-PCL triblock copolymer micelles using UV-23 spectrophotometric for solubility enhancement assay. *International Journal of Applied Pharmaceutics*. 2022; 14(1): 246-250. <https://doi.org/10.22159/ijap.2022v14i1.42961>.
 20. Fatmarahmi, Dhatmastuti C, Ratna A, Susudarti. Identification and quantification of metamizole in traditional herbal medicines using spectroscopy FTIR-ATR combined with chemometrics. *Research Journal of Pharmacy and Technology*. 2020; 14(8): 4413-4419. <https://doi.org/10.52711/0974-360X.2021.00766>.
 21. González-Domínguez R, Sayago A, Fernández-Recamales Á. An overview on the application of chemometrics tools in food authenticity and traceability. *Foods*. 2022; 11(23): 3940. <https://doi.org/10.3390/foods11233940>.
 22. 4. tiwi G. Ronny M. Retno M. Chitosan nanoparticle as a delivery system for polyphenols from meniran extract (*Phyllanthus niruri* L.): Formulation, optimization, and immunomodulatory activity. *International Journal of Applied Pharmaceutics*. 2019; 11(2): 50-58. <https://doi.org/10.22159/ijap.2019v11i2.29999>
 23. Kanwal T, Kawish M, Maharjan R, Ghaffar I, Ali HS, Imran M, Perven S, Saifullah S, Simjee SU, Shah MR. Design and development of permeation enhancer containing self-nanoemulsifying drug delivery system (SNEDDS) for ceftriaxone sodium improved oral pharmacokinetics. *Journal of Molecular Liquids*. 2019; 289: 1-11. <http://dx.doi.org/10.1016/j.molliq.2019.111098>
 24. Syukri Y 5. Martien R, Lukitaningsih E, Nugroho AE. Novel selfnano emulsifying drug delivery system (SNEDDS) of andrographolide isolated from *Andrographis paniculata* Nees: Characterization, in-vitro/invivo assessment. *Drug Delivery Science and Technology*. 2018; 47: 514-520. <http://dx.doi.org/10.1016/j.jddst.2018.06.014>
 25. Durak T, Depciuch J. Effect of plant sample preparation and 25. suring methods on ATR-FTIR spectra results. *Environmental and Experimental Botany*. 2020; 169. <http://dx.doi.org/10.1016/j.en.vexbot.2019.103915>
 26. Junges CH, Guerra CC, Gomes AA, Ferrão MF. Green analytical methodology for grape juice classification using FTIR spectroscopy combined with chemometrics. *Talanta Open*. 2022; 6: 100168. <https://doi.org/10.1016/j.talo.2022.100168>.
 27. Hoang VD. Chemometrics-assited spectrophotometric determination of ciprofloxacin and naphazonline in eye drops. *Asian Journal of Research in Chemistry*. 2014; 7(5): 461-465.
 28. 8. ndhi V, Santos, Pratiidnya S. Chemometric-assisted UV spectrophotometric method for determination of cefixime trihydrate and cloxacillin sodium in pharmaceutical dosage form. *Asian Journal of Research in Chemistry*. 2018; 11(4): 705-709. <https://doi.org/10.5958/0974-4150.2018.00124.4>.
 29. Rizt A. Public service motivation and performance: A critical perspective. Evidence-based HRM: a Global Forum for Empirical Scholarship. 2014; 2: 57- 79. <http://dx.doi.org/10.1108/EBHRM-07-2013-0020>.
 30. 6. hman A, Windarsih A. The Application of molecular spectroscopy in combination with chemometrics for halal authentication analysis: A Review. *International Journal of Molecular Sciences*. 2020; 21(14): 5155. <https://doi.org/10.3390/ijms21145155>.
 31. Patil P, Mahajan VR. Self-emulsifying drug delivery systems (SEDDS): A Brief Review. *Research Journal of Pharmaceutical Dosage Forms and Technology*. 2014; 6(2): 134-139.
 32. 18. anrao BM, Sundar PS, Nagsen S. Oral bioavailability enhancement of a poor water soluble drug by co-surfactant free self-emulsifying drug delivery system (SEDDS). *Research Journal of Pharmacy and Technology*. 2011; 4(10): 1557-1562.
 33. Shiyani S, Zubaidah, Pratiwi G. Chemometric approach to assess response correlation and its classification in simplex centroid design for pre-optimization stage of catechin-SNEDDS. *Research Journal of Pharmacy and Technology*. 2021; 14(11): 5863-5870. <https://doi.org/10.52711/0974-360X.2021.01020>.
 34. Shiyani S, Suryani RP, Mulyani LN, Pratiwi G. Stability study of 26. r saturable catechin-self 9. no emulsifying drug delivery system as antidiabetic therapy. *Biointerface Research in Applied Chemistry*. 2022; 12(5): 5811-5820. <https://doi.org/10.33263/BRIAC125.58115820>.

Preparation and FTIR-ATR combined with chemometrics analysis of selfemulsifying loaded sungkai extract from *Peronema canescens*

ORIGINALITY REPORT

8%

SIMILARITY INDEX

7%

INTERNET SOURCES

6%

PUBLICATIONS

3%

STUDENT PAPERS

PRIMARY SOURCES

1	fr.scribd.com Internet Source	1%
2	ethesis.nitrkl.ac.in Internet Source	1%
3	www.pnrjournal.com Internet Source	1%
4	Submitted to Udayana University Student Paper	<1%
5	gsconlinepress.com Internet Source	<1%
6	Faten Farouk, Shereen Essam, Amany Abdel-Motaleb, Rana El-Shimy, Wolfgang Fritzsche, Hassan Mohamed El-Said Azzazy. "Fast detection of Bacterial Contamination in Fresh Produce using FTIR and Spectral Classification", <i>Spectrochimica Acta Part A: Molecular and Biomolecular Spectroscopy</i> , 2022	<1%

7 jurnal.ugm.ac.id <1 %
Internet Source

8 ajrconline.org <1 %
Internet Source

9 Aziz Aynaou, Boubaker Youbi, Youssef Lghazi, Mohammed Ait Himi, Chaimaa El Haimer, Jihane Bahar, Ahmed Sahlaoui, Itto Bimaghra. "Nucleation, Growth and Electrochemical Performances of Polyaniline Electrodeposited on ITO Substrate", Journal of The Electrochemical Society, 2022
Publication

10 Manuel Alexandru Gitea, Simona Gabriela Bungau, Daniela Gitea, Bianca Manuela Pasca et al. "Evaluation of the Phytochemistry-Therapeutic Activity Relationship for Grape Seeds Oil", Life, 2023
Publication

11 Yandi Syukri, Annisa Fitria, Suci Hanifah, Muthiah Idrati. "Development of New Indonesian Propolis Extract-Loaded Selfemulsifying: Characterization, Stability and Antibacterial Activity", Advanced Pharmaceutical Bulletin, 2020
Publication

12 japsonline.com
Internet Source

<1 %

13

link.springer.com

Internet Source

<1 %

14

Losso, Jack. "Development and Delivery of Anti-Angiogenic Functional Food Products : Opportunities and Challenges", *Nutraceutical Science and Technology*, 2007.

Publication

<1 %

15

spgykj.com

Internet Source

<1 %

16

www.myfoodresearch.com

Internet Source

<1 %

17

Rao, S.V.R.. "Self-nanoemulsifying drug delivery systems (SNEDDS) for oral delivery of protein drugs", *International Journal of Pharmaceutics*, 20081001

Publication

<1 %

18

Submitted to University of Philadelphia - Jordan

Student Paper

<1 %

19

iopscience.iop.org

Internet Source

<1 %

20

www.science.gov

Internet Source

<1 %

- 21 apb.tbzmed.ac.ir Internet Source <1 %
-
- 22 www.researchgate.net Internet Source <1 %
-
- 23 Palash Sanphui, V. Kusum Devi, Deepa Clara, Nidhi Malviya, Somnath Ganguly, Gautam R. Desiraju. "Cocrystals of Hydrochlorothiazide: Solubility and Diffusion/Permeability Enhancements through Drug-Coformer Interactions", Molecular Pharmaceutics, 2015
Publication <1 %
-
- 24 Tasmina Kanwal, Muhammad Kawish, Rukesh Maharjan, Iqra Ghaffar et al. "Design and development of permeation enhancer containing self-nanoemulsifying drug delivery system (SNEDDS) for ceftriaxone sodium improved oral pharmacokinetics", Journal of Molecular Liquids, 2019
Publication <1 %
-
- 25 horizonpublishing.com Internet Source <1 %
-
- 26 siepub.unsri.dev Internet Source <1 %
-

Exclude bibliography On

Preparation and FTIR-ATR combined with chemometrics analysis of selfemulsifying loaded sungkai extract from Peronema canecens

GRADEMARK REPORT

FINAL GRADE

/100

GENERAL COMMENTS

Instructor

PAGE 1

PAGE 2

PAGE 3

PAGE 4

PAGE 5

PAGE 6

PAGE 7

RUBRIC: 6TH-8TH SCIENCE ARGUMENT (CER)

CLAIM

Take an arguable position on the scientific topic and develop the essay around that stance.

ADVANCED	The essay introduces a precise, qualitative and/or quantitative claim based on the scientific topic or text(s), regarding the relationship between dependent and independent variables. The essay develops the claim and counterclaim fairly, distinguishing the claim from alternate or opposing claims.
PROFICIENT	The essay introduces a clear, qualitative and/or quantitative claim based on the scientific topic or text(s), regarding the relationship between dependent and independent variables. The essay effectively acknowledges and distinguishes the claim from alternate or opposing claims.
DEVELOPING	The essay attempts to introduce a qualitative and/or quantitative claim, based on the scientific topic or text(s), but it may be somewhat unclear or not maintained throughout the essay. The essay may not clearly acknowledge or distinguish the claim from alternate or opposing claims.
EMERGING	The essay does not clearly make a claim based on the scientific topic or text(s), or the claim is overly simplistic or vague. The essay does not acknowledge or distinguish counterclaims.

EVIDENCE

Include relevant facts, definitions, and examples to back up the claim.

ADVANCED	The essay supplies sufficient relevant, accurate qualitative and/or quantitative data and evidence related to the scientific topic or text(s) to support its claim and counterclaim.
PROFICIENT	The essay supplies relevant, accurate qualitative and/or quantitative data and evidence related to the scientific topic or text(s) to support its claim and counterclaim.
DEVELOPING	The essay supplies some qualitative and/or quantitative data and evidence, but it may not be closely related to the scientific topic or text(s), or the support that is offered relies mostly on summary of the source(s), thereby not effectively supporting the essay's claim and counterclaim.
EMERGING	The essay supplies very little or no data and evidence to support its claim and counterclaim, or the evidence that is provided is not clear or relevant.

REASONING

Explain how or why each piece of evidence supports the claim.

ADVANCED	The essay effectively applies scientific ideas and principles in order to explain how or why the cited evidence supports the claim. The essay demonstrates consistently logical reasoning and understanding of the scientific topic and/or text(s). The essay's explanations anticipate the audience's knowledge level and concerns about this scientific topic.
----------	--

PROFICIENT	The essay applies scientific reasoning in order to explain how or why the cited evidence supports the claim. The essay demonstrates logical reasoning and understanding of the scientific topic and/or text(s). The essay's explanations attempt to anticipate the audience's knowledge level and concerns about this scientific topic.
DEVELOPING	The essay includes some reasoning and understanding of the scientific topic and/or text(s), but it does not effectively apply scientific ideas or principles to explain how or why the evidence supports the claim.
EMERGING	The essay does not demonstrate clear or relevant reasoning to support the claim or to demonstrate an understanding of the scientific topic and/or text(s).

FOCUS

Focus your writing on the prompt and task.

ADVANCED	The essay maintains strong focus on the purpose and task, using the whole essay to support and develop the claim and counterclaims evenly while thoroughly addressing the demands of the prompt.
PROFICIENT	The essay addresses the demands of the prompt and is mostly focused on the purpose and task. The essay may not acknowledge the claim and counterclaims evenly throughout.
DEVELOPING	The essay may not fully address the demands of the prompt or stay focused on the purpose and task. The writing may stray significantly off topic at times, and introduce the writer's bias occasionally, making it difficult to follow the central claim at times.
EMERGING	The essay does not maintain focus on purpose or task.

ORGANIZATION

Organize your writing in a logical sequence.

ADVANCED	The essay incorporates an organizational structure throughout that establishes clear relationships among the claim(s), counterclaims, reasons, and evidence. Effective transitional words and phrases are included to clarify the relationships between and among ideas (i.e. claim and reasons, reasons and evidence, claim and counterclaim) in a way that strengthens the argument. The essay includes an introduction and conclusion that effectively follows from and supports the argument presented.
PROFICIENT	The essay incorporates an organizational structure with clear transitional words and phrases that show the relationship between and among ideas. The essay includes a progression of ideas from beginning to end, including an introduction and concluding statement or section that follows from and supports the argument presented.
DEVELOPING	The essay uses a basic organizational structure and minimal transitional words and phrases, though relationships between and among ideas are not consistently

clear. The essay moves from beginning to end; however, an introduction and/or conclusion may not be clearly evident.

EMERGING

The essay does not have an organizational structure and may simply offer a series of ideas without any clear transitions or connections. An introduction and conclusion are not evident.

LANGUAGE

Pay close attention to your tone, style, word choice, and sentence structure when writing.

ADVANCED

The essay effectively establishes and maintains a formal style and objective tone and incorporates language that anticipates the reader's knowledge level and concerns. The essay consistently demonstrates a clear command of conventions, while also employing discipline-specific word choices and varied sentence structure.

PROFICIENT

The essay generally establishes and maintains a formal style with few possible exceptions and incorporates language that anticipates the reader's knowledge level and concerns. The essay demonstrates a general command of conventions, while also employing discipline-specific word choices and some variety in sentence structure.

DEVELOPING

The essay does not maintain a formal style consistently and incorporates language that may not show an awareness of the reader's knowledge or concerns. The essay may contain errors in conventions that interfere with meaning. Some attempts at discipline-specific word choices are made, and sentence structure may not vary often.

EMERGING

The essay employs language that is inappropriate for the audience and is not formal in style. The essay may contain pervasive errors in conventions that interfere with meaning, word choice is not discipline-specific, and sentence structures are simplistic and unvaried.