

In vivo study of antioxidant test of ethanolic extract of *Chromolaena odorata* Linn. leaves

By Indah Solihah



Original Article

In vivo study of the antioxidant test of ethanolic extract of *Chromolaena odorata* Linn. leaves

Indah Solihah¹, Herlina, Inayatul Munawwaroh, Riana Sari Puspita Rasyid²

¹ *Departement of Pharmacy Faculty of Mathematics and Natural Sciences, Universitas Sriwijaya, South Sumatera, Indonesia.*

² *Departement of Medicine, Faculty of Medicine, Universitas Sriwijaya, South Sumatera, Indonesia.*

ARTICLE INFORMATION

9

Received: February 00, 00

Revised: March 00, 00

Available online: April 00, 00

KEYWORDS

Chromolaena; Plant Leaves; Antioxidants; Quercetin; Flavonoids

CORRESPONDENCE

Phone: +6281277781119

E-mail: indahsolihah@mipa.unsri.ac.id

ABSTRACT

Background: *Chromolaena odorata* L. leaf was reported to contain phenolic group compounds, such as flavonoids. Flavonoid compounds have been reported to have antioxidant activity. Based on in vitro study, *Chromolaena odorata* L. leaves have potent antioxidant activity. However, in vivo, studies with dose variations have not been reported.

Purpose: This study evaluates the antioxidant activity with various doses of ethanolic extract of *Chromolaena odorata* L. leaves against male Wistar rats induced by paracetamol.

Methods: Flavonoid contents were measured spectrophotometrically based on the formation of a complex flavonoid-aluminum. Quercetin was used to make a calibration curve. In vivo test was used TBARS method carried out by measured malondialdehyde (MDA) level in male Wistar rats induced by paracetamol 2g/Kg BW. The test was carried out on extracts with doses of 125, 250, and 500 mg/Kg BW. Vitamin C with dose 6,5mg/Kg BW used as a positive control, and 1% of Na CMC used as the negative control. Histopathology assessment of liver used Hema-toxylin Eosin Stain.

Results: Ethanolic extract of *Chromolaena odorata* L. leaves contain flavonoid 126.459±0.163 mg/g extract as quercetin equivalent. Intoxication paracetamol on rats increased MDA serum level significantly different (p-value < 0.005) with control normal. Treatment of ascorbic acid and extracts decreased MDA serum level significantly different (p-value < 0.005) with control negative and improved the histological structure of hepatocytes.

Conclusion: Ethanolic extract of *Chromolaena odorata* Linn. dose 500 mg/Kg BW was the best treatment with exhibited 58.974% reduction of MDA serum level and better improve the histological structure hepatocytes than other doses.

INTRODUCTION

Degenerative diseases, such as diabetes, cardiovascular diseases, inflammation, cancer, aging, neurodegenerative diseases, and immunosuppression, are developed from the excess free radical molecules^{1,2}. The primary role of free radical molecules causes of these diseases is related to lipid peroxidation³. Free radical⁷ molecules are formed by standard metabolic action and have been reported to be formed by radiation, bacterial and viral toxin, smoking, alcohol, and psychological or emotional stress⁴.

Antioxidant compounds from medicinal plants play an essential role in slowing or preventing free radical molecules' oxidation. Phenolic compounds, alkaloids, organic sulfur compounds, α-tocopherol, and β-carotene are phytochemical compounds reported⁴ antioxidant properties⁵. *Chromolaena odorata* Linn. is one of the tropical plants used as traditional medicines for diabetes in Indonesia's regions. Most of the *Chromolaena* gen⁴ contains the flavonoids group. The previous study had reported that about 40 flavonoids had been identified from the *Chromolaena* genus and some of these flavonoid groups have strong antioxidant properties^{6,7}.

ht²://doi.org/10.30595/xxxxx

©(2020) by the Medisains Journal. Readers may use this article as long as the work is properly cited, the use is educational and not for profit, and the work is not altered. More information is available at [Attribution-NonCommercial 4.0 International](https://creativecommons.org/licenses/by-nc/4.0/).

In vitro, antioxidant activities of *Chromolaena odorata* L. leaves have been reported widely. Ethanol extract of *Chromolaena odorata* L. leaves with concentration 300 µg/mL has 59,89±0,002% DPPH scavenging activity⁸. Ethanol extract of *Chromolaena odorata* L. contains 7,39±0,40g chlorogenic acid in 100g of dry leaves showed the DPPH radical scavenging activity with IC50 of 72,23 µg/mL and 35,61% antioxidant activity of β-Carotene bleaching⁹. Methanol extract of *Chromolaena odorata* L. leaf has IC50 scavenging of DPPH radical at 82,18 µg/mL¹⁰. Ethanol extract of *Chromolaena odorata* L. leaves has the highest DPPH radical scavenging activity than n-hexane and ethyl acetate extract¹¹.

In vivo study using the TBARS method reported that ethanol extract of *Chromolaena odorata* L. leaf has lower lipid peroxidation at 20mg/Kg BW compared with the control group. However, this value was not significantly different (p>0,05) statistically¹². However, antioxidant in vivo studies use the TBARS method with dose variations that have not been reported. Currently, this research aims to evaluate the antioxidant activity with various doses of ethanol extract of *Chromolaena odorata* L. leaves against male Wistar rats induced by paracetamol.

METHOD

Study Design

This is an true-experimental randomized pretest-posttest controlled trial.

Study Site

Pharmacological and Biological Pharmacy laboratories, Faculty of Mathematics and Natural Sciences, Sriwijaya University. Dyatnitalis anatomical pathology laboratory, Palembang

Materials

The dried leaves of *Chromolaena odorata* Linn., ethanol, aqua dest, methanol, Quercetin (Sigma-Aldrich®), Sodium acetate, AlCl₃, aquabidest, Paracetamol standard (Dexa Medica), thiobarbituric acid (TBA) (Sigma-Aldrich®), trichloro acetate (TCA) (Sigma-Aldrich®), tetra-ethoxy-propane (TEP) (Sigma-Aldrich®), Na CMC, Ascorbic acid standard (Dexa Medica Pharm. Industry). Hematoxylin Eosin stain, Buffered formalin, Paraffin wax.

Plant Extraction

The dried leaves of *Chromolaena odorata* Linn. obtained from Belitung, South Sumatera, Indonesia. The 2 Kg of dried leaves were ground into powdered form then soak with ethanol 96% (1:10). The maceration process was kept in the amber bottle at room temperature for 72 hours and shaken occasionally. The macerate filtered then evaporated with a rotary evaporator at 60°C until getting a thick extract.

Total Flavonoid Content Assay

Total flavonoid content was measured spectrophotometrically based on the formation of complex flavonoid-aluminum¹³. Quercetin was used to make a calibration curve. Quercetin was prepared by concentration 5, 10, 15, 20, and 25 µg/mL in methanol solution. The extract was prepared by concentration 1000 µg/mL in methanol solution. One milliliter of sample solution was added with 1 mL of AlCl₃ 10% solution and 1 mL of sodium acetate 5%. The sample solution was incubated for 30 minutes, then measured absorbance value at λ 438 nm. The total flavonoid content was expressed as quercetin equivalents in mg/g (QE mg/g) extract.

In Vivo Procedure

Animal Preparation

Male Wistar rats (150-200 g) were used in this study, and seventh days were acclimated in the laboratory facility. All animals were maintained with approved animal care operating procedures consistently.

Experimental procedure

In vivo, the antioxidant assay was measured with the thiobarbituric acid reactive substance (TBARS) method¹⁴. The rats were divided into six groups, each comprising six rats. Group 1 (standard control) had free access to food. Group 2 (Negative control) received 2 g/Kg BW of paracetamol suspension p.o once a day for six days. Group 3 (Positive control) received 6.5 mg/Kg BW of ascorbic acid solution (in aquabidest) once a day six days later after receiving the same treatment of negative control. Group 4-6 (extract treatment) were received 125, 250, and 500 mg/Kg BW, respectively, once a day for six days later after received the same treatment of negative control. Blood serum was collected on the seventh and 14th day at sinus orbitalis. The MDA serum levels were obtained spectrophotometrically at 532 nm. Percent reduction of MDA level calculated with equation (1).

$$\text{Percent reduction MDA level (\%)} = \frac{\text{MDA before treatment} - \text{MDA after treatment}}{\text{MDA before treatment}} \times 100 \% \quad (1)$$

Histopathology of Liver

The histopathological assessment used the Hematoxylin Eosin stain as described before¹⁵. Liver tissues in each group were collected in 10% neutral buffered formalin. These tissues were processed and embedded in paraffin wax. Sections were cut used microtome with 5 µm of thickness and stained with hematoxylin and eosin (H&E). The sections were examined microscopically for the evaluation of histopathological changes.

Statistical Analysis

Results are expressed as mean ± standard deviation and analyzed using one-way ANOVA followed by Tukey's posthoc test and paired student T-test for serum MDA level

before and after treatment. The p values were two-tailed, and $p < 0.05$ was regarded as significant. Histopathological data were analyzed descriptively.

Ethical Consideration

The Health Research Review Committee approved this study of Mohammad Hoesin Central General Hospital and Faculty of Medicine Universitas Sriwijaya University (182/kepkrsmhfunksri/2019).

RESULTS

Plant Extraction and Total Flavonoid Measurement

Chromolaena odorata L. leaves were extracted with 96% ethanol to produce a 321.79 g viscous extract. The extract yield was 16.08% of the dry weight. Measurement of the total flavonoid levels of the extract used a quercetin calibration curve. Calibration curve equation obtained $y = 0.0188x - 0.0057$ with a value of $R^2 = 0.9998$. The quercetin calibration curve is shown in Figure 1. Based on this equation, the total flavonoid of ethanolic extract of *Chromolaena odorata* L. leaves express as quercetin equivalent was 126.459 ± 0.163 mg/g.




Figure 1. Quercetin Calibration Curve

In Vivo Test

Figure 2 shows that the negative, positive, and extract groups have been increased MDA levels after being intoxicated by 2g/Kg BW of paracetamol once a day for six days. In contrast, the standard control group was not induced by paracetamol as a reference for normal MDA levels. ANOVA analysis shows significantly different (p -value < 0.05) in the MDA level of each group. Based on Tukey's posthoc analysis, there is known that negative control, positive control, and 500mg/Kg BW extract groups have a significantly different value (p -value < 0.05) of MDA levels

with the standard control group. However, the MDA level on all group treatment had no significant difference (p -value > 0.05) with the negative control. It is indicated that paracetamol has been increased the MDA serum level of rats.

After the rats were treated for seven days, each group show a decrease in MDA levels, except for the standard and opposing control groups. Figure 3 shows that the lively control group and each extract group have significantly different (p -value $< 0,05$) MDA levels from the negative control group. Table 1 shows the percent reduction value of MDA level after treatment.

Table 1. Percent Reduction in MDA Level

Group	MDA level after Paracetamol intoxication (nmol/mL)	MDA level after treatment (nmol/mL)	Percent Reduction (%)
Negative control	2.654±0.599	2.510±0.434	5.426
Positive control	3.516±0.874	1.120±0.131*	68.146
125 mg/Kg BW of extract	2.471±0.261	1.490±0.093*	39.701
250 mg/Kg BW of extract	2.364±0.407	1.224±0.176*	48.223
500 mg/Kg BW of extract	2.729±0.899	1.120±0.318*	58.974

(* = significantly different (p -value < 0.05) of MDA before and after treatment with paired student T-test analysis)

A paired student T-test had used to analyze the differences of MDA level after paracetamol intoxication and MDA level after treatment. Table 1 found that a lively group and extract groups significantly reduce MDA level after treatment (p -value < 0.05). Positive control has the highest reduction of MDA level compared to other treatment groups. While in extract groups, 500 mg/Kg BW of ethanolic extract of *Chromolaena odorata* L. leaves was the best dose of an extract with 58.974% reduction of MDA level. Based on Tukey's posthoc analysis, all extract groups treatment had no significant difference (p -value > 0.05) in MDA level with the lively control group. It can be concluded that the ethanolic extract of *Chromolaena odorata* L. leaves has the same effectiveness as ascorbic acid.

Histopathology of Liver

There are differences in hepatocyte structure in the paracetamol-induced group of rats compared to the standard control. Paracetamol induction causes inflammation and necrosis of the hepatocyte cells (Figure 4) ¹⁵.




Figure 2. Serum MDA Level After Paracetamol Intoxication (* = significantly different with normal control (p-value < 0.05))




Figure 3. Serum MDA Level After Treatment (* = significantly different with negative control (p-value < 0.001))




Figure 4. Photograph of rat liver section shows A). Normal control shows the normal histological structure of hepatocyte (H) and sinusoid (S), B). Negative control shows more inflammatory infiltration, necrosis (arrow), and binucleated cell (BN) of hepatocyte, C). Positive control shows less necrosis (arrow) and binucleated cell (BN) of hepatocyte, D). Extract dose I shows inflammatory infiltration, necrosis (arrow), pyknotic nuclei (PK), and binucleated cells (BN) of hepatocyte, E). Extract dose II shows necrosis (arrow), pyknotic nuclei (PK), and binucleated cells (BN) of hepatocyte, F). Extract dose III shows necrosis (arrow), pyknotic nuclei (PK), and binucleated cells (BN) of the hepatocyte. (1) magnification 40x, (2) magnification 400x

DISCUSSION

Chromolaena odorata Linn. is a member of the family Eupatoriae (Asteraceae) ⁷. In Belitung, South Sumatera region, Indonesia, *Chromolaena odorata* Linn is known as "merdekaan," a popular folk medicine for diabetes treatment. Ethanolic extract of *Chromolaena odorata* Linn. leaves contain flavonoids, alkaloids, tannins, phenolics, saponins, steroids, and triterpenoids (data not shown). Some phenolic group compounds have been found in *Chromolaena odorata* L. leaf; there are p-coumaric acid, p-hydroxybenzoic acid, protocatechuic acid, protocatechuic acid, vanillic acid ¹⁶. Odoratenin, isosakuranetin, and

subscandenin, a new flavanone compound, in *Chromolaena odorata* L. leaf was found by Putri ⁶. High phenolic content is beneficial as the phenolic compound quench primary oxidants or free radicals ¹⁷. Balamurugan⁹ was investigated that indirect ethanolic extraction of *Chromolaena odorata* L. leaves has the highest total phenolic compound and total antioxidant content compare to other solvents in sequential extraction. Ethanol is a useful solvent to extract phenolic and flavonoid compounds and is safe for human consumption ¹⁸. Flavonoid and phenolic compounds have free radical scavenging activity by donating their free electron pairs to radical molecules ⁹. This study was measured the total flavonoid of ethanolic extract of *Chromolaena odorata* L. leaves; it has 126.459±0.163 mg QE/g extract.

Chromolaena odorata L. leaves have been shown as a potent antioxidant in various in vitro assays^{8-11,13,16,19-24}. However, in vivo assays of antioxidant activity of this plant are limited. Uhegbu¹² was investigating the activity of Chromolaena odorata L., Ageratum conyzoides L., and their combination to reduce lipid peroxidation in rats. The evaluation shows that ethanolic extract of Chromolaena odorata L. leaf has lower lipid peroxidation (TBARS) at 20mg/Kg BW than the control group. However, this value was not significantly different (p-value > 0.005) statistically. To our knowledge, in vivo study with various doses of Chromolaena odorata L. leaves has not been evaluated. This study was aimed to evaluate the antioxidant activity in system organisms used various doses of the extract. We used Wistar strain male rats as the test animal. The study used six groups: standard control, negative control, positive control, 125 mg/Kg BW, 250 mg/Kg BW, and 500 mg/Kg BW of ethanolic extract Chromolaena odorata Linn. leaves. Except for standard control, the five groups received 2mg/Kg BW of Paracetamol once a day for six days to elevate the MDA serum level.

Paracetamol with toxic dose can significantly elevate Malonaldehyde (MDA) level serum of rats. The elevation of Malonaldehyde (MDA) level serum indicates of increase in lipid peroxidation²⁵. Paracetamol had reported disturbing the balance between free radical molecules production and antioxidant protection, especially in the liver. N-acetyl-p-benzoquinone imine (NAPQI) formation by cytochrome P450 (CYP), a highly reactive toxic electrophile, is a toxic compound produced by paracetamol. When the rate of NAPQI formation exceeds the detoxification rate by glutathione, it is oxidized tissue macromolecules such as lipid or -SH group of proteins. Lipid peroxidation is an autocatalytic process, which is a common consequence of cell death²⁶. MDA is one of the end products in the lipid peroxidation process. The increased MDA content might have resulted from an increase of free radical molecules due to stress due to Paracetamol intoxication²⁷.

Serum MDA level of negative control, positive control, 125 mg/Kg BW, 250 mg/Kg BW, and 500 mg/Kg BW of ethanolic extract of Chromolaena odorata Linn. leaves after paracetamol intoxication has significantly higher (p-value < 0.05) than standard control (Figure 2). The reduction of MDA level serum in the positive control group, 125 mg/Kg BW, 250 mg/Kg BW, and 500 mg/Kg BW of ethanolic extract of Chromolaena odorata Linn. leaves have significantly different (p-value < 0.05) with negative control (Figure 3). Ascorbic acid in the positive control group has the highest percent reduction in MDA level (68.146%). Besides that, 500mg/Kg BW of ethanolic extract has a 58.974% reduction of MDA level, higher than other doses. Ascorbic acid as a standard drug, flavonoid, and phenolic compound in the extract has antioxidant activity that inhibits the enzyme involved in free radical molecules formation²⁸. Peluso²⁹ has been reported that flavonoids could

interfere with drugs' bioavailability through competition with cytochrome P₄₅₀ (CYP) enzymes. So, the metabolism of paracetamol into NAPQI form can be reduced.

Figure 4, A1-A2 liver sections of the standard control group showed the typical histological structure of hepatocytes with well-preserved cytoplasm and well-defined nucleus. The negative control group (Figure 4, B1-B2) showed disarrangement of normal hepatocytes, necrosis of cells, and inflammatory infiltration. There were binucleated cells and less necrosis of hepatocytes (Figure 4, C1-C2). Apart from necrosis of cells and binucleated cells, pyknotic nuclei were also found in extract treatment groups (Figure 4, D-F1-2). In our present study, there were histopathological changes in response to paracetamol. Indication of marked changes in the liver's overall histoarchitecture could be due to its toxic effect by the generation of radical molecules that damage the various membrane components of the cell—the necrotic condition in rats mostly characterized by pyknosis cytoplasm³⁰. The formation of highly reactive radicals because of paracetamol intoxication can cause liver function failure³¹. Lipid peroxidation can disturb cellular membranes' integrity, leading to the leakage of cytoplasmic enzymes due to severe histopathological damages^{32,33}. The phenolic compound, such as quercetin, was proven to possess hepatoprotective effect in an animal test against liver damage induced by lipid peroxidation³⁴. Phenolic compounds prevent paracetamol-induced lipid peroxidation was due to antioxidant activities. Chromolaena odorata L. leaves have strong antioxidant properties and inhibit lipid peroxidation due to hepatoprotective effect in rats against liver damage induced by paracetamol.

CONCLUSIONS AND RECOMMENDATION

This study concludes that ethanolic extract of Chromolaena odorata Linn. leaves contain a flavonoid of 126.459±0.163 mg QE/g extract. Extract with a 500 mg/Kg BW dose was the best treatment with exhibited 58.974% reduction of MDA serum level. Ascorbic acid and all extract treatments showed improvement in the histological structure of hepatocytes. Therefore, the next study should be directed toward using extract rich flavonoids to reduce doses and increase the activity. The small extract dosage will be more acceptable to make dosage preparations for human usage.

Acknowledgment

We wish to acknowledge Sriwijaya University that supported this research through SATEKS grant contract no.0684/UN9/SK.BUK.KP/2020.

REFERENCES

In vivo study of antioxidant test of ethanolic extract of *Chromolaena odorata* Linn. leaves

ORIGINALITY REPORT

7%

SIMILARITY INDEX

PRIMARY SOURCES

- 1** text-id.123dok.com Internet 42 words — 1%
- 2** Stefanus Mendes Kiik, Muhammad Saleh Nuwa. "Quality of life of the elderly: a comparison between community-dwelling elderly and in social welfare institutions", MEDISAINS, 2020 Crossref 42 words — 1%
- 3** Srinivasan, R.. "Antioxidant activity of *Caesalpinia digyna* root", Journal of Ethnopharmacology, 20070905 Crossref 25 words — 1%
- 4** Devi Anggraini Putri, Sri Fatmawati. " A New Flavanone as a Potent Antioxidant Isolated from L. Leaves ", Evidence-Based Complementary and Alternative Medicine, 2019 Crossref 22 words — 1%
- 5** www.apjtb.com Internet 21 words — 1%
- 6** Afolabi C. Akinmoladun, Kehinde O. Oguntunde, Lawrence O. Owolabi, Omotayo B. Ilesanmi et al. "Reversal of acetaminophen-generated oxidative stress and concomitant hepatotoxicity by a phytopharmaceutical product", Food Science and Human Wellness, 2017 Crossref 21 words — 1%
- 7** citeseerx.ist.psu.edu Internet 17 words — 1%

M. Sakti La Ore, Karna Wijaya, Wega Trisunaryanti, Wahyu Dita

8 Saputri et al. "The synthesis of SO₄/ZrO₂ and Zr/CaO catalysts via hydrothermal treatment and their application for conversion of low-grade coconut oil into biodiesel", Journal of Environmental Chemical Engineering, 2020
Crossref 16 words — < 1%

9 repository.uin-suska.ac.id
Internet 11 words — < 1%

10 www.intechopen.com
Internet 10 words — < 1%

EXCLUDE QUOTES OFF
EXCLUDE BIBLIOGRAPHY ON

EXCLUDE MATCHES OFF