

Title of Manuscript: **Cancer Treatment in Islamic Traditional
Medicine**

1. Proofread document received (September 22th, 2021)
-Document from proofreading service
2. Submitted to the journal "Arkus" (September 28th, 2021)
3. Peer Reviewer results: Revision Required (October 6th, 2021)
4. Revised version received by journal (October 16th, 2021)
5. Paper Accepted for publication (October 18th, 2021)
6. Galley proof (October 20th, 2021)
7. Paper published (October 21th, 2021)

September 22th, 2021

HM Publisher

Jl Sirnaraga No 99, 8 Ilir, Ilir Timur 3, Palembang, South Sumatra, Indonesia

CONFIDENTIAL
September 22th, 2021

Certificate Service Confirmation

To whom it may concern,
HM Publisher provided comprehensive editing services for manuscript entitled
Cancer Treatment in Islamic Traditional Medicine. The edit has achieved Grade
A: priority publishing; no language polishing required after editing. Should you
require any additional information, please do not hesitate to contact me.

Regards,



Khrishna Murti, PhD
Head of Language Institute-HM Publisher
Email: khrishnamurti@gmail.com

Submitted to the journal "Arkus" (September 28th, 2021)

Cancer Treatment in Islamic Traditional Medicine

Rachmat Hidayat^{1*}

¹ Department of Biology, Faculty of Medicine, Universitas Sriwijaya, Palembang, Indonesia

*Corresponding author email: dr.rachmat.hidayat@gmail.com

Abstract

Although significant progress has been made in cancer treatment in recent decades, the effectiveness of modern therapeutic approaches is often limited by their toxic effects on other organs. In addition, many people in the world have limited or unequal access to cancer treatment services. Therefore, utilizing information from traditional medicine systems to identify alternative methods to prevent and control cancer. The use of traditional medicine can lead to the discovery of new bioactive compounds and drugs that are available, safe and affordable. In Islamic medicine, cancer is known as a disease with many treatment options. Traditional Islamic medicine suggests several strategies for cancer control and treatment. Surgical manipulation, venesection, dietary adjustment, and use of natural medicines including solid, semisolid, and liquid dosage forms by oral and external routes of administration are among these methods. This literature study will presents cancer from the perspective of traditional Islamic medicine, its aetiology and management.

Keywords. Cancer Treatment, Complementary Medicine, Traditional Medicine, Islamic Medicine.

Introduction

Islamic traditional medicine (ITM) is a holistic system of medicine that developed during the Islamic Golden Age (750-1258 AD). It is practised and taught throughout the Islamic world. At that time, Muslim and non-Muslim medical scholars, especially Iranian doctors and pharmacists, translated the scientific knowledge they inherited from ancient Greek and Iranian (Persian) culture. Strives to revive and develop the knowledge, remove myths and errors, and impart authentic medicine knowledge. Although most of ITM's scientists are not Arabs, most of the books are written in Arabic. Razi (Razhes; 865-925 AD), Ahwazi (Haly Abbas; 930-994 AD), Avicenna (Ibnu Sina; 980-1037 AD), Jorjani (1042-1136 AD), and Aqili Khorasani (18th century) were prominent Iranian physicians who have the most significant contribution to the promotion of ITM.¹

Although significant progress has been made in cancer treatment in recent decades, the effectiveness of modern therapeutic approaches is often limited by their toxic effects on other organs. In addition, many people in the world have limited or unequal access to cancer treatment services. Therefore, utilizing information from traditional medicine systems to identify alternative methods to prevent and control cancer. In addition, the use of traditional medicine can lead to the discovery of new bioactive compounds and drugs that are available, safe and affordable. In Islamic medicine, cancer is known as a disease with many treatment options. This literature study will discuss cancer, its aetiology and management from the perspective of traditional Islamic medicine.²

Definition of cancer from ITM perspective

ITM is based on humour knowledge, which holds that the human body comprises four essential fluids, namely humour. The four fluids are blood, phlegm, yellow bile, and black bile, and each corresponds to a paired quality: hot and wet blood, cold and wet phlegm, hot and dry yellow bile, and cold and dry black bile. The right balance of humour is essential for maintaining good health. Thus, all diseases and disabilities, including cancer, are caused by excess or decay of this fluid.^{3,4}

According to ITM, cancer is a type of swelling of black bile that is accompanied by pain, throbbing, inflammation, and angiogenesis. The blood vessels around the tumour are full and stiff and contain dark, black blood. The designation "cancer" is due to the similarity between cancerous tumours that affect organs and crabs that catch their prey. The round shape, the darkness of the tumour and the origin of the blood vessels from its environment describe a crab's leg image. Cancer often affects soft and porous organs, and for this reason, cancer mainly involves the breast and innervated organs (such as the uterus) in women and the throat, larynx, testes and penis in men.⁵

Initially, the cancer is the size of a pea or smaller, complex, round, mobile, dark, and slightly warm. Then it will start to grow gradually and reach the size of a walnut or larger. It may be curable during the early stages of its development, but diagnosis is difficult at this stage. On the other hand, cancer treatment will be difficult or even impossible with the development and emergence of clinical manifestations.

Cancers of the eye, nasal cavity, breast, uterus, liver, and other visceral organs and skin are the most frequently mentioned cancers in the ITM text. Cancer is divided into two main types: ulcerative and nonulcerative cancer. Some cancerous tumours ulcerate easily, but some do not. Cancer sores usually spread centripetally, and the discharge is purulent. Appropriate use of drugs can prevent ulceration of susceptible tumours. On the other hand, some cancerous tumours that are not prone to ulceration will ulcerate after inappropriate drug administration.

Advanced and large tumours are susceptible and painful, with a characteristic red to yellow colour and a burning, stinging pain. Such tumours may erupt spontaneously, purulent and bloody fluid may appear on the wound surface. The resulting wound is susceptible and can corrode the surrounding tissue.

Cancer aetiology

According to the ITM, excessive accumulation of abnormal black bile in any part of the body is a significant cause of cancer. Ageing, prolonged exposure to stress, cold and dry foods and hard work are the leading causes of increased black bile production. In some cases, bleeding (such as menstruation, abnormal uterine bleeding, or bleeding haemorrhoids) is a defence mechanism against the accumulation of bad humour in the body. Therefore, stopping the bleeding completely by surgical procedures and other medications can accumulate black bile and increase the risk of cancer and other diseases stemming from excessive black bile (including cancer, melancholia, liver problems, psoriasis).⁶

Cancer management

Traditional Islamic medicine suggests several strategies for cancer control and treatment. Surgical manipulation, venesection, dietary adjustment, and use of natural

medicines including solid, semisolid, and liquid dosage forms by oral and external routes of administration are among these methods.⁷

Surgery and manipulation

Surgery is used to eradicate tumours in the early stages of their development. Small tumours distant from vital organs are good candidates for surgery. The tumour should be excised from its origin, and some of the adjacent unaffected tissue should also be excised. In addition, the bleeding should be allowed to continue until a large volume of blood is expelled, and the surrounding tissue must be compressed to expel the blood mixed with black bile thoroughly. After that, the injured site must be saved or burned. However, suppose the cancerous tumour is located in the vicinity of sensitive and vital organs. In that case, surgical procedures and cauterization will be hazardous and turn cancer into a non-healing ulcer. Regular venesection is also advised during the early stages of cancer to draw the blood of black bile from the body.⁸

Nutritional therapy

In traditional Islamic medicine, dietary recommendations have been proposed to slow the progression of advanced tumours that cannot be manipulated due to metastatic problems. In this case, nutritional treatment will increase the patient's longevity. Foods with a wet temper that produce high-quality blood such as almond oil, fresh small fish, soft boiled egg yolks, lamb, bird meat, ripe and sweet apples, sweet plums, bananas, blackcurrant (*Vigna mungo* (L.) Hepper), spinach (*Spinacia oleracea* L.), pumpkin, light wine, and fresh cow's milk and dough were given. On the other hand, excessive intake of foodstuffs can trigger the production of black bile in the body (such as eggplant (*Solanum melongena* L.), lentils (*Lens esculenta* Moench), dates (*Phoenix dactylifera* L.), cabbage (*Brassica oleracea* L.), meat beef, black and thick wine, and meat preserved with salt is strictly limited.⁹

Pharmacotherapy

From the point of view of traditional Islamic medicine, an excess of abnormal black bile in various parts of the body can lead to cancer formation. In treating cancer, black bile must be removed from the body using appropriate laxatives and prevent the formation and accumulation of black bile in the vessels as far as possible. For this purpose, many single and compound drugs of plant, animal and mineral origin have been recommended. Of the 107 plant species introduced in Iranian Islamic medicine for cancer treatment, 59 plants or their chemical compounds have been shown to have cytotoxic and antitumor activities in recent investigations, and several have entered clinical trials. Their effectiveness has been evaluated in humans.⁷⁻⁸

An essential point in cancer and tumour pharmacotherapy is avoiding caustic and irritant drugs to prevent further stimulation and ulceration. Cancer drugs can be administered internally (oral, enema, vaginal douche) or applied topically (topical oils, liniments, lotions, powders). Local anticancer drugs have the following objectives: cancer eradication, prevention of metastasis, prevention of ulceration, and tumour healing ulceration.

Mechanism of action of anticancer drugs

Anticancer drugs work through different mechanisms, namely as black bile laxative, antiulcer in cancerous wounds, wound healing drugs, and analgesics. Black bile laxatives are common anticancer drugs that can eliminate abnormal black bile from the rest of the body. Laxatives should be given frequently. From ITM's point of view, dodder clover (*Cuscuta epithimum Murr.*) is the most valuable black bile laxative, commonly used to treat diseases caused by excess or imbalance of this humour as all types of cancer, melancholia, leprosy, and vitiligo. For this purpose, a mixture of clover dodder with cheese whey or hydromel should be given frequently. A decoction of the plant in oxymel is also prescribed. Common polypody (*Polypodium vulgare L.*), French lavender (*Lavandula stoechas L.*), colocynth (*Citrullus colocynthis (L.) Schrad.*], and black hellebore (*Helleborus niger L.*) are other potent black bile laxatives.^{4,5}

Antiulcer drugs can inhibit tumour ulceration. *Urtica pilulifera L.* and *Aloe vera (L.) Burm.f.* is an example of such a plant. Wound healing drugs accelerate the healing of cancerous wounds. *Althaea officinalis L.*, *Brassica oleracea L.*, and *Viola odorata L.* have wound healing activity. In addition, analgesic drugs from plants can relieve cancer pain. *Parietaria officinalis L.* and *Solanum nigrum L.* are plants that have analgesic effects.

Conclusion

In conclusion, traditional Islamic medicine for cancer has been widely proven in modern research. Of the 107 plant species introduced in Iranian Islamic medicine for cancer treatment, 59 plants or their chemical compounds have been shown to have cytotoxic and antitumor activities in recent investigations, and several have entered clinical trials.

This finding shows the deep insight of Islamic doctors about cancer treatment. Despite the lack of modern facilities and developed equipment on their era, they introduced anticancer plants, showing cytotoxic properties in the latest research. The correlation between these findings signifies the originality of experience and studies, representing valuable funds and valuable knowledge dating back more than twelve centuries. This legacy is based on thousands of years of experience by ancient Greek, Indian and Iranian doctors and relies on many clinical trials on thousands of people. In addition, the application of traditional medicine knowledge reinterpreted by modern data can lead to more effective and evidence-based use of medicinal plants, which can contribute to therapeutic decisions in various diseases.

References

1. Jurenka J. Therapeutic applications of pomegranate (*Punica granatum L.*): a review. *Alternative Medicine Review*, 2008. 13 (2): 128-144.
2. E. P. Lansky, W. Jiang, H. Mo et al., Possible synergistic prostate cancer suppression by anatomically discrete pomegranate fractions," *Investigational New Drugs*, vol. 23, no. 1, pp. 11–20, 2005.
3. A. Malik and H. Mukhtar, "Prostate cancer prevention through pomegranate fruit," *Cell Cycle*, vol. 5, no. 4, pp. 371–373, 2006.
4. L. S. Adams, N. P. Seeram, B. B. Aggarwal, Y. Takada, D. Sand, and D. Heber, "Pomegranate juice, total pomegranate ellagitannins, and punicalagin suppress inflammatory cell signaling in colon cancer cells," *Journal of Agricultural and Food Chemistry*, vol. 54, no. 3, pp. 980–985, 2006.

5. N. Khan, F. Afaq, M. H. Kweon, K. Kim, and H. Mukhtar, "Oral consumption of pomegranate fruit extract inhibits growth and progression of primary lung tumors in mice," *Cancer Research*, vol. 67, no. 7, pp. 3475–3482, 2007.
6. V. M. Adhami, N. Khan, and H. Mukhtar, "Cancer chemoprevention by pomegranate: laboratory and clinical evidence," *Nutrition and Cancer*, vol. 61, no. 6, pp. 811–815, 2009.
7. E. Hornung, C. Pernstich, and I. Feussner, "Formation of conjugated Δ^{11} Δ^{13} -double bonds by Δ^{12} -linoleic acid (1,4)-acyl-lipid-desaturase in pomegranate seeds," *European Journal of Biochemistry*, vol. 269, no. 19, pp. 4852–4859, 2002.
8. R. F. Wang, W. D. Xie, Z. Zhang et al., "Bioactive compounds from the seeds of *Punica granatum* (pomegranate)," *Journal of Natural Products*, vol. 67, no. 12, pp. 2096–2098, 2004.
9. Behjat J., Milad I., Seyed A.E. Anticancer plants in islamic traditional medicine. *Complementary Therapies for the Body, Mind and Soul*. IntechOpen. 2015.

Submitted to the journal "Arkus" (September 28th, 2021)

Arkus
Multidisciplinary Journal



Submission acknowledgement

Dear author(s),

Rachmat Hidayat* has submitted the manuscript "Cancer Treatment in Islamic Traditional Medicine" to Arkus. The paper will be screened by editor and reviewed by peer review.

Cordially,

A handwritten signature in black ink, appearing to be "P. Magnano", is positioned to the left of the editor's name.

Prof. Paula Magnano, PhD

Editor **HM Publisher**



(*) Corresponding author

Peer Review Results

Dear author(s),

Rachmat Hidayat* has submitted the manuscript “Cancer Treatment in Islamic Traditional Medicine” to Arkus. The decision : Revision Required.

Cordially,



Prof. Paula Magnano, PhD

Editor



HM Publisher

(*) Corresponding author

Reviewer 1

Cancer Treatment in Islamic Traditional Medicine →1

Rachmat Hidayat^{1*}

¹ Department of Biology, Faculty of Medicine, Universitas Sriwijaya, Palembang, Indonesia

*Corresponding author email: dr.rachmat.hidayat@gmail.com

Abstract →3

Although significant progress has been made in cancer treatment in recent decades, the effectiveness of modern therapeutic approaches is often limited by their toxic effects on other organs. In addition, many people in the world have limited or unequal access to cancer treatment services. Therefore, utilizing information from traditional medicine systems to identify alternative methods to prevent and control cancer. The use of traditional medicine can lead to the discovery of new bioactive compounds and drugs that are available, safe and affordable. In Islamic medicine, cancer is known as a disease with many treatment options. Traditional Islamic medicine suggests several strategies for cancer control and treatment. Surgical manipulation, venesection, dietary adjustment, and use of natural medicines including solid, semisolid, and liquid dosage forms by oral and external routes of administration are among these methods. This literature study will presents cancer from the perspective of traditional Islamic medicine, its aetiology and management.

Keywords. Cancer Treatment, Complementary Medicine, Traditional Medicine, Islamic Medicine. →2

Introduction →4

Islamic traditional medicine (ITM) is a holistic system of medicine that developed during the Islamic Golden Age (750-1258 AD). It is practised and taught throughout the Islamic world. At that time, Muslim and non-Muslim medical scholars, especially Iranian doctors and pharmacists, translated the scientific knowledge they inherited from ancient Greek and Iranian (Persian) culture. Strives to revive and develop the knowledge, remove myths and errors, and impart authentic medicine knowledge. Although most of ITM's scientists are not Arabs, most of the books are written in Arabic. Razi (Razhes; 865-925 AD), Ahwazi (Haly Abbas; 930-994 AD), Avicenna (Ibnu Sina; 980-1037 AD), Jorjani (1042-1136 AD), and Aqili Khorasani (18th century) were prominent Iranian physicians who have the most significant contribution to the promotion of ITM.¹

Although significant progress has been made in cancer treatment in recent decades, the effectiveness of modern therapeutic approaches is often limited by their toxic effects on other organs. In addition, many people in the world have limited or unequal access to cancer treatment services. Therefore, utilizing information from traditional medicine systems to identify alternative methods to prevent and control cancer. In addition, the use of traditional medicine can lead to the discovery of new bioactive compounds and drugs that are available, safe and affordable. In Islamic medicine, cancer is known as a disease with many treatment options. This literature study will discuss cancer, its aetiology and management from the perspective of traditional Islamic medicine.²

Definition of cancer from ITM perspective

ITM is based on humour knowledge, which holds that the human body comprises four essential fluids, namely humour. The four fluids are blood, phlegm, yellow bile, and black bile, and each corresponds to a paired quality: hot and wet blood, cold and wet phlegm, hot and dry yellow bile, and cold and dry black bile. The right balance of humour is essential for maintaining good health. Thus, all diseases and disabilities, including cancer, are caused by excess or decay of this fluid.^{3,4}

According to ITM, cancer is a type of swelling of black bile that is accompanied by pain, throbbing, inflammation, and angiogenesis. The blood vessels around the tumour are full and stiff and contain dark, black blood. The designation "cancer" is due to the similarity between cancerous tumours that affect organs and crabs that catch their prey. The round shape, the darkness of the tumour and the origin of the blood vessels from its environment describe a crab's leg image. Cancer often affects soft and porous organs, and for this reason, cancer mainly involves the breast and innervated organs (such as the uterus) in women and the throat, larynx, testes and penis in men.⁵

Initially, the cancer is the size of a pea or smaller, complex, round, mobile, dark, and slightly warm. Then it will start to grow gradually and reach the size of a walnut or larger. It may be curable during the early stages of its development, but diagnosis is difficult at this stage. On the other hand, cancer treatment will be difficult or even impossible with the development and emergence of clinical manifestations.

Cancers of the eye, nasal cavity, breast, uterus, liver, and other visceral organs and skin are the most frequently mentioned cancers in the ITM text. Cancer is divided into two main types: ulcerative and nonulcerative cancer. Some cancerous tumours ulcerate easily, but some do not. Cancer sores usually spread centripetally, and the discharge is purulent. Appropriate use of drugs can prevent ulceration of susceptible tumours. On the other hand, some cancerous tumours that are not prone to ulceration will ulcerate after inappropriate drug administration.

Advanced and large tumours are susceptible and painful, with a characteristic red to yellow colour and a burning, stinging pain. Such tumours may erupt spontaneously, purulent and bloody fluid may appear on the wound surface. The resulting wound is susceptible and can corrode the surrounding tissue.

Cancer aetiology

According to the ITM, excessive accumulation of abnormal black bile in any part of the body is a significant cause of cancer. Ageing, prolonged exposure to stress, cold and dry foods and hard work are the leading causes of increased black bile production. In some cases, bleeding (such as menstruation, abnormal uterine bleeding, or bleeding haemorrhoids) is a defence mechanism against the accumulation of bad humour in the body. Therefore, stopping the bleeding completely by surgical procedures and other medications can accumulate black bile and increase the risk of cancer and other diseases stemming from excessive black bile (including cancer, melancholia, liver problems, psoriasis).⁶

Cancer management

Traditional Islamic medicine suggests several strategies for cancer control and treatment. Surgical manipulation, venesection, dietary adjustment, and use of natural

medicines including solid, semisolid, and liquid dosage forms by oral and external routes of administration are among these methods.⁷

Surgery and manipulation

Surgery is used to eradicate tumours in the early stages of their development. Small tumours distant from vital organs are good candidates for surgery. The tumour should be excised from its origin, and some of the adjacent unaffected tissue should also be excised. In addition, the bleeding should be allowed to continue until a large volume of blood is expelled, and the surrounding tissue must be compressed to expel the blood mixed with black bile thoroughly. After that, the injured site must be saved or burned. However, suppose the cancerous tumour is located in the vicinity of sensitive and vital organs. In that case, surgical procedures and cauterization will be hazardous and turn cancer into a non-healing ulcer. Regular venesection is also advised during the early stages of cancer to draw the blood of black bile from the body.⁸

Nutritional therapy

In traditional Islamic medicine, dietary recommendations have been proposed to slow the progression of advanced tumours that cannot be manipulated due to metastatic problems. In this case, nutritional treatment will increase the patient's longevity. Foods with a wet temper that produce high-quality blood such as almond oil, fresh small fish, soft boiled egg yolks, lamb, bird meat, ripe and sweet apples, sweet plums, bananas, blackcurrant (*Vigna mungo* (L.) Hepper), spinach (*Spinacia oleracea* L.), pumpkin, light wine, and fresh cow's milk and dough were given. On the other hand, excessive intake of foodstuffs can trigger the production of black bile in the body (such as eggplant (*Solanum melongena* L.), lentils (*Lens esculenta* Moench), dates (*Phoenix dactylifera* L.), cabbage (*Brassica oleracea* L.), meat beef, black and thick wine, and meat preserved with salt is strictly limited.⁹

Pharmacotherapy

From the point of view of traditional Islamic medicine, an excess of abnormal black bile in various parts of the body can lead to cancer formation. In treating cancer, black bile must be removed from the body using appropriate laxatives and prevent the formation and accumulation of black bile in the vessels as far as possible. For this purpose, many single and compound drugs of plant, animal and mineral origin have been recommended. Of the 107 plant species introduced in Iranian Islamic medicine for cancer treatment, 59 plants or their chemical compounds have been shown to have cytotoxic and antitumor activities in recent investigations, and several have entered clinical trials. Their effectiveness has been evaluated in humans.⁷⁻⁸

An essential point in cancer and tumour pharmacotherapy is avoiding caustic and irritant drugs to prevent further stimulation and ulceration. Cancer drugs can be administered internally (oral, enema, vaginal douche) or applied topically (topical oils, liniments, lotions, powders). Local anticancer drugs have the following objectives: cancer eradication, prevention of metastasis, prevention of ulceration, and tumour healing ulceration.

Mechanism of action of anticancer drugs

Anticancer drugs work through different mechanisms, namely as black bile laxative, antiulcer in cancerous wounds, wound healing drugs, and analgesics. Black bile laxatives are common anticancer drugs that can eliminate abnormal black bile from the rest of the body. Laxatives should be given frequently. From ITM's point of view, dodder clover (*Cuscuta epithymum* Murr.) is the most valuable black bile laxative, commonly used to treat diseases caused by excess or imbalance of this humour as all types of cancer, melancholia, leprosy, and vitiligo. For this purpose, a mixture of clover dodder with cheese whey or hydromel should be given frequently. A decoction of the plant in oxymel is also prescribed. Common polypody (*Polypodium vulgare* L.), French lavender (*Lavandula stoechas* L.), colocynth (*Citrullus colocynthis* (L.) Schrad.), and black hellebore (*Helleborus niger* L.) are other potent black bile laxatives.^{4,5}

Antiulcer drugs can inhibit tumour ulceration. *Urtica pilulifera* L. and *Aloe vera* (L.) Burm.f. is an example of such a plant. Wound healing drugs accelerate the healing of cancerous wounds. *Althaea officinalis* L., *Brassica oleracea* L., and *Viola odorata* L. have wound healing activity. In addition, analgesic drugs from plants can relieve cancer pain. *Parietaria officinalis* L. and *Solanum nigrum* L. are plants that have analgesic effects.

Conclusion→5

In conclusion, traditional Islamic medicine for cancer has been widely proven in modern research. Of the 107 plant species introduced in Iranian Islamic medicine for cancer treatment, 59 plants or their chemical compounds have been shown to have cytotoxic and antitumor activities in recent investigations, and several have entered clinical trials.

This finding shows the deep insight of Islamic doctors about cancer treatment. Despite the lack of modern facilities and developed equipment on their era, they introduced anticancer plants, showing cytotoxic properties in the latest research. The correlation between these findings signifies the originality of experience and studies, representing valuable funds and valuable knowledge dating back more than twelve centuries. This legacy is based on thousands of years of experience by ancient Greek, Indian and Iranian doctors and relies on many clinical trials on thousands of people. In addition, the application of traditional medicine knowledge reinterpreted by modern data can lead to more effective and evidence-based use of medicinal plants, which can contribute to therapeutic decisions in various diseases.

References→6

1. Jurenka J. Therapeutic applications of pomegranate (*Punica granatum* L.): a review. *Alternative Medicine Review*, 2008. 13 (2): 128-144.
2. E. P. Lansky, W. Jiang, H. Mo et al., Possible synergistic prostate cancer suppression by anatomically discrete pomegranate fractions," *Investigational New Drugs*, vol. 23, no. 1, pp. 11–20, 2005.
3. A. Malik and H. Mukhtar, "Prostate cancer prevention through pomegranate fruit," *Cell Cycle*, vol. 5, no. 4, pp. 371–373, 2006.
4. L. S. Adams, N. P. Seeram, B. B. Aggarwal, Y. Takada, D. Sand, and D. Heber, "Pomegranate juice, total pomegranate ellagitannins, and punicalagin suppress inflammatory cell signaling in colon cancer cells," *Journal of Agricultural and Food Chemistry*, vol. 54, no. 3, pp. 980–985, 2006.

5. N. Khan, F. Afaq, M. H. Kweon, K. Kim, and H. Mukhtar, "Oral consumption of pomegranate fruit extract inhibits growth and progression of primary lung tumors in mice," *Cancer Research*, vol. 67, no. 7, pp. 3475–3482, 2007.
6. V. M. Adhami, N. Khan, and H. Mukhtar, "Cancer chemoprevention by pomegranate: laboratory and clinical evidence," *Nutrition and Cancer*, vol. 61, no. 6, pp. 811–815, 2009.
7. E. Hornung, C. Pernstich, and I. Feussner, "Formation of conjugated Δ^{11} Δ^{13} -double bonds by Δ^{12} -linoleic acid (1,4)-acyl-lipid-desaturase in pomegranate seeds," *European Journal of Biochemistry*, vol. 269, no. 19, pp. 4852–4859, 2002.
8. R. F. Wang, W. D. Xie, Z. Zhang et al., "Bioactive compounds from the seeds of *Punica granatum* (pomegranate)," *Journal of Natural Products*, vol. 67, no. 12, pp. 2096–2098, 2004.
9. Behjat J., Milad I., Seyed A.E. Anticancer plants in islamic traditional medicine. *Complementary Therapies for the Body, Mind and Soul*. IntechOpen. 2015.

Reviewer Comment:

- 1→ Title of Manuscripts should be explained main review and declared type of literature review: narrative or systematic review.
- 2→ Keywords should be showed the main words of the study, the authors can use MeSH to develop keywords.
- 3→ Abstract should be showed the main of background, main of review and conclusion of study.
- 4→ Introduction should be showed the urgency of study (epidemiology data), biological plausibility concept, and lack of knowledge in the study.
- 5→ Conclusion should more specific and not more showed more review.
- 6→ Authors must check the references for make update references. References should no more than 10 years.

Reviewer 2

Cancer Treatment in Islamic Traditional Medicine →1

Rachmat Hidayat^{1*}

¹ Department of Biology, Faculty of Medicine, Universitas Sriwijaya, Palembang, Indonesia

*Corresponding author email: dr.rachmat.hidayat@gmail.com

Abstract →3

Although significant progress has been made in cancer treatment in recent decades, the effectiveness of modern therapeutic approaches is often limited by their toxic effects on other organs. In addition, many people in the world have limited or unequal access to cancer treatment services. Therefore, utilizing information from traditional medicine systems to identify alternative methods to prevent and control cancer. The use of traditional medicine can lead to the discovery of new bioactive compounds and drugs that are available, safe and affordable. In Islamic medicine, cancer is known as a disease with many treatment options. Traditional Islamic medicine suggests several strategies for cancer control and treatment. Surgical manipulation, venesection, dietary adjustment, and use of natural medicines including solid, semisolid, and liquid dosage forms by oral and external routes of administration are among these methods. This literature study will presents cancer from the perspective of traditional Islamic medicine, its aetiology and management.

Keywords. Cancer Treatment, Complementary Medicine, Traditional Medicine, Islamic Medicine. →2

Introduction →4

Islamic traditional medicine (ITM) is a holistic system of medicine that developed during the Islamic Golden Age (750-1258 AD). It is practised and taught throughout the Islamic world. At that time, Muslim and non-Muslim medical scholars, especially Iranian doctors and pharmacists, translated the scientific knowledge they inherited from ancient Greek and Iranian (Persian) culture. Strives to revive and develop the knowledge, remove myths and errors, and impart authentic medicine knowledge. Although most of ITM's scientists are not Arabs, most of the books are written in Arabic. Razi (Razhes; 865-925 AD), Ahwazi (Haly Abbas; 930-994 AD), Avicenna (Ibnu Sina; 980-1037 AD), Jorjani (1042-1136 AD), and Aqili Khorasani (18th century) were prominent Iranian physicians who have the most significant contribution to the promotion of ITM.¹

Although significant progress has been made in cancer treatment in recent decades, the effectiveness of modern therapeutic approaches is often limited by their toxic effects on other organs. In addition, many people in the world have limited or unequal access to cancer treatment services. Therefore, utilizing information from traditional medicine systems to identify alternative methods to prevent and control cancer. In addition, the use of traditional medicine can lead to the discovery of new bioactive compounds and drugs that are available, safe and affordable. In Islamic medicine, cancer is known as a disease with many treatment options. This literature study will discuss cancer, its aetiology and management from the perspective of traditional Islamic medicine.²

Definition of cancer from ITM perspective

ITM is based on humour knowledge, which holds that the human body comprises four essential fluids, namely humour. The four fluids are blood, phlegm, yellow bile, and black bile, and each corresponds to a paired quality: hot and wet blood, cold and wet phlegm, hot and dry yellow bile, and cold and dry black bile. The right balance of humour is essential for maintaining good health. Thus, all diseases and disabilities, including cancer, are caused by excess or decay of this fluid.^{3,4}

According to ITM, cancer is a type of swelling of black bile that is accompanied by pain, throbbing, inflammation, and angiogenesis. The blood vessels around the tumour are full and stiff and contain dark, black blood. The designation "cancer" is due to the similarity between cancerous tumours that affect organs and crabs that catch their prey. The round shape, the darkness of the tumour and the origin of the blood vessels from its environment describe a crab's leg image. Cancer often affects soft and porous organs, and for this reason, cancer mainly involves the breast and innervated organs (such as the uterus) in women and the throat, larynx, testes and penis in men.⁵

Initially, the cancer is the size of a pea or smaller, complex, round, mobile, dark, and slightly warm. Then it will start to grow gradually and reach the size of a walnut or larger. It may be curable during the early stages of its development, but diagnosis is difficult at this stage. On the other hand, cancer treatment will be difficult or even impossible with the development and emergence of clinical manifestations.

Cancers of the eye, nasal cavity, breast, uterus, liver, and other visceral organs and skin are the most frequently mentioned cancers in the ITM text. Cancer is divided into two main types: ulcerative and nonulcerative cancer. Some cancerous tumours ulcerate easily, but some do not. Cancer sores usually spread centripetally, and the discharge is purulent. Appropriate use of drugs can prevent ulceration of susceptible tumours. On the other hand, some cancerous tumours that are not prone to ulceration will ulcerate after inappropriate drug administration.

Advanced and large tumours are susceptible and painful, with a characteristic red to yellow colour and a burning, stinging pain. Such tumours may erupt spontaneously, purulent and bloody fluid may appear on the wound surface. The resulting wound is susceptible and can corrode the surrounding tissue.

Cancer aetiology

According to the ITM, excessive accumulation of abnormal black bile in any part of the body is a significant cause of cancer. Ageing, prolonged exposure to stress, cold and dry foods and hard work are the leading causes of increased black bile production. In some cases, bleeding (such as menstruation, abnormal uterine bleeding, or bleeding haemorrhoids) is a defence mechanism against the accumulation of bad humour in the body. Therefore, stopping the bleeding completely by surgical procedures and other medications can accumulate black bile and increase the risk of cancer and other diseases stemming from excessive black bile (including cancer, melancholia, liver problems, psoriasis).⁶

Cancer management

Traditional Islamic medicine suggests several strategies for cancer control and treatment. Surgical manipulation, venesection, dietary adjustment, and use of natural

medicines including solid, semisolid, and liquid dosage forms by oral and external routes of administration are among these methods.⁷

Surgery and manipulation

Surgery is used to eradicate tumours in the early stages of their development. Small tumours distant from vital organs are good candidates for surgery. The tumour should be excised from its origin, and some of the adjacent unaffected tissue should also be excised. In addition, the bleeding should be allowed to continue until a large volume of blood is expelled, and the surrounding tissue must be compressed to expel the blood mixed with black bile thoroughly. After that, the injured site must be saved or burned. However, suppose the cancerous tumour is located in the vicinity of sensitive and vital organs. In that case, surgical procedures and cauterization will be hazardous and turn cancer into a non-healing ulcer. Regular venesection is also advised during the early stages of cancer to draw the blood of black bile from the body.⁸

Nutritional therapy

In traditional Islamic medicine, dietary recommendations have been proposed to slow the progression of advanced tumours that cannot be manipulated due to metastatic problems. In this case, nutritional treatment will increase the patient's longevity. Foods with a wet temper that produce high-quality blood such as almond oil, fresh small fish, soft boiled egg yolks, lamb, bird meat, ripe and sweet apples, sweet plums, bananas, blackcurrant (*Vigna mungo* (L.) Hepper), spinach (*Spinacia oleracea* L.), pumpkin, light wine, and fresh cow's milk and dough were given. On the other hand, excessive intake of foodstuffs can trigger the production of black bile in the body (such as eggplant (*Solanum melongena* L.), lentils (*Lens esculenta* Moench), dates (*Phoenix dactylifera* L.), cabbage (*Brassica oleracea* L.), meat beef, black and thick wine, and meat preserved with salt is strictly limited.⁹

Pharmacotherapy

From the point of view of traditional Islamic medicine, an excess of abnormal black bile in various parts of the body can lead to cancer formation. In treating cancer, black bile must be removed from the body using appropriate laxatives and prevent the formation and accumulation of black bile in the vessels as far as possible. For this purpose, many single and compound drugs of plant, animal and mineral origin have been recommended. Of the 107 plant species introduced in Iranian Islamic medicine for cancer treatment, 59 plants or their chemical compounds have been shown to have cytotoxic and antitumor activities in recent investigations, and several have entered clinical trials. Their effectiveness has been evaluated in humans.⁷⁻⁸

An essential point in cancer and tumour pharmacotherapy is avoiding caustic and irritant drugs to prevent further stimulation and ulceration. Cancer drugs can be administered internally (oral, enema, vaginal douche) or applied topically (topical oils, liniments, lotions, powders). Local anticancer drugs have the following objectives: cancer eradication, prevention of metastasis, prevention of ulceration, and tumour healing ulceration.

Mechanism of action of anticancer drugs

Anticancer drugs work through different mechanisms, namely as black bile laxative, antiulcer in cancerous wounds, wound healing drugs, and analgesics. Black bile laxatives are common anticancer drugs that can eliminate abnormal black bile from the rest of the body. Laxatives should be given frequently. From ITM's point of view, dodder clover (*Cuscuta epithimum Murr.*) is the most valuable black bile laxative, commonly used to treat diseases caused by excess or imbalance of this humour as all types of cancer, melancholia, leprosy, and vitiligo. For this purpose, a mixture of clover dodder with cheese whey or hydromel should be given frequently. A decoction of the plant in oxymel is also prescribed. Common polypody (*Polypodium vulgare L.*), French lavender (*Lavandula stoechas L.*), colocynth (*Citrullus colocynthis (L.) Schrad.*), and black hellebore (*Helleborus niger L.*) are other potent black bile laxatives.^{4,5}

Antiulcer drugs can inhibit tumour ulceration. *Urtica pilulifera L.* and *Aloe vera (L.) Burm.f.* is an example of such a plant. Wound healing drugs accelerate the healing of cancerous wounds. *Althaea officinalis L.*, *Brassica oleracea L.*, and *Viola odorata L.* have wound healing activity. In addition, analgesic drugs from plants can relieve cancer pain. *Parietaria officinalis L.* and *Solanum nigrum L.* are plants that have analgesic effects.

Conclusion→5

In conclusion, traditional Islamic medicine for cancer has been widely proven in modern research. Of the 107 plant species introduced in Iranian Islamic medicine for cancer treatment, 59 plants or their chemical compounds have been shown to have cytotoxic and antitumor activities in recent investigations, and several have entered clinical trials.

This finding shows the deep insight of Islamic doctors about cancer treatment. Despite the lack of modern facilities and developed equipment on their era, they introduced anticancer plants, showing cytotoxic properties in the latest research. The correlation between these findings signifies the originality of experience and studies, representing valuable funds and valuable knowledge dating back more than twelve centuries. This legacy is based on thousands of years of experience by ancient Greek, Indian and Iranian doctors and relies on many clinical trials on thousands of people. In addition, the application of traditional medicine knowledge reinterpreted by modern data can lead to more effective and evidence-based use of medicinal plants, which can contribute to therapeutic decisions in various diseases.

References→6

1. Jurenka J. Therapeutic applications of pomegranate (*Punica granatum L.*): a review. *Alternative Medicine Review*, 2008. 13 (2): 128-144.
2. E. P. Lansky, W. Jiang, H. Mo et al., Possible synergistic prostate cancer suppression by anatomically discrete pomegranate fractions," *Investigational New Drugs*, vol. 23, no. 1, pp. 11–20, 2005.
3. A. Malik and H. Mukhtar, "Prostate cancer prevention through pomegranate fruit," *Cell Cycle*, vol. 5, no. 4, pp. 371–373, 2006.
4. L. S. Adams, N. P. Seeram, B. B. Aggarwal, Y. Takada, D. Sand, and D. Heber, "Pomegranate juice, total pomegranate ellagitannins, and punicalagin suppress inflammatory cell signaling in colon cancer cells," *Journal of Agricultural and Food Chemistry*, vol. 54, no. 3, pp. 980–985, 2006.

5. N. Khan, F. Afaq, M. H. Kweon, K. Kim, and H. Mukhtar, "Oral consumption of pomegranate fruit extract inhibits growth and progression of primary lung tumors in mice," *Cancer Research*, vol. 67, no. 7, pp. 3475–3482, 2007.
6. V. M. Adhami, N. Khan, and H. Mukhtar, "Cancer chemoprevention by pomegranate: laboratory and clinical evidence," *Nutrition and Cancer*, vol. 61, no. 6, pp. 811–815, 2009.
7. E. Hornung, C. Pernstich, and I. Feussner, "Formation of conjugated Δ^{11} Δ^{13} -double bonds by Δ^{12} -linoleic acid (1,4)-acyl-lipid-desaturase in pomegranate seeds," *European Journal of Biochemistry*, vol. 269, no. 19, pp. 4852–4859, 2002.
8. R. F. Wang, W. D. Xie, Z. Zhang et al., "Bioactive compounds from the seeds of *Punica granatum* (pomegranate)," *Journal of Natural Products*, vol. 67, no. 12, pp. 2096–2098, 2004.
9. Behjat J., Milad I., Seyed A.E. Anticancer plants in islamic traditional medicine. *Complementary Therapies for the Body, Mind and Soul*. IntechOpen. 2015.

Reviewer Comment:

- 1→ Title of Manuscripts should be explained main review and declared type of literature review: narrative or systematic review.
- 2→ Keywords should be showed the main words of the study, the authors can use MeSH to develop keywords.
- 3→ Abstract should be showed the main of background, main of review and conclusion of study.
- 4→ Introduction should be showed the urgency of study (epidemiology data), biological plausibility concept, and lack of knowledge in the study.
- 5→ Conclusion should more specific and not more showed more review.
- 6→ Authors must check the references for make update references. References should no more than 10 years.



ARKUS

Journal Homepage:

<https://hmpublisher.com/index.php/arkus>



Cancer Treatment in Islamic Traditional Medicine

Rachmat Hidayat^{1*}

¹ Department of Biology, Faculty of Medicine, Universitas Sriwijaya, Palembang, Indonesia

ARTICLE INFO

Keywords:

Cancer treatment
Complementary medicine
Traditional medicine
Islamic medicine

Corresponding author:

Rachmat Hidayat

E-mail address:

dr.rachmat.hidayat@gmail.com

The author has reviewed and approved the final version of the manuscript.

<https://doi.org/10.37275/arkus.v7i2.93>

ABSTRACT

Although significant progress has been made in cancer treatment in recent decades, the effectiveness of modern therapeutic approaches is often limited by their toxic effects on other organs. In addition, many people in the world have limited or unequal access to cancer treatment services. Therefore, utilizing information from traditional medicine systems to identify alternative methods to prevent and control cancer. The use of traditional medicine can lead to the discovery of new bioactive compounds and drugs that are available, safe and affordable. In Islamic medicine, cancer is known as a disease with many treatment options. Traditional Islamic medicine suggests several strategies for cancer control and treatment. Surgical manipulation, venesection, dietary adjustment, and use of natural medicines including solid, semisolid, and liquid dosage forms by oral and external routes of administration are among these methods. This literature study will presents cancer from the perspective of traditional Islamic medicine, its aetiology and management.

1. Introduction

Islamic traditional medicine (ITM) is a holistic system of medicine that developed during the Islamic Golden Age (750-1258 AD). It is practised and taught throughout the Islamic world. At that time, Muslim and non-Muslim medical scholars, especially Iranian doctors and pharmacists, translated the scientific knowledge they inherited from ancient Greek and Iranian (Persian) culture. Strives to revive and develop the knowledge, remove myths and errors, and impart authentic medicine knowledge. Although most of ITM's scientists are not Arabs, most of the books are written in Arabic. Razi (Razhes; 865-925 AD), Ahwazi (Haly Abbas; 930-994 AD), Avicenna (Ibnu Sina; 980-1037 AD), Jorjani (1042-1136 AD), and

Aqli Khorasani (18th century) were prominent Iranian physicians who have the most significant contribution to the promotion of ITM.¹

Although significant progress has been made in cancer treatment in recent decades, the effectiveness of modern therapeutic approaches is often limited by their toxic effects on other organs. In addition, many people in the world have limited or unequal access to cancer treatment services. Therefore, utilizing information from traditional medicine systems to identify alternative methods to prevent and control cancer. In addition, the use of traditional medicine can lead to the discovery of new bioactive compounds and drugs that are available, safe and affordable. In



<https://hmpublisher.com/index.php/arkus>

Islamic medicine, cancer is known as a disease with many treatment options. This literature study will discuss cancer, its aetiology and management from the perspective of traditional Islamic medicine.²

Definition of cancer from ITM perspective

ITM is based on humour knowledge, which holds that the human body comprises four essential fluids, namely humour. The four fluids are blood, phlegm, yellow bile, and black bile, and each corresponds to a paired quality: hot and wet blood, cold and wet phlegm, hot and dry yellow bile, and cold and dry black bile. The right balance of humour is essential for maintaining good health. Thus, all diseases and disabilities, including cancer, are caused by excess or decay of this fluid.^{3,4}

According to ITM, cancer is a type of swelling of black bile that is accompanied by pain, throbbing, inflammation, and angiogenesis. The blood vessels around the tumour are full and stiff and contain dark, black blood. The designation "cancer" is due to the similarity between cancerous tumours that affect organs and crabs that catch their prey. The round shape, the darkness of the tumour and the origin of the blood vessels from its environment describe a crab's leg image. Cancer often affects soft and porous organs, and for this reason, cancer mainly involves the breast and innervated organs (such as the uterus) in women and the throat, larynx, testes and penis in men.⁵

Initially, the cancer is the size of a pea or smaller, complex, round, mobile, dark, and slightly warm. Then it will start to grow gradually and reach the size of a walnut or larger. It may be curable during the early stages of its development, but diagnosis is difficult at this stage. On the other hand, cancer treatment will be difficult or even impossible with the development and emergence of clinical manifestations.

Cancers of the eye, nasal cavity, breast, uterus, liver, and other visceral organs and skin are the most

frequently mentioned cancers in the ITM text. Cancer is divided into two main types: ulcerative and nonulcerative cancer. Some cancerous tumours ulcerate easily, but some do not. Cancer sores usually spread centripetally, and the discharge is purulent. Appropriate use of drugs can prevent ulceration of susceptible tumours. On the other hand, some cancerous tumours that are not prone to ulceration will ulcerate after inappropriate drug administration.

Advanced and large tumours are susceptible and painful, with a characteristic red to yellow colour and a burning, stinging pain. Such tumours may erupt spontaneously, purulent and bloody fluid may appear on the wound surface. The resulting wound is susceptible and can corrode the surrounding tissue.

Cancer aetiology

According to the ITM, excessive accumulation of abnormal black bile in any part of the body is a significant cause of cancer. Ageing, prolonged exposure to stress, cold and dry foods and hard work are the leading causes of increased black bile production. In some cases, bleeding (such as menstruation, abnormal uterine bleeding, or bleeding haemorrhoids) is a defence mechanism against the accumulation of bad humour in the body. Therefore, stopping the bleeding completely by surgical procedures and other medications can accumulate black bile and increase the risk of cancer and other diseases stemming from excessive black bile (including cancer, melancholia, liver problems, psoriasis).⁶

Cancer management

Traditional Islamic medicine suggests several strategies for cancer control and treatment. Surgical manipulation, venesection, dietary adjustment, and use of natural medicines including solid, semisolid, and liquid dosage forms by oral and external routes of administration are among these methods.⁷



Surgery and manipulation

Surgery is used to eradicate tumours in the early stages of their development. Small tumours distant from vital organs are good candidates for surgery. The tumour should be excised from its origin, and some of the adjacent unaffected tissue should also be excised. In addition, the bleeding should be allowed to continue until a large volume of blood is expelled, and the surrounding tissue must be compressed to expel the blood mixed with black bile thoroughly. After that, the injured site must be saved or burned. However, suppose the cancerous tumour is located in the vicinity of sensitive and vital organs. In that case, surgical procedures and cauterization will be hazardous and turn cancer into a non-healing ulcer. Regular venesection is also advised during the early stages of cancer to draw the blood of black bile from the body.⁸

Nutritional therapy

In traditional Islamic medicine, dietary recommendations have been proposed to slow the progression of advanced tumours that cannot be manipulated due to metastatic problems. In this case, nutritional treatment will increase the patient's longevity. Foods with a wet temper that produce high-quality blood such as almond oil, fresh small fish, soft boiled egg yolks, lamb, bird meat, ripe and sweet apples, sweet plums, bananas, blackcurrant (*Vigna mungo* (L.) Hepper), spinach (*Spinacia oleracea* L.), pumpkin, light wine, and fresh cow's milk and dough were given. On the other hand, excessive intake of foodstuffs can trigger the production of black bile in the body (such as eggplant (*Solanum melongena* L.), lentils (*Lens esculenta* Moench), dates (*Phoenix dactylifera* L.), cabbage (*Brassica oleracea* L.), meat beef, black and thick wine, and meat preserved with salt is strictly limited.⁹

Pharmacotherapy

From the point of view of traditional Islamic medicine, an excess of abnormal black bile in various parts of the body can lead to cancer formation. In treating cancer, black bile must be removed from the body using appropriate laxatives and prevent the formation and accumulation of black bile in the vessels as far as possible. For this purpose, many single and compound drugs of plant, animal and mineral origin have been recommended. Of the 107 plant species introduced in Iranian Islamic medicine for cancer treatment, 59 plants or their chemical compounds have been shown to have cytotoxic and antitumor activities in recent investigations, and several have entered clinical trials. Their effectiveness has been evaluated in humans.⁷⁻⁸

An essential point in cancer and tumour pharmacotherapy is avoiding caustic and irritant drugs to prevent further stimulation and ulceration. Cancer drugs can be administered internally (oral, enema, vaginal douche) or applied topically (topical oils, liniments, lotions, powders). Local anticancer drugs have the following objectives: cancer eradication, prevention of metastasis, prevention of ulceration, and tumour healing ulceration.

Mechanism of action of anticancer drugs

Anticancer drugs work through different mechanisms, namely as black bile laxative, antiulcer in cancerous wounds, wound healing drugs, and analgesics. Black bile laxatives are common anticancer drugs that can eliminate abnormal black bile from the rest of the body. Laxatives should be given frequently. From ITM's point of view, dodder clover (*Cuscuta epithimum* Murr.) is the most valuable black bile laxative, commonly used to treat diseases caused by excess or imbalance of this humour as all types of cancer, melancholia, leprosy, and vitiligo. For this purpose, a mixture of clover dodder with cheese whey or hydromel should be given frequently. A decoction of the plant in oxymel is also



prescribed. Common polypody (*Polypodium vulgare* L.), French lavender (*Lavandula stoechas* L.), colocynth (*Citrullus colocynthis* (L.) Schrad.), and black hellebore (*Helleborus niger* L.) are other potent black bile laxatives. ^{4,5}

Antiulcer drugs can inhibit tumour ulceration. *Urtica pilulifera* L. and *Aloe vera* (L.) Burm.f. is an example of such a plant. Wound healing drugs accelerate the healing of cancerous wounds. *Althaea officinalis* L., *Brassica oleracea* L., and *Viola odorata* L. have wound healing activity. In addition, analgesic drugs from plants can relieve cancer pain. *Parietaria officinalis* L. and *Solanum nigrum* L. are plants that have analgesic effects.

2. Conclusion

In conclusion, traditional Islamic medicine for cancer has been widely proven in modern research. Of the 107 plant species introduced in Iranian Islamic medicine for cancer treatment, 59 plants or their chemical compounds have been shown to have cytotoxic and antitumor activities in recent investigations, and several have entered clinical trials.

This finding shows the deep insight of Islamic doctors about cancer treatment. Despite the lack of modern facilities and developed equipment on their era, they introduced anticancer plants, showing cytotoxic properties in the latest research. The correlation between these findings signifies the originality of experience and studies, representing valuable funds and valuable knowledge dating back more than twelve centuries. This legacy is based on thousands of years of experience by ancient Greek, Indian and Iranian doctors and relies on many clinical trials on thousands of people. In addition, the application of traditional medicine knowledge reinterpreted by modern data can lead to more effective and evidence-based use of medicinal plants, which can contribute to therapeutic decisions in various diseases.

3. References

1. Jurenka J. Therapeutic applications of pomegranate (*Punica granatum* L.): a review. *Alternative Medicine Review*, 2008; 13(2): 128-144.
2. E. P. Lansky, W. Jiang, H. Mo et al., Possible synergistic prostate cancer suppression by anatomically discrete pomegranate fractions," *Investigational New Drugs*. 2005; 23(1): 11-20,
3. A Malik and H. Mukhtar, "Prostate cancer prevention through pomegranate fruit," *Cell Cycle*. 2006; 5(4): 371-373,
4. L. S. Adams, N. P. Seeram, B. B. Aggarwal, Y. Takada, D. Sand, and D. Heber, "Pomegranate juice, total pomegranate ellagitannins, and punicalagin suppress inflammatory cell signaling in colon cancer cells," *Journal of Agricultural and Food Chemistry*. 2006; 54(3): 980-985.
5. N. Khan, F. Afaq, M. H. Kweon, K. Kim, and H. Mukhtar, "Oral consumption of pomegranate fruit extract inhibits growth and progression of primary lung tumors in mice," *Cancer Research*. 2007; 67(7): 3475-3482,
6. V. M. Adhami, N. Khan, and H. Mukhtar, "Cancer chemoprevention by pomegranate: laboratory and clinical evidence," *Nutrition and Cancer*. 2009; 61(6): 811-815,
7. E. Hornung, C. Pernstich, and I. Feussner, "Formation of conjugated Δ 11 Δ 13-double bonds by Δ 12-linoleic acid (1,4)-acyl-lipid-desaturase in pomegranate seeds," *European Journal of Biochemistry*. 2002; 269(19): 4852-4859,
8. R. F. Wang, W. D. Xie, Z. Zhang et al., "Bioactive compounds from the seeds of *Punica granatum* (pomegranate)," *Journal*



of Natural Products. 2004; 67(12): 2096–2098,

9. Behjat J., Milad I., Seyed A.E. Anticancer plants in islamic traditional medicine.


Complementary Therapies for the Body, Mind and Soul. IntechOpen. 2015.



Letter of Acceptance

Manuscript "Cancer Treatment in Islamic Traditional Medicine" by Rachmat Hidayat*, has been accepted to publish in Arkus Vol 7 issue 2 in October 2021.

Cordially,



Prof. Paula Magnano, PhD

Editor



HM Publisher

(*) Corresponding author

The Corresponding Author can access the account in website :

<https://hmpublisher.com/index.php/arkus/login>

User: rachmat_hdyt

Password: 210587



ARKUS

Journal Homepage:

<https://hmpublisher.com/index.php/arkus>



Cancer Treatment in Islamic Traditional Medicine

Rachmat Hidayat^{1*}

¹ Department of Biology, Faculty of Medicine, Universitas Sriwijaya, Palembang, Indonesia

ARTICLE INFO

Keywords:

Cancer treatment
Complementary medicine
Traditional medicine
Islamic medicine

Corresponding author:

Rachmat Hidayat

E-mail address:

dr.rachmat.hidayat@gmail.com

The author has reviewed and approved the final version of the manuscript.

<https://doi.org/10.37275/arkus.v7i2.93>

ABSTRACT

Although significant progress has been made in cancer treatment in recent decades, the effectiveness of modern therapeutic approaches is often limited by their toxic effects on other organs. In addition, many people in the world have limited or unequal access to cancer treatment services. Therefore, utilizing information from traditional medicine systems to identify alternative methods to prevent and control cancer. The use of traditional medicine can lead to the discovery of new bioactive compounds and drugs that are available, safe and affordable. In Islamic medicine, cancer is known as a disease with many treatment options. Traditional Islamic medicine suggests several strategies for cancer control and treatment. Surgical manipulation, venesection, dietary adjustment, and use of natural medicines including solid, semisolid, and liquid dosage forms by oral and external routes of administration are among these methods. This literature study will presents cancer from the perspective of traditional Islamic medicine, its aetiology and management.

1. Introduction

Islamic traditional medicine (ITM) is a holistic system of medicine that developed during the Islamic Golden Age (750-1258 AD). It is practised and taught throughout the Islamic world. At that time, Muslim and non-Muslim medical scholars, especially Iranian doctors and pharmacists, translated the scientific knowledge they inherited from ancient Greek and Iranian (Persian) culture. Strives to revive and develop the knowledge, remove myths and errors, and impart authentic medicine knowledge. Although most of ITM's scientists are not Arabs, most of the books are written in Arabic. Razi (Razes; 865-925 AD),

Ahwazi (Haly Abbas; 930-994 AD), Avicenna (Ibnu Sina; 980-1037 AD), Jorjani (1042-1136 AD), and Aqili Khorasani (18th century) were prominent Iranian physicians who have the most significant contribution to the promotion of ITM.¹

Although significant progress has been made in cancer treatment in recent decades, the effectiveness of modern therapeutic approaches is often limited by their toxic effects on other organs. In addition, many people in the world have limited or unequal access to cancer treatment services. Therefore, utilizing information from traditional medicine systems to



<https://hmpublisher.com/index.php/arkus>

identify alternative methods to prevent and control cancer. In addition, the use of traditional medicine can lead to the discovery of new bioactive compounds and drugs that are available, safe and affordable. In Islamic medicine, cancer is known as a disease with many treatment options. This literature study will discuss cancer, its aetiology and management from the perspective of traditional Islamic medicine.²

Definition of cancer from ITM perspective

ITM is based on humour knowledge, which holds that the human body comprises four essential fluids, namely humour. The four fluids are blood, phlegm, yellow bile, and black bile, and each corresponds to a paired quality: hot and wet blood, cold and wet phlegm, hot and dry yellow bile, and cold and dry black bile. The right balance of humour is essential for maintaining good health. Thus, all diseases and disabilities, including cancer, are caused by excess or decay of this fluid.^{3,4}

According to ITM, cancer is a type of swelling of black bile that is accompanied by pain, throbbing inflammation, and angiogenesis. The blood vessels around the tumour are full and stiff and contain dark, black blood. The designation "cancer" is due to the similarity between cancerous tumours that affect organs and crabs that catch their prey. The round shape, the darkness of the tumour and the origin of the blood vessels from its environment describe a crab's leg image. Cancer often affects soft and porous organs, and for this reason, cancer mainly involves the breast and innervated organs (such as the uterus) in women and the throat, larynx, testes and penis in men.⁵

Initially, the cancer is the size of a pea or smaller, complex, round, mobile, dark, and slightly warm. Then it will start to grow gradually and reach the size of a walnut or larger. It may be curable during the early stages of its development, but diagnosis is difficult at this stage. On the other hand, cancer treatment will be difficult or even impossible with the

development and emergence of clinical manifestations.

Cancers of the eye, nasal cavity, breast, uterus, liver, and other visceral organs and skin are the most frequently mentioned cancers in the ITM text. Cancer is divided into two main types: ulcerative and nonulcerative cancer. Some cancerous tumours ulcerate easily, but some do not. Cancer sores usually spread centripetally, and the discharge is purulent. Appropriate use of drugs can prevent ulceration of susceptible tumours. On the other hand, some cancerous tumours that are not prone to ulceration will ulcerate after inappropriate drug administration.

Advanced and large tumours are susceptible and painful, with a characteristic red to yellow colour and a burning, stinging pain. Such tumours may erupt spontaneously, purulent and bloody fluid may appear on the wound surface. The resulting wound is susceptible and can corrode the surrounding tissue.

Cancer aetiology

According to the ITM, excessive accumulation of abnormal black bile in any part of the body is a significant cause of cancer. Ageing, prolonged exposure to stress, cold and dry foods and hard work are the leading causes of increased black bile production. In some cases, bleeding (such as menstruation, abnormal uterine bleeding, or bleeding haemorrhoids) is a defence mechanism against the accumulation of bad humour in the body. Therefore, stopping the bleeding completely by surgical procedures and other medications can accumulate black bile and increase the risk of cancer and other diseases stemming from excessive black bile (including cancer, melancholia, liver problems, psoriasis).⁶

Cancer management

Traditional Islamic medicine suggests several strategies for cancer control and treatment. Surgical



manipulation, venesection, dietary adjustment, and use of natural medicines including solid, semisolid, and liquid dosage forms by oral and external routes of administration are among these methods.⁷

Surgery and manipulation

Surgery is used to eradicate tumours in the early stages of their development. Small tumours distant from vital organs are good candidates for surgery. The tumour should be excised from its origin, and some of the adjacent unaffected tissue should also be excised. In addition, the bleeding should be allowed to continue until a large volume of blood is expelled, and the surrounding tissue must be compressed to expel the blood mixed with black bile thoroughly. After that, the injured site must be saved or burned. However, suppose the cancerous tumour is located in the vicinity of sensitive and vital organs. In that case, surgical procedures and cauterization will be hazardous and turn cancer into a non-healing ulcer. Regular venesection is also advised during the early stages of cancer to draw the blood of black bile from the body.⁸

Nutritional therapy

In traditional Islamic medicine, dietary recommendations have been proposed to slow the progression of advanced tumours that cannot be manipulated due to metastatic problems. In this case, nutritional treatment will increase the patient's longevity. Foods with a wet temper that produce high-quality blood such as almond oil, fresh small fish, soft boiled egg yolks, lamb, bird meat, ripe and sweet apples, sweet plums, bananas, blackcurrant (*Vigna mungo* (L.) Hepper), spinach (*Spinacia oleracea* L.), pumpkin, light wine, and fresh cow's milk and dough were given. On the other hand, excessive intake of foodstuffs can trigger the production of black bile in the body (such as eggplant (*Solanum melongena* L.), lentils (*Lens esculenta* Moench), dates (*Phoenix dactylifera* L.), cabbage (*Brassica oleracea* L.), meat

beef, black and thick wine, and meat preserved with salt is strictly limited.⁹

Pharmacotherapy

From the point of view of traditional Islamic medicine, an excess of abnormal black bile in various parts of the body can lead to cancer formation. In treating cancer, black bile must be removed from the body using appropriate laxatives and prevent the formation and accumulation of black bile in the vessels as far as possible. For this purpose, many single and compound drugs of plant, animal and mineral origin have been recommended. Of the 107 plant species introduced in Iranian Islamic medicine for cancer treatment, 59 plants or their chemical compounds have been shown to have cytotoxic and antitumor activities in recent investigations, and several have entered clinical trials. Their effectiveness has been evaluated in humans.⁷⁻⁸

An essential point in cancer and tumour pharmacotherapy is avoiding caustic and irritant drugs to prevent further stimulation and ulceration. Cancer drugs can be administered internally (oral, intravenous, vaginal douche) or applied topically (topical oils, liniments, lotions, powders). Local anticancer drugs have the following objectives: cancer eradication, prevention of metastasis, prevention of ulceration, and tumour healing ulceration.

Mechanism of action of anticancer drugs

Anticancer drugs work through different mechanisms, namely as black bile laxative, antiulcer in cancerous wounds, wound healing drugs, and analgesics. Black bile laxatives are common anticancer drugs that can eliminate abnormal black bile from the rest of the body. Laxatives should be given frequently. From ITM's point of view, dodder clover (*Cuscuta epithimum* Murr.) is the most valuable black bile laxative, commonly used to treat diseases caused by excess or imbalance of this



humour as all types of cancer, melancholia, leprosy, and vitiligo. For this purpose, a mixture of clover dodder with cheese whey or hydromel should be given frequently. A decoction of the plant in oxymel is also prescribed. Common polypody (*Polypodium vulgare* L.), French lavender (*Lavandula stoechas* L.), colocynth (*Citrullus colocynthis* (L.) Schrad.), and black hellebore (*Helleborus niger* L.) are other potent black bile laxatives.^{4,5}

Antiulcer drugs can inhibit tumour ulceration. *Urtica pilulifera* L. and *Aloe vera* (L.) Burm.f. is an example of such a plant. Wound healing drugs accelerate the healing of cancerous wounds. *Althaea officinalis* L., *Brassica oleracea* L., and *Viola odorata* L. have wound healing activity. In addition, analgesic drugs from plants can relieve cancer pain. *Parietaria officinalis* L. and *Solanum nigrum* L. are plants that have analgesic effects.

2. Conclusion

In conclusion, traditional Islamic medicine for cancer has been widely proven in modern research. Of the 107 plant species introduced in Islamic medicine for cancer treatment, 59 plants or their chemical compounds have been shown to have cytotoxic and antitumor activities in recent investigations, and several have entered clinical trials.

This finding shows the deep insight of Islamic doctors about cancer treatment. Despite the lack of modern facilities and developed equipment on their era, they introduced anticancer plants, showing cytotoxic properties in the latest research. The correlation between these findings signifies the originality of experience and studies, representing valuable funds and valuable knowledge dating back more than twelve centuries. This legacy is based on thousands of years of experience by ancient Greek, Indian and Iranian doctors and relies on many clinical trials on thousands of people. In addition, the application of traditional medicine knowledge

reinterpreted by modern data can lead to more effective and evidence-based use of medicinal plants, which can contribute to therapeutic decisions in various diseases.

3. References

1. Jurenka J. Therapeutic applications of pomegranate (*Punica granatum* L.): a review. *Alternative Medicine Review*, 2008; 13(2): 128-144.
2. E. P. Lansky, W. Jiang, H. Mo et al., Possible synergistic prostate cancer suppression by anatomically discrete pomegranate fractions," *Investigational New Drugs*. 2005; 23(1): 11-20,
3. A Malik and H. Mukhtar, "Prostate cancer prevention through pomegranate fruit," *Cell Cycle*, 2006; 5(4): 371-373, S. Adnan, M. P. Seeram, B. B. Aggarwal, Y. Nakada, D. Sand, and D. Heber, "Pomegranate juice, total pomegranate ellagitannins, and punicalagin suppress inflammatory cell signaling in colon cancer cells," *Journal of Agricultural and Food Chemistry*. 2006; 54(3): 980-985.
5. N. Khan, F. Afaq, M. H. Kweon, K. Kim, and H. Mukhtar, "Oral consumption of pomegranate fruit extract inhibits growth and progression of primary lung tumors in mice," *Cancer Research*. 2007; 67(7): 3475-3482,
6. V. M. Adhami, N. Khan, and H. Mukhtar, "Cancer chemoprevention by pomegranate: laboratory and clinical evidence," *Nutrition and Cancer*. 2009; 61(6): 811-815,
7. E. Hornung, C. Pernstich, and I. Feussner, "Formation of conjugated $\Delta 11$ $\Delta 13$ -double bonds by $\Delta 12$ -linoleic acid (1,4)-acyl-lipid-desaturase in pomegranate seeds," *European Journal of Biochemistry*. 2002; 269(19): 4852-4859,



8. R. F. Wang, W. D. Xie, Z. Zhang et al., "Bioactive compounds from the seeds of Punica granatum (pomegranate)," Journal of Natural Products. 2004; 67(12): 2096–2098,

9. Behjat J., Milad I., Seyed A.E. Anticancer plants in islamic traditional medicine. Complementary Therapies for the Body, Mind and Soul. IntechOpen. 2015.

Galey



CERTIFICATE

O F P U B L I C A T I O N

For the article titled:

Cancer Treatment in Islamic Traditional Medicine

Authored by;

Rachmat Hidayat

Published in

Arkus Volume 7 Issue 2 2021

Indexed in:

