

# Geothermal Study of the Airklinсар Geothermal Field Empat Lawang District, Sumatera Selatan Province, Indonesia

*by Frinsyah Virgo*

---

**Submission date:** 25-May-2023 08:18PM (UTC+0700)

**Submission ID:** 2101622590

**File name:** IJBAS-IJENS\_FV2022.pdf (256.51K)

**Word count:** 2576

**Character count:** 14484

# Geothermal Study of the Airklinsar Geothermal Field Empat Lawang District, Sumatera Selatan Province, Indonesia

Suharno<sup>1</sup>, F. Virgo<sup>2,3</sup>, and Wahyudi<sup>3</sup>

1. Geophysic Department of Lampung University

2. Physic Department Sriwijaya University

3. The geophysic of Physic Departmen of Gadjah Mada University

**Abstract-** The first geothermal exploration activities in Empat Lawang District, Soth Sumatera Province, Indonesia were carried out in the Airklinsar field, Ulu Musi Sub-District. The discharge test measurements show that wells Airklinsar spring produce predominantly from a single-phase, reservoir (liquid) with a temperature of 225°C and 223°C. The temperature analisis using silica and Na-Mg geothermometers. The chemical composition of the reservoir water was analyzed by standard methods and subsequently classified using Cl-SO<sub>4</sub>, HCO<sub>3</sub> and Cl-Li-B triangular diagrams. A Na-K-Mg triangular diagram was used to classify waters according to the state of equilibrium at given temperatures. The geothermal waters are of chloride type and from a relatively old hydrothermal system. Thermal fluid is in equilibrium with reservoir rocks. The chemical geothermometers were used to predict subsurface temperature. Silica geothermometers give temperature more close to measured temperature whereas cation geothermometers show significantly higher Silica geothermometers.

**Index Term--** geological, geochemical, geothermometers, geoinicator, Airklinsar geothermalfiel

## I. INTRODUCTION

The Airklinsar geothermal field is one of the prospective areas in Sumatera Selatan Province, Indonesia for geothermal exploration which is located on one of the Bukit Bukit barisan mountaneous. In the other hands, its also is located on one of the major tectonic belts of the Fault Sumatera System. The hot springs are located mainly on major active fractures and volcanic areas. The Airklinsar is one of the interisting areas in Empat Lawang District. The Airklinsar is one of the prospective areas in Empat Lawang District, Sumatera Selatan Province for geothermal exploration which was initiated by Virgo Team are the first in June 2012.

## II. MATERIAL AND METHOD

### A. Geology of Airklinsar

The Airklinsar geothermal field is situated on the major tectonic belts of the Fault Sumatera System. There are volcanic areas spread throughout the country. The country has many hot springs with a variety of temperatures ranging up to 98°C. The hot springs are located mainly on major active

fractures and volcanic areas one of which is Airklinsar. The Airklinsar hot spring is spreadout within the Ulu Musi and Talang Padang Sub-District of Empat Lawang District. The geological of the Ulu Musi and Talang Padang Sub-District extends elongated the Fault Sumatera System between the Sumatera Selatan and Bengkulu Provinces, bisides of Bukit Barisan mountainous. The Fault Sumatera System runs the length of the Sumatera Island and transects the Conozoic volcanic arc [1]. The activity of magma was prominent during middle Miocene extend to Pliocene [2].

The Airklinsar geothermal field is spreadout as shown in Figure 1. The lithologies such as Qa is Aluvium, QTV is Rhyoandesite Volcanic Unit, Tmbt is Baturaja Foration, Tma is Air Benakat Formation, Tomt is Talang Akar Formation, and Tpok is Kikim Formation. The rocks of the Airklinsar geothermal field are divided into six zones that are characterized in Figure 1 modified from [3].




Fig. 1. The Geology of Airklinsar geothermal field within the Ulu Musi and Talang Padang Sub-Districts. Qa is Aluvium, QTV is Rhyoandesite Volcanic Unit, Tmbt is Baturaja Foration, Tma is Air Benakat Formation, Tomt is Talang Akar Formation, and Tpok is Kikim Formation.

### B. Cemical Analysis

The geochemical study is based on discharge water samples collected from the hot spring Airklinsar discharge. The water samples collected from the discharge of two hot water springs. These samples were collected in June 2012. Samples were untreated and included acidified water. Chemical analyses of Na, K, Ca, Mg, B, Li, and SO<sub>4</sub> were

carried out in the site laboratory of Lampung University (Bandar Lampung), and the  $\text{SiO}_2$  pH, Cl,  $\text{HCO}_3$ , in laboratory of Sriwijaya University (Palembang).

The relative Cl,  $\text{SO}_4$  and  $\text{HCO}_3$  contents thermal waters. The composition area the where ionic solute geothermal should be indicated reservoir condition. Li, Cl and B are conservative elements in the geothermal system. They are fixed in fluid phases and have not equilibrated. The conservative elements are the best geoindicators of the origin of the geothermal system. B/Cl ratio and Cl-Li-B ternary diagram were used to indicate the source of the fluid.

Temperature calculation based on silica geothermometer, Na-K geothermometers (Fournier and Giggenbach) [4, 5], K-Mg geothermometer [6]. Besides, the ternary diagram Na-K and K-Mg also use for evaluating equilibration temperatures.

### III. RESULT

The average chemical compositions of the geothermal water from the Airklinsar hot spring discharge are presented in Table I. A plot of the relative concentrations of Cl,  $\text{SO}_4$  and  $\text{HCO}_3$  is illustrated in Figure 2. A plot of the relative concentrations of Cl, Li and B is shown in Figure 3. A plot of the Na-K-Mg ternary diagram shows the equilibrium between the geothermal fluids and rock and reservoir temperature is shown in Figure 4. The subsurface temperatures calculated using geothermometer presented in Table II.

TABLE I  
THE AVERAGE CHEMICAL COMPOSITIONS OF THE GEOTHERMAL WATER FROM AIRKLINSAR HOT SPRING DISCHARGE

No	Sample	(ppm)
1	Na	32
2	K	9.1
3	Ca	133
4	Mg	1.5
5	B	17
6	Li	0.223
7	$\text{SO}_4$	340
8	$\text{HCO}_3$	61
9	$\text{SiO}_2$	50
10	Cl	664

### IV. DISCUSSION

The Airklinsar possesses great potentiality for the utilization of geothermal energy. The region has been centre of attraction to a number of visiting national scientists, encouraging them to collect and analyse geothermal water samples at different localities on a sporadic basis. One of such studies has determined the temperature and reservoir characteristic.

#### A. Classification of the thermal fluids

The average chemical compositions of the geothermal water from the Airklinsar are presented in Table 1. The discharge water from Airklinsar hot spring are of the chlorate type with alkaline pH (6.2-7.8) and with total dissolved solids in range of 500-750 mg/kg. A Cl and Ca are predominant cations with concentration of 664 and 133 mg/l respectively, whereas Mg is present in about (1.5 mg/l). The chemical compositions of the waters were classified on the basis of major ions using the Cl- $\text{SO}_4$ - $\text{HCO}_3$  triangular diagram of Giggenbach (see Figure 2) [7, 8]. The sample plot to area of immature waters and can be classified as Cl-rich geothermal water.




Fig. 2. Airklinsar hot spring situation in the ternary diagram of Cl- $\text{SO}_4$  and  $\text{HCO}_3$ , due to characterised the reservoir.  $\text{HCO}_3/\text{Cl}$  closed to 0.09 with %Cl is 62.3% and % $\text{HCO}_3$  is 5.7%.

Cl and B are conservative elements in the geothermal system. They are fixed in fluid phase and have not equilibrated. The conservative elements are the best geoindicators of the origin of the geothermal system. B/Cl ratio and Cl-Li-B ternary diagram were used to indicate the source of the fluid. A plot of the relative concentrations of Cl, Li and B is shown in Figure 3. The geothermal waters have high Cl content relative to Li and B, indicating that they are from an old hydrothermal system and that fluid migrated from the old basement rock.

The B/Cl ratio of the water from airklinsar hot spring is 0.03 (see Figure 3) [8]. The B/Cl ratio should be representative of andesitic reservoir and rhyolitic system or due to absorption into alteration clays along the geothermal flow. The geothermal waters have high Cl content relative to Li and B, indicating that sample come from an old hydrothermal system and that fluid migrated from the old basement rock.




Fig. 3. Airklinsar hot spring situation in the ternary diagram of Cl-Li and B. B/Cl closed to 0.03 with %Cl and %B are 60% and 40% respectively.

By considering the diagram presented in Figure 4, the Airklinsar hot springs plot in the immature water part, so using the chemical geothermometers is not reliable according to the theory used to establish the diagram. Presumably, fast circulation of fluid through the rock fractures, the sample have not gained equilibrium with rock, (see Na-K-Mg diagram, Figure 4). This causes the water to be immature, considering the ion exchange processes that, equilibrium has not been reached yet with rock minerals because of circulation flow. The sample plot to area of immature waters and can be classified as Cl-rich geothermal water which formed by the interaction of geothermal fluids with the host rock and dilution with low salinity water at depth [9].

**B. Subsurface Temperature Estimation**

Chemical geothermometers are used in order to estimate the reservoir temperature. The important criteria for chemical geothermometer application to thermal spring are the pH, temperature and discharge rate of the spring. Both silica and cation geothermometers are applied to the Airklinsar hot springs. Some give unreliable results such as either lower than spring temperature or extremely high temperature. These equations are based on geothermometers for quartz, which assume that these minerals used in geothermometers, are not in equilibrium with rock – water interaction in reservoir.




Fig. 4. Airklinsar hot spring situation in the ternary diagram of K-Na and Mg, with %Na is 2,4% and %Mg is 91%.

TABLE II  
TEMPERATURE CALCULATION OF THE AIRKLINSAR HOT SPRING CHEMISTRY GEOTHERMOMETER

No.	Geothermometes	Temperatures
1	SiO <sub>2</sub> no steam loss	102,0 <sup>o</sup> C
2	SiO <sub>2</sub> max steam loss	102,7 °C
3	Na-K (Fournier)	325,8 °C
4	Na-K (Giggenbach)	331,5 °C
5	K-Mg	86,8 °C

The silica and cation geothermometers were used for the evaluation of subsurface temperature for well discharges (Table 2). The Na-K-Ca geothermometers of and Giggenbach (1988) were used. Source of temperature equations: T (Na-K) [5, 10], T-SiO<sub>2</sub> no steam loss, T-SiO<sub>2</sub>-maxsteam loss [4, 8, 11, 12]. These give reservoir temperatures ranging from 86<sup>o</sup>C to 330 °C. In comparison with SiO geothermometer, the temperature calculated by the quartz geothermometer [4] is about 102<sup>o</sup>C. It may be inferred that quartz controls the dissolved silica concentration in the deep reservoir [13]. The temperatures obtained from cation geothermometers are significantly higher than from silica geothermometers. The Na-K geothermometer of [4] and [5] suggested reservoir temperature in the range of 326-331<sup>o</sup>C which is higher than SiO geothermometers range of 102-103<sup>o</sup>C [8].

The Na-K-Mg triangular diagram shows the equilibrium between the geothermal fluids and rock and reservoir temperature (Figure 4). Figure 4 shows that samples from studied Airklinsar hot spring geothermal field fall on the immature water, suggested attainment of the water rock equilibrium [8]. By considering the diagram presented in Figure 4, the Airklinsar hot springs plot in the immature water part, so using the chemical geothermometers is not reliable according to the theory used to establish the diagram. In this Na-K-Mg diagram the sample have not gained equilibrium

with rock, presumably due to fast circulation of fluid through the rock fractures. This causes the water to be immature, equilibrium has not been reached yet with rock minerals because of circulation flow. Figure 4 shows that samples from studied Airklinsar hot spring geothermal fields.

### C. Reservoir Estimation

Even though the above table suggests that none of the springs have a huge mass flow rate, a number of springs emerging in the vicinity could have lowered the flow rate. Water containing chloride concentration about 650 ppm from Airklinsar does meet the mixing characteristics since their pH values do lie between 6 and 8. The waters at Airklinsar hot spring lying in this region have relatively high chloride, suggesting that the chloride waters as indicated by the Giggenbach's diagram of concentrations of the major anions, Cl, SO<sub>4</sub> and HCO<sub>3</sub> (Figure 2). The characteristic the reservoir represented the HCO<sub>3</sub>/Cl closed to 0.09 with %Cl is 62.3% and %HCO<sub>3</sub> is 5.7%.

### D. Mixing Characteristics of Thermal Waters

The following characteristics of mixing are described by [8], (1) A huge mass flow rate of spring, (2) Calcite unsaturation, (3) A high concentration of silica relative to discharge temperature. Calcite is either supersaturated, suggesting that there is mixing. No high concentration of silica is observed relative to discharge temperature in the spring waters. It leads to conclude that the waters are mixed.

### V. CONCLUSIONS

The geothermal waters discharged from Airklinsar geothermal field of the Empat Lawang is of chlorate type with a high concentration of Cl. Chemical compositions of reservoir waters indicate that the reservoir is located in the liquid dominant zone and geothermal waters come from an old geothermal system. Thermal fluid is in equilibrium with reservoir rocks, which can be the product of water-rock interaction at high temperature. Common geothermometer have been used for estimating the subsurface temperature. The results from Na-K geothermometers and Na-K-Mg geoindicator, indicate that the reservoir temperature. The calculated temperatures using Na-K geothermometers indicate at a range 326°C to 331°C. However the Na-K-Mg geoindicator is not indicate the clear temperature.

Common geothermometers have been used for estimating the subsurface temperature. The results from silica geothermometers indicate that quartz can control silica concentration in the reservoir or quartz geothermometers better indicate the reservoir temperature. The calculated temperatures using quartz thermometer is 102°C and 102.7°C.

Cation geothermometers give temperature significantly higher than quartz geothermometers.

### ACKNOWLEDGMENT

The authors thank to Virgo Team that prepared the data and gave consent to publish this paper. Gratefully acknowledged to Prof. PRL Browne of the Honorary Research Scientist Institute of Earth Science and Engineering, Auckland, New Zealand for his help.

### REFERENCES

- [1] Fitch, F. J., 1972. Plate Convergence, transcurent faults and internal deformation adjacent to Southeast Asia and the western Pacific. *J. Geophys. Res.* 77, p.4432-60.
- [2] de Coster, G.L., 1974. The Geology of central Sumatera Basin. Proceedings of the Indonesian Assotoin of Geologists, 3<sup>rd</sup> Annual Convention, 77-110.
- [3] Gafoer, S., T. C. Amin and R. Pardede, 1994. Geology of Baturaja Quadrangle, Sumatera. Geological Research and Development Centre, Directorate General of Geology and Mineral Resources.
- [4] Fournier, R. O., 1977. Chemical Geothermometers and Mixing Models for Geothermal System, *Geothermics*, 5, (1977), 41-50.
- [5] Giggenbach, W. F., 1988. Geothermal Gas Equilibria. *Geochemica Cosmochimica Acta* 52, (1988), 2749-2765.
- [6] Giggenbach, W.F., 1991. Chemical Techniques in Geothermal Exploration. In D'Amore, F (coordinator), Application of geochemistry in geothermal reservoir development. UNITAR/UNDP publication, Rome, (1991), 119-142.
- [7] Giggenbach, W. F., 1980. Geothermal Solute Equilibria. Derivation of Na-K-Mg-Ca geoindicators. *Geochemica Cosmochimica Acta* 52 (1980).
- [8] Simmons, F. W., 1998. Lecturer Note Book the Geothermal Institute the University of Auckland New Zealand.
- [9] White, D.E. and L.G. Muffler, 1971. Vapour-dominated Hydrothermal System Compared with Hot Water System. *Economic Geology*, 66, (1971), 75-97.
- [10] Fournier, R., and Truesdell A., 1973. An Empirical Na-K-Ca Geoindicators, *Geochim. Cosmochim. Acta*, 37, (1973), 1255-1275. Giggenbach, W.: Geothermal Solute Equilibria. Derivation of Na-K-Mg-Ca Geoindicators, *Geochim. Cosmochim. Acta*, 52, (1988), 2749-2765.
- [11] Virgo, F., Wahyudi, Suharno, A. Zaenudin dan W. Suryanto, 2012. Magnetic Gradient Temperature and Geochemistry Surveys Within Pasema Air Keruh Geothermal Area Empat Lawang District, South Sumatera Province Indonesia. 3rd Joghha International Conference on Physics 2012, Yogyakarta, 18-19 September 2012.
- [12] Arnorsson, S., E. Gunnlaugsson and H. Svavarsson, 1983. The Chemistry of Geothermal Waters in Iceland III. Chemical Geothermometry in Geothermal Investigations, *Geochim. Cosmochim. Acta*, 47, (1983), 567-577.
- [13] D'Amore, F., 1991. Application of Geochemistry in Geothermal Reservoir Development. UNITAR/UNDP publication, Rome, (1991), 93-95.
- [14] Arnorsson, S. (ed), 2000. Isotopic and Chemical Techniques in Geothermal Exploration, Development and Use. Sampling Method, Data Handling, Interpretation. International Atomic Energy Agency, Vienna, (2000), 351 pp.

# Geothermal Study of the Airklinsar Geothermal Field Empat Lawang District, Sumatera Selatan Province, Indonesia

---

ORIGINALITY REPORT

---

**11** %  
SIMILARITY INDEX

**7** %  
INTERNET SOURCES

**5** %  
PUBLICATIONS

**5** %  
STUDENT PAPERS

---

MATCH ALL SOURCES (ONLY SELECTED SOURCE PRINTED)

---

2%  
★ Submitted to University of Edinburgh  
Student Paper

---

Exclude quotes Off  
Exclude bibliography Off

Exclude matches Off