

# The Risk Factors Analysis Occurrence of Chlamydia Infection to Direct Female Seks Workers (DFSW) in Indonesia

*by* Rini Mutahar

---

**Submission date:** 23-May-2023 11:38PM (UTC+0700)

**Submission ID:** 2100183853

**File name:** The\_Risk\_Factors\_Analysis\_Occurrence\_of.pdf (1.53M)

**Word count:** 3790

**Character count:** 18696

## 8 The Risk Factors Analysis Occurrence of Chlamydia Infection to Direct Female Seks Workers (DFSW) in Indonesia

Masayu Gemala Rabiah<sup>1</sup>, Rini Mutahar<sup>1\*</sup> and Rico Januar Sitorus<sup>1</sup>

<sup>22</sup>  
<sup>2</sup> Faculty of Public Health, Universitas Sriwijaya, Ogan Ilir, Sumatera Selatan, Indonesia

6  
**Abstract.** Chlamydia infection is one of the most common sexual 19  
transmitted infections worldwide. Based on data of IBBS 2015 in  
Indonesia, the highest prevalence of chlamydia occurred in direct female  
sex worker group (32.21%). This study a STIs to determine the risk factors  
for chlamydia infection in direct sex workers. This research uses Cross-  
Sectional study design. The population of this study were all direct female  
sex workers as many as 3,789 people with samples in accordance with  
inclusion and exclusion criteria as many as 3.114 people. The prevalence  
2 direct female sex workers with chlamydia infection was 1.018 (32.4%).  
The result of the analysis showed that there was a relationship between age  
(PR: 0.541 95% CI: 0.452-0.643) and length of work as FDSW (PR: 1.,60,  
95% CI: 1.087-1.512) chlamydia infection in female sex workers in  
Indonesia. most dominant variable has an effect on the incidence of  
chlamydia infection that is long wo 2 ng as FDSW after controlled by age  
(PR 1.282 95% CI: 1.087-1.,12). It is expected for the government to  
improve the implementation of prevention program of chlamydia not only  
among female sex worker but also proactive to customer.

### 1 Introduction

Chlamydia Infection is the infection that is caused by more than 30 types of bacteria, viruses, parasites, and fungi which is transmitted through sexual relations both vaginal, anal and oral. However, transmission can also occur from mother to the fetus in the womb or at birth, through blood products, or medical contaminated transfer. STIs can be classified in two type based on how it is cured, the first one is STIs that can be cured such as syphilis, 15  
gonorrhea, chlamydia, trichomoniasis. The other one is STIs that can not be cured like hepatitis B, herpes, Human Immunodeficiency Virus/HIV, dan Human Papiloma Virus/HPV.[1,2] 13

The Data from World Health Organization (WHO) shows that more than 1 million STIs are obtained every day and there are an estimated 357 million STIs (Chlamydia, Gonorrhea, Syphilis and Trichomoniasis), 131 million of which are the incidence of Chlamydia infections that occurred every year in the age group 15-49 year.[3] This sexually transmitted

5  
\*Corresponding author: rini\_mutahar@fkm.unsri.ac.id

infection (STI) is one of the causes of health, social and economic problems in many countries including Indonesia.

Chlamydia infection is one of the most common sexually transmitted infections worldwide.[4.5] In America more than 3 million new cases are reported annually, of which 70-90% are asymptomatic, because of the high rate of infection that is asymptomatic the true prevalence of infection can be higher.[6] Based on the Biological and Behavioral Survey 2015, in Indonesia the highest prevalence of chlamydia was in the DFSW group (32.21%), followed by the IFSW group (30.29%), MSM (18.53%) and Transgender (16.78% ) From these data shows that sexual chlamydia infection is the most common among direct female sex workers (DFSW).[1]

Female sex workers is a core group in STIs transmission. Through the development of the epidemic, pathogens spread from the core group, female sex workers, directly to the customer population. Population this customer will transmit the disease to sexual partner in the common population, thus forming the chain of transmission of chlamydia infection.[1] This infection is very influenced by behavior and style patterns someone's life [7] Some other risk factors are new or multiple sexual partners. Inconsistent contraceptive use, commercial sex workers, low education status and unmarried status[8]

The purpose of this research is to know the factors that influence the incidence of chlamydia infection in direct female sex workers in several major cities in Indonesia with analysis secondary data Integrated Biological and Behavioral Survey (IBBS) 2015 was carried out by the Ministry of poor living conditions Health Affairs.

## 2 Materials and Methods

The design of this research was cross sectional study by using secondary data sources from the 2015 Integrated Biological and Behavioral Survey (IBBS). The independent variables in this study were age, marital status, education level, age at the first sex, number of customers, frequency of sex, consistency of condom used, consistency of vaginal douching, and long working as a direct female sex worker while the dependent variable is the incidence of Chlamydia Infection. The target population in this study were all direct female sex workers in Indonesia.

The inclusion criteria in this study were direct female sex workers who were 15 years of age or older who had commercial sex with at least one customer in the last month and were at the survey site at the time of the survey team's visit , while the exclusion criteria in this study namely incomplete or missing data. The minimum sample size is 260 respondents, after the exclusion criteria were carried out, the total sample in this study was 3.143 respondents. Statistics data analysis using computer software included univariate, bivariate and multivariate (regression logistics prediction factor model ).

### 3 Results and Discussions

Table 1. Dependent and Independent Variable Frequency Distribution

Variable	Amount (n)	Percentage (%)
<b>Chlamydia Infection</b>		
Yes	1018	32.4
No	2125	67.6
<b>Age</b>		
<25 years	781	24.8
≥ 25 years	2362	75.2
<b>Marital Status</b>		
Single	488	15.5
Married	527	16.8
Divorced	1819	57.9
Death divorce	307	9.8
No answer	2	1
<b>Level of Education</b>		
Never go to school	130	4.1
Not completing elementary school / equivalent	482	15.3
Graduated from elementary school	774	24.6
Not graduating from junior high school / equivalent	358	11.4
Graduated from junior high school / equivalent	823	26.2
Not graduating from high school or equivalent	188	6.0
Graduated from high school / equivalent	365	11.6
Academy / College	23	7
<b>First time sex</b>		
< 20 years	2500	79.5
≥ 20 years	643	20.5
<b>Number of Customers</b>		
≥ 5 people	1913	60.9
<5 people	1230	39.1
<b>Frequency of Sex</b>		
≥ 5 times	2129	67.7
<5 times	1014	32.3
<b>Consistency of Condom Use</b>		
Inconsistent	1971	62.7
Consistent	1172	37.3
<b>Vaginal Douching Consistency</b>		
Consistent	1999	63.6
Inconsistent	1144	36.4
<b>Working Period of DFSW</b>		
≥ 4 years	1143	36.4
<4 years	2000	63.6

Based on Table 1. direct female sex workers (DFSW) who were infected with chlamydia were 32.4 %, the majority of DFSW were more than 25 years old at 75.2 %, the majority of respondents were divorced as much as 57.9 %, the majority of direct female sex worker education level graduated from junior high school as much as 26.2 %, most DFSW have sex for the first time at the age of less than 20 years as much as 79,5%, the majority DFSW get customers in the past one week more than 5 people by 60.9%. Most of the direct female sex workers had sex frequency in the past week more than 5 times, namely 67.7%. The majority of direct female sex workers have not consistently used condoms in the past

week as much as 62.7%. A total of 63.6% of female sex workers were immediately consistent in performing *vaginal douching* in the past week. The majority of respondents work as DFSW less than 4 years, which is 63.6% (Table 1).

**Table 2 .** Relationship of Independent Variable with Chlamydia Infections in Direct Female Sex Workers in Indonesia

Chlamydia Infection								
Variable	9 Yes		No		Total N	%	p- value	PR 95% CI
	n	%	n	%				
<b>Age</b>								
< 25 years	344	44.0	437	56.0	781	100	<0.0001*	1.544
≥ 25 years	674	28.5	1688	71.5	2362	100		1.394-1.709
<b>Marital status:</b>								
Single	870	33.3	1746	66.7	2616	100	0.024*	1.184
Married	148	28.1	379	71.9	527	100		1.022-1.372
<b>Level of education</b>								
Low (<High school)	871	31.6	1884	68.4	2755	100	0.016*	0.834
High (> High School)	147	37.9	241	62.1	388	100		0.726-0.959
<b>First Time Sex</b>								
< 20 years	828	33.1	1672	66.9	2500	100	0.093	1.121
≥ 20 years	190	29.5	453	70.5	643	100		0.983-1.279
<b>Number of Customers</b>								
≥ 5 people	627	32.8	1286	67.2	1913	100	0.590	1.031
< 5 people	391	31.8	839	68.2	1230	100		0.929-1.144
<b>Frequency of Sex</b>								
≥ 5 times	716	33.6	1413	66.4	2129	100	0.035*	1.129
< 5 times	302	29.8	712	70.2	1014	100		1.010-1.263
<b>Consistency of Condoms Use</b>								
Inconsistent	657	33.3	1314	66.7	1971	100	0.154	1.082
Consistent	361	30.8	811	69.2	1172	100		0.973-1.203
<b>Vaginal Douching Consistency</b>								
Consistent	638	31.9	1361	68.1	1999	100	0.478	0.961
Inconsistent	380	33.2	764	66.8	1144	100		0.866-1.066
<b>Working Period of DFSW</b>								
≥ 4 years	310	27.1	833	72.9	1143	100	<0.0001*	0.766
< 4 years	708	35.4	1292	64.9	2000	100		0.685-0.885

\* Significant at 5% alpha

10  
 The results of the bivariate analysis show that the variables associated with the incidence of chlamydia infection in direct female sex workers included Age (PR = 1.544), Marital Status (PR = 1.184), Level of Education (PR = 0.834), Frequency of Sex (PR = 1.129), and the duration worked as a DFSW (PR = 0.766). While the unrelated variables were the age of first sex, number of customers, consistency of condom use, and consistency of vaginal douching (Table 2).

**Table 3.** Multivariate Analysis Results

Variable	Early Modeling of Multivariate Analysis			Final Model of Multivariate Analysis		
	p-value	PR	95% CI	p-value	PR	95% CI
Age	<0.0001	0.562	0.471-0.670	<0.0001	0.541	0.456-0.643
Marital status	0.117	0.845	0.685-1.043	-	-	-
Education Level	0.051	1,253	0.999-1.570	-	-	-
First time sex	0.589	0.947	0.779-1.152	-	-	-
Frequency of Sex	0.233	0.904	0.767-1.067	-	-	-
Consistency of Condoms	0.256	0.912	0.779-1.069	-	-	-
Working period as a DFSW	0.006	1.260	1.067-1488	0.003	1.282	1.087-1.512

The results of the analysis show that the most dominant variable influencing the incidence of chlamydia infection in direct female sex workers was the duration of work as DFSW after being controlled by the age variable. The result of the analysis is known by the table above that the time working variable as DFSW obtained the largest PR value of 1.282 (95% CI: 1.087-1.512). Direct female sex workers who work more than 4 years 1.282 times higher risk of chlamydia infection compared to direct female sex workers who work less than 4 years (Table 3).

The data which used by researchers in analyzing the risk factors for the incidence of chlamydia infection in direct female sex workers in Indonesia are data from the 2015 Integrated Biology and Behavior Survey (IBBS) with cross sectional design in order to collect data of STBP respondent. The deficiency of this research is the study design is difficult seeing the causal relationship or in explaining which conditions are first of a condition of health problems studied because of the measurement of variables carried out at the same time. Another deficiency in this study is using secondary data that have different purposes from the research purposes.

### 3.1 Occurrence of Chlamydia Infection

Chlamydia infection is the most common sexually transmitted infection in the world. This infection occurs without symptoms that typically distinguish cervicitis due to Chlamydia trachomatis and cervicitis due to other organisms. The high level of asymptomatic infection is one of the factors that influence the prevalence of infections that can actually be higher [3].

Direct female sex worker is a core group in STI's transmission.[1] Factor of risk for chlamydia infection in female sex workers directly, among others, the young (<25 years), sexual partner with urethritis, multiple sex partners, swab endocervical cause bleeding, presence of secretions endocervical mucopurulent, contraceptives such as condoms inconsistently or not using condoms during sex, low educational status and unmarried status[9] The prevalence of direct female sex workers who experienced chlamydia infection was 1.018 respondents ( 32.4 %) and who did not experienced chlamydia infection 2.125 respondents ( 67.6 %).

### 3.2 Age of Direct Female Sex Workers

In this study shows that the majority of of DFSW aged < 25 years with chlamydia infection was 44%. Based on the *chi-square* calculation, the *p-value* is <0.0001 ( *p-value*

<0.05). Respondents with age less than 25 years has an opportunity of 1.544 times to be at risk of developing chlamydia infection compared with respondents over the age of 25 years.

The results of this study are supported by Puspita (2017) which states that there is a significant relationship between age with the incidence of chlamydia infection (p-value = 0.012) OR value of 3.683. [9] Young age (less than 25 years) is the most powerful factor associated with Chlamydia infection there are direct female sex workers.[10,15] This can be explained by anatomical differences in the cervix of a younger woman, where the junction is *squamo-columnar*, the main host target for *C. trachomatis* which is everted and thus more open.[8,19] Marcone et.al research in Rome states that age factors are associated with the incidence of Chlamydia Infection. Women with age less than 25 years were 12% at risk of developing chlamydia infection compared to women aged more than 25 years (OR: 2.33 95% CI: 1.02- 5.31; p = 0.04)[11].

### 3.3 Marital Status

The result of this study indicate that status of marriage is a risk factor for the incidence of chlamydia infection. On status variables marriage obtained a PR value of 1.184 means that respondents who are not married / divorced have a risk of 1.184 times for the risk of developing chlamydia infection compared with respondents with married status (95% CI : 1.04-1.372) [12].

This research is also in line with research conducted in Indonesia in 2017 which states there is a relationship between marital status and the incidence of chlamydia infection (p-value = 0.035) with an OR value of (3.189 95% CI: 1.1-8.5) [9] Supported by research in India in 2013 which showed that status divorced and unmarried related with sexual risk OR 3.24, 95% CI: 1.63 -6.43 [13].

Unmarried status can increase the risk of getting STIs includes chlamydia infection compared to married respondents.[23,24] Unmarried status affect sexual needs to be higher compared to someone who has already get married, so it increases the risk of not having sex safe with a partner who can transmit chlamydia infection [9].

### 3.4 Level of Education

A person who is educated has a wider possibility to be able to find information, receive and access various information, especially about the importance and whether or not prevention of chlamydia infection. [14,25] The results of this study indicate that majority of respondents with low education experienced an incidence of chlamydia infection (31.6%). This is evidenced by the results of statistical tests with p-value (0.016 <0.05).

The result of this study is in line with the research conducted by [9] which states that there is a relationship between the level of education and the incidence of chlamydia infection (p-value = 0.040). The value of OR is 3.193( 95% CI: 1.16-74).[9] However, the result of this study contrast with research in Indonesia which states that there is no relationship between the level of education and the incidence of chlamydia infection (p-value 0.727 OR 1.06 95% CI : 0.74-1.51)[16].

Education related to someone's ability to accept and to respond information. In the health side, education factor is classified as individual predisposing factors to utilize health facilities that are due to differences in things knowledge of health, value, and the individual's attitude. Therefore, the sensitivity in facing and overcoming a health problem will be better and ripe than someone with a low education [25].

### 3.5 Frequency of Sex

Frequency of sexual intercourse DFSW have done in the past week. The higher frequency of sex in the past week, the greater the risk of STIs. This is because of the increased chance of contracting STIS at each sexual frequency. [14,21] This study shows that the frequency of having sex more than 5 times per week increases the risk of chlamydia infection

1.129 times higher. However, this research contrasts with [20] study which states that there is no relationship between the frequency of having sex with the incidence of chlamydia infection (OR = 1.21; 95% CI = 0.51 to 2.86; p = 0.661)[18-22].

However, based on the result of the multivariate analysis, it was obtained to see the relationship between the frequency of sex and the incidence of chlamydia infection by analyzing other covariates. Statistical test result showed that there was no relationship between the frequency of having sex with the incidence of chlamydia infection (p-value 0.233; PR 0.904; 95% CI: 0.767-1.067).

### 3.6 Working Period as a Direct Female Sex Worker (DFSW)

The duration of work as DFSW is one of the important factors related to the incidence of chlamydia infection. The longer the service period of a DFSW, the higher possibility serving customers who have been infected with chlamydia. [17,20] The risk of transmission can increase along with the length of time to become DFSW. Chlamydia infection transmission can continue to increase along with the start of work to become DFSW. Becoming a sex worker may not be a living well job, there is the time when DFSW will stop and the number of their customers decreases.

Length work as DFSW is a protective factor against the incidence of chlamydia infection. In the working time variable as DFSW obtained the value of PR 0.766 means that the respondents with a working period of more than 4 years prevents 0.766 times to experience chlamydia infection compared to respondents with less than 4 years of work.

Based on the result of the multivariate analysis obtained to see the old working relationship as DFSW with incidence of chlamydia infection by analyzing other covariates. The results of statistical tests showed that the duration of work as DFSW was the most dominant factor affecting the incidence of chlamydia infection (p-value = 0.003; PR = 1.228 95% CI : 1.087-1.512). Direct female sex workers who work less than 4 years 1,282 times higher risk for chlamydia infection compared to direct female sex workers working directly over 4 years.

## 4 Conclusions

4  
There is a significant relationship between age, marital status, level of education, frequency of sex, working period as DFSW and no relationship between the first age of 10, number of customers, consistency of condoms and consistency of vaginal douching with the incidence of chlamydia infection in direct sex workers in Indonesia. The most dominant variable effect on the incidence of chlamydia infection in women direct sex workers are working period as Direct Female Sex Worker (DFSW) with PR value 1.282 (95% CI: 1.087-1.512).

For the government, it is better to improve the implementation of prevention and control programs for chlamydia infections to reduce the prevalence of chlamydia infections not only among women who are direct sex workers but also need to be carried out proactively to customers in order to increase awareness in women of direct sex workers at the risk of



chlamydia infection, provide capacity building training programs and increase the awareness of the importance of education and provide facilities health services for diagnosis and developing STI clinics at the location / localization of sex workers.

## References

1. Kemenkes RI., *Pedoman Penatalaksanaan Infeksi Menular Seksual* (Dirjen Pengendalian Penyakit dan Penyehatan Lingkungan, Jakarta, 2015)
2. WHO, *Factsheet of Sexually Transmitted Infections* (2013)
3. WHO, *Global Health Sector Strategy On Sexually Transmitted Infections 2016-2021* (2016)
4. Paavonen J., *Ann Med. J. state of the art* **44**, 1 (2012)
5. WHO, *Prevalence and incidence of selected sexually transmitted infections, Chlamydia trachomatis, Neisseria gonorrhoeae, syphilis and Trichomonas vaginalis: methods and results used by WHO to generate 2005 estimates* (Geneva, WHO, 2011)
6. Murtiastutik D., *Infeksi Chlamydia pada wanita. Buku ajar infeksi menular seksual* (Airlangga University Press, Surabaya, 2008)
7. Widodo, E., *J. Jpki*, **2** (2009)
8. Novak M., Novak D., *J. Srh* **4**, 4 (2013)
9. Puspita, Linda, *Jik* **2**, 1 (2017)
10. Darville T., *J. Spid* **1**, 2 (2008)
11. Marcone, V *et al.*, *Es*, **17**, 25 (2012)
12. Lanjouw, E *et al.*, **0**, 0 (2015)
13. Malhotra, Meenakshi et al. *Ijmr, J. Genital Chlamydia Trachomatis : An Update* **30**, 2 (2013)
14. Notoadmodjo, S., *Promosi Kesehatan dan Perilaku Kesehatan*. (Rineka Cipta , Jakarta, 2012)
15. Arifin, Fitriana N, Praba G, Ari U **1**, 2 (2012)
16. Widiastuti, Yani, et al., *Kesehatan Reproduksi* (Yogyakarta, 2009)
17. Karmila, Nelva, *Infeksi Chlamydia Trachomatis* (Fakultas Kedokteran, Universitas Sumatera Utara, 2011)
18. Kusnan, Adius. *Jk*. **4**, 2 (2013)
19. Aridawarni, Yulica, *Jos.A* **2**, 1 (2014)
20. Nugrahaeni, Anita et al., *Jeph, J. Risk Factor of Gonnorrhoea Among Female Indirect Sex Worker* **2**, 3 (2017)
21. Setyawulan, *Hubungan praktek Pencegahan Penyakit Menular Seksual dengan Kejadian Penyakit Menular* (2007)
22. Pane, M., Bppt, *Aspek Klinis dan Epidemiologi Penyakit* (2008)
23. Depkes RI., *Surveilans Terpadu Biologi dan Perilaku* (STBP, Jakarta, 2011)
24. Luong, Me-Linh et al., *Jogc, Vaginal Douching, Bacterial vaginosis, adn Spontaneous Pretetm Birth* **32**, 4 (2010)
25. Mitchellc et al., *Std* **38**, 12 (2011)

# The Risk Factors Analysis Occurrence of Chlamydia Infection to Direct Female Seks Workers (DFSW) in Indonesia

## ORIGINALITY REPORT

13%

SIMILARITY INDEX

10%

INTERNET SOURCES

8%

PUBLICATIONS

4%

STUDENT PAPERS

## PRIMARY SOURCES

1	<a href="http://ims.pnzgu.ru">ims.pnzgu.ru</a> Internet Source	2%
2	<a href="http://www.semanticscholar.org">www.semanticscholar.org</a> Internet Source	2%
3	Submitted to Liberty University Student Paper	1%
4	Hardisman, Lusiana El Sinta Bustami, Lailani Najrani. "DETERMINANTS OF EXCLUSIVE BREASTFEEDING IN AIR DINGIN PRIMARY HEALTH CENTER IN 2018", Walter de Gruyter GmbH, 2020 Publication	1%
5	<a href="http://pdfs.semanticscholar.org">pdfs.semanticscholar.org</a> Internet Source	1%
6	<a href="http://www.tulipgroup.com">www.tulipgroup.com</a> Internet Source	1%
7	Submitted to Universitas Indonesia Student Paper	1%
8	<a href="http://healthdocbox.com">healthdocbox.com</a> Internet Source	1%
9	<a href="http://journal.unnes.ac.id">journal.unnes.ac.id</a> Internet Source	1%
10	<a href="http://www.aisyah.journalpress.id">www.aisyah.journalpress.id</a> Internet Source	<1%
11	Hudila Rifa Karmia, Meilinda Agus, Shinta Maya Sari. "FACTORS RELATED TO PREGNANT	<1%

# WOMAN PARTICIPATION IN ATTENDING PREGNANCY CLASS IN WORK AREA OF LUBUK KILANGAN HEALTH CENTER IN 2017", Walter de Gruyter GmbH, 2020

Publication

12	<a href="https://eurosurveillance.org">eurosurveillance.org</a> Internet Source	<1 %
13	<a href="http://www.afro.who.int">www.afro.who.int</a> Internet Source	<1 %
14	<a href="http://www.balimedicaljournal.org">www.balimedicaljournal.org</a> Internet Source	<1 %
15	<a href="http://balitbangkota.palangkaraya.go.id">balitbangkota.palangkaraya.go.id</a> Internet Source	<1 %
16	<a href="http://ejournal.umpri.ac.id">ejournal.umpri.ac.id</a> Internet Source	<1 %
17	<a href="http://researchonline.lshtm.ac.uk">researchonline.lshtm.ac.uk</a> Internet Source	<1 %
18	<a href="http://eprints.ulm.ac.id">eprints.ulm.ac.id</a> Internet Source	<1 %
19	Hanny Nilasari, Roro Inge Ade Krisanti, Yeva Rosana, Fitri Azizah. "Diagnostic value of the QuickStripe™ chlamydia rapid test among high-risk women in Jakarta", International Journal of STD & AIDS, 2022 Publication	<1 %
20	<a href="http://bmcinfectdis.biomedcentral.com">bmcinfectdis.biomedcentral.com</a> Internet Source	<1 %
21	<a href="http://medicopublication.com">medicopublication.com</a> Internet Source	<1 %
22	<a href="http://sportdocbox.com">sportdocbox.com</a> Internet Source	<1 %
23	Linda Puspita. "Analisis Faktor yang Berhubungan dengan Kejadian Infeksi	<1 %

Menular Seksual pada Wanita Pekerja  
Seksual", Jurnal Aisyah : Jurnal Ilmu  
Kesehatan, 2017

Publication

---

24

Muhammad Asrol, Hakara Warid,  
Ardhianiswari Diah Ekawati. "Triple Bottom  
Line Analysis and Assessment Towards  
Sustainable Palm Oil Plantation", E3S Web of  
Conferences, 2023

Publication

---

<1 %

---

Exclude quotes Off

Exclude matches Off

Exclude bibliography On