PAPER • OPEN ACCESS

The dynamics of rainfall and temperature on peatland in South Sumatra during the 2019 extreme dry season

To cite this article: M Irfan et al 2021 J. Phys.: Conf. Ser. 1940 012030

View the article online for updates and enhancements.

You may also like

- <u>What is the rate of groundwater level</u> <u>decline on peatlands in South Sumatera</u> <u>during the 2019 extreme dry season?</u> M Irfan, O C Satya, Arsali et al.
- Food security prospects of rural community in the change and degraded peatland landscape of South Sumatra B Winarno, S Lestari, Ramawati et al.
- Determination of soil moisture reduction rate on peatlands in South Sumatera due to the 2019 extreme dry season
 M Irfan, E Koriyanti, Awaluddin et al.



245th ECS Meeting

San Francisco, CA May 26–30, 2024

PRIME 2024 Honolulu, Hawaii October 6–11, 2024 Bringing together industry, researchers, and government across 50 symposia in electrochemistry and solid state science and technology

Learn more about ECS Meetings at http://www.electrochem.org/upcoming-meetings



Save the Dates for future ECS Meetings!

The dynamics of rainfall and temperature on peatland in South Sumatra during the 2019 extreme dry season

M Irfan^{1,2,*}, F Virgo¹, M Y N Khakim¹, M Ariani¹, A Sulaiman³, I Iskandar^{1,2}

¹Department of Physics, Faculty of Mathematics and Natural Sciences, Sriwijaya University, Indonesia;

²Graduate School of Sciences, Faculty of Mathematics and Natural Sciences, Sriwijaya University, Indonesia;

³Agency for Assessment and Application of Technology, Jakarta, Indonesia

*irfplg@yahoo.com

Abstract. During the extreme dry season of 2019 massive fires broke out on peatlands in South Sumatera. This study examines the dynamics of rainfall and temperature in the peatlands of South Sumatera in the 2019 dry season as one of the fire disaster mitigation efforts. The data used are in situ measurement data by the Peatland Restoration Agency's measurement stations on two peatlands in South Sumatera. The results of this study indicate that rainfall in July until October 2019 was very minimal even in one of the study sites in August there was no rain. This shows that rainfall anomaly has occurred along with massive fires so that rainfall is possible to be used as one of the fire control parameters in peatlands. The lack of rainfall in South Sumatera during this period was due to the positive Indian Ocean Dipole phenomenon that occurred in the Indian Ocean. The results of this study also showed that temperature did not have a clear pattern of relationship with fire events on peatlands.

1. Introduction

Indonesia is a country that has very large tropical peatlands, which are around 20 million hectares. These peatlands are spread across almost all parts of Indonesia, especially on the islands of Sumatera, Papua and Kalimantan. The area of peatland on the island of Sumatera is around 6 million hectares, mainly in South Sumatera, Riau and Jambi provinces^[1-6].

In every dry season, peatlands burn so that prevention is necessary. In the extreme dry season of 2019, there were massive fires on peatlands in South Sumatera which burned 328,457 hectares of peatlands ^[7]. One of the prevention methods is controlling the parameters related to peatland fire events. Parameters estimated to be closely related to peatland fires include rainfall and temperature. Therefore, it is necessary to study the dynamics of rainfall and temperature through an accurate measurement system for prevention of future fires.

So far, rainfall and temperature data on peatlands only consist of remote sensing data from satellite measurements ^[8-13]. In-situ measurements are believed to be closer to the true value when compared to the results of remote sensing measurements. The Indonesian government, through the Peatland Restoration Agency, has set up several stations to measure in-situ parameters related to peatland fires. The parameters are rainfall, temperature, groundwater level, and soil moisture [14,15]. In South Sumatera, several of these stations have been established since July 2017. This study uses rainfall and

Content from this work may be used under the terms of the Creative Commons Attribution 3.0 licence. Any further distribution of this work must maintain attribution to the author(s) and the title of the work, journal citation and DOI. Published under licence by IOP Publishing Ltd 1

temperature data from the measurement results of these stations to assess whether these two parameters have changed significantly during the dry season in 2019.

2. Methodology

2.1. Data

The data used in this study came from the results of in-situ measurements at two stations belonging to the Indonesian Peatland Restoration Agency (BRG) which are located on peatlands in South Sumatera. These data are hourly rainfall and temperature measured for the period 1 January 2019 to 31 December 2019. Station names and coordinates are shown in Table 1.

Table 1. Name of the BRG meas	suring station and its coordinates
-------------------------------	------------------------------------

Station Name	Coordinate
Sungai Saleh-1 (SS1)	-2.911, 105.082
Sungai Lumpur-1 (SL1)	-2.677, 105.143

2.2. Data analysis

Rainfall and temperature data that have been obtained are processed to obtain daily average data and monthly average data. Daily average data is displayed in the form of time series graph, and monthly average data is shown in tabular form. Analysis was carried out on the time series graphs and tables obtained to assess whether there was a significant change in the value of rainfall and temperature during the extreme dry season in 2019. If there is a significant change in value, it can be concluded that these parameters have a close relationship with fire events on peatlands in South Sumatera. This parameter can be used as one of the controlling parameters for fire prevention in peatlands.

3. Result and Discussion

3.1. The dynamics of rainfall

The study of the dynamics of daily rainfall is carried out through analysis of daily rainfall time series graph and monthly rainfall data table. The time series graph of the dynamics of daily rainfall at two measurement stations SS1 and SL1 during 2019 is shown in Figure 1. In Figure 1, it can be seen that daily rainfall decreased drastically in July until October 2019.




Figure 1. Time series of daily rainfall at (a) SS1 and (b) SL1

Table 2 shows the monthly rainfall at these 2 locations. In Table 2, it can be seen that the monthly rainfall in July until October 2019 is much lower than the other months, even in August there was almost no rain at all. This is what causes the peatlands to become very dry, which causes massive fires on the peatlands. The period from July to October 2019 is called the extreme dry season.

1940 (2021) 012030 doi:10.1088/1742-6596/1940/1/012030


Figure 1 and Table 2 show a close relationship between rainfall and fire events on peatlands in South Sumatera. If the parameters of rainfall can be controlled, it can be used as an alternative effort to prevent fires in peatlands.

This extreme drought has had a negative impact on the agriculture, water resources, forestry and environment sectors. This extreme drought is triggered by the anomalous phenomenon of sea surface temperature (SST) in the Indian Ocean. In this phenomenon, the sea surface temperature in East Africa is warmer than the sea surface temperature in Southwest Sumatera. This phenomenon is called the Indian Ocean Dipole (IOD) which strengthened from April 2019 to December 2019 and caused very low rainfall in the dry season period from July to October 2019^[14, 16-20].

	•	
Month	Rainfall (mm)	
	SS1	SL1
January	90.5	243
February	196	176
March	197.5	369
April	151.5	215.5
May	80	71
June	91.5	106
July	39.5	64
August	0	3
September	2.5	14
October	4	48.5
November	166.5	113.5
December	257	359.5

Table 2. Monthly rainfall

In general, the dry season in 2019 shows drier conditions than the 2018 dry season and the climatological normal reference for 1981-2010, although not drier than the dry season conditions in 2015 when there was a strong El Nino phenomenon at that time.



Indian Ocean Diploe (IOD) is an ocean-atmosphere phenomenon in the equatorial region of the Indian Ocean that affects the climate in Indonesia and other countries around the Indian Ocean basin. As the name implies, IOD is characterized by an anomaly of Sea Surface Temperature (SST) between the 'two poles' of the Indian Ocean, namely the West Indian Ocean (50E-70E, 10S-10N) and Southeast (90E-110E, 10S-0S). Horizontal temperature variations at sea level are generally influenced by the position (radiation) of the sun and water mass.

The difference in SST anomaly between the two regions in the West and Southeast Indian Ocean is called the Dipole Mode Index (DMI), and is used to measure the strength of the IOD itself. The period in which the DMI is positive is generally referred to as the positive IOD period (IOD +), and vice versa, when the DMI is negative it is called the negative IOD period (IOD-) ^[21, 22]. DMI in 2019 has a high value as shown in Figure 2 ^[18].

In the IOD + period, the waters in the Southeast Indian Ocean were generally colder (temperatures are lower than average), whereas the waters in the western Indian Ocean would be warmer (temperatures higher than average). As a result, convection (which is the initial process of cloud and rain formation) will shift from the East Indian Ocean towards the West, and bring a lot of rain to the eastern part of the African continent. On the other hand, areas of the Eastern Indian Ocean that are "left behind" by convection (such as Indonesia) will suffer from drought.

The IOD- period characteristic is the opposite of IOD +. SST in the Southeast Indian Ocean will be warmer, while in the West it will be colder. This will result in drought in Eastern Africa and increased rainfall in Indonesia, especially West Indonesia which is adjacent to the Indian Ocean.

3.2. The dynamics of temperature

The temperature data used is the hourly average T data which is processed into average temperature per day and per month originating from 2 BRG stations, namely: SL1 and SS1 for the period January 1, 2019 until December 31, 2019. The daily average temperature in the form of a time series graph is shown in Figure 3, and the monthly average temperature is shown in Table 3.




Figure 3. Time series of average daily temperature at (a) SS1 and (b) SL1

In Figures 3 (a) and (b), it can be seen that in the dry season from July to October 2019 there was a slight decrease in the daily average temperature. In Table 3, it can be seen that at the SS1 location in July to September 2019 the temperature was slightly lower than the other months, while at the SL1 location there was no significant change in temperature during the extreme dry season.

If we look for the relationship between temperature and the IOD + event, which is from July to October 2019, the temperature of the water surface in the Indian Ocean (SST) in the Southeast is lower than the temperature in the West, so it is estimated that the lower SST temperature is related to the low daily average temperature of peatlands in South Sumatera.

If temperature is to be used as a fire control parameter in peatlands, it is quite difficult to apply because the pattern of the relationship between temperature and the incidence of fire in peatlands is not clear. The relationship between temperature dynamics and fire events on peatlands in South Sumatera is not significant.

1940 (2021) 012030 doi:10.1088/1742-6596/1940/1/012030

IOP Publishing

	Temperature (°C)	
SS1	SL1	
28.04	29.21	
27.72	29.13	
28.68	30.10	
28.38	30.47	
27.91	30.17	
27.21	29.71	
26.70	29.10	
26.43	28.50	
26.45	27.89	
28.48	29.18	
29.28	28.98	
28.45	27.96	
	28.04 27.72 28.68 28.38 27.91 27.21 26.70 26.43 26.45 28.48 29.28	

Table 3. Average monthly temperature

4. Conclusion

The analysis that has been carried out on the dynamics of rainfall and temperature in the extreme dry season of 2019 shows that rainfall has a very significant change in value, while temperature has also changed but the pattern of change is unclear. This very significant change in rainfall indicates a close relationship between rainfall and the occurrence of fires on peatlands during the extreme dry season of 2019. The change in the value of these two parameters is due to the IOD + phenomenon. If it is used to control fires on peatlands, the rainfall parameter is more likely than temperature because rainfall dynamics have a significant relationship with fire events on peatlands.

Acknowledgements

We thank to Sriwijaya University for supporting this research activity through the *Hibah Unggulan Kompetitif* 2019 for the first author and a *Hibah Unggulan Profesi* 2020 for the last author. We also thank to the Agency for Peatland Restoration (BRG) for providing rainfall and temperature data.

References

- [1] Kajita R 2019 Historical precipitation data in Sumatera and Kalimantan from 1879 to 1900, by using Dutch colonial materials *IOP Conf. Ser. Earth Environ. Sci.* **361** (1) p. 012003.
- [2] Yule C M 2010 Loss of biodiversity and ecosystem functioning in Indo-Malayan peat swamp forests *Biodivers Conserv.* **19** (2) pp. 393–409.
- [3] Takeuchi W, Hirano T, Roswintiarti O 2015 Estimation model of groundwater table at peatland in Central Kalimantan, Indonesia *Trop. Peatl. Ecosyst* pp. 445–453
- [4] Kobayashi S 2015 Peatland and peatland forest in Brunei Darussalam *Tropical Peatland Ecosystem* pp.75-89
- [5] Osaki M and Tsuji N 2015 Tropical peatland ecosystems Mitsuru Osaki (Sapporo Japan: Spinger) pp. 1–651
- [6] Irfan M, Kurniawati N, Ariani M, Sulaiman I, and Iskandar I 2020 Study of groundwater level and its correlation to soil moisture on peatlands in South Sumatera," J. Phys. Conf. Ser. 1568 (1).
- [7] Osawa T and Kajita R 2020 News of Indonesian fires in 2019 Newsletter, Toward the Regeneration Tropical Peatland Societies Project **8** pp.8-10.
- [8] Tekeli A E and Fouli H 2017 Reducing False Flood Warnings of TRMM Rain Rates Thresholds over Riyadh City, Saudi Arabia by Utilizing AMSR-E Soil Moisture Information *Water Resour. Manag.* **31** (4) pp. 1243–1256.

- [9] Li Z, Lin X and Cai W 2017 Realism of modelled Indian summer monsoon correlation with the tropical Indo-Pacific affects projected monsoon changes *Sci. Rep.* **7** (1) pp. 1–7.
- [10] Gunawan, Sihombing C, Nugraheni E, and Sudarsono W 2018 the Relationship Between Rainfall, Air Temperature and Wind Speed Effects Dengue Hemorrhagic Fever Case in Bengkulu City At 2009-2014 Januari 2018 JKD 7 (1), pp. 366–380.
- [11] Hidayat, Teuling A J, Vermeulen B, Taufik M, Kastner K 2017 Hydrology of inland tropical lowlands: The Kapuas and Mahakam wetlands *Hydrol. Earth Syst. Sci.* 21 (5) pp. 2579–2594.
- [12] Fistikoglu O, Gunduz O, Simsek C 2016 The Correlation Between Statistically Downscaled Precipitation Data and GWL Records in North-Western Turkey *Water Resour. Manag.* **30** (15) pp. 5625–5635.
- [13] Hoffmann L 2019 From ERA-Interim to ERA5: The considerable impact of ECMWF's nextgeneration reanalysis on Lagrangian transport simulations Atmos. Chem. Phys. 19 (5) pp. 3097–3214.
- [14] Irfan M, Satya O C, Virgo F, Sutopo, Ariani M, Sulaiman A, and Iskandar I 2020 Is there a correlation between rainfall and soil moisture on peatlands in South Sumatra? J. Phys. Conf. Ser. 1572 p. 012040.
- [15] Irfan M, Mardiansyah W, Ariani M, Sulaiman A, and Iskandar I 2019 Is TRMM product good proxy for gauge precipitation over peat land area of the South Sumatera? J. Phys. Conf. Ser. 1282 p. 012021
- [16] Preethi B, Sabin T P, Adedoyin J A, and Ashok K 2015 Impacts of the ENSO Modoki and other tropical indo-pacific climate-drivers on African rainfall Sci. Rep. 5.
- [17] Lim E P and Hendon H H 2017 Causes and Predictability of the Negative Indian Ocean Dipole and Its Impact on la Niña during 2016 Sci. Rep. 7 (1) pp. 1–12
- [18] Doi T, Behera S K, Yamagata, T 2020 Predictibality of the super IOD event in 2019 and its link with El Nino Modoki *Geophysical Research Letters* pp. 1-9.
- [19] Muhammad F R , Lubis S W, Tiarni I, and Setiawan S 2019 Influence of the Indian Ocean Dipole (IOD) on Convectively Coupled Kelvin and Mixed Rossby-Gravity Waves IOP Conf. Ser. Earth Environ. Sci. 284 (1).
- [20] Behera S K and Ratnam J V 2018 Quasi-asymmetric response of the Indian summer monsoon rainfall to opposite phases of the IOD *Sci. Rep.* **8** (1) pp. 1–9.
- [21] Lestari D O, Sutriyono E, Sabaruddin, and Iskandar I 2018 Severe Drought Event in Indonesia Following 2015/16 El Niño/positive Indian Dipole Events J. Phys. Conf. Ser. 1011 (1) pp. 1-5.
- [22] Iskandar I, Irfan M, Syamsuddin F, Johan A, and Poerwono P 2013Trend in Precipitation Over Sumatera Under the Warming Earth *Int. J. Remote Sens. Earth Sci.* 8 (1)