

# Photographic evidence of freshwater whipray *Urogymnus* *dalyensis*(Myliobatiformes: Dasyatidae) in Indonesian waters

*By* Universitas Sriwijaya



Ecologica Montenegrina  
ISSN 2336-0744 (online) ISSN 2337-0173 (print)

*Ecologica Montenegrina* 22: 166-170 (2019)  
This journal is available online at: [www.biotaux.org/em](http://www.biotaux.org/em)

Article

## Photographic evidence of freshwater whipray *Urogymnus dalyensis* (Myliobatiformes: Dasyatidae) in Indonesian waters

YUANITA WINDUSARI<sup>1</sup>, LAILA HANUM<sup>1</sup>, DONI SETIAWAN<sup>1</sup> & MUHAMMAD IQBAL<sup>2\*</sup>

<sup>1</sup> Department of Biology, Faculty of Science, Sriwijaya University, Jalan Raya Palembang-Prabumulih km 32, Indralaya, Sumatera Selatan 30662, Indonesia. Email: [ywindusari@yahoo.com](mailto:ywindusari@yahoo.com)

<sup>2</sup> Conservation Biology Program, Faculty of Science, Sriwijaya University, Jalan Padang Selasa 524, Palembang, Sumatera Selatan 30139, Indonesia. E-mail: [kpbsos26@yahoo.com](mailto:kpbsos26@yahoo.com)

\* Corresponding author [[kpbsos26@yahoo.com](mailto:kpbsos26@yahoo.com)]

Received 13 January 2019 | Accepted by V. Pešić: 18 September 2019 | Published online 29 September 2019.

### Abstract

On 24 December 2017, an individual of *Urogymnus dalyensis* (c. 90–110 cm) was caught and photographed from Lampu Satu beach, Merauke, Indonesia. Lampu Satu beach is an estuarine habitat on sandy substrate surrounded by settlement with mainly coconut trees, and approximately 5 km of Maro river (a muddy substrate surrounded by mangrove vegetation). This finding constitutes the first record of *Urogymnus dalyensis* for Indonesia.

**Key words:** First record; Dasyatidae; *Urogymnus dalyensis*; West Papua; Merauke; estuarine.

### Introduction

The Stingrays (family Dasyatidae) are small to massive rays (adult disc width from 23 cm to at least 220 cm), with the largest estimated to weigh up to 600 kg, making them amongst the largest of all fishes (Last et al., 2016a). Members of the family are found in fresh, brackish, and also marine (continental and insular shelves and uppermost slopes; one species oceanic) waters, and are known from the Atlantic (including Mediterranean Sea), Indian and Pacific oceans (Nelson et al., 2016). Stingray taxonomy is made difficult by the similarity in appearance between many species, and also because they are often poorly represented in museum collections due to their large size (Last and Compagno, 1999; Last et al., 2016a).

There are up to 52 species of stingrays of the family Dasyatidae that are known to enter or to live permanently in freshwaters in the Indo-Pacific region (Last and Stevens, 1994; Last et al., 2016b; White et al. 2017). One of them is the giant freshwater whipray *Urogymnus dalyensis*. This species was previously described as *Himantura chaophraya* (Monkolprasit & Roberts, 1990; Last & Stevens, 1994). Specimens from Australia and Papua New Guinea were then split to become *Himantura dalyensis* (Last and Manjajii-Matsumoto, 2008). More recent revisions have been made within the Dasyatidae family, and the species is now called *Urogymnus dalyensis* (Last et al. 2016a; Last et al., 2016b; Froese and Pauly 2018).

The presence of a single individual of *Urogymnus dalyensis* based photographic evidences, as described in this paper, is the first known record of this species from Indonesian waters.

## Materials and Methods

On 24 December 2017, a single large dasyatid ray was caught and photographed by local people (Fig. 1-2) at Lampu Satu beach ( $08^{\circ}30'47.81''S$ ,  $140^{\circ}22'18.71''E$ ), Merauke district, Papua province, Indonesia (Fig. 3). Lampu Satu beach is an estuarine habitat on sandy substrate surrounded by settlement and mainly coconut trees. This site is 5 km east of the mouth of the Maro River (largest river in Merauke district). The mouth of Maro river is muddy substrate surrounded by mangrove vegetation.

This large dasyatid ray was estimated from photographs to have a disc width of between 90 and 110 cm. Unfortunately the specimen was not preserved, nor detailed morphometric measurements or meristic counts taken. Identification is therefore based on the features that are visible in the photographs.

The ray were immediately determined as *Urogymnus dalyensis* by combination of major features and its distributional range.



**Figure 1.** The specimen of *Urogymnus dalyensis* caught on 24 December 2017 at Lampu Satu beach, Merauke district, Papua province, Indonesia (Photo: Lham Jr.).

## Results and Discussion

The fish was initially identified as *Urogymnus dalyensis* using the following specific characteristics: large body; dorsal surface yellowish brown; plain coloured whipray with oval to subcircular disc with an enlarged triangular apical lobe and tail whip-like without skin folds (as per description by Last and Manjaji-Matsumoto (2008), Last et al. (2016a) and White et al. (2017)) (see Fig. 1-2).

The species is distinguished from other *Urogymnus* species (*Urogymnus acanthobothrium*, *U. asperimus*, *U. granulatus* and *U. lobistoma*) which occur in Australia and Papuan waters by its disc pattern (Last and Stevens, 1994; Last and Manjaji-Matsumoto, 2008; Last et al., 2016a; Last et al., 2016c; White et al., 2017). Our preliminary identification as *Urogymnus dalyensis* was confirmed by Dr. William White and

PHOTOGRAPHIC EVIDENCE OF UROGYMNUS DALYENSIS IN INDONESIAN WATERS


Dr. Mabel Manjaji-Matsumoto, the leading expert for dasyatids of the Indo-Pacific region (*Pers. Comm.* via Muhammad Iqbal). *U. polylepis* is the only similar species to *U. dalyensis* (Last et al., 2016b). The main characters of the upper surface to distinguish *U. dalyensis* from *U. polylepis* are the proportion of the tail length (1.9–2.2 vs. 2.8–3.1), and the size of the mid-shoulder denticles (2–4 greatly enlarged denticles vs 1–6 small or inconspicuous). Unfortunately these characters are not clear in the photographs. However, *U. dalyensis* has a combination of yellowish brown dorsal surface, and length of orbit and spiracle 2.6–3.2 in snout length (Fig. 1-2); vs greyish or brownish dorsal surface, and length of orbit and spiracle 3.3–4.4 in snout length (Last et al. 2016b).



**Figure 2.** Another side of *Urogymnus dalyensis* caught on 24 December 2017 at Lampu Satu beach, Merauke district, Papua province, Indonesia (Photo: Lham Jr.).

*Urogymnus dalyensis* is typically found in the estuarine and freshwater environments of northern Australia, although it may also occur in the Fly River Basin of Papua New Guinea (Last and Stevens, 1994; Last and Manjaji-Matsumoto, 2008; White et al., 2017). Range has not been fully mapped, and it has been suggested that this species may occur in most large tropical rivers of northern Australia (Last and Stevens, 1994; Kyne, 2016).

The record of *U. dalyensis* in Indonesian waters provides additional information that extends the known distribution of this large whipray, although this extension is not that surprising, since it is from about 75 km from Papua New Guinean waters territory, and about 150–200 km from the Fly River of Papua New Guinea (closest area previously known to have *U. dalyensis*). The absence of previous records of *U. dalyensis* from Indonesian Papua waters is likely to be a lack of local experts to identify and document stingray records. Recent studies show a few species of stingrays in Indonesian waters that are present locally but which have been overlooked previously (e.g. Iqbal and Yustian, 2016; Iqbal et al., 2017; Iqbal et al., 2018; Windusari et al., 2019). Genetic samples of *U. dalyensis* from Indonesian Papua would be useful to compare relationships between this Indonesian population, and other known populations from Papua New Guinea and Australian waters (William White *Pers. Comm.*).



**Figure 3.** Map showing location of *Urogymnus dalyensis* (yellow circle) in Lampu Satu beach, Merauke district, Papua province, Indonesia.

#### Acknowledgements

We thank Mr. Lham Jr who shared his photo of a large whipray from Merauke with us. We are very grateful to Dr. William White (CSIRO Marine & Atmospheric Research, Australia) and Dr. Mabel Manjaji-Matsumoto (Borneo Marine Research Institute, Universiti Malaysia Sabah) who confirmed our identification. Thanks also to Dr. William White and Dr. Mat Gilfedder who kindly agreed to proof-read the manuscript. Finally, we would like to thank anonymous reviewers for their useful comments which have improved this paper.

#### References

- Froese, R. & Pauly, D. (eds). (2017) *Urogymnus dalyensis* (Last & Manjaji-Matsumoto, 2008). <https://www.fishbase.de/summary/Urogymnus-dalyensis> [accessed 20/June/2018].
- Iqbal, M. & Yustian, I. (2016) Occurrence of the giant freshwater stingray *Urogymnus polylepis* in Sumatra, Indonesia (Chondrichthyes: Dasyatidae). *Ichthyological Exploration of Freshwaters*, 27(4), 333-336.
- Iqbal, M., Setiawan, D. & Ajiman. (2017) Presence of *Fluvitrygon oxyrhynchus* in Sumatra, Indonesia (Chondrichthyes: Dasyatidae). *Ichthyological Exploration of Freshwaters*, 28, 85-87.

*PHOTOGRAPHIC EVIDENCE OF UROGYMNUS DALYENSIS IN INDONESIAN WATERS*

- Iqbal, M., Setiawan, D. & Ajiman (2018) New data on the distribution of the endangered white-edge freshwater whipray *Fluvitrygon signifer* (Chondrichthyes: Dasyatidae). *Ichthyological Exploration of Freshwaters*, 28(2), 171-176.
- Kyne, P.M. (2016) *Urogymnus dalyensis*. The IUCN Red List of Threatened Species 2016: e.T195319A104250402. <http://dx.doi.org/10.2305/IUCN.UK.2016-3.RLTS.T195319A104250402.en> [accessed 20/June/2018].
- Last, P.R. & Compagno, L.J.V. (1999) Dasyatidae (Stingrays) In: Carpenter K.E. & Niem, V. (Eds.), *FAO species identification guide for fishery purposes. The living marine resources of the Western Central Pacific. Volume 3. Batoid fishes, chimaeras and bony fishes part 1 (Elopidae to Linophrynidae)*. FAO, Rome, pp. 1479-1510.
- Last, P.R. & Manjaji-Matsumoto, B.M. (2008) *Himantura dalyensis* sp. nov., a new estuarine whipray (Myliobatoidei: Dasyatidae) from northern Australia. In: Last, P.R., White, W.T. & Pogonoski, J.J. (Eds.) *Descriptions of new Australian Chondrichthyans*. CSIRO Marine and Atmospheric Research, Hobart, pp. 283-291.
- Last, P.R., Manjaji-Matsumoto, B.M., Naylor, G.J.P. & White, W.T. (2016a) Stingrays (Family Dasyatidae) In: Last, P.R., William, W.T., de Carvalho, M.R., Sérét, B., Stehmann, M.F.W. & Naylor, G.J.P. (Eds.), *Rays of the world*. Cornell University Press, Ithaca, pp. 521-618.
- Last, P.R., Naylor, G.J.P. & Manjaji-Matsumoto, B.M. (2016b) A revised classification of the family Dasyatidae (Chondrichthyes: Myliobatiformes) based on new morphological and molecular insights. *Zootaxa*, 4139, 345-368.
- Last, P.R. & Stevens, J.D. (1994) Sharks and rays of Australia. CSIRO, Australia, 513 pp + 84 pls.
- Last, P.R., White, W.T. & Kyne, P.M. (2016c). *Urogymnus acanthobothrium* sp. nov., a new euryhaline whipray (Myliobatiformes: Dasyatidae) from Australia and Papua New Guinea. *Zootaxa*, 4147(2), 162-176.
- Monkolprasit, S. & Roberts, T.R. (1990) *Himantura chaophryaya*, a new giant freshwater stingray from Thailand. *Japanese Journal of Ichthyology*, 37(3), 203-208.
- Nelson, J.S., Terry, C., Grande, T.C. & Wilson, M.V.H. (2016) Fishes of the World. 5<sup>th</sup>Edit., John Wiley & Sons, Hoboken, 752 pp.
- White, W.T., Baje, L., Sabub, B., Appleyard, S.A., Pogonoski, J.J. & Mana, R.R. (2017) *Sharks and rays of Papua New Guinea*. Australian Centre for International Agricultural Research, Canberra. 327 pp.
- Windusari, Y., Iqbal, M., Hanum, L., Zulkifli, H. & Yustian, I. (2019) Contemporary distribution records of the giant freshwater stingray *Urogymnus polylepis* in Borneo (Chondrichthyes: Dasyatidae). *Ichthyological Exploration of Freshwaters* (in press).

# Photographic evidence of freshwater whipray *Urogymnus dalyensis*(Myliobatiformes: Dasyatidae) in Indonesian waters

---

ORIGINALITY REPORT

---

5%  
SIMILARITY INDEX

---

MATCH ALL SOURCES (ONLY SELECTED SOURCE PRINTED)

---

★M. S. Imanudin, E. Armanto, R. H. Susant, S. M. Bernas. "The study water table fluctuation in tidal lowland for developing agricultural water management strategies: (A case study for corn cultivation after rice)", 2010 International Conference on Chemistry and Chemical Engineering, 2010 1%

[Crossref](#)

---

EXCLUDE QUOTES ON  
EXCLUDE BIBLIOGRAPHY ON

EXCLUDE SOURCES < 1%  
EXCLUDE MATCHES < 10 WORDS

**LEMBAR  
HASIL PENILAIAN SEJAWAT SEBIDANG (PEER REVIEW)  
KARYA ILMIAH: JURNAL ILMIAH**

Judul Artikel Ilmiah	: Photographic evidence of freshwater whipray Urogymnus dalyensis (Myliobatiformes: Dasyatidae) in Indonesian waters
Penulis Artikel Ilmiah	: Yuanita Windusari, Laila Hanum, Doni Setiawan, dan Muhammad Iqbal
Identitas Jurnal Ilmiah	: <ul style="list-style-type: none"> <li>a. Nama Jurnal : Ecologica Montenegrina</li> <li>b. ISSN/ISBN : 2336-9744</li> <li>c. Nomor/Volume/Hal : 1 / 22 / 166 - 170</li> <li>d. Edisi (Bulan/Tahun) : 2019</li> <li>e. Penerbit : Center for Biodiversity of Montenegro</li> <li>f. Jumlah Halaman : 5</li> <li>g. Jurnal terindeks di : Scimagojr (Scopus) dan Clarivate Analytics</li> </ul>

Kategori Publikasi Jurnal Ilmiah :  Jurnal Internasional Bereputasi(terindeks database internasional dan berfaktor dampak)  
 (Beri √ pada kategori yang tepat)  Jurnal Internasional terindeks pada basis data internasional bereputasi  
 Jurnal Internasional terindeks pada basis data internasional  
 Jurnal Nasional Terakreditasi peringkat 1 dan 2  
 Jurnal Nasional Terakreditasi peringkat 3 dan 4

Hasil Penilaian Validasi:

No.	ASPEK	URAIAN/KOMENTAR PENILAIAN
1	Indikasi Plagiasi	27%
2	Linieritas	Linier

I. Hasil Penilaian Peer Review:


Komponen Yang Dinilai	Nilai Maksimal Jurnal Ilmiah =				Nilai Akhir Yang Diperoleh
	Internasional Bereputasi(terindeks database internasional dan berfaktor dampak) Maks=40	Internasional terindeks pada basis data internasional bereputasi Maks=30	Internasional terindeks pada basis data internasional Maks=20	Nasional Terakreditasi peringkat 3&4	
Kelengkapan dan Kesesuaian unsur isi jurnal (10 %)	4				4
Ruang Lingkup dan kedalaman pembahasan (30 %)	12				12
Kecukupan dan Kemutahiran data/Informasi dan metodologi (30 %)	12				12
Kelengkapan unsur dan Kualitas penerbit (30 %)	12				12
Total = (100 %)	40				40
Kontribusi Pengusul (Penulis Pertama/Anggota Utama)	Anggota	$0,133 \times 40 = 5,33$			

**KOMENTAR/ULASAN PEER REVIEW**

- Kelengkapan dan Kesesuaian Unsur	Unsur-unsur jurnal lengkap dan sesuai
- Ruang Lingkup dan Kedalaman Pembahasan	Pembahasan
- Kecukupan dan Kemutakhiran Data &Metodologi	Data dan metodologi
- Kelengkapan Unsur & Kualitas Penerbit	

Inderalaya,  
Penilai

Juni 2020



Nama : Dr. Harry Widjajanti, M.Si.  
NIP : 196112121987102001  
Unit Kerja : Fakultas MIPA  
Instansi : Universitas Sriwijaya  
Bidang Ilmu : Biologi  
Jabatan/Pangkat : Lektor Kepala / Pembina (IV/a)

**LEMBAR  
HASIL PENILAIAN SEJAWAT SEBIDANG (PEER REVIEW)  
KARYA ILMIAH: JURNAL ILMIAH**

Judul Artikel Ilmiah : Photographic evidence of freshwater whipray Urogymnus dalyensis (Myliobatiformes: Dasyatidae) in Indonesian waters

Penulis Artikel Ilmiah : Yuanita Windusari, Laila Hanum, Doni Setiawan, dan Muhammad Iqbal

Identitas Jurnal Ilmiah :

- a. Nama Jurnal : Ecologica Montenegrina
- b. ISSN/ISBN : 2336-9744
- c. Nomor/Volume/Hal : 1 / 22 / 166 - 170
- d. Edisi (Bulan/Tahun) : 2019
- e. Penerbit : Center for Biodiversity of Montenegro
- f. Jumlah Halaman : 5
- g. Jurnal terindeks di : Scimagojr (Scopus) dan Clarivate Analytics

Kategori Publikasi Jurnal Ilmiah :  Jurnal Internasional Bereputasi(terindeks database internasional dan berfaktor dampak)  
(Beri √ pada kategori yang tepat)

Jurnal Internasional terindeks pada basis data internasional bereputasi  
 Jurnal Internasional terindeks pada basis data internasional  
 Jurnal Nasional Terakreditasi peringkat 1 dan 2  
 Jurnal Nasional Terakreditasi peringkat 3 dan 4

Hasil Penilaian Validasi:

No.	ASPEK	URAIAN/KOMENTAR PENILAIAN
1	Indikasi Plagiasi	27 %
2	Linieritas	Linier

I. Hasil Penilaian Peer Review:

Komponen Yang Dinilai	Nilai Maksimal Jurnal Ilmiah =				Nilai Akhir Yang Diperoleh
	Internasional Bereputasi(terindeks database internasional dan berfaktor dampak) Maks=40	Internasional terindeks pada basis data internasional bereputasi Maks=30	Internasional terindeks pada basis data internasional Maks=20	Nasional Terakreditasi peringkat 3&4	
Kelengkapan dan Kesesuaian unsur isi jurnal (10 %)	4				4
Ruang Lingkup dan kedalaman pembahasan (30 %)	12				12
Kecukupan dan Kemutahiran data/Informasi dan metodologi (30 %)	12				12
Kelengkapan unsur dan Kualitas penerbit (30 %)	12				12
Total = (100 %)	40				40
Kontribusi Pengusul (Penulis Pertama/Anggota Utama)	Anggota	$10,133 \times 40 = 5,33$			

**KOMENTAR/ULASAN PEER REVIEW**

- Kelengkapan dan Kesesuaian Unsur	Unsur - unsur jurnal lengkap dan sesuai
- Ruang Lingkup dan Kedalaman Pembahasan	Pembahasan
- Kecukupan dan Kemutakhiran Data &Metodologi	Data dan metodeologi
- Kelengkapan Unsur & Kualitas Penerbit	

Inderalaya, Juni 2020  
Penilai



Nama : Dr. Salni, M.Si.  
NIP : 196608231993031002  
Unit Kerja : Fakultas MIPA  
Instansi : Universitas Sriwijaya  
Bidang Ilmu : Biologi  
Jabatan/Pangkat : Lektor Kepala / Pembina (IV/a)