

# B9-Turnitin-Insecticidal Activity and Phytochemical Profiles of *Avicennia marina*

*by Universitas Sriwijaya Unsri*

---

**Submission date:** 14-Jun-2023 09:13AM (UTC+0700)

**Submission ID:** 2115624180

**File name:** B9-Insecticidal\_Activity\_and\_Phytochemical\_Profiles-2023.pdf (430.33K)

**Word count:** 7581

**Character count:** 40036

## Insecticidal Activity and Phytochemical Profiles of *Avicennia marina* and *Excoecaria agallocha* Leaves Extracts

Rozirwan<sup>1\*</sup>, Muhtadi<sup>2</sup>, Tengku Zia Ulqodry<sup>1</sup>, Redho Yoga Nugroho<sup>2</sup>, Nadila Nur Khotimah<sup>2</sup>, Fauziyah<sup>1</sup>, Wike Ayu Eka Putri<sup>1</sup>, Riris Aryawati<sup>1</sup>, Che Abd Rahim Mohamed<sup>3</sup>

<sup>1</sup>Department of Marine Science, Faculty of Mathematics and Natural Sciences, Sriwijaya University  
Indralaya 30862, South Sumatra, Indonesia

<sup>2</sup>Environmental Management Study Program, Graduate Program, Sriwijaya University  
Palembang 30139, Indonesia

<sup>3</sup>Faculty of Science and Technology, Universiti Kebangsaan Malaysia  
43600 UKM Bangi, Selangor, Malaysia  
Email: rozirwan@unsri.ac.id

### Abstract

Toxic bioactive compounds can be obtained from various mangrove plants that have the potential to be developed in agriculture as bioinsecticides. The coast of South Sumatra has relatively abundant *Avicennia marina* and *Excoecaria agallocha* vegetation. This study was conducted to analyze the bioactivity of insecticides and the phytochemical profiles of mangrove leaves *A. marina* and *E. agallocha* was extracted with methanol, ethyl acetate, and n-hexane from the most toxic extract fraction. Samples were obtained from the Barong River in Sembilang National Park, Banyuasin Regency. The insecticidal activity of the samples was carried out on *Gryllus bimaculatus* and *Tenebrio molitor*. Furthermore, the phytochemical profile was analyzed using GC-MS. Based on the results of insecticidal activity on *G. bimaculatus*, the respective LC<sub>50</sub> values of *A. marina* and *E. agallocha* extracts for the hexane fraction were 12,562 mg.L<sup>-1</sup> and 15,464 mg.L<sup>-1</sup>, ethyl acetate 9,986 mg.L<sup>-1</sup> and 10,292 mg.L<sup>-1</sup>, methanol 6,454 mg.L<sup>-1</sup> and 6,969 mg.L<sup>-1</sup>. Whereas in *T. molitor*, the LC<sub>50</sub> values for the hexane fraction were 10,682 mg.L<sup>-1</sup> and 11,070 mg.L<sup>-1</sup>, ethyl acetate 9,065 mg.L<sup>-1</sup> and 9,269 mg.L<sup>-1</sup>, methanol 4,799 mg.L<sup>-1</sup>, and 5,408 mg.L<sup>-1</sup>. The results of GC-MS analysis on the methanol extract of *A. marina* leaves which contained phytochemical compounds such as alcohol, lauric acid, myristic, linoleic, elaidate, stearate, endogenous, olead, phthalic ester, and siloxane. Based on the insecticide toxicity category, the insecticidal activity of both *A. marina* and *E. agallocha* leaves extracts were low and non-toxic. Further research is needed regarding variations in anti-insecticide of mangrove extract measurements in the future studies.

**Keywords:** Insecticidal, Mangrove, Phytochemical, Toxicity

### Introduction

One of the main ways to eliminate food and environmental poverty in the world is to make agriculture more environmentally friendly to create a balance between the production of materials for population growth and food security (Zegeye et al., 2022). Problems are the main obstacle to achieving higher crop production (Souto et al., 2021). So far, the control of agricultural pests is still very dependent on synthetic insecticides for land protection and post-harvest crops (Bertomeu-Sánchez, 2019). The use of synthetic insecticides can cause failure in pest control due to unwanted side effects on plants and the rapid evolution of insecticide resistance in agroecosystems (Alfaro-Tapia et al., 2021). It not only impacts plants and the environment, but also humans who are in direct contact with synthetic insecticides that can cause cancer, hypertension, kidney failure, lung damage, and neurological

disorders (Krishnamurthy et al., 2020). Therefore, discoveries related to new natural pesticides are urgently needed to prevent damage to crops and not to the environment (Rahman et al., 2021).

The knowledge of scientists about the actual harmful effects of the massive use of synthetic insecticides has increased and is changing the direction of research to find new solutions. The solution that can be given is to look for alternatives to synthetic insecticides that are more natural, environmentally friendly, effective, high efficiency, and have low levels of environmental degradation (Zhu et al., 2022). Alternatives to synthetic insecticides are vegetable insecticides that can be obtained from natural materials such as plants, microorganisms, animals, and certain minerals as natural substances that can control pests with non-toxic and low toxicity mechanisms (De Bernardi et al., 2022). More than 2000 plant species have potential insecticidal activity (Souza et al., 2017).

\* ) Corresponding author  
© Ilmu Kelautan, UNDIP

<https://ejournal.undip.ac.id/index.php/ijms>  
DOI: 10.14710/ik.ijms.28.2.148-160

Received : 22-12-2022  
Accepted : 24-05-2023

Mangrove is one of the plants that have been widely used in the health sector because it has the potential various biological activity from the bioactive compounds it produces (Ulmursida *et al.*, 2017; Sopalun *et al.*, 2021). These perfectly vegetated plants grow in intertidal or brackish water and greatly adapt to environmental changes (Abdul-Azeez *et al.*, 2022). These natural processes will encourage plants to produce natural compounds in response to environmental changes. These compounds are called secondary metabolites, which are unique and structurally diverse and attract significant attention in organic chemistry and pharmacology (Chen *et al.*, 2022). This bioactive compound in mangrove plants has a toxic effect. Toxicity is a reaction caused by natural components extracts that can cause biological damage, including death in the test organism (Rozirwan *et al.*, 2022). Toxicity is most often manifested in certain parts of consumption because the toxic constituents in plants generally come from organic compounds and are rarely found in minerals (Elouardi *et al.*, 2022). Several researchers have considered the toxic effects of mangroves as potential future drug development, such as raw materials for anti-inflammatory, anticancer, antioxidant, and antimicrobial (Parthiban *et al.*, 2021).

The application of secondary metabolites in mangroves to overcome problems in agriculture is an exciting study. One of the existing problems is fighting agricultural pests. Plants with potential for bioactivity usually contain bioactive compounds such as essential oils, fatty acids, alkaloids, glycosides, flavonoids, and esters (Sakul, 2017). These compounds are known to have chemical properties that can affect insects through antifeedant mechanisms, contact toxicity, and reduction of fecundity and fertility (Tak *et al.*, 2017). Mangroves of *A. marina* and *E. agallocha* are known to have toxic activity and potential as insecticides. Several previous studies revealed that testing of *A. marina* leaf extract against mosquito larvae revealed a *Aedes aegypti* of 0.164 mg.L<sup>-1</sup> and *Culex quinquefasciatus* LC<sub>50</sub> of 0.197 mg.L<sup>-1</sup> (Karthi *et al.*, 2020). Furthermore, tests of *E. agallocha* Pradeepa *et al.* (2015) stating that the toxic methanol and *E. agallocha* leaf extract were tested on *A. aegypti* and *C. quinquefasciatus* larvae, which resulted in 100% mortality. A methanol extract of *E. agallocha* leaves with a concentration of 0.05% was tested against the larval pest *Crocidolomia pavonana*, causing a mortality of 86.25%. Based on these studies, the leaf extract of *A. marina* and *E. agallocha* strongly affect mosquito larvae, and caterpillar larvae (Melo *et al.*, 2015). The toxicity of mangrove leaves of *A. marina* and *E. agallocha* leaves, as evidenced against these test organisms can be predicted to be that the extract is also toxic to test insects, which is greater than

larvae. So far, there have been no studies using insects as test animals of Crickets (*Gryllus bimaculatus*) and caterpillars (*Tenebrio molitor*) (Irwan *et al.*, 2021). Sembilang National Park which is located on the coast of the Banyuasin regency South Sumatera, is an area that has various types of mangroves (Rozirwan *et al.*, 2022a; Rozirwan *et al.*, 2022c), including the two most dominating *A. marina* and *E. agallocha*. Research on the insecticide toxicity of mangrove leaves extracts of *A. marina* and *E. agallocha* has never been carried out. This research is an interesting topic to be studied to prove its activity as an insecticide. In the future, its potential can be developed as an alternative source of insecticides from natural materials to replace synthetic insecticides.

## Materials and Methods

Samples of *A. marina* and *E. agallocha* leaves were taken in September 2021 at the Barong River, Sembilang National Park, Banyuasin Regency, South Sumatra (Figure 1.). This sampling location had coordinated on 2.160000 S° and 104.896389 E°. the location had also much biodiversity, such as mangrove vegetation, macrozoobenthos community, and migratory birds (Rozirwan *et al.*, 2022a; Fitriah *et al.*, 2023). Mangrove guidebooks referred to (Giesen *et al.*, 2007). *E. agallocha* and *A. marina* samples were plucked from the fresh leaves using the purposive sampling method (Rahman *et al.*, 2020).

### Environmental quality measurement

Environmental parameters were measured at the sampling location. These parameters include temperature, salinity, dissolved oxygen, and acidity (Ramses *et al.*, 2020; Almaniar *et al.*, 2021). All environmental parameters were measured using portable tools.

### Plant maceration and extraction

Stratified maceration had been used in this study with solvents polarity: methanol (polar), ethyl acetate (semi-polar), and hexane (non-polar) (Pontes *et al.*, 2020). Mangrove leaves were placed in a glass jar and macerated with the three solvents in a ratio of 1:4 (w/v) the first maceration with hexane for 2 x 24 h and the next were ethyl acetate and methanol. The filtrate was filtered and the solvent was replaced using a new hexane solvent every 24 h (Rozirwan *et al.*, 2022b). The final filtrate of each solvent obtained was evaporated using a rotary evaporator at a temperature of 40°C (Rozirwan *et al.*, 2023).

### Insecticide activity test

The test insects used to refer to (Irwan *et al.*, 2021), using crickets (*G. bimaculatus*) and

caterpillars (*T. molitor*); each test treatment was ten tails. Furthermore, the concentration of the insecticide test solution modified the research (Khan *et al.*, 2017). Insecticide activity test used stock solution 10,000 mg.L<sup>-1</sup>. This solution was prepared from 1 g of each fraction and dissolves it in 1 mL of dimethyl sulfoxide, then adding aquades up to 99 mL. The test concentration was 10,000, 8,000, 6,000, 4,000, 2,000 and 1,000 mg.L<sup>-1</sup>. The jar containing insects was sprayed with 5 mL extract solution (Table 1.).

**Gas chromatography-mass spectroscopy analysis**

The components of the phytochemical profile contained in the leaf extract of *A. marina* could be analyzed using GC-MS. The data bank in the Wiley 7 library had been used as a comparison data for the results of the analysis graph spectrum (Roziwan *et al.*, 2022b)

**Percentage of yield value and extract weight**

The percentage of yield value and extract samples using the formula (1), and formula (2) used for the calculation of the percentage of weight.

$$\% \text{ Yield value} = \frac{\text{Final weight} - \text{Initial weight}}{\text{Initial weight}} \times 100 \quad (1)$$

$$\% \text{ Extract weight} = \frac{\text{Final weight}}{\text{Initial weight}} \times 100 \quad (2)$$

**Percentage of mortality**

The mortality categories according to Roziwan *et al.* (2022b) were non-toxic (50%), moderately toxic (50%-75%), and highly toxic (75%-100%). The percentage of deaths was analyzed based on the number of test insects that died after 24 h using the formula (3).

$$\% \text{ Mortality} = \frac{\text{Number of dead individuals}}{\text{number of test individuals}} \times 10 \quad (3)$$

**Percentage of LC50**

LC<sub>50</sub> values were obtained from data analysis of test concentrations, total individual trials, and mortality. There was to know before determining of LC<sub>50</sub> value was the log 10 concentration test and mortality probit value. The standard linear regression equation was used to analyze LC<sub>50</sub> values with Microsoft Excel 2019 (4). Based on Roziwan *et al.* (2022b), LC<sub>50</sub> value categories included non-toxic (> 1000), weak toxic (500–1000), moderate toxic (100–500), and very toxic (< 100).

$$y = a + bx \quad (4)$$




Figure 1. Map of sampling locations for *A. marina* and *E. agallocha*



## Results and Discussion

### Description of mangrove *A. marina* and *E. agallocha* leaves

Leaves morphological observations were done to determine the characteristics between *A. marina* and *E. agallocha* leaves (Figure 2.). Barong Kecil River is one of the areas included in the Sembilang National Park and is located in wetlands, and is dominated by mangrove forests (Ratmoko *et al.*, 2021). In this region, mangrove plants of various genera, such as *E. agallocha* and *A. marina*. Mangrove habitat is very decisive for growth and development such a muddy substrate conditions (Saputra *et al.*, 2021) and it was also reported that sedimentary mangroves also has antibacterial potential (Lalitha *et al.*, 2021). A selection of old leaves is more appropriate to be sampled than young leaves. According to Sumartini *et al.* (2022), the type and level of leaf maturity had an influence on phytochemical compounds, such as the old leaves of *Rhizophora* sp. contain more phenolic, and flavonoid compounds that function as antioxidants.

### Environmental characteristics


The mangroves *A. marina* and *E. agallocha*, discovered at the sampling site, coexist. Both were

associated with other mangrove species and grow on muddy substrates. Ecologically, mangrove growth was influenced by several parameters of the aquatic environment, such as temperature, salinity, pH, and DO water (Table 2.).

Mangrove growth in an area is strongly influenced by environmental conditions. Based on (Table 2.) the parameters of the water quality of the sampling location of this study were still within normal limits, good mangrove growth is in the water salinity does not exceed 34 psu, DO >5 mg.L<sup>-1</sup>, temperature range of 28-32°C, and pH 7-8.5. The temperature obtained is optimal for mangrove growth because mangroves were intolerant of temperatures below 0 °C and high temperatures, which could cause delays in the tree deposition process (Noor *et al.*, 2015). Salinity is also an important component in the mangrove growth and development system because if there is excessive salt content, it would certainly inhibit the growth and development of mangroves and could even cause death of mangroves (Sudhir *et al.*, 2022). In addition, the pH value obtained was classified as neutral and suitable for the sustainability of mangrove life (Wahyuni *et al.*, 2018). The dissolved oxygen parameter indicates the value that low at low tide and rising on the high tide. Almaniar *et al.* (2021) the quality of environmental parameters in mangrove forests was influenced by nutrient and anthropogenic sources.

**Table 1.** Preparation of test solution

Concentration (mg.L <sup>-1</sup> )	Extract solution (mL)	Saline water (mL)	Final volume (mL)
10,000	100	0	100
8,000	80	20	100
6,000	75	25	100
4,000	67	33	100
2,000	50	50	100
1,000	50	50	100
0 as control (-)	-	50	100



**Figure 2.** Morphology of mangrove leaves, (1A) *A. marina*, (2B) *E. agallocha*.

**Extract characteristics of *A. marina* and *E. agallocha* mangrove leaves**

The weight loss percentage of the wet and dry sample leaves was 67.9% for the *A. marina* sample and 72.4% for the *E. agallocha* sample (Table 3.). These results indicated that the moisture content of the leaves of *A. marina* was lower than *E. agallocha*.

The removal of the water content in the sample was carried out by drying until the water content was completely lost because the range of compounds contained in the sample would be better if the sample is dry. Meanwhile, the extraction process of *A. marina* and *E. agallocha* leaf samples was carried out using different solvents (Table 4.). The selection of this stratified maceration aims to obtain the best extract results with high toxicity. Maceration with different samples and solvents would certainly affect the value of the weight percentage of the extract obtained. According to Gori *et al.* (2021), graded maceration is very effective and accurate for used in exploration activities for bioactivity compounds.

Different solvents were used in the leaves extraction processes of *A. marina* and *E. agallocha*. The results showed that the weight of the extract produced by the samples of *A. marina* and *E.*

*agallocha* leaves was the highest in the methanol extract at 7.08 g and 5.21 g, respectively. Each type of mangrove extracted using graded solvent had a different yield value (%). This yield value indicated the percentage of material remaining from the extraction and determined the effectiveness of the solvent in attracting and separating bioactive compounds. Based on (Table 6.), which showed the increasingly polar properties of solvents, the higher the percentage yield value. This statement was following the results of the extraction where the methanol extract in polar solvents in *A. marina* was 2.83%, and in *E. agallocha* it was 2.08%. Generally, secondary metabolites of *A. marina* were polar, so the methanol extract percentage was quite large. Based on these properties, the active compound is dissolved in a solvent that has the same polarity (Soedirga *et al.*, 2020).

**Determination of mortality**

The percentage values of mortality testing of *A. marina* and *E. agallocha* leaves extracts against *G. bimaculatus* (Table 5.) for the methanol, ethyl acetate, and n-hexane fractions, respectively, were at a concentration of 10,000 mg.L<sup>-1</sup>, 8,000 mg.L<sup>-1</sup>, 6,000 mg.L<sup>-1</sup>, 4,000 mg.L<sup>-1</sup>, 2,000 mg.L<sup>-1</sup>, 1000 mg.L<sup>-1</sup>, and negatif control (0 mg.L<sup>-1</sup>).

**Table 2.** Environmental quality parameters

Environmental quality	Score	Measuring Instrument
Salinity (psu)	20.30	Handrefractometer
Dissolved Oxygen (mg.L <sup>-1</sup> )	9.63	DO/Temp Meter Gondo
Temperature (°C)	33.00	DO/Temp Meter Gondo
Acidity (pH)	7.50	pH Meter HI 83141

**Table 3.** Depreciation percentage of weight

Sample leaves	Sample weight (g)		Depreciation percentage (%)	Weight percentage (%)
	Wet	Dry		
<i>A. marina</i>	1,000	321	67.9	32.1
<i>E. agallocha</i>	1,000	276	72.4	27.6

**Table 4.** Percentage of extract

Sample leaves	Solution	Extract weight (g)		Depreciation percentage (%)	Weight percentage (%)
		Dry powder	Crude extract		
<i>A. marina</i>	Methanol	250	7.08	97.2	2.83
	Ethyl acetate	250	5.23	97.9	2.09
	N-hexane	250	2.63	98.9	1.05
<i>E. agallocha</i>	Methanol	250	5.21	97.9	2.08
	Ethyl acetate	250	4.56	98.2	1.82
	N-hexane	250	3.87	98.5	1.55

*A. marina* and *E. agallocha* leaves were extracted using graded solvents of methanol, ethyl acetate, and hexane, which were then tested on *G. bimaculatus* with 6 test concentrations showing different mortality for each extract fraction. The methanol fraction showed a high mortality value compared to the ethyl acetate and hexane fractions in both *A. marina* and *E. agallocha*. Subsequent mortality tests were also carried out on *T. molitor* test insects for leaves extracts of *A. marina* and *E. agallocha* (Table 6).

Mortality in *T. molitor* was carried out as comparison data on whether there was a significant difference or directly proportional to the mortality of *G. bimaculatus*. Based on the results of the *T. molitor* mortality test for 24 hours of treatment with the same extract and test concentration, the percentage results were directly proportional to the mortality of *G. bimaculatus*. The methanol extract fraction showed the highest mortality compared to the fractions of ethyl acetate and n-hexane. The percentage mortality in *G. bimaculatus* methanol extract with a concentration of 10,000 mg.L<sup>-1</sup> showed the highest results in both *A. marina* and *E. agallocha* (Table .5). Likewise with *T. molitor* which showed the percentage results were directly proportional to the mortality of

*G. bimaculatus* (Table 6). According to (Permatasari et al., 2021), the greater the concentration of extracts, the higher the mortality rate and this illustrates that each different concentration affects the percentage of deaths produced. Based on the category of percentage mortality, this study was included in the moderate toxic. Then it was clarified by (Permatasari and Asri, 2020), the standard of botanical pesticides for biological control was stated to be effective toxic if the mortality value was 80%, if the mortality value was 50%, the extract was not effectively used as a vegetable insecticide.

#### Determination of LC<sub>50</sub>

The LC<sub>50</sub> value is basically obtained from linear regression analysis. The LC<sub>50</sub> value test was carried out to determine the concentration value that was able to cause the death of *G. bimaculatus* insects by 50 % of the total tested insects (Table 7.).

Based on the test results in this study, in addition to testing the *G. bimaculatus*, extracts were also tested on *T. molitor* (Table 8.), it was carried out as comparison data to determine whether there was a difference in the mortality and the LC<sub>50</sub> obtained.

**Table 5.** Percentage mortality of *G. bimaculatus*

Test insects	Concentration (mg.L <sup>-1</sup> )	<i>A. marina</i> (%)			<i>E. agallocha</i> (%)		
		Methanol	Ethyl acetate	N-hexane	Methanol	Ethyl acetate	n-hexane
<i>G. bimaculatus</i>	10,000	53±1.15	30±1.73	17±1.15	47±0.58	27±1.15	10±1.73
	8,000	47±1.53	23±1.53	13±0.58	40±1.00	20±0.00	7±0.58
	6,000	40±1.00	13±1.53	7±0.58	33±0.58	13±1.15	3±0.58
	4,000	23±0.58	7±0.58	3±0.58	20±1.00	7±1.15	3±0.58
	2,000	10±1.00	3±0.58	0±0.00	10±1.00	3±0.58	0±0.00
	1,000	0±0.00	0±0.00	0±0.00	0±0.00	0±0.00	0±0.00
	0	0±0.00	0±0.00	0±0.00	0±0.00	0±0.00	0±0.00

**Table 6.** Mortality percentage of *T. molitor*

Test insects	Concentration (mg.L <sup>-1</sup> )	<i>A. marina</i> (%)			<i>E. agallocha</i> (%)		
		Methanol	Ethyl acetate	N-hexane	Methanol	Ethyl acetate	n-hexane
<i>T. molitor</i>	10,000	77±1.53	30±1.73	30±1.73	73±1.53	30±1.00	27±1.53
	8,000	70±1.73	23±1.15	20±1.00	67±1.15	23±1.53	17±1.53
	6,000	60±1.73	17±1.15	10±0.00	53±1.53	17±1.53	10±1.00
	4,000	37±0.58	13±0.58	7±0.58	33±0.58	10±1.00	7±1.15
	2,000	27±1.15	7±0.58	3±0.58	23±1.53	3±0.58	3±0.58
	1,000	23±0.58	0±0.00	0±0.00	20±1.73	0±0.00	0±0.00
	0	0±0.00	0±0.00	0±0.00	0±0.00	0±0.00	0±0.00

**Table 7.** Classification and determination of LC<sub>50</sub> (*G. bimaculatus*)

Sample leaves	Solution	Linear regression			LC <sub>50</sub> (mg.L <sup>-1</sup> )	Category
		a	B	R <sup>2</sup>		
<i>A. marina</i>	Methanol	-14,1142	5,0170	0,8238	6,454	Moderate
	Ethyl acetate	-11,9575	4,2400	0,8331	9,986	Moderate
	n-hexane	-15,4042	4,9778	0,8888	12,562	Non-toxic
<i>E. agallocha</i>	Methanol	-13,3975	4,7870	0,8011	6,969	Moderate
	Ethyl acetate	-11,7252	4,1682	0,8231	10,292	Non-toxic
	n-hexane	-13,9887	4,5326	0,8637	15,464	Non-toxic

**Table 8.** Classification and determination of LC<sub>50</sub> (*T. molitor*)

Sample leaves	Solution	Linear regression			LC <sub>50</sub> (mg.L <sup>-1</sup> )	Category
		a	B	R <sup>2</sup>		
<i>A. marina</i>	Methanol	0,1611	1,3144	0,8677	4,799	Moderate
	Ethyl acetate	-11,7095	4,2223	0,7606	9,065	Moderate
	n-hexane	-11,4992	4,0954	0,8134	10,682	Non-toxic
<i>E. agallocha</i>	Methanol	-0,1807	1,3878	0,8665	5,408	Moderate
	Ethyl acetate	-11,6561	4,1986	0,7643	9,269	Moderate
	n-hexane	-11,2438	4,0165	0,8016	11,070	Non-toxic

Based on the tests in this study, referring to the LC<sub>50</sub> toxicity category on (Rozirwan *et al.*, 2022b) the leaves of *A. marina* and *E. agallocha* against *G. bimaculatus*, the smallest value was in the methanol fraction (Table 7). Meanwhile, for the insect *T. molitor* (Table 8) the smallest LC<sub>50</sub> value was also in the methanol fraction. These results showed that the higher the LC<sub>50</sub> value indicated the ability of the extract to be non-toxic. Bioactive compounds are toxic if they can cause a deadly effect that can cause the death of more than 50 % in a short time. *G. bimaculatus* and *T. molitor* belong to the major insect groups and are more substantial than test organisms that are often used in toxicity research.

#### GC-MS analysis of methanol extract *A. marina*

GC-MS analysis was taken based on the higher mortality fraction and LC<sub>50</sub> values. Based on the test results, the extract with a substantial value is in the methanol fraction of *A. marina* leaves. The graph obtained as many as 14 peak points. The detected compounds were alcohols and organic fatty acids. The secondary metabolites contained in the extract affected the insecticide toxicity in the test (Figure 3.). The compounds identified based on the height of the chromatogram peaks are presented in full in (Table 9). The mass spectrum on the chromatogram graph of the results of the analysis that has been matched is identical to the mass spectrum in the base data library: WILEY.

GC-MS chromatogram analysis of the methanol fraction of *A. marina* leaves clearly showed the presence of 14 secondary metabolites based on the data bank library: WILEY 7. Based on the peaks detected that the main groups of compounds included alcohol, lauric acid, myristic, palmitic, linoleic, elaidate groups, stearate, oleic, phthalate ester, and siloxane. The results of the highest level of toxicity were found in the methanol extract of *A. marina* with the lowest LC<sub>50</sub> value indicating that there were bioactive compounds that affected the results. Based on the results of GC-MS analysis (Table 9), most of the methanol extracts of *A. marina* leaves contain fatty acid compounds such as Dodecanoic acid, methyl ester, Dodecanoic acid, Tridecanoic acid, 12-methyl-, methyl ester, Hexadecanoic acid, methyl ester, 7,10-Octadecadienoic acid, methyl ester, 13-Octadecenoic acid, methyl ester, Heptadecanoic acid, 16-methyl-, methyl ester, 9-Octadecenamide, (Z)-, 1,4 Benzenedicarboxylic acid, bis (2-ethylhexyl) ester, Octasilaxane, 1,1,3,5,5,7,7,9,9,11,11,13,13,15,15-hexadecame, and Phthalic acid, di(2 propylpentyl) ester.

Fatty acids can be found in green plants and were produced in chloroplasts. In addition to the leaves of *A. marina*, Jaritkhuan and Suanjit (2018) also revealed that the leaves of the mangroves of *Aurantiochytrium mangrovei* and *Aurantiochytrium Limacinum* contains fatty acids. Several studies had revealed that fatty acids reported as antifungal,

















antibacterial, antioxidant, and anti-inflammatory activities (Sohaib *et al.*, 2022). Furthermore on Hexadecanoic acid, methyl ester (palmitic acid) has antibacterial activity (Karunia *et al.*, 2017). 9-Octadecenamamide, (Z)- is oleic acid which is lethal to *A. aegypti* mosquito larvae, these compounds include oleoamides which can cause motor dysfunction and are commonly used as sedatives. 1,4 Benzenedicarboxylic acid, bis(2-ethylhexyl) ester is a flatalic acid ester reported as antifungal activity. Phthalic acid, di(2 propylpentyl) ester compound has antimicrobial and anti-inflammatory activity (Chakraborty *et al.*, 2022). Hexadecanamide has anti-inflammatory, antifungal and anti-nociceptive activity (Rahbar *et al.*, 2012). Dodecanoic acid, methyl ester and dodecanoic acid are included in lauric acid. Lauric acid is a saturated fatty acid that can be used as an antimicrobial activity (Li *et al.*, 2022). Octasiloxane,1,1,3,3,5,5,7,7,9,9,11,11,13,13,15,15-hexadecamethyl is a compound that reported as insecticidal activity (Farag *et al.*, 2021). As for the

organic acids compounds in Heptadecanoic acid, 16-methyl-, methyl ester of stearic acid, Tridecanoic acid, 12-methyl-, methyl ester of myristic acid, 7,10-Octadecadienoic acid, methyl ester linoleic acid and Trans-13-Octadecenoic acid, methyl ester of elaidic acid reported as antibacterial (Ravichandran *et al.*, 2022). Stearic, oleic, myristic, linoleic and elaidic acids reported as antibacterial (Sani *et al.*, 2022).

Some of the statements above reveal that phytochemical profile of *A. marina* leaves extract has a potential bioactivity. The results of the percentage mortality and LC<sub>50</sub> test showed that the methanol extract of *A. marina* leaves was a fraction that had a higher mortality and toxicity level even though it was in the very low toxic category compared to the fractions of ethyl acetate and n-hexane. The extract methanol of *A. marina* leaf produces the most dominant bioactive compound is organic fatty acids and shows low and non-toxic activity against test insects.

**Table 9.** Retention time, peak area, compound name, formula, and structure compound

Ret. time	Peak Area%	Name of Compounds	Formula	Structure Compounds
10.31	3,83	2-(2-Butoxyethoxy)ethanol	C <sub>8</sub> H <sub>18</sub> O <sub>3</sub>	
13.36	2,43	Dodecanoic acid, methyl ester	C <sub>13</sub> H <sub>26</sub> O <sub>2</sub>	
13.65	2,24	Dodecanoic acid	C <sub>12</sub> H <sub>24</sub> O <sub>2</sub>	
14.66	2,82	Tertbutyloxyformamide, N-methyl- N-[4-(1 pyrrolidinyl)- 2 butynyl]	C <sub>14</sub> H <sub>24</sub> N <sub>2</sub> O <sub>2</sub>	
15.10	3,10	Tridecanoic acid, 12-methyl-, methyl ester	C <sub>15</sub> H <sub>30</sub> O <sub>2</sub>	
17.85	10,16	Hexadecanoic acid, methyl ester	C <sub>17</sub> H <sub>34</sub> O <sub>2</sub>	
20.23	2,33	7,10-Octadecadienoic acid, methyl ester	C <sub>19</sub> H <sub>34</sub> O <sub>2</sub>	
20.31	9,29	Trans-13-Octadecenoic acid, methyl ester	C <sub>19</sub> H <sub>36</sub> O <sub>2</sub>	
20.60	7,16	Heptadecanoic acid, 16-methyl-, methyl ester	C <sub>19</sub> H <sub>38</sub> O <sub>2</sub>	
21.25	4,25	Hexadecanamide	C <sub>16</sub> H <sub>33</sub> NO	
23.30	3,45	9-Octadecenamamide, (Z)-	C <sub>18</sub> H <sub>35</sub> NO	
24.90	9,06	Phthalic acid, di(2 propylpentyl) ester	C <sub>24</sub> H <sub>38</sub> O <sub>4</sub>	
26.49	15,82	1,4-Benzenedicarboxylic acid, bis(2-ethylhexyl) ester	C <sub>24</sub> H <sub>38</sub> O <sub>4</sub>	
28.28	2,51	Octasiloxane,1,1,3,3,5,5,7,7,9,9,11,11,13,13,15,15-hexadecamethyl-	C <sub>16</sub> H <sub>50</sub> O <sub>7</sub> Si <sub>8</sub>	

\*Source 2D structure and Graph image : (Kim *et al.*, 2022).




Figure 3. GC-MS chromatogram of methanol extract *A.marina*.

## Conclusion

The insecticidal potency of *A. marina* and *E. agallocha* leaves extracts was in the methanol extract fraction in the very low toxic category. This is evidenced by the results of the  $LC_{50}$  value of *A. marina* methanol extract testing against *G. bimaculatus* ( $6.454 \text{ mg.L}^{-1}$ ) and *T. molitor* ( $4.799 \text{ mg.L}^{-1}$ ). Then testing the methanol extract of *E. agallocha* against *G. bimaculatus* and *T. molitor* of  $6.969 \text{ mg.L}^{-1}$  and  $5.408 \text{ mg.L}^{-1}$ . The results of the GC-MS analysis of the most potential extract as an insecticide based on the high toxicity value were in the methanol extract of *A. marina* leaves which were suspected to contain alcohol and organic fatty acids.

## Acknowledgement

The part of this research was funded by Ministry of Research, Technology, and Higher Education of the Republic of Indonesia under the "Penelitian Dasar Unggulan Perguruan Tinggi" (Grant Number: 059/E5/PG.02.00.PL/2023).

## References

Abdul Azeez, S., Gnanappazham, L., Muraleedharan, K. R., Revichandran, C., John, S., Seena, G. & Thomas, J. 2022. Multi-decadal changes of

mangrove forest and its response to the tidal dynamics of thane creek, Mumbai. *J. Sea Res.*, 180: 102162. <https://doi.org/10.1016/J.SEARES.2021.102162>

Alfaro-Tapia, A., Alvarez-Baca, J.K., Fuentes-Contreras, E. & Figueroa, C. C. 2021. Biological control may fail on pests applied with high doses of insecticides: Effects of sub-lethal concentrations of a pyrethroid on the host-searching behavior of the aphid parasitoid *Aphidius colemani* (hymenoptera, braconidae) on aphid pests. *Agric.*, 11(6): 1–14. <https://doi.org/10.3390/AGRICULTURE11060539>

Almaniar, S., Rozirwan & Herpandi. 2021. Abundance and diversity of macrobenthos at Tanjung Api-Api waters, South Sumatra, Indonesia. *AACL Bioflux*, 14: 1486–1497.

Bertomeu-Sánchez, J.R. 2019. Introduction. Pesticides: Past and Present. *HoST - J. Hist. Sci. Technol.*, 13(1): 1–27. <https://doi.org/10.2478/HOST-2019-0001>

Bibi, S. N., Gokhan, Z., Rajesh, J. & Mahomoodally, M. F. 2020. Fungal endophytes associated with mangroves – Chemistry and biopharmaceutical potential. *South African J. Bot.*, 134: 187–212. <https://doi.org/https://doi.org/10.1016/j.sajb.2019.12.016>

- Chakraborty, B., Kumar, R.S., Almansour, A.I., Perumal, K., Nayaka, S. & Brindhadevi, K. 2022. *Streptomyces filamentosus* strain KS17 isolated from microbiologically unexplored marine ecosystems exhibited a broad spectrum of antimicrobial activity against human pathogens. *Process Biochem.*, 117: 42–52. <https://doi.org/10.1016/J.PROCBIO.2022.03.010>
- Chen, S., Cai, R., Liu, Z., Cui, H. & She, Z. 2022. Secondary metabolites from mangrove-associated fungi: source, chemistry and bioactivities. *Nat. Prod. Rep.*, 39(3): 560–595. <https://doi.org/10.1039/D1NP00041A>
- Dahibhate, N.L., Saddhe, A.A. & Kumar, K. 2019. Mangrove plants as a source of bioactive compounds: A review. *Nat. Prod. J.*, 9(2): 86–97.
- De Bernardi, A., Marini, E., Casucci, C., Tiano, L., Marcheggiani, F., Ciani, M., Comitini, F., Taskin, E., Puglisi, E. & Vischetti, C. 2022. Ecotoxicological effects of a synthetic and a natural insecticide on earthworms and soil bacterial community. *Environ. Adv.*, 8: 100225. <https://doi.org/10.1016/J.ENVADV.2022.100225>
- Elouardi, M., Zair, T., Mabrouki, J., Fattah, G., Benchrifa, M., Qisse, N. & El Belghiti, M. A. 2022. A review of botanical, biogeographical phytochemical and toxicological aspects of the toxic plants in Morocco. *Toxicol. Anal. Clin.*, 34(4): 215–228. <https://doi.org/10.1016/J.TOXAC.2022.04.003>
- Farag, S.M., Essa, E.E., Alharbi, S. A., Alfarraj, S. & Abu El-Hassan, G.M.M. 2021. Agro-waste derived compounds (flax and black seed peels): Toxicological effect against the West Nile virus vector, *Culex pipiens* L. with special reference to GC-MS analysis. *Saudi J. Biol. Sci.*, 28(9): 5261–5267. <https://doi.org/10.1016/J.SJBS.2021.05.038>
- Fitria, Y., Rozirwan, Fitriani, M., Nugroho, R. Y., Fauziyah & Putri, W.A.E. 2023. Gastropods as bioindicators of heavy metal pollution in the Banyuasin estuary shrimp pond area, South Sumatra, Indonesia. *Acta Ecol. Sin.*, <https://doi.org/https://doi.org/10.1016/j.chn.aes.2023.05.009>
- Giesen, W., Wulffraat, S., Zieren, M. & Scholten, L. 2007. Mangrove guidebook for Southeast Asia. In *Mangrove guidebook for Southeast Asia*. FAO Regional Office for Asia and the Pacific.
- Gori, A., Boucherle, B., Rey, A., Rome, M., Fuzzati, N. & Peuchmaur, M. 2021. Development of an innovative maceration technique to optimize extraction and phase partition of natural products. *Fitoterapia*, 148: 104798. <https://doi.org/10.1016/j.fitote.2020.104798>
- Hazra, D.K., Karmakar, R., Poi, R., Bhattacharya, S. & Mondal, S. 2017. Recent advances in pesticide formulations for eco-friendly and sustainable vegetable pest management: A review. *Arch. Agric. Environ. Sci.*, 2(3): 232–237.
- Irwan, Z., Kamarudin, W.F.W., Korish, U.A.S.A., Z.N.A., Rusli, A.S. & Sallehuddin, S. 2021. Effectiveness of *Annona Squamosa* and *Annona Muricata* Seed Extracts as Ingredients in Bio-pesticides Spray. *IOP Conf. Ser. Mater. Sci. Eng.*, 1176(1): 012007. <https://doi.org/10.1088/1757-899x/1176/1/012007>
- Jarikhuan, S. & Suanjit, S. 2018. Species diversity and polyunsaturated fatty acid content of thraustochytrids from fallen mangrove leaves in Chon Buri province, Thailand. *Agric. Nat. Resour.*, 52(1): 24–32. <https://doi.org/10.1016/J.ANRES.2018.05.002>
- Karthi, S., Vinothkumar, M., Karthic, U., Manigandan, V., Saravanan, R., Vasantha-Srinivasan, P., Kamaraj, C., Shivakumar, M.S., De Mandal, S., Velusamy, A., Krutmuang, P. & Senthil-Nathan, S. 2020. Biological effects of *Avicennia marina* (Forssk.) vierh. extracts on physiological, biochemical, and antimicrobial activities against three challenging mosquito vectors and microbial pathogens. *Environ. Sci. Pollut. Res.*, 27(13): 15174–15187. <https://doi.org/10.1007/S11356-020-08055-1>
- Karunia, S.D., Supartono & Sumarni, W. 2017. Analisis Sifat Anti Bakteri Ekstrak Biji Srikaya (*Annona squamosa* L) dengan Pelarut Organik. *Indones. J. Chem. Sci.*, 6(1): 56–60.
- Khan, K.M., Collier, A.D., Meshalkina, D.A., Kysil, E.V., Khatsko, S.L., Kolesnikova, T., Morzherin, Y.Y., Warnick, J.E., Kalueff, A.V. & Echevarria, D.J. 2017. Zebrafish models in neuropsychopharmacology and CNS drug discovery. *Br. J. Pharmacol.*, 174(13): 1925–1944. <https://doi.org/10.1111/bph.13754>
- Kim, S., Cheng, T., He, S., Thiessen, P.A., Li, Q., Gindulyte, A. & Bolton, E.E. 2022. PubChem Protein, Gene, Pathway, and Taxonomy Data Collections: Bridging Biology and Chemistry through Target-Centric Views of PubChem Data. *J. Mol. Biol.*, 434(11): 167514. <https://doi.org/10.1016/J.JMB.2022.167514>
- Krishnamurthy, S., Veerasamy, M. & Karruppaya, G. 2020. A Review on Plant Sources for Nano

- Biopesticide Production. *Lett. Appl. Nano-BioScience*, 9(3): 1348–1358. <https://doi.org/10.33263/lianbs93.13481358>
- Lalitha, P., Parthiban, A., Sachithanandam, V., Purvaja, R. & Ramesh, R. 2021. Antibacterial and antioxidant potential o
- A., Murugan, K. & Sridhar, R. 2021. An integrative review on bioactive compounds from Indian mangroves for future drug discovery. *South African J. Bot.*, 149: 899–915.
- Permatasari, G.A., Hariani, N. & Trimurti, S. 2020. Armyworm Mortality Test (*Spodoptera litura* F.) Against Mother-in-law's Tongue Plant Extract (*Sansevieria trifasciata* Prain). *J. Bioterdidik*, 8(3): 56–67. <https://doi.org/10.23960/jbt.v8.i3.07>
- Permatasari, S.C. & Asri, M.T. 2021. The Effectiveness of Ethanol Extract of Kirinyuh Leaves (*Eupatorium odoratum*) Against *Spodoptera litura* Larvae Mortality. *LenteraBio Berk. Ilm. Biol.*, 10(1): 17–24. <https://doi.org/10.26740/LENERABIO.V10N1.P17-24>
- Pontes, A.L. da S., Mesquita, V.C., Chaves, F. de O., da Silva, A.J.R., Kaplan, M.A.C. & Fingolo, C.E. 2020. Phthalates in *Avicennia schaueriana*, a mangrove species, in the State Biological Reserve, Guaratiba, RJ, Brazil. *Environ. Adv.*, 2: 100015. <https://doi.org/10.1016/J.ENVADV.2020.100015>
- Pradeepa, P., Subalakshmi, K., Saranya, A., Dinesh, P., Raj, V.A. & Ramanathan, T. 2015. Milky mangrove *Excoecaria agallocha* L. Plant as a source for potential mosquito larvicides. *J. Appl. Pharm. Sci.*, 5(3): 102–105. <https://doi.org/10.7324/JAPS.2015.50317>
- Quiroz, S., Cespedes, C.L., Alderete, J.B. & Alarcon, J. 2015. Ceanothane and oleanane-type triterpenes from *Talguenea quinquenervia* have insecticidal activity against *Cydia pomonella*, *Tenebrio molitor* and *Drosophila melanogaster*. *Ind. Crops Prod.*, 74: 759–766. <https://doi.org/10.1016/J.INDCROP.2015.06.003>
- Rahbar, N., Shafaghat, A. & Salimi, F. 2012. Antimicrobial activity and constituents of the hexane extracts from leaf and stem of *Origanum vulgare* L. ssp. *Viride* (Boiss.) Hayek. growing wild in Northwest Iran. *J. Med. Plants Res.*, 6(13): 2681–2685. <https://doi.org/10.5897/JMPR11.1768>
- Rahman, K.S., Islam, M.N., Ahmed, M.U., Bosma, R. H., Debrot, A.O. & Ahsan, M.N. 2020. Selection of mangrove species for shrimp based silvo-aquaculture in the coastal areas of Bangladesh. *J. Coast. Conserv.* 2020 245, 24(5): 1–13. <https://doi.org/10.1007/S11852-020-00770-8>
- Rahman, M.A., Paul, R.R., Biswas, C., Akter, H., Rouf, R., Nath, S., Shilpi, J.A., Nahar, L., Sarker, S.D. & Uddin, S.J. 2021. Pesticidal Activity of Sundarban Mangrove Plant Extracts against *Sitophilus* Pests and Identification of Active Constituents Using LC-MS. *Adv. Pharmacol. Pharm. Sci.*, 2021:.. <https://doi.org/10.1155/2021/1540336>
- Ramses, I., Amelia, F., Rozirwan & Suheryanto. 2020. Diversity and abundance of polychaetes in the west coast waters of Batam Island, Kepulauan Riau Province-Indonesia. *AACL Bioflux*, 13(1): 381–391.
- Ratmoko, I.A., Budianta, D., Ridho, R., Wildayana, E., Idrus, Z. & Nuryamsasni, N. 2021. Analytic hierarchy process for zonation review of Sembilang National Park, Banyuasin Regency, South Sumatera Province, Indonesia. *IOP Conf. Ser. Earth Environ. Sci.*, 737(1): 12019.
- Ravichandran, A.T., Rathika, R. & Kumaresavanji, M. 2022. Non-linear optical behaviour and antibacterial activity of semi-organic picolinic acid zinc sulphate single crystal. *Mater. Today Proc.*, 68:502-505. <https://doi.org/10.1016/J.MATPR.2022.07.432>
- Rozirwan, F., Nugroho, R.Y., Melki, Ulqodry, T.Z., Agustriani, F., Ningsih, E.N., Putri, W.A.E., Absori, A. & Iqbal, M. 2022a. An ecological assessment of crab's diversity among habitats of migratory birds at berbak-sembilang national park indonesia. *Int. J. Conserv. Sci.*, 13(3): 961–972.
- Rozirwan, N., Nugroho, R.Y., Diansyah, G., Muhtadi, Fauziyah, Putri, W.A.E. & Agussalim, A. 2023. Phytochemical composition, total phenolic content and antioxidant activity of *Anadara granosa* (Linnaeus, 1758) collected from the east coast of South Sumatra, Indonesia. *Baghdad Sci. J.*, <https://doi.org/10.21123/bsj.2023.6941>
- Rozirwan, Nugroho, R.Y., Hendri, M., Fauziyah, Putri, W.A.E. & Agussalim, A. 2022b. Phytochemical profile and toxicity of extracts from the leaf of *Avicennia marina* (Forssk.) Vierh. collected in mangrove areas affected by port activities. *South African J. Bot.*, 150: 903–919. <https://doi.org/10.1016/jsajb.2022.08.037>
- Rozirwan, Nugroho, R. Y., Wulandari, P. I., Aryawati, R., Fauziyah, Putri, W. A. E., Agussalim, A. & Isnaini.



- 2022c. Bacillariophyceae Distribution and Water Quality in Estuarine-Mangrove Environments: The Commonest Phytoplankton in Musi Estuary, Indonesia. *J. Hunan Univ. Nat. Sci.*, 49(12): 78–88.
- Sakul, E.H. (Ernest). 2017. Impact Of Botanical Insecticides Derived From *Pangium edule* Reinw And *Annona muricata* L Seed Extracts On The “Gay Gantung” Diamondback Moth, *Plutella Xylostella* L. *Agrotech J.*, 2(2): 27–35.
- Sani, M.S.A., Bakar, J., Azid, A. & Iqbal, M.J. 2022. Chemometrics-based evaluation on the effect of sonication, contact time and solid-to-solvent ratio on total phenolics and flavonoids, free fatty acids and antibacterial potency of *Carica papaya* seed against *S. enteritidis*, *B. cereus*, *V. vulnificus* and *P. mirabilis*. *Food Chem. Adv.*, 1: p.100033. <https://doi.org/10.1016/J.FOCHA.2022.100033>
- Saputra, A., Nugroho, R.Y., Isnaini, R. & Rozirwan. 2021. A review: The potential of microalgae as a marine food alternative in Banyuasin Estuary, South Sumatra, Indonesia. *Egypt. J. Aquat. Biol. Fish.*, 59: 1053–1065. <https://doi.org/10.21608/EJABF.2021.170654>
- Soedirga, L.C., Hardoko & Prameswari, G. 2020. Determination of solvent and ratio sample-solvent towards the production of oligo-glucosamine obtained from fermented tiger shrimp (*Penaeus monodon*) shell's chitin by using precipitation method. *Food Res.*, 4(6): 2163–2168. [https://doi.org/10.26656/fr.2017.4\(6\).255](https://doi.org/10.26656/fr.2017.4(6).255)
- Sohaib, M., Al-Barakah, F.N.I., Migdadi, H.M. & Husain, F.M. 2022. Comparative study among *Avicennia marina*, *Phragmites australis*, and *Moringa oleifera* based ethanolic-extracts for their antimicrobial, antioxidant, and cytotoxic activities. *Saudi J. Biol. Sci.*, 29(1): 111–122. <https://doi.org/10.1016/J.SJBS.2021.08.062>
- Sopalun, K., Laosripaiboon, W., Wachirachai, A. & Iamtham, S. 2021. Biological potential and chemical composition of bioactive compounds from endophytic fungi associated with thai mangrove plants. *South African J. Bot.*, 141: 66–76. <https://doi.org/10.1016/j.sajb.2021.04.031>
- Souto, A.L., Sylvestre, M., Tölke, E.D., Tavares, J.F., Barbosa-Filho, J.M. & Cebrián-Torrejón, G. 2021. Plant-Derived Pesticides as an Alternative to Pest Management and Sustainable Agricultural Production: Prospects, Applications and Challenges. 26(16): 1–34. <https://doi.org/10.3390/MOLECULES26164835>
- Souza, C.M., Baldin, E.L.L., Ribeiro, L.P., Silva, I.F., Morando, R., Bicalho, K.U., Vendramim, J.D. & Fernandes, J.B. 2017. Lethal and growth inhibitory activities of Neotropical Annonaceae-derived extracts, commercial formulation, and an isolated acetogenin against *Helicoverpa armigera*. *J. Pest Sci. (2004)*, 90(2): 701–709.
- Sudhir, S., Arunprasath, A. & Sankara Vel, V. 2022. A critical review on adaptations, and biological activities of the mangroves. *J. Nat. Pestic. Res.*, 1: 1–11. <https://doi.org/10.1016/J.NAPER.2022.100006>
- Sumartini, P. W., Ratrinia & Hutabarat, R. F. 2022. The effect of mangrove types and leave maturity on the mangrove leaves (*Sonneratia alba*) and (*Rhizophora mucronata*) tea powder - IOPscience. *IOP Conf. Ser. Earth Environ. Sci.*, 1–11. <https://doi.org/10.1088/1755-1315/967/1/012018>
- Tak, J.H., Jovel, E. & Isman, M.B. 2017. Synergistic interactions among the major constituents of lemongrass essential oil against larvae and an ovarian cell line of the cabbage looper, *Trichoplusia ni*. *J. Pest Sci. 2017* 902, 90(2): 735–744. <https://doi.org/10.1007/S10340-016-0827-7>
- Wahyuni Baderan, D.K., Utina, R. & Lapolo, N. 2018. Vegetation structure, species diversity, and mangrove zonation patterns in the Tanjung Panjang Nature Reserve Area, Gorontalo, Indonesia. *Int. J. Appl. Biol.*, 2(2): 1–12. <https://doi.org/10.20956/ijab.v2i2.5752>
- Wahyuni, D. 2015. New Bioinsecticide Granules Toxin from Extract of Papaya (*Carica Papaya*) Seed and Leaf Modified Against *Aedes aegypti* Larvae. *Procedia Environ. Sci.*, 23: 323–328. <https://doi.org/10.1016/J.PROENV.2015.01.047>
- Winthachai, S.; Vaithanomsat, P.; Punsuvon, V. 2015. Solubility Behaviour of Baicalin in Polar and Non-Polar Solvents: Solubility Parameters Approach. *Asian J. Chem.*, 27(9): 3507–3510.
- Zegeye, M.B., Meshesha, G.B. & Shah, M.I. 2022. Measuring the poverty reduction effects of adopting agricultural technologies in rural Ethiopia: findings from an endogenous switching regression approach. 8(5): 1–12. <https://doi.org/10.1016/J.HELIVON.2022.E09495>
- Zhang, W.J., Zhang, Z., Chen, Z.Y., Liang, J.Y., Geng, Z.F., Guo, S.S., Du, S.S. & Deng, Z.W. 2017.

- Chemical Composition of Essential Oils from Six *Zanthoxylum* Species and Their Repellent Activities against Two Stored-Product Insects. *J. Chem.*, 2017: 1–7. <https://doi.org/10.1155/2017/1287362>
- Zhu, X., Chen, S., Zheng, Y., Zhang, Y., Hsiang, T., Huang, R., Qi, J., Gan, T., Chang, Y. & Li, J. 2022. Antifungal and insecticidal activities of rhein derivatives: synthesis, characterization and preliminary structure–activity relationship studies. *Nat. Prod. Res.*, 36(16): 4140–4146. <https://doi.org/10.1080/14786419.2021.1977804>
- Zikankuba, V.L., Mwanyika, G., Ntwenya, J.E. & James, A. 2019. Pesticide regulations and their malpractice implications on food and environment safety. *Cogent Food Agric.*, 5(1): 1601544.

# B9-Turnitin-Insecticidal Activity and Phytochemical Profiles of Avicennia marina

---

## ORIGINALITY REPORT

---

5%

SIMILARITY INDEX

4%

INTERNET SOURCES

5%

PUBLICATIONS

1%

STUDENT PAPERS

---

## MATCH ALL SOURCES (ONLY SELECTED SOURCE PRINTED)

---

2%

★ Ajib Setyo Arifin, Rafifa Maricha Putri Erisa, Muhammad Suryanegara. "Sea and Ship Waves Spectrum Measurement Using Wireless Sensor Network in Java Sea", 2018 Second World Conference on Smart Trends in Systems, Security and Sustainability (WorldS4), 2018

Publication

---

Exclude quotes On

Exclude matches < 1%

Exclude bibliography On