

Antioxidant and Antibacterial Activities of Endophytic Fungi Extracts of *Syzygium zeylanicum*

by Hary Widjajanti

Submission date: 21-Jun-2023 01:58PM (UTC+0700)

Submission ID: 2120183248

File name: terial_Activities_of_Endophytic_Fungi_Extracts_STI_JULI_2022.pdf (221.43K)

Word count: 6755

Character count: 37443

Antioxidant and Antibacterial Activities of Endophytic Fungi Extracts of *Syzygium zeylanicum*

Syarifah^{1,2}, Elfita^{3*}, Hary Widjajanti⁴, Arum Setiawan⁴, Alfia Rahma Kurniawati²

¹Graduate School of Sciences, Faculty of Mathematics and Natural Sciences, Universitas Sriwijaya, Palembang, 30139, Indonesia

²Department of Biology, Faculty of Science and Technology, Universitas Islam Negeri Raden Fatah, Palembang, 30126, Indonesia

³Department of Chemistry, Faculty of Mathematics and Natural Sciences, Universitas Sriwijaya, Palembang, 30862, Indonesia

⁴Department of Biology, Faculty of Mathematics and Natural Sciences, Universitas Sriwijaya, Palembang, 30862, Indonesia

*Corresponding author: elfita69@gmail.com

Abstract

Syzygium zeylanicum, one of the other therapeutic plants found in Indonesia, is used to treat various ailments, such as antimicrobial, anti-inflammatory, antioxidant, arthritis, antidiabetic, mosquitocidal, antitumor, and anti-rheumatic agents. The massive use of plant extract has caused the development of isolation of bioactive compounds thru their endophytic. The present research aimed to obtain endophytic fungal isolates from the stem bark and leaves of Jambu nasi-nasi (*S. zeylanicum*) and analyze endophytic fungal extracts' antioxidant and antibacterial activity. Endophyte identification was performed morphologically, and isolates with high biological activity were molecularly characterized. The antibacterial activity was evaluated by the disc diffusion method, and the antioxidant activity was evaluated by the DPPH method. In total, 10 endophytic fungi were isolated and identified as *Phialemonium* sp. (Code SZT3), *Acremonium* sp. (Code SZT4), *Trichoderma aureoviridae* (Code SZT5), *Trichoderma koningi* (Code SZT7), *Phytium torulosum* (Code SZL1), *Phytium zingiberum* (Code SZL2), *Septonema* sp. (Code SZL3), *Lasiodiplodia pseudotheobromae* (Code SZL4), *Volutella ciliata* (SZL5), and *Trichophyton mentagrophytes* (Code SZL7). Isolate SZL4 gave activity the highest antioxidant (IC₅₀ = 3.30 µg/mL) and strong antibacterial activity against four pathogens bacterial (*S. thypi*, *B. subtilis*, *S. aureus*, and *E. coli*). The potential endophytic fungi based on molecular analysis was *Lasiodiplodia pseudotheobromae* with accession number OK668257. These endophytic fungi can be developed into new sources of antibacterial and antioxidant bioactive compounds through further studies.

Keywords

Antibacterial Activity, Antioxidant Activity, Endophytic Fungi, *Syzygium zeylanicum*

Received: 9 February 2022, Accepted: 5 June 2022

<https://doi.org/10.26554/sti.2022.7.3.303-312>

1. INTRODUCTION

The search for sources of biologically active compounds is constantly being carried out by many emerging diseases, including infectious diseases, cancer, and other dangerous diseases (Segaran and Sathivelu, 2019; Teodoro, 2019; Zhao et al., 2010). Endophytic fungi have great potential in searching for new drug sources as they can produce bioactive compounds that can be processed into drugs (Bhardwaj and Agrawal, 2014; Lu et al., 2012; Omoareloje and Van Staden, 2020; Uzma et al., 2019). Beneficial endophytic fungi are easy to grow, have a short life cycle, and can produce large amounts of bioactive compounds by cultivation methods (Kumar et al., 2019).

Improving resistance through pathogenic microorganisms to business pills is a applicable trouble confronted by health-care offerings and has become an extreme problem worldwide (Mane et al., 2018). Factors contributing to this resistance

include the widespread and often inappropriate use of antibiotics, poor sanitation, the constant movement of tourists, and the growing number of immunocompromised patients and late diagnosis of infections. Fungi are easy to grow and have a short lifespan, and cultivation methods can produce many biologically active compounds (Manyi-Loh et al., 2018).

Accordingly, there is a need for an intensive search for novel antibacterial agents from various natural sources, including endogenous fungi (Kaul et al., 2012). In a recent study, antibacterial activity was measured against *Salmonella thypi* and *Escherichia coli* (gram-negative pathogenic bacteria); *Bacillus subtilis* and *Staphylococcus aureus* (gram-positive pathogenic bacteria).

Bacterial resistance can be severe when the cell body in damaged condition. The integrity of the cell can destroy by radicals in the body, and it can be solved using antioxidant agents (Pandey et al., 2014). Degenerative diseases such as cancer, atherosclerosis, diabetes, rheumatoid arthritis, and decreased

immune response are caused by free radicals (Phaniendra et al., 2015; Sharma et al., 2018). Correlation damaged cell, free radical and bacterial resistance was the reason to be needed to find a new source with antibacterial and antioxidant potential.

Leaves of *S. zeylanicum* plant have been used by people in Indonesia as a medicinal plant related to pathogenic bacterial infections and the effects of free radicals in the body (Anoop and Bindu, 2014; Anoop and Bindu, 2015; Deepika et al., 2014; Vinodkumar, 2015). Endophytes, especially those that inhabit medicinal plant tissues, are often used as sources of bioactive compounds. Some plants can degrade the bioactive compounds they contain to endophytic microbes that grow in their tissues so that these endophytic microbes can produce the same compounds as their hosts. Our previous research Syarifah et al. (2021) found the antibacterial compound p-hydroxybenzaldehyde in endophytic fungi *Penicillium brefeldianum* (isolated from *S. zeylanicum* root bark). This p-hydroxybenzaldehyde compound is also produced by its host. To continue our research series, in this article, *S. zeylanicum* reports the potential bioactivity of endophytes from other parts of the plant, the bark of leaves and stems. Stem bark and leaves have higher presence and yield than flowers and seed (Figueiredo et al., 2008). Leaves and stems are the main organs responsible for accumulating active components with important medicinal value. Leaves are places for photosynthesis and play an important role in the plant's life. The leaves and stem can also be used as a synthetic storage organs for secondary metabolites (Li et al., 2020). To the best of our knowledge, this paper is the first report to expose the potential of the endophytic fungal extract from stem bark and leaves of *S. zeylanicum* as antibacterial and antioxidant.

2. EXPERIMENTAL SECTION

2.1 Plant Materials

The stem bark and leaves of *S. zeylanicum* (L.) were collected from the PALI regency (Penukal Abab Lematang Ilir), South Sumatra. The plant has been registered in the botanical field of the BRIN (Badan Riset dan Inovasi Nasional) Cibinong Research Center for Biology with the number B-417/V/DI.05.07/10/2021. Sampling was carried out in new state in July 2020. Sampling was carried out in a new state in July 2020. Isolation of endophytic fungi using young leaves and stem bark tissue. The leaves used are leaves in the third leaves position from the tip of the branch. The stem bark used is the bark from the central unit.

2.2 Isolation of Endophytic Fungi

Isolation of endophytic fungi begins by disinfecting the surface of leaves and stem bark of *S. zeylanicum*. Fresh plant offal was washed under running water until clean for ± 5 min. Then soak in alcohol for ± 3 min. Then rinse with sterile distilled water for ± 1 min, and then soak in 3% (w/v) sodium hypochlorite (NaOCl) for 1 min (Habisukan et al., 2021; Hanin and Fitri-asari, 2019). Surface sterilized leaves were cut aseptically by ± 2 cm, while the stem bark was cut $\pm 3 \times 0.5$ cm. The samples

were inoculated in PDA (Potato Dextrose Agar) plate, and incubated for 3-14 days at room temperature. Inspections were made every day until the fungi were visible. Fungal colonies grown on PDA plates with different morphological characteristics (color, size, and texture) were further purified. Purification was performed by transferring colonies to fresh PDA plate with single-spore isolation and then incubated at room temperature for 2×24 h. Purified fungal colonies are then transferred to the culture medium (in petri dishes) and stock cultures (in vitro) by culturing them on a PDA medium (Herlinda et al., 2021).

2.3 Endophytic Fungi Identification

Phenotypic characters both macroscopically and microscopically were used for endophytic fungi identification. Observation of colony characteristics included: colony color surface and reverse side; colony texture (cottony, granular, powdery, slimy); presence of exudate drops; presence of radial lines; presence of concentric circles. Microscopic characterization analysis using slide culture methods and observation of the hyphae, spores, color, and other specific characteristics under microscope until 500 x magnification. Both macroscopic and microscopic characterization then compared with fungi identification literature such as identification keys Pictorial Atlas of Soil and Seed Fungi (Watanabe, 2002), Fungi and Food Spoilage (Pitt and Hocking, 2009), Larone's medically necessary fungi (Walsh et al., 2018) and other fungi identification journal.

2.4 Endophytic Fungi Diversity Analysis

Diversity analysis of endophytic fungi from the leaves and stem bark of *S. zeylanicum* was estimated using the Simpson diversity index (Simpson, 1949) and Shannon-Weiner diversity index (Shannon and Weiner, 1949). The ecological interrelationships between the endophytic fungi species and the different organ were interpreted using Paleontological Statistics (PAST) Software to construct the Principal Component Analysis (PCA) (Hammer et al., 2001).

2.5 Molecular Identification

Potential endophytic fungi isolate continues with molecular identification based on the Internal Transcribed Spacer (ITS) area of DNA (rDNA). Primer ITS1 (5'-TCCTCCGCTTATTGATATG C-3') and ITS 4 (5'-TCCTCCGCTTATTGATATG C-3') was used for the amplification process. Assembly of DNA sequences forward and reverse primers were compiled using the Bioedit program to cut the unnecessary sequences. The sequence assembly was then entered into the Basic Local Alignment Search Tool (BLAST) at the website address <http://blast.ncbi.nlm.nih.gov/Blast.cgi>. Next, the sequences from sample and database were aligned with the CLUSTAL W method using the MEGA11 program, and a phylogenetic tree was constructed using the Neighbor-joining tree method with a bootstrap value of 1000 (Tamura et al., 2021).

2.6 Extraction and Cultivation

All endophytic fungi isolates were cultured using 6 blocks of pure cultured agar (± 6 mm in diameter) in 300 mL potato

dextrose broth (PDB) medium. Each isolate was refined in an erlenmeyer flask up to 300 mL×5. The cultures were then incubated for 4 weeks at room temperature under static conditions. The medium and biomass were separated using filter paper. Then ethyl acetate solvent was added to the culture medium (1:1) and extracted by partition (repeated three times). The ethyl acetate extract was separated using a rotary evaporator to obtain the extract (Budiono et al., 2019; Habisukan et al., 2021). The extract was concentrated using an oven at 40°C. The concentrated extract was weighed on an analytical balance.

2.7 Antibacterial Activity

Antibacterial activity is analysed using Kirby-Bauer method with NA (Natrium Agar) medium. Bacterial tests used 2 gram-negative bacteria (*Escherichia coli* InaCCB5 and *Salmonella thypi* ATCC1048) and 2-gram-positive bacteria (*Staphylococcus aureus* InaCCB4 and *Bacillus subtilis* InaCCB1204). The blank disc paper was dripped to endophytic fungal extract with a concentration of 400 µg/disc. Dilution of fungi extracts using Dimethylsulfoxide (DMSO). A positive control using Tetracycline with a concentration 30 µg/disc. Disc paper test was placed on the MHA (Muller Hinton Agar) plate that had been inoculated with the bacteria. Plate then incubated for 1×24 hours at incubation with setting 37°C, then the inhibition zone was observed. The diameter of the inhibition zone formed was measured with a ruler. The criteria for determining the antibacterial activity of the test sample and the diameter of the inhibition zone were determined by the following formula (Elfita et al., 2019; Syarifah et al., 2021):

$$\text{Strong} : \frac{A}{B} \times 100\% > 70\%;$$

$$\text{Moderate} : 50\% < \frac{A}{B} \times 100\% < 70\%;$$

$$\text{Weak} : \frac{A}{B} \times 100\% < 50\%$$

A: Inhibition zone (mm) of samples

B: Inhibition zone (mm) of antibiotic standard

2.8 Antioxidant Activity Analysis

Antioxidant activity was analyzed using the DPPH (2,2-diphenyl-1-picryl-hydrazyl-hydrate) free radical method. Each endophytic fungi ethyl acetate extract was dissolved in methanol to concentrations of 1000, 500, 250, 125, 62.5, 31.25, 15.625 g/mL (three replications). At 0.2 mL of each concentration 3.8 mL of 0.5 mM DPPH solution was added. The mixture became homogenized and left in a darkish tube for 30 minutes. Absorption was measured the use of a UV-Vis spectrophotometer at λ max 517 nm (Fadhillah et al., 2019). In this study, ascorbic acid was used as the standard antioxidant. Antioxidant activity was calculated from the inhibition rate of DPPH absorption and the IC₅₀ value.

$$\% \text{Inhibition} = \frac{A_k - A_s}{A_s}$$

A_k = Absorbance of control

A_s = Absorbance of samples

3. RESULTS AND DISCUSSION

3.1 Endophytic Fungi of *Syzygium zeylanicum*

Isolation of endophytic fungi was characterized by the presence of fungal mycelium around the surface sterilized plant organs. The leaves and stem bark of *S. zeylanicum* dominated by white fungal colony (Figure 1). The frequency of endophyte establishment by whether young, mature old segments (Disanayake et al., 2015). Young leaves have lower concentrations of antifungal and anti-herbivorous substances compared to mature stems (stem bark), which may be the reason for the lower leaf colonization rate than stem bark. A total of 16 endophytic fungi were isolated from stem bark (4 isolates) and leaves (6 isolates) of *S. zeylanicum*.






Figure 1. Endophytic Fungi from *Syzygium zeylanicum*. (A) Leaves (B) Stem Bark. Isolation Using PDA Medium after 7 Days Incubation in Room Temperature

Colony morphology of the 10 endophytic fungi isolates varied in shape and color (Figure 2) along with the shape of the hyphae and conidia which showed their characteristics (Figure 3). Colonies of endophytic fungus isolates showed various physical appearances. Large-scale hyphal growth occurred in bark and leaf samples, with predominant green, white, and black colonies. The results of the isolation of endophytic fungi from the bark obtained five isolates with codes SZT3 to SZT5 and SZT7 (Figure 2: A-D), and from leaf organs, seven isolates were obtained with codes SZL1 to SZL7 (Figure 2: E-J). The population structure of endophytes can be significantly affected by factors such as host's genetic culture, age, and environmental conditions (Jia et al., 2016).

Endophytic fungi are facultative biotrophs, so they can complete their life cycle outside their host. The role of endophytic fungi can support the accumulation of bioactive compounds by helping to increase the production of secondary metabolites in their host plants (Khare et al., 2018; Sharma et al., 2018; Yuan et al., 2019). Endophytic fungi can produce bioactive compounds exclusively for their host plants, compounds can be the same or different from their hosts, this is very important to improve the adaptability of endophytic fungi and their host

Table 1. Colony Characteristics of Endophytic Fungi of Jambu Nasi-Nasi

Code	Surface Color	Reverse Color	Texture	Elevation	Pattern	Exudate Drops	Radial Line	Concentric Circle
SZT3	Yellowish white	Clear yellow	Cottony	Flat	Spread	-	-	√
SZT4	Grayish white	Unpigmented	Cottony	Flat	Spread	-	-	√
SZT5	Yellowish green	Clear yellow	Powdery	Flat	Spread	-	√	√
SZT7	Yellowish white	Clear yellow	Cottony	Flat	Spread	-	-	√
SZL1	Milky white	White	Cottony	Flat	Zonate	-	-	-
SZL2	Dusty white	Unpigmented	Cottony	Umbonate	Spread	-	-	√
SZL3	Cream and white	Cream	Velvety	Umbonate	Flowery	-	-	√
SZL4	Gray	Gray	Cottony	Raised	Spread	-	-	-
SZL5	Greenish gray	Dark green	Cottony	Flate	Zonate	-	-	√
SZL7	Milky white	White	Velvety	Flat	Zonate	-	-	√

**Figure 2.** Colony Morphology of Endophytic Species; from Stem Bark (SZT3 (A); SZT4 (B); SZT5 (C); SZT7 (D)); and from Leaves (SZL1 (E); SZL2 (F); SZL3(G); SZL4 (H); SZL5 (I); SZL7 (J)). Colony Growth on Potato Dextrose Agar Medium after 5-7 Days**Figure 3.** Microscopic Characterization of Endophytic Fungal Isolates; from Stem Bark (SZT3 (A); SZT4 (B); SZT5 (C); SZT7 (D)); and from Leaves (SZL1 (E); SZL2 (F); SZL3 (G); SZL4 (H); SZL5 (I); SZL7 (J)). Observations Were Made on after 5-7 Days on PDA Medium Under Digital Microscope Magnification 400×

plants, such as ¹¹tolerance to biotic and abiotic stresses (Jia et al., 2016).

3.2 Morphological Identification of Endophytic Fungi

Fungal identification begins by comparing morphologically known species. Observations can be made with the naked eye (macroscopic) or with a compound microscope (microscopic). Table 1 shows the characteristics of colonies from color to growth type. On the other hand, Table 2 shows the microscopic features from the shape of the spores to the specific features.

SZL4 isolate had colonies that proliferated throughout the disc with a cottony texture and grayish-white (Figure 2L; Table 1). The microscopic data supported identification where the isolates had sub-ovoid to ellipsoid conidia with wide rounded apex, 1 insulated and thick-walled (Figure 3L; Table 2). Based on these characteristics, SZL4 is close to the genus *Lasidopodia*, which is by what was reported by Kapoor and Saxena (2014).

Endophytic fungi isolate SZT5 and SZT7 had the same colony shape with the yellowish-green colony and when compared with microscopic morphology (Figures 3C & 3D), both had setae like hyphae with globose-shaped conidia (Table 2). Phialid in SZT2 & SZT7 has a verticillate shape so it is close to

the morphology of *Trichoderma koningi*. Meanwhile, the isolate SZT5 had verticillate, short, and thin phialid (Table 2). And the conidia from this isolate were in the form of hyaline and ovate phialosporus, so they were identified as *Trichoderma aureoviridae*.

Isolates SZL1, SZL2, and SZL3 had yellowish-white colonies with a flowery pattern (Figure 2E-G). The isolate SZL2 (Figure 3F) showed coenocytic hyphae with lobate sporangiospores and coiling around the oogoniumphores (Table 2) so that SZL2 was identified as *Phytium zingiberum*. While SZL1 has a sporangia lobate form that forms vesicle formations so that this fungus is close to *Phytium torulosum*. And in SZL3 isolate the conidia were not completely differentiated, where the conidia grew directly on the hyphae and were blastosporous so that the SZL3 isolate was closer to *Septonema* sp.

SZT3 isolate had yellowish-white colonies that spread throughout the disc with a cotton-like texture (Figure 2A). SZT3 isolate was identified as *Phialemonium* sp. because microscopically (Figure 3A) it has septate hyphae and there are single short conidia along the hyphae. SZT4 colonies (Figure 2B) grew all over the disc, grayish-white in color with a cottony

Table 2. Microscopic Characteristics of Jambu Nasi-Nasi Endophytic Fungi

Isolate Code	Type of Spore	Shape of Spore	Hyphae	Specific Characteristic	Species of Identification
SZT3	Conidia	Peglike	Septate	Single-celled, shirt and peglike	<i>Phialemonium</i> sp.
SZT4	Conidia	Oblong	Septate	Conidia are oblong, easily disrupted clusters	<i>Acremonium</i> sp.
SZT5	Conidia	Globose	Septate	Conidiospores branches, conidia phialosporous	<i>Trichoderma aureoviridae</i>
SZT7	Conidia	Globose	Septate	Conidiospores branches, conidia phialosporous	<i>Trichoderma koningi</i>
SZL1	Oospore	Lobate	Coenocytic	Vesicle formation from lobate sporangia	<i>Phytium torulosum</i>
SZL2	Antheridium spore	Lobate	Coenocytic	Coiling around oogoniumphores	<i>Phytium zingiberum</i>
SZL3	Conidia	Blastosporous	Septate	Conidia is not well differentiated, truncate both ends	<i>Septonema</i> sp.
SZL4	Conidia	Cylindrical	Septate	Bearing monoverticillate penicilla	<i>Lasiodiplodia pseudotheobromae</i>
SZL5	Conidia	Cylindrical	Septate	Sporodochia subglobose, conidiophores phialosporous	<i>Volutella cilliata</i>
SZL7	Conidia	Subglobose	Septate	Ovate conidia, short phialides, densely arranged	<i>Trichophyton mentagrophytes</i>

texture. The microscopic characteristics of SZT4 (Figure 3B; Table 2) have a single erect and hyaline phialid, and oblong conidia and conidia form easily disturbed clusters at the ends of the phialides. The isolate SZT4 was identified as *Acremonium* sp. based on its morphological characteristics.

Table 3. Endophytic Fungi Diversity from *S. zeylanicum*

Taxon (Genera)	Tissues of <i>Syzygium zeylanicum</i>		
	Leaves	Stem Bark	Total
Phytium	2	0	0
Septonema	1	0	1
Lasiodiplodia	1	0	1
Volutella	1	0	1
Trichophyton	1	0	1
Trichoderma	0	2	2
Phialemonium	0	1	1
Acremonium	0	1	1
No. of total fungal isolates	6	4	10
Simpson's index (D)	0.1837	0.4400	0.1389
Simpson's index of diversity (1-D)	0.8163	0.5600	0.8611
Shannon index of diversity (H')	1.7480	0.9503	2.0950

SZL5 isolates had greenish-gray colonies and dark reverse color (Figure 2I) supported microscopically (isolate 3I; Table 2) showing subglobose sporodochia and phialosporous and cylindrical conidiophores. Based on the macroscopic and

microscopic morphological characteristics, it was close to the isolate *Volutella ciliate*. SZL7 colonies were milky white with a velvety texture (Figure 2J) with microscopic morphology having septate hyphae and branching conidiophores and macroconidia attached to the hyphae (Figure 3J) so that SZL7 isolate was identified as *Trichophyton mentagrophytes*.

3.3 Endophytic Fungi Diversity Analysis

The stem bark of *S. zeylanicum* obtained four isolates and the leaves obtained six isolates. The diversity index values (Shannon and Weiner, 1949; Simpson, 1949) for these *S. zeylanicum* tissues are listed in Table 3. The α -diversity values were acquired thru the PAST software (Sadeghi et al., 2019). The Shannon-Weiner and Simpson variety index values had been maximum in leaves ($H'=1.748$ and $1-D=0.8163$, respectively) and lowest in root bark ($H'=0.9503$ and $1-D=0.56$, respectively). In addition, the general variety values of the endophytic fungal populations had been $H'=2.095$ and $1-D=0.8611$ (Table 3).

The two principal components or axes (1-2) obtained by PCA accounted for 78% of the total variability of the fungus (Figure 4). Analysis showed that some isolates showed affinity for specific tissues. There was no common genus of endophytes from the bark of leaves and stems, but the genus *Trichoderma* was found in the bark of the roots of *S. zeylanicum* (Syarifah et al., 2021). Some isolated endophytic taxa prefer a particular tissue type. Some endophytic can invade the tree host through the root system and move from the soil to new niches within the plant (Gazis and Chaverri, 2010).

Table 4. Antibacterial Activity Percentage of Endophytic Fungi Extracts of *S. zeilanicum* (at a Concentration of 400 µg/disc) Compared to Tetracycline 30 µg/disc and Antioxidant Activity with Ascorbic Acid Antioxidant Standard and Weight of The Extract from Endophytic Fungi Cultivation in PDB Medium (5×300 mL)

Isolate Code	Genus/Species of Identification	Fungal Extract Weight (Gram)	% Antibacterial Activity				Antioxidant Activity IC50 (µg/mL)	Antioxidant Activity IC50 (µg/mL)
			<i>E. coli</i>	<i>B. subtilis</i>	<i>S. thypi</i>	<i>S. aureus</i>		
SZT3	<i>Phialemonium</i> sp.	5.2	81.0±3.19 ***	72.8±5.40 ***	80.8±2.56 ***	77.7±0.23 ***	28.5 ***	
SZT4	<i>Acronium</i> sp.	5.5	68.5±1.82 **	62.4±3.20 **	69.5±1.27 **	71.6±0.95 ***	3.56 ****	
SZT5	<i>Trichoderma</i> sp.	5.3	63.8±3.45 **	57.4±4.20 **	74.4±1.63 ***	70.6±1.71 ***	10.25 ***	
SZT7	<i>Trichoderma koningi</i>	4.3	75.9±2.16 ***	81.0±3.43 ***	78.7±2.29 ***	84.0±1.61 ***	98.41 ***	
SZL1	<i>Phytium</i> sp.	4.7	63.1±0.78 **	57.6±5.27 **	74.9±2.09 ***	78.8±1.07 ***	9.15 ****	
SZL2	<i>Phytium zingiberum</i>	5.6	34.7±0.82 *	40.6±1.97 *	58.5±0.68 **	65.4±4.20 **	4.31 ****	
SZL3	<i>Septonema</i> sp.	5.1	74.6±2.12 ***	64.7±4.93 **	73.4±0.71 ***	82.0±0.96 ***	6.79 ****	
SZL4	<i>Lasiodiplodia pseudotheobromae</i>	8.8	76.7±1.90 ***	78.3±1.85 ***	76.3±1.43 ***	76.0±1.71 ***	3.30 ****	
SZL5	<i>Volutella ciliata</i>	4.6	59.1±1.86 ***	50.6±0.66 ***	57.2±0.35 ***	62.5±0.85 ***	6.18 ****	
SZL7	<i>Trichophyton mentagrophytes</i>	4.7	46.6±1.48 *	42.1±6.05 *	75.3±1.72 ***	69.1±5.58 **	27.69 ***	
	Positive control		Tetracycline 100±1.40 ***	Tetracycline 100±1.28 ***	Tetracycline 100±1.12 ***	Tetracycline 100±1.70 ***	Ascorbic Acid 2.73 ****	

Note: Antibacterial activity percentage: *** strong (≥70%), **moderate (50-70%), and *weak (<50%) [15,42]. Antioxidant activity IC50 (µg/mL): ****⁶very strong <10 µg/mL; ***strong <100 µg/mL; **moderat 100-500 µg/mL; *weak >500 µg/mL (Mbekou et al., 2021; Metasari et al., 2020).




Figure 4. PCA of Endophytic Fungi Isolated from Leaves and Stem Bark of *S. zeylanicum*. Components 1 and 2 Described 78% of The Total Variation of The Fungi. The Percentages Shown Are Percentages of Variation Explained by The Components. The Data was Processed Using The PAST Software

3.4 Antibacterial and Antioxidant Activity of Endophytic Fungi Extract

Biological activity screening for potential endophytic fungus *S. zeylanicum* was carried out by analyzing its antibacterial and antioxidant activity. Antioxidant activity was carried out by DPPH activity to remove free radicals. Upon receiving an electron from an antioxidant compound, the DPPH radical is reduced to DPPH (Budiono et al., 2019; Praptiwi et al., 2018). The purple color of the DPPH radical changes to yellow (Pavithra and Vadivukkarasi, 2015). Overall endophytic fungi from leaves and bark showed vital to extreme antioxidant activity.

Endophytic fungi ethyl acetate extract from *S. zeylanicum* has potential as an antioxidant and antibacterial as shown in Table 4. There are 7 out of 10 extracts of endophytic fungi showing strong antibacterial activity (>70% inhibition) against *S. typhi* and *S. aureus* bacteria. While the antibacterial activity against *E. coli* showed as many as 5 fungal extracts that had strong activity and against *B. subtilis* bacteria as many as 4 isolates which had strong activity. All endophytic fungi extracts showed very strong antioxidant values (IC₅₀ <10 µg/mL).

The data in Table 4 shows that all endophytic fungal extracts could inhibit the growth of the bacteria which was indicated by the formation of a clear zone. The majority of extracts gave moderate to strong antibacterial activity against the four tested bacteria. SZL1, SZL2 and SZL7 extract provided weak antibacterial activity. The same thing can be seen from the ability of endophytic fungi extracts to reduce DPPH free radicals. All extracts provide very strong antioxidant activity. Isolate SZL4 and SZL5 endophytic fungi from the leaves and stem bark of *S. zeylanicum* had very strong antioxidant and strong antibacterial against four pathogens bacterial. In this study, only one isolate

was selected to be identified to the molecular stage with potential antioxidant and antimicrobial activity with a high yield of extract.

A total of 7 out of 10 extracts of the endophytic fungus *S. zeylanicum* showed a positive correlation with having strong antioxidant and antibacterial activity. The development of compounds that have dual activity as antioxidant and antibacterial is beneficial in the treatment of chronic infections such as Alzheimer's- infection is commonly associated with gram-negative anaerobic bacteria (Dioguardi et al., 2020), Cystic Fibrosis, and other complex multigenic diseases (Martelli and Giacomini, 2018; Oset-Gasque and Marco-Contelles, 2018). We suggest that bioactive compounds with antioxidant and antibacterial activity include beta-lactam-based compounds, Dihydroquinolines, Quinolines, Piperidone-hydrazides, Isatin-thiosemicarbazones, Barbiturates, Indolophanes, Triazoles and substituted 3-indoles (Martelli and Giacomini, 2018).

Antibacterials and antioxidant can be positively or negatively correlated. 70% *Syzygium zeylanicum* endophytic fungi extract showed a positive correlation where the antioxidant and antibacterial activity values were equally strong. In the study of Li et al. (2020) stated that natural flavonoids (i.e. quercetin, baicalin and rutin) have high antibacterial and antioxidant activity.

Not all endophytic fungi extract of *S. zeylanicum* showed a positive correlation, where 3 fungal extracts showed strong antioxidant activity but weak to moderate antibacterial activity. The negative correlation between antioxidant and antibacterial was also mentioned in another study where several tetradecane compounds, -palmitolactone and ethyl hydrocinnamate contributed to this correlation (Bittencourt et al., 2015). With the antibacterial and antioxidant screening of *S. zeylanicum* fungal extract, it can be used as a reference in obtaining alternatives to produce potential active compounds.

The low yield of bioactive substances is an obstacle in the development of these bioactive substances as candidates for medicinal ingredients because to go through a long research stage, many substances are needed. In this study, SZL4 isolates of endophytic fungi were found that produce twice the yield of other endophytic fungi, and have strong antibacterial and antioxidant activities. This discovery is very valuable so further research needs to be designed to make endophytic fungi as source of future medicinal raw materials.

3.5 Molecular Identification

The endophytic fungi isolates selected for the molecular identification were SZL4 fungi isolated from leaves. Extracts SZL4 fungi produce more yield than other extracts Fungi. The potential for endophytic fungi to be developed as a new source of medicinal raw materials, apart from the ability to biological activity, can also be viewed from the yield of the resulting extract. Molecular test results are presented in the form of a phylogenetic tree (Figure 5) to assist in the identification process.

Isolate SZL4 has a similarity of 100% and is in the same

- Dissanayake, A. J., W. Zhang, L. Mei, E. Chukeatirote, J. Y. Yan, X. Li, and K. D. Hyde (2015). *Lasiodiplodia pseudotheobromae* Causes Pedicel and Peduncle Discolouration of Grapes in China. *Australasian Plant Disease Notes*, **10**(1); 1–5
- Elfita, E., Mardiyanto, Fitriya, J. E. Larasati, Julinar, H. Widjajanti, and Muharni (2019). Antibacterial Activity of *Cordyline fruticosa* Leaf Extracts and its Endophytic Fungi Extracts. *Biodiversitas Journal of Biological Diversity*, **20**(12); 3804–3812
- Fadhillah, F., Elfita, Muharni, H. Yohandini, and H. Widjajanti (2019). Chemical Compound Isolated from Antioxidant Active Extract of Endophytic Fungus *Cladosporium tenuissimum* in *Swietenia mahagoni* Leaf Stalks. *Biodiversitas Journal of Biological Diversity*, **20**(9); 2645–2650
- Figueiredo, A. C., J. G. Barroso, L. G. Pedro, and J. J. Scheffer (2008). Factors Affecting Secondary Metabolite Production in Plants: Volatile Components and Essential Oils. *Flavour and Fragrance Journal*, **23**(4); 213–226
- Gazis, R. and P. Chaverri (2010). Diversity of Fungal Endophytes in Leaves and Stems of Wild Rubber Trees (*Hevea brasiliensis*) in Peru. *Fungal Ecology*, **3**(3); 240–254
- Habisukan, U. H., E. Elfita, H. Widjajanti, A. Setiawan, and A. R. Kurniawati (2021). Diversity of Endophytic Fungi in *Syzygium aqueum*. *Biodiversitas Journal of Biological Diversity*, **22**(3); 1129–1137
- Hammer, Ø., D. A. Harper, and P. D. Ryan (2001). PAST: Paleontological Statistics Software Package for Education and Data Analysis. *Palaeontologia Electronica*, **4**(1); 9
- Hanin, N. A. and P. D. Fitriyari (2019). Identification of Endophytic Fungi from Fruits and Seeds of Jambolana (*Syzygium cumini* L.) Skeels. In *IOP Conference Series: Earth and Environmental Science*, **276**; 012060
- Herlinda, S., M. Gustianingtyas, S. Suwandi, R. Suharjo, J. M. P. Sari, and R. P. Lestari (2021). Endophytic Fungi Confirmed as Entomopathogens of The New Invasive Pest, The Fall Armyworm, *Spodoptera frugiperda* (JE Smith)(Lepidoptera: Noctuidae), Infesting Maize in South Sumatra, Indonesia. *Egyptian Journal of Biological Pest Control*, **31**(1); 1–13
- Jia, M., L. Chen, H. L. Xin, C. J. Zheng, K. Rahman, T. Han, and L. P. Qin (2016). A Friendly Relationship Between Endophytic Fungi and Medicinal Plants: a Systematic Review. *Frontiers in Microbiology*, **7**; 906
- Kapoor, N. and S. Saxena (2014). Potential Xanthine Oxidase Inhibitory Activity of Endophytic *Lasiodiplodia pseudotheobromae*. *Applied Biochemistry and Biotechnology*, **173**(6); 1360–1374
- Kaul, S., S. Gupta, M. Ahmed, and M. K. Dhar (2012). Endophytic Fungi from Medicinal Plants: a Treasure Hunt for Bioactive Metabolites. *Phytochemistry Reviews*, **11**(4); 487–505
- Khare, E., J. Mishra, and N. K. Arora (2018). Multifaceted Interactions Between Endophytes and Plant: Developments and Prospects. *Frontiers in Microbiology*, **9**; 2732
- Kumar, V., R. Soni, L. Jain, B. Dash, and R. Goel (2019). Endophytic Fungi: Recent Advances in Identification and Explorations. *Advances in Endophytic Fungal Research*; 267–281
- Li, Y., D. Kong, Y. Fu, M. R. Sussman, and H. Wu (2020). The Effect of Developmental and Environmental Factors on Secondary Metabolites in Medicinal Plants. *Plant Physiology and Biochemistry*, **148**; 80–89
- Lu, Y., C. Chen, H. Chen, J. Zhang, and W. Chen (2012). Isolation and Identification of Endophytic Fungi from *Actinidia macrosperma* and Investigation of Their Bioactivities. *Evidence-Based Complementary and Alternative Medicine*, **2012**; 1–8
- Mane, R. S., P. M. Paarakh, and A. B. Vedamurthy (2018). Brief Review on Fungal Endophytes. *International Journal of Secondary Metabolite*, **5**(4); 288–303
- Manyi-Loh, C., S. Mamphweli, E. Meyer, and A. Okoh (2018). Antibiotic Use in Agriculture and its Consequential Resistance in Environmental Sources: Potential Public Health Implications. *Molecules*, **23**(4); 795
- Martelli, G. and D. Giacomini (2018). Antibacterial and Antioxidant Activities for Natural and Synthetic Dual-Active Compounds. *European Journal of Medicinal Chemistry*, **158**; 91–105
- Mbekou, M. I. K., D. Dize, V. L. Yimgang, F. Djague, R. M. K. Toghueo, N. Sewald, B. N. Lenta, and F. F. Boyom (2021). Antibacterial and Mode of Action of Extracts from Endophytic Fungi Derived from *Terminalia mantaly*, *Terminalia catappa*, and *Cananga odorata*. *BioMed research international*, **2021**
- Metasari, S., E. Elfita, M. Muharni, and H. Yohandini (2020). Antioxidant Compounds from The Stem Bark of *Syzygium samarangense* L. *Molekul*, **15**(3); 175–183
- Nei, M. and S. Kumar (2000). *Molecular Evolution and Phylogenetics*. Oxford University Press, USA
- Omoarelojic, L. O. and J. Van Staden (2020). Plant-Endophytic Fungi Interactions: a Strigolactone Perspective. *South African Journal of Botany*, **134**; 280–284
- Oset-Gasque, M. J. and J. Marco-Contelles (2018). Alzheimer's Disease, The "One-Molecule, One-Target" Paradigm, and The Multitarget Directed Ligand Approach. *ACS Chemical Neuroscience*, **9**(3); 401–403
- Pandey, G., K. Verma, and M. Singh (2014). Evaluation of Phytochemical, Antibacterial and Free Radical Scavenging Properties of *Azadirachta indica* (Neem) Leaves. *International Journal of Pharmacy and Pharmaceutical Sciences*, **6**(2); 444–447
- Pavithra, K. and S. Vadivukkarasi (2015). Evaluation of Free Radical Scavenging Activity of Various Extracts of Leaves from *Kedrostis foetidissima* (Jacq.) Cogn. *Food Science and Human Wellness*, **4**(1); 42–46
- Phaniendra, A., D. B. Jestadi, and L. Periyasamy (2015). Free Radicals: Properties, Sources, Targets, and Their Implications in Various Diseases. *Indian Journal of Clinical Biochemistry*, **30**(1); 11–26
- Pitt, J. I. and A. D. Hocking (2009). The Ecology of Fungal Food Spoilage. In *Fungi and Food Spoilage*. Springer;3–9

- Praptiwi, R. M., D. Wulansari, A. Fathoni, and A. Agusta (2018). Antibacterial and Antioxidant Activities of Endophytic Fungi Extracts of Medicinal Plants from Central Sulawesi. *Journal of Applied Pharmaceutical Science*, **8**(08); 069–074
- Sadeghi, F., D. Samsampour, M. A. Seyahooei, A. Bagheri, and J. Soltani (2019). Diversity and Spatiotemporal Distribution of Fungal Endophytes Associated with *Citrus reticulata* Cv. Siyadoo. *Current Microbiology*, **76**(3); 279–289
- Segaran, G. and M. Sathiavelu (2019). Fungal Endophytes: a Potent Biocontrol Agent and a Bioactive Metabolites Reservoir. *Biocatalysis and Agricultural Biotechnology*, **21**; 101284
- Shannon, C. and V. Weiner (1949). A Mathematical Theory of Communication University Press. *Illinois Urban*, **1**; 101–107
- Sharma, G. N., G. Gupta, and P. Sharma (2018). A Comprehensive Review of Free Radicals, Antioxidants, and Their Relationship with Human Ailments. *Critical Reviews™ in Eukaryotic Gene Expression*, **28**(2)
- Simpson, E. H. (1949). Measurement of Diversity. *Nature*, **163**(4148); 688–688
- Syarifah, S., E. Elfita, H. Widjajanti, A. Setiawan, and A. R. Kurniawati (2021). Diversity of Endophytic Fungi from The Root Bark of *Syzygium zeylanicum*, and The Antibacterial Activity of Fungal Extracts, and Secondary Metabolite. *Biodiversitas Journal of Biological Diversity*, **22**(10); 4572–4582
- Tamura, K., G. Stecher, and S. Kumar (2021). MEGA11: Molecular Evolutionary Genetics Analysis Version 11. *Molecular Biology and Evolution*, **38**(7); 3022–3027
- Teodoro, A. J. (2019). Bioactive Compounds of Food: Their Role in The Prevention and Treatment of Diseases
- Uzma, F., C. D. Mohan, C. N. Siddaiah, and S. Chowdappa (2019). Endophytic Fungi: Promising Source of Novel Bioactive Compounds. In *Advances in Endophytic Fungal Research*. Springer; 243–265
- Vinodkumar, T. G. (2015). Leaf Essential Oil Composition of Six *Syzygium* Species from The Western Ghats, South India. *Records of Natural Products*, **9**(4); 592
- Walsh, T. J., R. T. Hayden, and D. H. Larone (2018). *Larone's Medically Important Fungi: a Guide to Identification*. John Wiley & Sons
- Watanabe, T. (2002). *Pictorial Atlas of Soil and Seed Fungi: Morphologies of Cultured Fungi and Key to Species*. CRC press
- Wei, W., N. Jiang, Y. N. Mei, Y. L. Chu, H. M. Ge, Y. C. Song, S. W. Ng, and R. X. Tan (2014). An Antibacterial Metabolite from *Lasiodiplodia pseudotheobromae* F2. *Phytochemistry*, **100**; 103–109
- Yuan, Z., Y. Tian, F. He, and H. Zhou (2019). Endophytes from *Ginkgo biloba* and Their Secondary Metabolites. *Chinese Medicine*, **14**(1); 1–40
- Zhao, J., L. Zhou, J. Wang, T. Shan, L. Zhong, X. Liu, and X. Gao (2010). Endophytic Fungi for Producing Bioactive Compounds Originally from Their Host Plants. *Curr Res, Technol Educ Trop Appl Microbiol Microbial Biotechnol*, **1**; 567–576

Antioxidant and Antibacterial Activities of Endophytic Fungi Extracts of *Syzygium zeylanicum*

ORIGINALITY REPORT

14%

SIMILARITY INDEX

7%

INTERNET SOURCES

13%

PUBLICATIONS

3%

STUDENT PAPERS

PRIMARY SOURCES

- 1 dspace.nm-aist.ac.tz 1 %
Internet Source
- 2 Flor N. Rivera-Orduña, Roberto A. Suarez-Sanchez, Zoila R. Flores-Bustamante, Jorge N. Gracida-Rodriguez et al. "Diversity of endophytic fungi of *Taxus globosa* (Mexican yew)", *Fungal Diversity*, 2010 1 %
Publication
- 3 Yanqun Li, Dexin Kong, Ying Fu, Michael R. Sussman, Hong Wu. "The effect of developmental and environmental factors on secondary metabolites in medicinal plants", *Plant Physiology and Biochemistry*, 2020 1 %
Publication
- 4 www.semanticscholar.org 1 %
Internet Source
- 5 pdfs.semanticscholar.org 1 %
Internet Source
- 6 Submitted to Universitas Airlangga 1 %
Student Paper

7	academic-accelerator.com Internet Source	1 %
8	Riri Novita Sunarti, Sri Budiarti, Marieska Verawaty, Bayo Alhusaeri Siregar, Poedji Loekitowati Hariani. "Diversity of Antibiotic-Resistant <i>Escherichia coli</i> from Rivers in Palembang City, South Sumatra, Indonesia", AIMS Environmental Science, 2022 Publication	<1 %
9	Giulia Martelli, Daria Giacomini. "Antibacterial and antioxidant activities for natural and synthetic dual-active compounds", European Journal of Medicinal Chemistry, 2018 Publication	<1 %
10	"Advances in Endophytic Fungal Research", Springer Nature, 2019 Publication	<1 %
11	"Advances in Endophytic Research", Springer Nature, 2014 Publication	<1 %
12	Yi-Jie Zhai, Guang-Ming Huo, Qiang Zhang, Ding Li, Da-Cheng Wang, Jian-Zhao Qi, Wen-Bo Han, Jin-Ming Gao. " Phaeosphaones: Tyrosinase Inhibitory Thiodiketopiperazines from an Endophytic ", Journal of Natural Products, 2020 Publication	<1 %

13	tjnpr.org Internet Source	<1 %
14	www.thaiscience.info Internet Source	<1 %
15	Mohamed K. Abdel-Rafei, Noura M. Thabet, Mohamed M. Amin. "Concerted regulation of OPG/RANKL/ NF-κB/MMP-13 trajectories contribute to ameliorative capability of prodigiosin and/or low dose γ-radiation against adjuvant- induced arthritis in rats", <i>International Immunopharmacology</i> , 2022 Publication	<1 %
16	Shubhpriya Gupta, Preeti Chaturvedi, Manoj G. Kulkarni, Johannes Van Staden. "A critical review on exploiting the pharmaceutical potential of plant endophytic fungi", <i>Biotechnology Advances</i> , 2020 Publication	<1 %
17	jurnal.uns.ac.id Internet Source	<1 %
18	Meyyappan Narayanan, Divya Baskaran, Vasumathi Sampath. " Experimental design of hydrotropic extraction for recovery of bioactive limonin from lemon (L.) seeds ", <i>Separation Science and Technology</i> , 2021 Publication	<1 %

19 Sharmilla Ashokhan, Sujatha Ramasamy, Saiful Anuar Karsani, Rashidi Othman, Jamilah Syafawati Yaacob. " Analysis of bioactive pigments in coloured callus of for possible use as functional natural colourants ", Pigment & Resin Technology, 2019
Publication

20 "NCSBN's Environmental Scan COVID-19 and Its Impact on Nursing and Regulation", Journal of Nursing Regulation, 2021
Publication

21 Submitted to Universitas Tidar
Student Paper

22 Submitted to University of Liverpool
Student Paper

23 Hui Wang, Tianxing Liu, Zhihong Xin. "A new glucitol from an endophytic fungus Fusarium equiseti Salicorn 8", European Food Research and Technology, 2014
Publication

24 Martín-Rodríguez, Alberto, Fernando Reyes, Jesús Martín, Juan Pérez-Yépez, Milagros León-Barrios, Alan Couttolenc, César Espinoza, Ángel Trigos, Víctor Martín, Manuel Norte, and José Fernández. "Inhibition of Bacterial Quorum Sensing by Extracts from

Aquatic Fungi: First Report from Marine Endophytes", Marine Drugs, 2014.

Publication

25

Prabukumar Seetharaman, Sathishkumar Gnanasekar, Rajkuberan Chandrasekaran, Gobinath Chandrakasan et al. "Isolation of limonoid compound (Hamisonine) from endophytic fungi *Penicillium oxalicum* LA-1 (KX622790) of *Limonia acidissima* L. for its larvicidal efficacy against LF vector, *Culex quinquefasciatus* (Diptera: Culicidae)", Environmental Science and Pollution Research, 2017

Publication

26

Rosario del Carmen Flores-Vallejo, Jorge Luis Folch-Mallol, Ashutosh Sharma, Alexandre Cardoso-Taketa et al. "ITS2 ribotyping, in vitro anti-inflammatory screening, and metabolic profiling of fungal endophytes from the Mexican species *Crescentia alata* Kunth", South African Journal of Botany, 2020

Publication

27

www.ijsk.org
Internet Source

28

Prasad G. Jamkhande, Vikas A. Suryawanshi, Tukaram M. Kaylankar, Shailesh L. Patwekar. "Biological activities of leaves of ethnomedicinal plant, *Borassus flabellifer*

<1 %

<1 %

<1 %

<1 %

Linn. (Palmyra palm): An antibacterial, antifungal and antioxidant evaluation", Bulletin of Faculty of Pharmacy, Cairo University, 2016

Publication

29

Yawei Que, Donghai Huang, Shuangjun Gong, Xuejiang Zhang et al. "Indole-3-Carboxylic Acid From the Endophytic Fungus Lasiodiplodia pseudotheobromae LPS-1 as a Synergist Enhancing the Antagonism of Jasmonic Acid Against Blumeria graminis on Wheat", Frontiers in Cellular and Infection Microbiology, 2022

Publication

<1 %

30

dokumen.pub
Internet Source

<1 %

31

e-journal.unair.ac.id
Internet Source

<1 %

32

"Antimicrobial compounds from microorganisms: Production, characterization, and applications", Antimicrobials, 2015.

Publication

<1 %

33

"Pharmaceuticals from Microbes", Springer Nature, 2019

Publication

<1 %

34

"Postharvest Pathology", Springer Science and Business Media LLC, 2021

Publication

<1 %

35

Satish K. Verma, Surendra K. Gond, Ashish Mishra, Vijay K. Sharma et al. "Impact of environmental variables on the isolation, diversity and antibacterial activity of endophytic fungal communities from *Madhuca indica* Gmel. at different locations in India", *Annals of Microbiology*, 2013

Publication

<1 %

36

krishikosh.egranth.ac.in

Internet Source

<1 %

37

Anupma Dahiya, Kavita Chahar, Satyavir S. Sindhu. "The rhizosphere microbiome and biological control of weeds: A review", *Spanish Journal of Agricultural Research*, 2020

Publication

<1 %

38

Gurleen Kaur Sodhi, Sanjai Saxena. "Chapter 12 Role of Endophytic Fungi in Promoting Plant Health", Springer Science and Business Media LLC, 2023

Publication

<1 %

39

Jelly Milinia Puspita Sari, Siti Herlinda, Suwandi Suwandi. "Endophytic fungi from South Sumatra (Indonesia) in seed-treated corn seedlings Affecting development of the fall armyworm, *Spodoptera frugiperda* J.E. Smith (Lepidoptera: Noctuidae)", *Egyptian Journal of Biological Pest Control*, 2022

Publication

<1 %

40

Libo Xiang, Shuangjun Gong, Lijun Yang, Jianjun Hao, MinFeng Xue, FanSong Zeng, Xuejiang Zhang, WenQi Shi, Hua Wang, Dazhao Yu. "Biocontrol potential of endophytic fungi in medicinal plants from Wuhan Botanical Garden in China", Biological Control, 2016

Publication

<1 %

41

Norma Tiku Kambuno, Melva Louisa, Puspita Eka Wuyung, Taniawati Supali. "Impact of Ultrasonic Assisted Extraction on the Protein Yield from Moringa Oleifera Lam Leaves and Its Functional Characterization", Research Square Platform LLC, 2023

Publication

<1 %

42

Sandip Chowdhury, Suvranil Ghosh, Surendra K. Gond. " Anti - MRSA and clot lysis activities of isolated from Roxb ", Journal of Basic Microbiology, 2022

Publication

<1 %

43

Supaphon Preuttiorn, Preedanon Sita. "Antimicrobial and antioxidant activities of endophytic fungi extracts isolated from Carissa carandas", African Journal of Microbiology Research, 2019

Publication

<1 %

44

Vishnu Sharma, Tarun Kumar Kumawat, Ruchi Seth, Anima Sharma, Subhash Chandra. "A

<1 %

study on predominance of keratinophilic flora in soil of Jaipur, India", Journal of King Saud University - Science, 2020

Publication

45

Mara L.F. Bittencourt, Paulo R. Ribeiro, Rosana L.P. Franco, Henk W.M. Hilhorst, Renato D. de Castro, Luzimar G. Fernandez. "Metabolite profiling, antioxidant and antibacterial activities of Brazilian propolis: Use of correlation and multivariate analyses to identify potential bioactive compounds", Food Research International, 2015

Publication

<1 %

46

Saranya Jayaram, Soma Biswas, Indhu Philip, Mridul Umesh, Suma Sarojini. "Differential Laccase Production among Diverse Fungal Endophytes in Aquatic Plants of Hulimavu Lake in Bangalore, India", Journal of Pure and Applied Microbiology, 2023

Publication

<1 %

47

ojs.umrah.ac.id

Internet Source

<1 %

48

"Endophytes", Springer Science and Business Media LLC, 2021

Publication

<1 %

Exclude quotes On

Exclude matches Off

Exclude bibliography On