

13

The Rate of Metabolism and Performance of Broiler Who Gets Skin Treatment Rations Containing Cassava Flour and Injection Solution Thyroid Gland

Yuanita Windusari, Meisji Liana Sari, Sofia Sandi S., and Agus Budiyanasyah

Abstract—The aims of this research was to find out the effect of the thyroid gland flour given by injection and skin cassava flour given in the diet on metabolic rate and performance of broiler chickens. Feeds were arranged with protein content (18%) and metabolism energy (2900 kcal/kg) in accordance with recommendations of Nutrition Research Council (NRC). As many as 35 were one week old chicks used in this study. Completely random design was used in this research, with 5 treatments and 4 replicates. Each of these repeats was placed 3 ducks and kept for 3 months. The treatment consisted of P1 (control diet), P2 (70% basal diet + 30% cassava peel without steam), P3 (100% basal ration with treatment of the thyroid gland injection of starch solution as much as 10 mg per day in subcutaneous), P4 (100% basal ration with treatment of the thyroid gland flour solution injection of 20 mg per day in subcutaneous). The results showed that metabolic activity of broilers increased concurrently with the increasing age characterized by increased oxygen consumption, body heat production and basal metabolic rate.

Index Terms—Countaining casava flour, solution tyroid gland, chicken.


Yuanita Windusari is with the Department of Biology, Faculty of Mathematics and Natural Sciences, Sriwijaya University, South Sumatera, Indonesia (e-mail: ywindusari@yahoo.com).

Meisji Liana Sari and Sofia Sandi S. are with the Department of Animal Science, Faculty of Agriculture, Sriwijaya University, South Sumatera, Indonesia (e-mail: meisji@yahoo.com).

Agus Budiyanasyah is with Faculty of Animal Science, Jambi University, Indonesia (e-mail: budiyanasyah_agus@yahoo.com).

[PDF]

Cite: Yuanita Windusari, Meisji Liana Sari, Sofia Sandi S., and Agus Budiyanasyah, "The Rate of Metabolism and Performance of Broiler Who Gets Skin Treatment Rations Containing Cassava Flour and Injection Solution Thyroid Gland," *International Journal of Chemical Engineering and Applications* vol. 5, no. 4 pp. 287-290, 2014.



The Rate of Metabolism and Performance of Broiler Who Gets Skin Treatment Rations Containing Cassava Flour and Injection Solution Thyroid Gland

Yuanita Windusari, Meisji Liana Sari, Sofia Sandi S., and Agus Budiaryansyah

Abstract—The aims of this research was to find out the effect of the thyroid gland flour given by injection and skin cassava flour given in the diet on metabolic rate and performance of broiler chickens. Feeds were arranged with protein content (18%) and metabolism energy (2900 kcal/kg) in accordance with recommendations of Nutrition Research Council (NRC). As many as 35 were one week old chicks used in this study. Completely random design was used in this research, with 5 treatments and 4 replicates. Each of these repeats was placed 3 ducks and kept for 3 months. The treatment consisted of P1 (control diet), P2 (70% basal diet + 30% cassava peel without steam), P3 (100% basal ration with treatment of the thyroid gland injection of starch solution as much as 10 mg per day in subcutaneous), P4 (100% basal ration with treatment of the thyroid gland flour solution injection of 20 mg per day in subcutaneous). The results showed that metabolic activity of broilers increased concurrently with the increasing age characterized by increased oxygen consumption, body heat production and basal metabolic rate.

Index Terms—Countaining casava flour, solution thyroïd gland, chicken.

I. INTRODUCTION

Endocrine system in the body serves to unify and harmonize the various processes and systems in the body, thus allowing the changes according to environmental conditions. This integration by hormone produced by certain glands in the body and transported to target cells with circulation. Thyroid hormone is one of the endocrine systems in the body. In the active form of thyroid hormone is as tri-iodothyronine (T3) and thyroxine (T4) has an important role in the physiology of the body, such as in growth, reproduction, metabolism and other foods. Thyroid hormone stimulates protein synthesis in tissues that synthesize RNA at the core level [1].

Therefore it can be said to affect thyroid hormone metabolism controls the general. Synthetic thyroid hormone is a substance that supposedly could provide benefit when given to chickens. This is because the nature of thyroid hormone like noted above, so that could be expected to increase the rate of metabolism and performance of broiler

chickens, as well as synthetic thyroid hormone can increase the activity of the thyroid gland to produce thyroid hormones in broiler chickens. However, data on the number or doses that can be given to broiler chickens is not known for sure. Therefore, the research about the right amount or dose of a synthetic thyroid hormone use in broiler chickens needs to be known.

On the other hand the use of certain materials that are goitrogenik can inhibit mineral absorption of iodine by the thyroid gland, resulting in the formation of hormones from the thyroid gland is inhibited. It is known that the formation of the mineral iodine is a component of thyroid hormones. Raw foods are goitrogenic among others are derived from cassava (including bark, tubers and cassava leaves). This is due to the material containing cyanide (HCN). Thiocyanate, the results of cyanide ion detoxification of HCN in the heart will inhibit absorption (uptake) in the blood iodine by the thyroid gland.

This study was conducted to determine the effect of the thyroid gland flour given by injection and skin cassava flour given in the diet on metabolic rate and performance of broiler chickens.

II. MATERIALS AND METHODS

A. Materials Research

A total of 35 were one week old chicks used in this study. Another material used is the thyroid gland flour dissolved in distilled water with a concentration of 10 mg and 20 mg per 0.5 cc for one injection every day and calculated based on the need for injection for 7 days. Ration ration used was compiled based on the needs of the chicks initial period under the recommendations of the Nutrition Research Council 1994 were compiled using materials yellow corn, fish meal, coconut meal, fine bran, soybean meal, bone meal and premix and feed ingredients cassava peel which is steamed and dried with the treatment without steamed. Weighing both the chicks and feed treatments using Ohaus digital scale brands with a level of accuracy 0.0001 grams.

Use measure oxygen consumption Calorimeter chamber made of jar equipped with a rubber hose and a syringe to measure the volume of incoming air, pipe manometer scale U-shaped letters made of glass and NaOH solution as the absorbent. To measure oxygen consumption when used timekeeping device (timer).

B. Research Methods

A total of 10 chicks aged one week to judge the rate of metabolism by measuring oxygen consumption and heat

Manuscript received July 8, 2013; revised December 20, 2013.

Yuanita Windusari is with the Department of Biology, Faculty of Mathematics and Natural Sciences, Sriwijaya University, South Sumatera, Indonesia (e-mail: ywindusari@yahoo.com).

Meisji Liana Sari and Sofia Sandi S. are with the Department of Animal Science, Faculty of Agriculture, Sriwijaya University, South Sumatera, Indonesia (e-mail: meisji@yahoo.com).

Agus Budiaryansyah is with Faculty of Animal Science, Jambi University, Indonesia (e-mail: budiaryansyah_agus@yahoo.com).

production of the body by using the system to determine the rate of metabolism beginning of broiler chickens. A total of 25 chicks then maintained for one week to be given five (5) kinds of treatment. Each treatment performed in 5 (five) chicks. These five kinds of treatment are:

- P1 : Chickens fed control diet (100% basal ration)
 P2 : Chickens fed rations containing dried cassava peel without steamed as much as 30% (70% basal diet + 30% cassava peel without steam) Chickens fed rations containing dried cassava peel without steamed as much as 30% (70% basal diet + 30% cassava peel without steam)
 P3 : Chickens fed rations containing skin-dried cassava steamed prior treatment by 30% (70% + 30% basal ration skin steamed cassava)
 P4 : Chickens fed control diet (100% basal ration) with treatment of the thyroid gland injection of starch solution as much as 10 mg per day in sub-cutaneous
 P5 : Chickens fed control diet (100% basal ration) with treatment of the thyroid gland flour solution injection of 20 mg per day in sub-cutaneous;

Basal ration composition and nutrient content of the diets are presented in Table I.

TABLE I: THE COMPOSITION OF FEED INGREDIENTS MAKING UP THE BASAL RATION

Composition of feed ingredients	(%)
Fine bran	8.5
Yellow corn	54.0
Soybean meal	15.5
Fish meal	13.4
Coconut cake	6.5
Vegetable oil	2.0
Top mix	0.1
Total	100
Metabolizable energy (kcal / kg)	3001
Crude protein (%)	21.2
Crude fat (%)	11.13
Crude fiber (%)	3.56
Calcium (%)	1.25

Before being given treatment, chicks were weighed beforehand to determine initial body weight. Weighing of body weight every day in the morning. Rationing and performed daily drinking water ad libitum. Daily feed intake was measured by weighing before being given rations and rations remaining after 24 hours of administration. Feed intake is calculated based on the difference given rations and rations rest. Body weight gain was calculated based on daily body weight gain. After one week, the chicks back measurements performed by measuring the metabolic rate of oxygen consumption and heat production of the body.

Measurement of oxygen consumption and Heat Production Rate Metabolism Oxygen consumption measurements conducted using a closed-circuit system using

a Calorimeter chamber in the form of jars and equipment. Put chicks into a jar, then sealed and airtight. Before the chicks put in a jar, tube manometer scale has been filled with water as a control tool that the volume of oxygen consumed is entered already. To absorb the carbon dioxide released by the oxidation of the rest of the chicks, at the bottom of the jar is placed in a bottle NaOH grain size of 10 cc. A number of oxygen volume syringe is inserted through a predetermined volume (initial measurement is 15 cc and 20 cc). One of the air intake causing the water level in the pipe manometer rose motivated by the pressure of air entering through the pipe. The beginning of time through the syringe air intake recorded, and after the water level in the capillary equal end time is recorded as the volume of oxygen consumed. During the measurement of oxygen consumption, air temperature (T1) and pressure (P1) is recorded, as well as the existing state of chicks observed in jars. Oxygen consumption (V1) standardized at 760 mm Hg pressure (P2) at a temperature of 273 °K (T2). Oxygen consumption (V2) is calculated with the formula $P1V1/T1 = P2V2/T2$. Heat production was calculated based on oxygen consumption multiplied by the equivalent calorific value. On chickens that are given feed with a mixture of carbohydrates, fat and protein, equivalent calorific value standardized to the value of 4.825 Kcal / liter of oxygen consumed. Chicks' basal metabolic rate is calculated based on the body's production of heat per gram of metabolic body weight per day (0.75 g BB / day).

Observed Variables Parameters observed in this penlitian include oxygen consumption (liters / head / day), body heat production (kcal/head day), basal metabolic rate (kcal / 0.75 g BB / day), feed intake (g / day), body weight gain (g / day) and feed conversion.

Data processing Research using completely randomized design (CRD) with 5 treatments and 5 replications. Data were processed using variance used in accordance with the design by using the SAS program. To test for differences between treatment used Duncan's multiple range test [2] and also using the SAS program.

III. RESULTS AND DISCUSSION

A. Effect of Treatment of Metabolic Rate

Treatment effect on oxygen consumption, body heat production and basal metabolic rate are presented in Table II. Results of analysis of variance showed that the treatment significantly ($P < 0.05$) on oxygen consumption and heat production of broiler chicken's body, but did not significantly affect the basal metabolic rate. Based on the results of Duncan's multiple range test (Table II) showed that the oxygen consumption and heat production in chickens treated with diet containing cassava peel as much as 30% (either steamed or without steamed) was significantly higher than that of chickens that received control treatment, but the consumption of oxygen and production of body heat from the chickens given the treatment of thyroid powder injection solution as much as 10 mg / head / day or 20 mg / head / day was not significantly different than the control treatment. This means cassava significantly increased oxygen consumption and heat

activity, some reports get mixed results, no one can improve body weight gain [5]-[7] but the reality weight gain not different from controls. Allegedly required a long time to be giving thyroid powder can provide metabolic stimulation.

The results get all the feed efficiency of the treatment was not significantly different from the control treatment, but demonstrated the highest efficiency in the treatment of thyroid flour administration of 20 mg / head / day. Allegedly giving thyroid flour as much as 20 mg / head / day can increase protein synthesis so that a more efficient use of feed. It is known that thyroid hormone stimulates the formation of protein networks, namely by increasing protein synthesis through increased RNA synthesis at the level of protein in the nucleus and cytoplasm [8]. Ration efficiency of the worst occurred in chickens treated diet containing cassava skin without steamed, and there kecendungan that the implementation of this diet containing cassava peel as much as 30% ration becomes inefficient. Allegedly there is a toxic effect of HCN that inhibit growth.

IV. CONCLUSION

Provision of treatment rations containing cassava peel as much as 30% and as much thyroid powder injection 10 mg / head / day and 20 mg / head / day increase oxygen consumption and heat production of the body, while the rate of basal metabolism is not influenced treatment.

Metabolic activity of broilers increased concurrently with the increasing age characterized by increased oxygen consumption, body heat production and basal metabolic rate.

Giving skin treatment diet containing 30% cassava flour and thyroid injection as much as 10 mg / head / day tended to increase feed intake and feed efficiency decrease, whereas thyroid powder injection of 20 mg / head / day will increase feed efficiency.

REFERENCES

- [1] H. A. Harper, V. W. Rodwell, and P. A. Mayes, *Review of Physiological Chemistry (Biochemistry)*, Translation by M. Muliawan, 17th ed. CV.EGC. Book Medical Publishers, Jakarta, 1979, pp. 376.
- [2] L. R. Steel and J. H. Torrie, *Principle and Procedure Statistics*, McGraw-Hill Book Company Inc., New York, 1980.
- [3] P. McDonald, R. A. Edward, and J. F. D. Greenhalgh, *Animal Nutrition*, 2nd ed, The English Language Book Society and Longman, London, 1977, pp. 479.
- [4] C. Devendra, "Source cassava as feed for ruminants," in *Cassava as Animal Feed*, B. Nestle and M. Graham Eds. IDRC, Canada, 1977, pp. 107-119.
- [5] A. Atomarsono, "Role of Thyroid Hormone in Relation to the Use of Protein Performances Broiler," dissertation, Graduate School of Bogor Agriculture Institute, Bogor, West Java, Indonesia, 1989, pp. 125-220.
- [6] J. D. May, "Effect of dietary thyroid hormone on growth and feed efficiency on broiler," *Poult. Sci.* vol. 59, pp. 882-892, 1980.
- [7] C. E. Leung, J. E. Taylor, and V. Inderstin, "Effect of thyroid hormone on growth, feed efficiency and plasma T3 and T4 in 4 Week Old Cockerel," *Poult. Sci.*, vol. 61, pp. 1500-1501, 1982.
- [8] H. Astier, *Thyroid Gland in Birds: Structure and Function*. In: *Avian Endocrinology*. A. Apple and M. H. Stetson, Eds. Academic Press, New York, 1980, pp. 167-189.



Yuanita Windusari was born on 14 September 1969 at Tanjungkarang, Province of Lampung, Indonesia, who received her bachelor's degree (Bachelor of Science) in the Department of Biology, Faculty of Mathematics and Natural Sciences, University of Sriwijaya in Palembang, South Sumatra, Indonesia in 1993. In 2002, she obtained postgraduate degree (M.Sc.) from Institute Technology of Bandung in Bandung, Indonesia in the field of Developmental Biology and Toxicology.

In 2009 she followed the Sandwich-like programme include studies of wetlands in the UNESCO-IHE, Delft, Netherland. In 2012, get the title of Doctor from Environmental Science Programme, University of Sriwijaya in Palembang, South Sumatra, Indonesia for the general study is wetlands, while the dissertation study on the patterns of vegetation succession on land of tailings. Until now she has been active as a lecturer, researcher, and several university activities. She is also active as a researcher at "Research Center of Sub-optimal land Development of Sriwijaya University (PUR - PLSO)". Many studies, symposium, seminar, training and workshops at the local, national and international have been followed. Was awarded as the best presentation and in 2013 was awarded the best presentation at the International Symposium of Lowland and Technology in Nusadua, Bali, Indonesia, organized the International Association of Lowland and Technology based in Saga University Japan. Recent research related to carbon stocks in forest vegetation in the city of Palembang, South Sumatra, Indonesia and the quality of the river waters. The results of these studies generally have been published in several journals nationally accredited.



Meisji Liana Sari was born at Palembang on 27 May 1970. She has got her bachelor degree from Jambi university, South Sumatra Indonesia in 1994. In 2002, she has got her master degree from Bogor Agriculture Institute (IPB), West Java Indonesia. She got her Doctoral degree also from Bogor Agriculture Institute (IPB), West Java Indonesia in 2012. She is a lecturer at Nutrition and Animal Feed Department, Sriwijaya University (UNSRI). She is also active as a researcher at "Research Center of Sub-optimal land Development of Sriwijaya University (PUR - PLSO)".