

# Length-weight Relationship and Condition Factor of *Trichopodus trichopterus* from Ogan Ilir Peat Swamp, South Sumatra, Indonesia

*by* Muslim Muslim

---


**Submission date:** 15-Jun-2024 05:58PM (UTC+0700)

**Submission ID:** 2402912134

**File name:** hopterus\_from\_Ogan\_Iilir\_Peat\_Swamp,\_South\_Sumatra,\_Indonesia.pdf (595.32K)

**Word count:** 3440

**Character count:** 18519



### 3

## Length-weight relationship and condition factor of *Trichopodus trichopterus* from Ogan Ilir peat swamp, South Sumatra, Indonesia

<sup>1</sup>Muslim Muslim, <sup>2</sup>Anita Karolina, <sup>2</sup>Reno Irawan, <sup>1</sup>Muhammad Rizky Ariansyah Putra

<sup>15</sup>  
<sup>1</sup> Aquaculture Study Program, Faculty of Agriculture, Universitas Sriwijaya, Indralaya, Indonesia; <sup>2</sup> Aquatic Resource Management Study Program, Faculty of Fisheries, Universitas Islam Ogan Komering Ilir Kayuagung, Indonesia. Corresponding author: M. Muslim, muslim\_bda@unsri.ac.id

**Abstract.** The study of the length-weight relationship and condition factor is a fundamental component of fisheries management. *Trichopodus trichopterus* (Pallas, 1770) is a native species of Indonesia, used as food and aquarium fish. The purpose of this study was to analyze the length-weight relationship and condition factor of *T. trichopterus* from Ogan Ilir peat swamp, South Sumatra, Indonesia. A total of 85 samples were used in this study. The results showed that the length-weight relationship of the *T. trichopterus*, the predictive model of the weight of the fish from the length of the fish is in the exponential form with the equation  $y=0.0207x^{2.8607}$  ( $R^2=0.84$ ,  $p<0.01$ ), with an a coefficient of 0.0207 and a b constant of 2.86. The current study hopes to provide the first baseline data about the length-weight relationship and condition factor of *T. trichopterus* from the Ogan Ilir peat swamp. The data obtained are very useful for the sustainable management of this resource.

**Key Words:** anabantiformes, fisheries management, floodplain, labyrinth fish, osphronemidae.

**Introduction.** The three-spot gourami *Trichopodus trichopterus* (Pallas, 1770) is a species of fish native to southeastern Asia but also introduced elsewhere (Low 2019). These fish live in swamps, marshes, lakes, canals, and lowland wetlands. *T. trichopterus* is an omnivore and requires both algae-based and meaty foods. An algae-based flake food, along with freeze-dried bloodworms, *Tubifex* sp. worms, and brine shrimp, provides these fish with the proper nutrition. Live foods such as mosquito larvae and *Daphnia* sp. are also beneficial. In Indonesia, this fish can be found on the islands of Sumatra and Kalimantan. The local people use it as food and ornamental fish.

The fish generally increase in length and weight during development. Factors that affect fish growth are internal and external. Internal factors include health status, life cycle, sexes, and parental inheritance, while external factors include physical, chemical, and biological parameters of the habitat. The length-weight relationship is crucial for ecological assessments and monitoring (Orlov & Binohlan 2009), population stock assessment studies (Augustina et al 2022), determining the growth type of fish (Ricker 1975; Erzini 1994), several biological aspects of the species (Freitas et al 2017), determining the life history (Ferdaushy & Alam 2015), determining the growth and body condition (Zuchi et al 2020), describing geographic and seasonal conditions (Lima et al 2021).

Some previous studies were conducted on the length-weight relationship of Indonesian fish species: *Rasbora tawarensis* and *Poropuntius tawarensis* (Muchlisin et al 2010), *Tor tambra* (Muchlisin et al 2015), *Crassostrea virginica*, *Crassostrea gigas*, *Crassostrea iridescens*, *Crassostrea angulata*, and *Ostrea edulis* (Octavina et al 2015), *Hemibagrus wyckii* (Aryani et al 2016), *Decapterus macrosoma* (Pattikawa et al 2017), *Barbonymus gonionotus* (Batubara et al 2019), *Channa striata* (Ahmadi 2018), *Anabas testudineus* (Nobe et al 2019), *Oryzias matanensis* (Rinandha et al 2020; Nugroho et al 2021), *Mystacoleucus padangensis* (Nasution et al 2021), *Pristolepis grootii* (Muslim et al

2022) and others. <sup>16</sup> This study aims to determine the length-weight relationships and condition factor of *T. trichopterus* from Ogan Ilir peat swamp, South Sumatra, Indonesia.

**Material and Method.** This study was conducted in Ogan Ilir peat swamp, South Sumatra, Indonesia, from September to December 2022 (rainy season). The samples were collected from two sampling stations: station 1 (3°12'21.16"S-104°38'50.25"E) and station 2 (3°12'18.92"S-104°38'49.758"E) (Figure 1).




Figure 1. Map of the Ogan Ilir peat swamp, South Sumatra, Indonesia; the red dots mark the location of sampling sites.

<sup>6</sup> The fish were collected using traditional fishing gear, namely traps and gillnets. Sampling was carried out twice in one month. *T. trichopterus* samples were transported to the Batanghari Sembilan Fish Breeding Unit, Ogan Ilir Regency, under live conditions for measurement. The measured samples were reared as ornamental aquarium fish. A total of 85 *T. trichopterus* were collected at the sampling stations. The <sup>12</sup> total length (TL, cm) and body weight (BW, g) were determined for each individual. <sup>13</sup> The TL of the fish was measured from the tip of the mouth to the tip of the caudal fin. TL was measured using a caliper (0.1 cm accuracy), and the BW of the sample fish was measured using a digital scale (0.01 g accuracy).

The analysis of the length-weight relationship used a linear allometric model for the calculation of parameters 'a' and 'b'. The calculation was done by measuring changes in weight and length. The bias correction that exists in the change in the average weight of the logarithmic unit is used as a weight prediction calculation on the length parameter according to the formula (De Robertis & Williams 2008):

$$BW = aTL^b$$

<sup>1</sup> Where: BW is the body weight of each fish (g); TL is the total length of each fish (cm); a, b are constants. <sup>2</sup>

The fish growth patterns are determined based on the value of b. When it equals 3, then the growth pattern is isometric <sup>2</sup> or weight gain is equivalent to the length of the fish, and when the value of b not equal to 3, then the growth pattern is allometric. These growth patterns are divided into two categories, namely positive and negative allometric. Whenever the value of b is less than 3, it is called positive allometric (the increase in length is less than the weight gain) (Kirankaya et al 2014). The significance of the length-weight relationship of the specimens was tested among stations by a t-test.

The relative weight (Wr) and condition factor coefficient (C) were used to evaluate the condition factor of each individual. The Wr was determined based on the equation (Rypel & Richter 2008):

$$Wr = \frac{BW}{Ws} \times 100$$

Where: Wr is the relative weight, BW is the body weight of each fish, and Ws is the standard weight predicted from the same sample because it is calculated from the combined length-weight regression through the distance between species.

The Fulton condition factor (K) was determined based on Okgerman (2005) with the following formula:

$$K = BW \times TL^{-3} \times 100$$

Where: K is the condition factor; BW is the body weight (g); TL is the total length (cm).

**Results and Discussion** The TL range was between 5.3 and 12.1 cm with an average of  $8.56 \pm 1.15$  cm, and the BW range was between 1.69 and 26.65 g with an average of  $10.21 \pm 4.28$  g (Figure 2). Eight length class intervals were obtained as the result of the TL frequency distribution analysis. The most frequent fish size classes are in the TL of 7–7.99 cm, 8–8.99 cm, and 9–9.99 cm classes. The least frequent fish size is in the 5–5.99 cm and 12–12.99 cm classes. *T. trichopterus* had a larger maximum TL compared to the TL of *T. trichopterus* in other studies (Table 1) and the same maximum TL as the fish reported by Cuadrado et al (2019) in Lake Esperanza, Agusan del Sur, Philippines, which was 12.1 cm. However, its size is smaller compared to *T. trichopterus* data recorded in FishBase, where the maximum length of *T. trichopterus* can reach 15 cm in natural waters (Froese & Pauly 2021).




Figure 2. Total length and weight of *Trichopodus trichopterus* (Pallas, 1770) from Ogan Ilir peat swamp, South Sumatra, Indonesia.

Length-weight relationships and condition factor of *Trichopodus trichopterus* (Pallas, 1770) in different regions

Location	Author	Length (cm)	a	b	R <sup>2</sup>	GP	K
Agusan Marsh	Jumawan & Seronay (2017)	4.9-11.5	0.029	3.04	0.96	A+	-
Martapura	Aminah & Ahmadi (2018)	3-11.1	0.0002	1.60	0.70	A-	1.88
Bendungan Lempake	Jusmaldi et al (2021)	3.16-10.35	0.0003	2.86	0.94	A-	1.05
Ciperet River	Wahyudewantoro et al (2021)	3.42-10.75	0.0006	2.40	0.99	A-	1.00
Muara Angke	Wahyudewantoro et al (2021)	3.54-10.33	0.0003	2.62	0.99	A-	1.00
Ogan Ilir, Peat Swamp	Current study	5.3-12.1	0.0207	2.86	0.84	A-	1.01

Note: a and b - constants; R<sup>2</sup> - coefficient of determination; GP - growth pattern; A+ - allometric positive; A- - allometric negative; K - condition factor.

20

The results of the analysis of the length-weight relationship are expressed in the regression equation  $BW=0.0207 \times TL^{2.8607}$  (Figure 3). The TL and BW of *T. trichopterus* show a positive association with a correlation coefficient (r) of 0.92 and a coefficient of determination (R<sup>2</sup>) of 0.84, indicating that TL influences 84% of the change in BW of *T. trichopterus*. The results of the t-test show that the regression coefficient (b) is significantly different from 3 (p>0.05). *T. trichopterus* has a b value of 2.86, meaning a negative allometric growth pattern (the increase in length is not followed by an increase in weight).




Figure 3. The length-weight relationship of *Trichopodus trichopterus* (Pallas, 1770) from Ogan Ilir peat swamp, South Sumatra, Indonesia.

In this study, the growth of *T. trichopterus* was negative allometric. Negative allometric *T. trichopterus* growth was also found in Martapura (Aminah & Ahmadi 2018), at the Lempake Dam (Jusmaldi et al 2021), in the Ciperet River, and in Muara Angke (Wahyudewantoro et al 2021). According to Mazumder et al (2016), factors that may cause allometric negative growth patterns in *T. trichopterus* are habitat conditions that are not suitable for growth. Mon et al (2020) explained that fish growth with  $b < 3$  indicates an aquatic environment that is not good for fish growth. Meanwhile,  $b > 3$  (positive allometric) indicates good

environmental factors. The cause of the difference in the value of  $b$  in fish is thought to be due to ecological differences such as temperature and food supply.

The  $K$  value is used to assess the health condition of fish (Ahmadi 2018; Basak & Hadiuzzaman 2019). The Fulton condition factor is generally used to assess the health of the fish, indicating the nutritional and physiological status of the fish (Muchlisin et al 2017; Oktaviani et al 2020). According to Fafioye & Ayodele (2018), a  $K$  value less than 1 indicates that the fish habitat is not fully suitable for fish health. The mean value of the condition factor of the *T. trichopterus* in the study was 1.01, which indicated the fish were in good condition.

**Conclusions.** The relationship between the length and weight of the *T. trichopterus* has a positive correlation, and the fish present a negative allometric growth pattern. The condition factor value of 1.01 indicated that the *T. trichopterus* population in the Ogan Ilir peat swamp was in good condition.

**Acknowledgements.** We would like to thank the Batanghari Sembilan fish breeding unit, Ogan Ilir. Thank you to Rais Pebri Sagita, Ripal Nur Hidayat, Ahmad Muslim Rosyidi, Aiverdaofi, Muhammad Aulia Hibatilla, Shafira Az-Zahra for their assistance.

**Conflict of Interest.** The authors declare that there is no conflict of interest.

## References

- Ahmadi A., 2018 The length-weight relationship and condition factor of the threatened snakehead (*Channa striata*) from Sungai Batang River, Indonesia. Polish Journal of Natural Sciences 33(4):607-623.
- Aminah S., Ahmadi A., 2018 Experimental fishing with led light traps for threespot gourami (*Trichogaster trichopterus*) in Martapura, Indonesia. International Journal of Fisheries and Aquatic Studies 6(1):37-42.
- Aryani N., Suharman I., Hasibuan S., 2016 Length-weight relationship and condition factor of the critically endangered fish of Geso, *Hemibagrus wyckii* (Bleeker, 1858) bagridae from Kampar Kanan River, Indonesia. Journal of Entomology and Zoology Studies 4(2):119-122.
- Augustina T. X., Sreeram M. P., Sreekumar K. M., Raju A. K., 2022 Length-weight relation and relative condition factor of six deep-sea fishes from the South-eastern Arabian Sea. Journal of the Marine Biological Association of India 64(2):118-122.
- Basak S. S., Hadiuzzaman M., 2019 Length-weight relationship, condition factor and relative condition factor of kalibaus fish *Labeo calbasu* (Hamilton, 1822) of Kaptai Lake, Rangamati, Bangladesh. International Journal of Fisheries and Aquatic Studies 7(5):231-235.
- Batubara A. S., Muchlisin Z. A., Efizon D., Elvyra R., Irham M., 2019 Length-weight relationships and condition factors of the naleh fish, *Barbonymus gonionotus* (Pisces, Cyprinidae) harvested from Nagan Raya Waters, Indonesia. Vestnik Zoologii 53(1):75-82.
- Cuadrado J. T., Lim D. S., Alcontin R. M. S., Calang J. L. L., Jumawan J. C., 2019 Species composition and length-weight relationship of twelve fish species in the two lakes of Esperanza, Agusan del Sur, Philippines. FishTaxa 4(1):1-8.
- De Robertis A., Williams K., 2008 Weight-length relationships in fisheries studies: the standard allometric model should be applied with caution. Transactions of the American Fisheries Society 137(3):707-719.
- Erzini K., 1994 An empirical study of variability in length-at-age of marine fishes. Journal of Applied Ichthyology 10(1):17-41.
- Fafioye O., Ayodele O., 2018 Length-weight relationship and condition factor of four commercial fish species of Oyan Lake, Nigeria. Examines in Marine Biology and Oceanography 2(2):227-230.
- Ferdaushy M. H., Alam M. M., 2015 Length-length and length-weight relationships and condition factor of nine freshwater fish species of Nageshwari, Bangladesh. International Journal of Aquatic Biology 3(3):149-154.

- Freitas T. M. D. S., Souza J. B. D. S., Prudente B. D. S., Montag L. F. D. A., 2017 Length-weight relationship in ten fish species from the Nhamundá River, the Amazon Basin, Brazil. *Acta Amazonica* 47(1):75-78.
- Jumawan J. C., Seronay R. A., 2017 Length-weight relationships of fishes in eight floodplain lakes of Agusan Marsh, Philippines. *Philippine Journal of Science*. 146(1):95-99.
- Jusmaldi J., Dianingrum A. R., Hariani N., 2021 [The growth pattern and condition factors of three spot gourami *Trichopodus trichopterus* (Pallas, 1770) from the Lempake Dam, East Kalimantan]. *Indonesian Journal of Ichthyology* 21(3):215-233. [In Indonesian]
- Kırankaya Ş. G., Ekmekçi F. G., Yalçın-Özdilek Ş., Yoğurtçuoğlu B., Gençoğlu L., 2014 Condition, length-weight and length-length relationships for five fish species from Hirfanlı Reservoir, Turkey. *Journal of Fisheries Sciences* 8(3):208-213.
- Lima J. S., da Costa I. D., Zalmon I. R., 2021 Length-weight relationship of fish species captured around an artificial offshore reef (northern Rio de Janeiro, Brazil). *Journal of Applied Ichthyology* 37(2):337-341.
- Mazumder S. K., Das S. K., Bakar Y., Ghaffar M. A., 2016 Effects of temperature and diet on length-weight relationship and condition factor of the juvenile Malabar blood snapper (*Lutjanus malabaricus* Bloch & Schneider, 1801). *Journal of Zhejiang University Science B* 17(8):580-590.
- Mon E. E., Swe T., Zin P. P., Dwe K. L., 2020 Length-weight relationship, condition factor and sex ratio of tade mullet (*Liza tade* Forsskal, 1775) from Mawlamyine, Mon state, Myanmar. *Journal of Aquaculture and Marine Biology* 9(4):107-112.
- Muchlisin Z. A., Batubara A. S., Azizah M. N. S., Adlim M., Hendri A., Fadli N., Muhammadar A. A., Sugianto S., 2015 Feeding habit and length weight relationship of keureling fish, *Tor tambra* Valenciennes, 1842 (Cyprinidae) from the western region of Aceh Province, Indonesia. *Biodiversitas* 16:89-94.
- Muchlisin Z. A., Fransiska V., Muhammadar A. A., Fauzi M., Batubara A. S., 2017 Length-weight relationships and condition factors of the three dominant species of marine fishes caught by traditional beach trawl in Ulelhee Bay, Banda Aceh City, Indonesia. *Croatian Journal of Fisheries* 75(3):104-112.
- Muchlisin Z. A., Musman M., Azizah M. N. S., 2010 Length-weight relationships and condition factors of two threatened fishes, *Rasbora tawarensis* and *Poropuntius tawarensis*, endemic to Lake Laut Tawar, Aceh Province, Indonesia. *Journal of Applied Ichthyology* 26(6):949-953.
- Muslim M., Wardani W., Sahusilawane H., Oktarina S., Rifa'i R., Heltonika B., 2022 Length-weight relationship and environmental parameters of Indonesian leaf fish (*Pristolepis grootii*, Bleeker 1852) in Kelekar River, South Sumatera, Indonesia. *International Journal of Progressive Sciences and Technologies* 31(1):110-117.
- Nasution S. H., Akhdiana I., Syawal M. S., Nurhidayat A., 2021 Length-weight relationship and condition factor of endemic fish bilih (*Mystacoleucus padangensis* Blkr) in Lake Singkarak, West Sumatra, Indonesia. *E3S Web of Conferences* 322:01036.
- Ndobe S., Masyahoro A., Serdiati N., Moore A. M., 2019 Meristic characters and length-weight relation of climbing perch (*Anabas testudineus*) from wetlands in Sigi District, Central Sulawesi, Indonesia. *IOP Conference Series: Earth and Environmental Science* 370:012001.
- Nugroho R. A., Florentino A. P., Lariman L., Aryani R., Rudianto R., Kusneti M., 2021 [Length-weight relationship and relative condition factors of five fish species in the Suwi Muara Ancalong River, East Kutai]. *Biota* 6(2):64-70. [In Indonesian].
- Octavina C., Yulianda F., Krisanti M., Muchlisin Z. A., 2015 Length-weight relationship of Ostreidae in the Kuala Gigieng estuary, Aceh Besar District, Indonesia. *AAFL Bioflux* 8(5):817-823.
- Okgerman H., 2005 Seasonal variations in the length-weight relationship and condition factor of rudd (*Scardinius erythrophthalmus* L.) in Sapanca Lake. *International Journal of Zoological Research* 1:6-10.
- Oktaviani D., Prianto E., Nugroho D., 2020 Length-weight, maturity, and condition factor of torpedo scads (*Megalaspis cordyla*, Linnaeus 1758) in the Java Sea, Indonesia. *Biodiversitas* 21(4):1527-1534.

- Orlov A., Binohlan C., 2009 Length–weight relationships of deep-sea fishes from the western Bering Sea. *Journal of Applied Ichthyology* 25(2):223-227.
- Pattikawa J. A., Tetelepta J. M., Ongkers O. T., Uneputty P. A., Lewerissa H., 2017 Size distribution, length-weight relationship and age group of *Decapterus macrosoma* in eastern waters of Ambon Island, Indonesia. *AAFL Bioflux* 10(4):969-976.
- Ricker W. E., 1975 Computation and interpretation of biological statistics of fish populations. *Bulletin of the Fisheries Research Board of Canada* 191:1-382.
- Rinandha A., Omar S. B. A., Tresnati J., Yanuarita D., Umar T., 2020 Length-weight relationship and condition factors of Matano medaka (*Oryzias matanensis* Aurich, 1935) in Towuti Lake, South Sulawesi, Indonesia. *ACCL Bioflux* 13(4):1946-1954.
- Rypel A. L., Richter T. J., 2008 Empirical percentile standard weight equation for the blacktail redhorse. *North American Journal of Fisheries Management* 28(6):1843-1846.
- Wahyudewantoro G., Haryono H., Sulistiono S., Dina R., 2021 Growth pattern, condition factor and reproductive aspects of three spot gourami *Trichopodus trichopterus* (Pallas, 1770) in mangrove waters of Muara Angke Jakarta and Ciperet Cilacap, Indonesia. *IOP Conference Series: Earth and Environmental Science* 744:012095.
- Zuchi N., Röpke C., Shibuya A., Farago T., Carmona M., Zuanon J., Amadio S., 2020 Length-weight relationship of fish species from Central Amazon floodplain. *Journal of Applied Ichthyology* 36(6):837–841.
- \*\*\* Froese R., Pauly D., 2021 FishBase. Available at: [www.fishbase.org](http://www.fishbase.org)
- \*\*\* Low B. W., 2019 *Trichopodus trichopterus*. The IUCN Red List of Threatened Species 2019: e.T187981A89805622. Available at: <https://dx.doi.org/10.2305/IUCN.UK.2019-2.RLTS.T187981A89805622.en>

Received: 12 April 2023. Accepted: 29 June 2023. Published online: 19 January 2024.

Authors:

Muslim Muslim, Aquaculture Study Program, Faculty of Agriculture, Universitas Sriwijaya, Jl. Palembang - Prabumulih Km. 32, 30662 Indralaya, South Sumatra, Indonesia, e-mail: muslim\_bda@unsri.ac.id  
 Anita Karolina, Aquatic Resource Management, Faculty of Fisheries, Universitas Islam Ogan Komering Ilir Kayuagung, Jl. Letnan Mughtar Saleh No. 1, 30611 Kayuagung, Ogan Komering Ilir Regency, South Sumatra, Indonesia, e-mail: karolina.uniski@gmail.com  
 Reno Irawan, Aquatic Resource Management, Faculty of Fisheries, Universitas Islam Ogan Komering Ilir Kayuagung, Jl. Letnan Mughtar Saleh No. 1, 30611 Kayuagung, Ogan Komering Ilir Regency, South Sumatra, Indonesia, e-mail: renoirawanpsdp@gmail.com  
 Muhammad Rizky Ariansyah Putra, Aquaculture Study Program, Faculty of Agriculture, Universitas Sriwijaya, Jl. Palembang - Prabumulih Km. 32, 30662 Indralaya, South Sumatra, Indonesia, e-mail: mrizkiariansyahputra@gmail.com

This is an open-access article distributed under the terms of the Creative Commons Attribution License, which permits unrestricted use, distribution and reproduction in any medium, provided the original author and source are credited.

How to cite this article:

Muslim M., Karolina A., Irawan R., Putra M. R. A., 2024 Length-weight relationship and condition factor of *Trichopodus trichopterus* from Ogan Ilir peat swamp, South Sumatra, Indonesia. *AAFL Bioflux* 17(1):91-97.



# Length-weight Relationship and Condition Factor of Trichopodus trichopterus from Ogan Ilir Peat Swamp, South Sumatra, Indonesia

---

## ORIGINALITY REPORT

---

**20%**  
SIMILARITY INDEX

%  
INTERNET SOURCES

**20%**  
PUBLICATIONS

%  
STUDENT PAPERS

---

## PRIMARY SOURCES

---

- 1** A W Perdana, A S Batubara, F M Nur. "LWRs, condition factors and isopod parasites infection of the sumbo fish Selar crumenophthalmus (Pisces: Carangidae) in Lampulo fishing port, Banda Aceh, Aceh Province, Indonesia", IOP Conference Series: Earth and Environmental Science, 2019  
Publication **2%**
- 2** N Auliyah. "Lenght-weight relationship and condition factor of huluu fish (Giuris margaritacea) in Limboto Lake", IOP Conference Series: Materials Science and Engineering, 2019  
Publication **2%**
- 3** Zainal A. Muchlisin, Vera Fransiska, Abdullah A. Muhammadar, Muhammad Fauzi, Agung S. Batubara. "Length-weight relationships and condition factors of the three dominant species of marine fishes caught by traditional **2%**

beach trawl in Ulelhee Bay, Banda Aceh City, Indonesia", Croatian Journal of Fisheries, 2017

Publication

---

4

D S Said, N Mayasari. "The development of adapted marmorated medaka *Oryzias marmoratus* (Aurich, 1935) at ex-situ habitat", IOP Conference Series: Earth and Environmental Science, 2021

Publication

---

1 %

5

Md. Arifur Rahman, Md. Rased Khan Manon, Md. Rahamat Ullah, Asraful Alam et al. "Biometric indices, growth pattern, and physiological status of captive-reared indigenous Yellowtail brood catfish, *Pangasius pangasius* (Hamilton, 1822)", Environmental Science and Pollution Research, 2023

Publication

---

1 %

6

Syafrialdi ., Dahelmi ., Dewi Imelda Roe, Hafrijal Syandri. "Length-Weight Relationship and Condition Factor of Two-Spot Catfish (*Mystus nigriceps* [Valenciennes, 1840]) (Pisces, Bagridae), from Kampar Kanan River and Kampar Kiri River in Indonesia", Pakistan Journal of Biological Sciences, 2020

Publication

---

1 %

7

Ayat N. Salman, Abdul-Razak M Mohamed. "Growth, Mortality and Yield-per-recruit of Nile Tilapia (*Oreochromis niloticus*) in

1 %

8

*Dagnew Mequanent, Minwyelet Mingist, Abebe Getahun, Wassie Anteneh, Benyam Hailu. "The newly built Ribb Reservoir fisheries, Tana Sub-basin, Ethiopia: New fishery establishment, diversity, production, challenges and management", Aquaculture, Fish and Fisheries, 2022*

*Publication*

---

1 %

9

*"Morphometric characteristic and growth patterns of climbing perch (anabas testudineus) from sungai batang river, Indonesi", International Journal of Hydrology, 2019*

*Publication*

---

1 %

10

*Manmatha Nath Sarker, Sana Ahmed, Shampa Mitra, Prosenjit Pramanick, Sufia Zaman, Abhijit Mitra. "Application of Condition Index to Evaluate Environmental Health of Fishes: A Case Study With Labeo Rohita", Bangladesh Journal of Zoology, 2023*

*Publication*

---

1 %

11

*HAUWAU SALELE ABUBAKAR, Armaya'u Hamisu Bichi, Akeem Babatunde Dauda. "SPECIES COMPOSITION, SIZE DISTRIBUTION, CONDITION FACTOR AND GROWTH PATTERN OF CICHLIDS FROM ZOBE RESERVOIR, DUTSIN-MA, KATSINA*

1 %

STATE, NIGERIA.", *FUDMA Journal of Agriculture and Agricultural Technology*, 2023

Publication

---

12

*Shishir Kumar Dey, Md Rayhan Hossain, Zoarder Faruque Ahmed, Kaniz Fatema, Sayeeda Sultana. "Length-length relationship: an index of growth pattern for lesser spine eel *Macragnathus Aculeatus* (Bloch, 1786) in Bangladesh", *Asian-Australasian Journal of Bioscience and Biotechnology*, 2016*

Publication

---

1 %

13

*Kousar Jan, Imtiaz Ahmed. " Comparative study of length-weight relationships and biological indices of Himalayan snow trout, , inhabiting two lotic water bodies in the Kashmir Valley ", *Fisheries & Aquatic Life*, 2021*

Publication

---

1 %

14

*M Dwirastina, Y C Ditya, Herlan. "Estimation of Fish Production Potential with Benthos Biomass Approach in Sumani and Ombilin River of Singkarak Lake West Sumatra", *IOP Conference Series: Earth and Environmental Science*, 2021*

Publication

---

1 %

15

*E Anggraini, T E Riyanti, C Irsan, H Hamidson, M Sefrila, A Kurnianingsih, S F Sitepu. "Insect pests in smallholding coconut plantation in Marga Sungsang Village, Banyuasin Regency, South*

1 %

*Sumatra, Indonesia", IOP Conference Series:  
Earth and Environmental Science, 2024*

*Publication*

---

16

*F Y Yalindua, A Faricha, P S Ibrahim, R Huwae.  
"Length-weight relationship of seven Apogonidae  
species in Lembeh Strait", IOP Conference Series:  
Earth and Environmental Science, 2022*

*Publication*

---

17

*George Isaac Nodza, Eze Tochukwu, Aramide  
Dolapo Igbari, Temitope Olabisi Onuminya,  
Oluwatoyin Temidayo Ogundipe. "Application of  
IUCN Red List Criteria for Regional Assessment of  
Some Northern Savanna Trees of Nigeria, West  
Africa", Research Square Platform LLC, 2024*

*Publication*

---

18

*Chinyama Makeche Mauris, Nhiwatiwa Tamuka,  
Chitondo Lufeyo, Kanyati Malvern et al.  
"Comparative study of growth rates, condition  
factors and natural mortality of Oreochromis  
niloticus fish from culture fisheries and capture  
fisheries at Lake Kariba, Zambia", International  
Journal of Fisheries and Aquaculture, 2023*

*Publication*

---

19

*L Kasmini, T A Barus, M A Sarong, M B Mulya. "  
Morphometric study of pacific oyster ( ) in the  
coastal area of Banda Aceh ", Journal of Physics:  
Conference Series, 2018*

*Publication*

---

1 %

1 %

1 %

1 %

20

*Pattikawa Jesaja Ajub, Mamesah Julieta Adriana Bertha, Tetelepta Johannes Marten Stephan, Natan Yuliana, Pietersz Janson Hans. "Biological aspects of roundscads (Decapterus spp.) inhabiting the waters of Southeast Maluku, Eastern Indonesia", Fisheries and Aquatic Sciences, 2023*

*Publication*

---

1%

21

*Yeasmin Akter, Md. Hafiz All Hosen, Md. Idris Miah, Zoarder Faruque Ahmed, Mousumi Sarker Chhanda, Sheik Istiak Md. Shahriar. "Impact of gonad weight on the length-weight relationships of river catfish (Clupisoma garua) in Bangladesh", The Egyptian Journal of Aquatic Research, 2019*

*Publication*

---

1%

*Exclude quotes*      *On*

*Exclude matches*      *< 1%*

*Exclude bibliography*      *On*