

1. Proofread document received (2023-09-10)
2. Cover letter submission to the journal "Baghdad Science Journal" (2023-10-12)
3. Submission (2023-10-19)
4. Correction round 1 (2024-01-20)
5. Correction round 2 (2024-01-22)
6. Correction round 3 (2024-01-23)
7. Primarily accept the submission (2024-01-23)
8. Final check on manuscripts (2024-03-12)
9. Acceptance letter (2024-05-02)
10. Published

## SEXUAL DIMORPHISM AND REPRODUCTIVE BIOLOGY OF BRONZE FEATHERBACK (*Notopterus notopterus*, Pallas 1769) FROM KELEKAR RIVER, OGAN ILIR, SOUTH SUMATRA

**Abstract:** *Sexual dimorphism and reproductive biology are fundamental aspects of fish breeding studies. The aim of this research was to analyze the sexual dimorphism and reproductive biology of *N. notopterus*. A total of 74 *N. notopterus* collected from the Kelekar River in Ogan Ilir Regency, consisting 38 males (TL: 18–23.6 cm; BW: 35.1–92.1 g) and 36 females (TL: 19.6–26.3 cm; BW: 49.4–133.8 g). Seventeen morphometric characters, three meristic characters, and five reproductive biological parameters were analyzed. The results showed the differences in the morphometric characteristics of male and female *N. notopterus* in the characters of DTBH, BH, AF-CFL, and AT. However, there was no difference in the meristic characters. The male gonadosomatic index ranged from 0.15 to 0.61%, and the female from 1.12 to 9.10%. The hepatosomatic index of males ranged from 0.19 to 1.38%, and that of females ranged from 0.15 to 1.23%, with a sex ratio of 1.056:1, fecundity of 1.007–3.901 eggs, ova diameters of 1.1–3.9 mm, and maturity levels of developed gonads to mature for males and mature to spawning for females.*

~~Sexual dimorphism and reproductive biology are fundamental aspects of fish breeding studies. The aim of this research was to analyze the sexual dimorphism and reproductive aspects of *N. notopterus*. The samples were collected from the Kelekar River in Ogan Ilir Regency. A total of 74 *N. notopterus* samples were used: 38 males (TL: 18–23.6 cm; BW: 35.1–92.1 g) and 36 females (TL: 19.6–26.3 cm; BW: 49.4–133.8 g). Seventeen morphometric characters, three meristic characters, and five reproductive biological parameters were analyzed. The results showed that there were differences in the morphometric characteristics of male and female *N. notopterus* in the characters of DTBH, BH, AF-CFL, and AT, but there was no difference in the meristic character. The male gonadosomatic index ranges from 0.15 to 0.61%, and the female from 1.12 to 9.10%. The hepatosomatic index of males ranged from 0.19 to 1.38%, and that of females ranged from 0.15 to 1.23%, with a sex ratio of 1.056:1, fecundity of 1.007–3.901 eggs, ova diameters of 1.1–3.9 mm, and maturity levels of developed gonads to mature for males and mature to spawning for females.~~

**Keywords:** *bronze featherback*; *knife fish*; reproductive biology; sexual dimorphism

### Introduction

Bronze featherback (*Notopterus notopterus*, Pallas 1769) is a freshwater fish species native to Indonesia. *N. notopterus* is one of the fish species of the Notopteridae family (Kottelat, 2013). The Notopteridae consist ~~of~~ approximately ten species in four genera: Chitala, Notopterus, Xenomystus, and Papyrocranus (Barby et al., 2018). The only recognized species of Notopterus at present is *N. notopterus* (Lavoué et al., 2020). The *N. notopterus* distribution in Asia includes Indonesia, India, Malaysia, Thailand, Bangladesh, Pakistan, and Vietnam (Barby et al., 2018; Hilton & Lavoué, 2018; Kottelat, 2013; Mustafa et al., 2014; Naeem et al., 2010; Srivastava, Singh, & Pandey, 2012). In Indonesia, this species ~~is~~

Formatted: Font: Italic

Formatted: Font: Italic

distributed in Java, Sumatra, and Kalimantan (Kottelat, 2013; Kottelat et al., 1996). *N. notopterus* was found in Kolong-Bendungan Simpung, Bangka Island (Gustomi et al., 2016), Ogan River, Kelekar River (tributary of Musi River, South Sumatra), waters in Bangka Island as well as Kota Bangun and Tanah Ulu (Mahakam River, East Kalimantan) (Wibowo et al., 2009), Sail River, Pekanbaru City, Riau Province (Purba et al., 2017), Sei Gesek Reservoir, Bintan, Riau Islands (Rapita et al., 2021), Kelekar River (Muslim & Syaifudin, 2022), and Lebung Karang fishery reservoir, Ogan Ilir (Muslim & Lestari, 2005).

Several studies on *N. notopterus* have been conducted, among others, on aspects of length-weight relationship, relative condition factor and fecundity (Kaur & Rawal, 2017; Kaushik, Sahu, & Nath, 2019; Parasmewaran & Sinha, 1966; Paul, Nath, & Dutta, 2022; Winn et al., 2021), genetic diversity (Wibowo et al., 2009), morphometrics and growth patterns (Purba et al., 2017), food and feeding habits (Rapita et al., 2021), reproductive biology (Gustomi et al., 2016), determination of condition factors, somatic condition, hepatosomatic index and gonadosomatic index (Sudarshan & Kulkarni, 2013), behavior and reproduction (Sukendi et al., 2020), potential biomarkers of oxidative stress (Mohanty & Samanta, 2016), embryo and larval development (Srivastava et al., 2012), relative condition factors (Achakzai et al., 2015; Kumar & Kiran, 2016), population dynamics (Mustafa et al., 2014), and relationship of ecological factors and commercial (Kyaw et al., 2020).

Fishery management mainly depends on knowledge of fish reproductive biology (Dinh & Tran, 2018; Parmaksiz et al., 2017). In adult fish, sexual dimorphism is a frequent and well-known phenomenon (Capasso et al., 2021). The majority of fishes are bisexual and have various sexually dimorphic characteristics in their secondary sexual characteristics (Chacko & Sekharan, 2022). Distinguishing characteristics for morphological qualities of many teleosts have been documented, including body color, ornamental patterns, breeding tubercles on the snout, pectoral fins and scales, fin hooks, body shape, and fin size and shape (Ambus & Moody, 2018; Garcia & Zuanon, 2019; Motomura & Babaran, 2020; Mainero et al., 2023; Uba, 2019; Ueng et al., 2022).

*N. notopterus* is one of the most valuable freshwater fish in South Sumatra, Indonesia. This fish utilized by the community in South Sumatra, especially for a daily side dish and a raw material for the culinary industries of pempek (fish cake) and kerupuk-kemplang (fish crackers). Therefore, this species has the potential for cultivation.

~~*N. notopterus* is one of the most valuable species of freshwater fish in South Sumatra, Indonesia. This fish is utilized by the community of South Sumatra as a daily side dish and also as a raw material for the culinary industries of pempek (fish cake) and kerupuk-kemplang (fish crackers). So this species has the potential to be cultivated~~ (Muslim et al., 2020).


There is relatively little known about sexual dimorphism and reproductive biology in *N. notopterus*. Such information is ~~very important as basic~~ crucial for initial data

Formatted: Font: Not Italic

~~infer~~ breeding of *N. notopterus*. The aim of the study was to analyze the sexual dimorphism and reproductive biology of *N. notopterus* from Sungai Kelekar, Ogan Ilir. The results of this study are ~~very useful~~ beneficial in the selection of *N. notopterus* broodstock candidates for breeding programs.

## Materials and Methods

~~The f~~ Fish collection was conducted in Kelekar River, Tanjung Baru Village, North Indralaya District, Ogan Ilir Regency, South Sumatra (3°14'11.8 "S, 104°36'58.3 "E) (Figure 1). Sampling was conducted in February–May 2023. The temperature, water acidity, and dissolved oxygen at the sampling site ranged from 24.4°C to 28.3°C, 5.5 to 6.4, and 4.9 to 6.4 mg/L, respectively. Fishing was conducted by fishermen using fish traps. A total of 74 *N. notopterus* samples were collected: 38 males (TL: 18–23.6 cm; BW: 35.1–92.1 g) and 36 females (TL: 19.6–26.3 cm; BW: 49.4–133.8 g). The *N. notopterus* samples were transported to the Fisheries Basic Laboratory, Aquaculture Study Program, Department of Fisheries, Faculty of Agriculture, Universitas Sriwijaya, for analysis of morphometric-meristic characters and also aspects of reproductive biology.





**Figure 1.** Map of *Notopterus notopterus* (Pallas 1769) sample collection in Kelekar River, Ogan Ilir Regency, South Sumatra

In the laboratory, fish were weighed using a digital scale (accuracy 0.01 gram). Then observe the fish morphologically and count the meristic characters

(number of rays of the dorsal fin, number of rays of the pectoral fin, and number of rays of the anal fin). Furthermore, the samples were measured for morphometric characters using a caliper (accuracy 0.01 cm) and a ruler (accuracy 0.1 cm). The morphometric characters measured were seventeen characters (Figure 2). Next, the fish were dissected, and the gonads and liver were taken, separated from other organs. Gonads and liver were weighed for the calculation of gonadosomatic index (IGS) = [gonad weight (g) / fish weight (g) x 100] and hepatosomatic index (IHS) = [liver weight (g) / fish weight (g) x 100] (Muddasir & Neelofar, 2017). Calculation of fecundity was done gravimetrically, namely, the fish gonads were taken partially, and then the number of eggs in the partial gonads was calculated. Fecundity = [number of sample gonad eggs x (total gonad weight/partial gonad weight)] (Bishai et al., 1974). Oocyte diameter was observed under a digital microscope. The maturity level of fish gonads was categorized based on the morphology and color of the gonads, IGS, and oocyte diameter. The sex ratio was calculated based on ~~the ratio of~~ the number of male and female fish. The data obtained were grouped into two categories, namely male and female fish. Data tabulation using the Microsoft Excel program, ~~then~~ ~~Furthermore, it was~~ analyzed descriptively to obtain a differences between male and female fish.

Formatted: Font: 12 pt





**Figure 2.** Morphometric character measurements of bronze featherback (*Notopterus notopterus*, Pallas 1769): TL (total length), SL (standard length), M-O (mouth to operculum distance), M-DF (mouth to dorsal fin distance), DF-C (dorsal fin to caudal distance), DFL (dorsal fin length), PFL (pectoral fin length), AF-CFL (anal fin to caudal fin length), BH (body height), HL (head length), HH (head height), DM (eye diameter), DTBH (Degree of Tilt of the Back of the Head), BT (body thickness), AT (abdominal thickness), HM (height of mouth), WM (width of mouth)

## Results and Discussion

A total of 74 *N. notopterus* samples, consisting of 38 males and 36 females, have been successfully measured for morphometric characters and calculated for meristic characters. The results of measurements and calculations of *N. notopterus* samples in this study are presented in Table 1. Morphological differences between male and female *N. notopterus* are presented in Figure 3.

**Table 1.** Morphometric and meristic characters of male and female *Notopterus notopterus* (Pallas, 1770) from Kelekar River, Ogan Ilir Regency, South Sumatra

Morphometric character	Male		TL/ character (%)	Female		TL/ character (%)
	Range (cm)	AVG±STD		Range (cm)	AVG±STD	
TL	18-23.6	20.39±1.39	-	19.6-26.3	22.46±1.56	-
SL	16-21.6	18.60±1.45	91.25	17.3-24.2	20.45±1.59	91.07
HL	1.3-3.5	2.07±0.51	10.18	1.6-3.7	2.16±0.47	9.64
HH	2.1-3.5	2.63±0.38	12.90	2.4-4.3	2.85±0.39	12.70
ED	0.5-0.7	0.64±0.06	3.16	0.8-1.0	0.86±0.06	3.87
BH	3.1-5.1	4.29±0.44	21.07	4.8-6.4	5.62±0.50	25.05
PFL	2.1-3.3	2.53±0.35	12.44	2.2-3.6	2.73±0.37	12.16
DFL	1.7-2.8	2.19±0.27	10.77	1.9-3.2	2.33±0.30	10.38
AF-CFL	13.2-18.4	15.42±1.22	75.62	14.0-19.1	16.44±1.36	73.20
M-O	3.0-4.7	3.65±0.40	17.91	3.0-4.9	3.90±0.44	17.40
M-DF	8.1-11.5	9.75±0.68	47.85	9.5-12.3	10.58±0.80	47.14
DF-C	8.0-11.6	9.85±1.02	48.32	8.5-12.5	10.94±0.98	48.72
BT	1.1-1.3	1.19±0.08	5.87	1.4-1.9	1.51±0.14	6.72

AT	0.6-1.0	0.89±0.11	4.40	1.1-1.6	1.26±0.13	5.61
MH	0.8-1.2	1.02±0.09	5.00	0.8-1.2	1.05±0.09	4.67
MW	0.8-1.1	0.96±0.10	4.73	0.8-1.2	1.00±0.11	4.47
DTBH (°)	32.0-50.0	42.36±3.51	207.76	47.0-63.0	52.6±3.06	234.35
<b>Meristic character</b>	Number of rays of the male fish		Number of rays of the female fish			
Pectoral (P)	10-14		11-14			
Dorsal (D)	6-9		6-9			
Anal+Caudal (A+C)	107-118		110-118			

The male and female *N. notopterus* were distinguished based on the difference in the proportion value of DTBH characters 26.59%; BH 3.98%; AF-CFL 2.42%; AT 1.21%. TL male fish 180-236 mm; female 196-263 mm; BW male 35.1-92.1 g; female 49.4-133.8 g; BH male 3.1-5.1 mm; female 4.8-6.4 mm; BT male 1.1-1.3 mm; female 1.4-1.9 mm; AT male 0.6-1.0 mm; female 1.1-1.6 mm. The total length of *N. notopterus* used in this study ranges from 18 cm to 23.6 cm; body weight ranges from 35.1 g to 92.1 g for male fish; The length ranges from 19.6 cm to 26.3 cm; and body weight ranges from 49.4 g to 33.8 g for female fish. According to the study results of Gustomi et al. (2016), the total length of female fish ranged from 130 mm to 249 mm, males from 120 mm to 232 mm, and the body weight of males was 22.17–97.17 g and females 38.98–120.47 g. The total length of *N. notopterus* ranged from 30 mm to 249 mm. The difference between male and female *N. notopterus* is based on the difference in the proportion value of DTBH characters 26.59%; BH 3.98%; AF CFL 2.42%; AT 1.21%. TL male fish 180-236 mm; female 196-263 mm; BW male 35.1-92.1 g; female 49.4-133.8 g; BH male 3.1-5.1 mm; female 4.8-6.4 mm; BT male 1.1-1.3 mm; female 1.4-1.9 mm; AT male 0.6-1.0 mm; female 1.1-1.6 mm. The total length of *N. notopterus* used in this study ranges from 18 cm to 23.6 cm; body weight ranges from 35.1 g to 92.1 g for male fish; length ranges from 19.6 cm to 26.3 cm; and body weight ranges from 49.4 g to 33.8 g for female fish. According to the study results of Gustomi et al., (2016), the total length of female fish ranged from 130 mm to 249 mm, males from 120 mm to 232 mm, and the body weight of males was 22.17-97.17 g and females 38.98-120.47 g. The total length of *N. notopterus* ranged from 30 mm to 249 mm. The total length of *N. notopterus* ranges from 401–950 mm (Wibowo et al., 2009). The male *N. notopterus* measured between 19.1 cm and 24.1 cm in length and weighed between 48.7 g and 133.4 g, while the female measured between 17.2 cm and 25.6 cm in length and weighed between 69.7 g and 151.2 g (Paul et al., 2022).

The dorsal fin of *N. notopterus* is fused with the caudal fin and anal fin. Each fin contains only rays and no spine. Meristic characters have been successfully counted. The results showed that P.10-14, D.6-9, and A+C.107-118. The results of the study by Rianti et al., (2021) showed that P.11-14, D.5-7, and A.98-111. According to the results of the Purba et al., (2017) study, the rays of the P. 11-14, D. 6-7, and A. 99-111. According to Mulyani & Budijono (2020), the number of rays on the P.13-14, D.7-9, and A.97-111. However, According to Sudarto (2011) stated that, D.7-9, and A.97-111. According to Kottelat et al., (1996), P.13-14, and A. 99-111.

One of the vital physiological processes that is essential to the life cycle of any organism, including fish, is reproduction (Muchlisin, 2014). Planning

improved conservation and management strategies for fishery resources requires a fundamental understanding of the reproductive cycle of fish (Parmaksiz et al., 2017; Radhakrishnan et al., 2020). Some aspects of *N. notopterus* reproductive biology observed in this study are presented in Table 2.

**Table 2.** Reproductive aspects of male and female *Notopterus notopterus* (Pallas, 1770) from Kelekar River, Ogan Ilir Regency, South Sumatra

Parameters	Sampling Period							
	February		March		April		May	
	<b>Male &amp; Female</b>							
GSI (%)	0.31±0.05	5.18±1.04	0.39±0.11	4.50±1.16	0.30±0.14	3.22±1.69	0.28±0.06	3.59±2.57
HSI (%)	0.65±0.22	0.53±0.10	0.52±0.19	0.51±0.22	0.79±0.23	0.80±0.36	0.79±0.27	0.73±0.19
Ova diameter (mm)	-	1.2-3.6	-	1.1-3.9	-	1.2-3.9	-	1.2-3.8
Fecundity	-	1361-3543	-	1455-3121	-	1060-1471	-	1007-3901
Sex ratio	1:1		1.14:1		1.5:1		0.86:1	

The GSI utilized to ascertain the fish's reproductive periodicity (Jewel et al., 2019). Table 2 shows that the GSI value of female fish was highest in February, and the male fish was highest in March. The GSI of female fish is higher than that of male fish. The GSI value is also related to the stage of gonad maturity. The GSI value increases with increasing gonad maturity. The increase in GSI indicates a developmental process in the gonads. The GSI value will reach a maximum value before spawning occurs. In addition, the GSI value is influenced by environmental factors related to the food availability as an energy source for somatic development and fish reproduction (Kasmi et al., 2017).

HSI values for female fish were highest in April and for males in May. HSI is the percentage value of the ratio between liver weight and total body weight. One of the organs that plays a role in fish reproduction is the liver. The liver plays an important role in the vitellogenesis process. This process triggered by the increasing concentration of estradiol in the blood. Estradiol is an important hormone produced by the ovaries in female fish, beneficial in the process of vitellogenesis. The higher the level of estradiol concentration in the blood, the earlier the liver will carry out the vitellogenesis process, and gonad maturation will be faster (Garnayak et al., 2013; Sullivan et al., 2018). The HSI indicated an energy reserve in the liver (Pandit & Gupta, 2019) and a bio-indicator of contaminant exposure (Araújo et al., 2018; Singh & Srivastava, 2015). The GSI is a well-known connection that is routinely utilized to ascertain the fish's reproductive periodicity (Jewel et al., 2019). Table 2 shows that the GSI value of female fish was highest in February, and that of male fish was highest in March. The GSI of female fish is higher than that of male fish. The GSI value is also related to the stage of gonad maturity. The GSI value increases with increasing gonad maturity. The increase in GSI indicates a developmental process in the gonads. The GSI value will reach a maximum value before spawning occurs. In addition, the GSI value is influenced by environmental factors related to the

Formatted: Justified



availability of food as an energy source for somatic development and fish reproduction (Kasmi et al., 2017).


HSI values for female fish were highest in April and for males in May. HSI is the percentage value of the ratio between liver weight and total body weight. One of the organs that plays a role in fish reproduction is the liver. The liver plays an important role in the vitellogenesis process. This process is triggered by the increasing concentration of estradiol in the blood. Estradiol is an important hormone produced by the ovaries in female fish that are in the process of vitellogenesis. The higher the level of estradiol concentration in the blood, the earlier the liver will carry out the vitellogenesis process, and gonad maturation will be faster (Garnayak et al., 2013; Sullivan et al., 2018). The HSI is an indicator of energy reserves in the liver (Pandit & Gupta, 2019) and a bio indicator of contaminant exposure (Araújo et al., 2018; Singh & Srivastava, 2015).

The fecundity of *N. notopterus* in this study ranged from 1.007 to 3.901 eggs per individual. The results of the study by Azadi et al., (1995) showed that fecundity values ranged from 53 to 748 and from 175 to 4.494 (Gupta & Ray, 2022). The value of fecundity is influenced by several factors, one of which is environmental. Variations in the number of fish fecundities are also caused by variations in the length and weight of fish. According to Parasmewaran & Sinha (1966), the range of ova per g body weight of the fish was 3-07 to 5-94, and there was a trend towards an increase as the fish grew larger.

The diameter of the ova of *N. notopterus* in this study ranged from 1.1 mm to 3.9 mm. The diameter of ova from a previous studies ranged from 1.05 to 2.2 mm (Gustomi et al., 2016). The oocyte diameter measured in this study is that of a categorized as mature ovary. The higher the oocyte diameter value, the higher the ovary's maturity. The results showed that the diameter of the oocyte in the ovary varied. This indicates that the development of oocytes in the ovary is not uniform. It is suspected that *N. notopterus* spawns gradually. This is in accordance with the opinion of Gustomi et al., (2016) that *N. notopterus* releases mature oocytes gradually. *N. notopterus* spawns more than once during a spawning season (Parasmewaran & Sinha, 1966).

The sex ratio is an important component in the study of fish reproduction (Purushottama et al., 2017; Radhakrishnan et al., 2020). The balance of the number of male and female fish in a body of water affects the continued existence of a species. The results of this study show that the number of male fish is higher than the number of female fish, except in the May sampling period. This result indicates that the population of male fish is greater than that of female fish.





**Figure 3.** The bronze featherback (*Notopterus notopterus*, Pallas 1769) from Kelekar River, Ogan Ilir Regency, South Sumatra (A) male and (B) female

### Conclusion

The results of this study found that there are differences in the morphology of male and female *N. notopterus*. There are differences in male and female morphometric characters in DTBH, BT, AF-CFL, and AT characters, and there are no differences in meristic characters. The gonadosomatic index of male fish ranged from 0.15% to 0.61%, and that of females ranged from 1.12% to 9.10%. The hepatosomatic index of male fish ranged from 0.19% to 1.38%, and females ranged from 0.15 to 1.23%; the sex ratio of the sample fish obtained was 1.056:1. The fecundity value of the sampled fish ranged from 1,007 to 3,901. The oocyte diameter of the sample fish ranged from 1.1 to 3.9 mm. The gonadal maturity stage of the sample fish was at the developing to mature stage for males and mature to spawning for females. The results of this study provide basic information for *N. notopterus* breeding studies.

This study denoted a differences in the morphology of male and female *N. notopterus*. There are differences in male and female morphometric characters in DTBH, BT, AF-CFL, and AT characters. However there are no differences in meristic characteristic. The gonadosomatic index of male fish ranged from 0.15% to 0.61%, and that of females ranged from 1.12% to 9.10%. The hepatosomatic index of male fish ranged from 0.19% to 1.38%, and females ranged from 0.15 to 1.23%; the sex ratio of the sample fish obtained was 1.056:1. The fecundity value of the sampled fish ranged from 1,007 to 3,901. The oocyte diameter of the sample fish ranged from 1.1 to 3.9 mm. The gonadal maturity stage of the sample fish was at the developing to mature stage for males and mature to spawning for females.

### Acknowledgements

We appreciate the fisherman (Mr. Nur and Mr. Lukman) who assisted with the fish collection. We also extend thanks to Universitas Sriwijaya's Rector and Head of LPPM. The research of this article was funded by DIPA of the Public Service Agency of Universitas Sriwijaya 2023. Nomor SP DIPA-

023.17.2.677515/2023, On November 30, 2022. In accordance with the Rector's Decree Number: 0188/UN9.3.1/SK/2023, On April 18, 2023.

## References

- Achakzai, W.M., Baloch, W.A., & Soomro, A.N. (2015). Length-weight relationship and condition factor of *Notopterus notopterus* (Pallas, 1769) from Manchar Lake Sindh, Pakistan. *Sindh University Research Journal (Science Series)*, 47(3), 515–518.
- Ambus, N., & Moody, E.K. (2018). Size-based sexual dimorphism of nitrogen excretion in livebearing fishes. *Ecology of Freshwater Fish*, 1–7. <https://doi.org/10.1111/eff.12445>
- Araújo, F.G., Morado, C.N., Parente, T.T.E., Paumgartten, F.J.R., & Gomes, I.D. (2018). Biomarkers and bioindicators of the environmental condition using a fish species (*Pimelodus maculatus* Lacepède, 1803) in a tropical reservoir in Southeastern Brazil. *Brazilian Journal of Biology*, 78(2), 351–359. <https://doi.org/10.1590/1519-6984.167209>
- Azadi, M., Islam, M., Nasiruddin, M., & Quader, M. (1995). Reproductive biology of *Notopterus notopterus* (Pallas) in Kaptai reservoir, Bangladesh. *Bangladesh Journal of Zoology*, 23, 215–220.
- Barby, F., Ráb, P., Lavoué, S., Ezaz, T., Bertollo, L., Kilian, A., ... Cioffi, M. (2018). From chromosomes to genome: Insights into the evolutionary relationships and biogeography of old. *Genes*, 9(6), 306. <https://doi.org/10.3390/genes9060306>
- Bishai, H.M., Ishak, M.M., & Labib, W. (1974). Fecundity of the mirror carp *Cyprinus carpio* L at the serow fish farm (Egypt). *Aquaculture*, 4, 257–265.
- Capasso, L., Carmela, M., Re, D.E.L., Bravi, S., & Taverne, L.P. (2021). Sexual dimorphism in a population of Pycnodontiformes of the Aptian (Lower Cretaceous) from Profeti (Province of Caserta, Southern Apennines, Italy). *Bollettino Del Museo Civico Di Storia Naturale Di Verona*, 45, 31–41.
- Chacko, J., & Sekharan, M. (2022). Sexual dimorphism in structures, size and shape of the cyprinid Nilgiri melon barb, *Haludaria fasciata*. *Fisheries and Aquatic Life*, 30, 138–148. <https://doi.org/10.2478/aopf-2022-0013>
- Dinh, Q.M., & Tran, N.T.T. (2018). Reproductive biological traits of the goby *Stigmatogobius pleurostigma* (Bleeker, 1849) from the Mekong Delta, Vietnam. *Indian Journal of Fisheries*, 65(1), 20–25. <https://doi.org/10.21077/ijf.2018.65.1.68188-04>
- Garcia, E.Q., & Zuanon, J. (2019). Sexual dimorphism in the electric knifefish, *Gymnorhamphichthys rondoni* (Rhamphichthyidae: Gymnotiformes). *Acta Amazonica*, 49, 213–220. <https://doi.org/http://dx.doi.org/10.1590/1809-4392201804392> ORIGINAL
- Garnayak, S.K., Mohanty, J., Rao, T.V., Sahoo, S.K., & Sahoo, P.K. (2013). Vitellogenin in Asian catfish, *Clarias batrachus*: Purification, partial characterization and quantification during the reproductive cycle by ELISA. *Aquaculture*, 392–395, 148–155. <https://doi.org/10.1016/J.AQUACULTURE.2013.02.020>
- Gupta, S., & Ray, A. (2022). Review on biology and culture potential of

- Notopterus notopterus* (Pallas, 1769). *International Journal of Aquatic Biology*, 10(2), 78–91.
- Gustomi, A., Sulistiono, S., & Yonvitner. (2016). Reproductive biology featherback (*Notopterus notopterus* Pallas, 1769) in Simpung Reservoir, Bangka island. *Jurnal Ilmu Pertanian Indonesia*, 21(1), 56–62. <https://doi.org/10.18343/jipi.21.1.56>
- Hilton, E.J., & Lavoué, S. (2018). A review of the systematic biology of fossil and living bony-tongue fishes, Osteoglossomorpha (Actinopterygii: Teleostei). *Neotropical Ichthyology*, 16(3), e180031. <https://doi.org/10.1590/1982-0224-20180031>
- Jewel, A.S., Haque, A., Ferdous, M.S., Khatun, M.S., Khanom, D.A., & Hasan, J. (2019). Gonadosomatic index and fecundity of threatened reba carp, *Cirrhinus reba* (Hamilton), in the Ganges River (northwest Bangladesh). *Fisheries and Aquatic Life*, 27, 80–85. <https://doi.org/10.2478/aopf-2019-0009>
- Jir, E.S.D., Motomura, H., & Babaran, R.P. (2020). Sexual dimorphism in the spotted stinger *Inimicus sinensis* (Valenciennes, 1833 ) and notes on pore structures on the body. *Philippine Journal of Systematic Biology Sexual*, 12(2), 1–11.
- Kasmi, M., Hadi, S., & Kantun, W. (2017). Reproductive biology of Indian mackerel, *Rastreliger kanagurta* (Cuvier, 1816) in Takalar Coastal Waters, South Sulawesi. *Jurnal Iktiologi Indonesia*, 17(3), 259–271.
- Kaur, V., & Rawal, Y.K. (2017). Length-weight relationship (LWR) in *Notopterus notopterus* (Pallas) from Sukhna Lake, Chandigarh. *IOSR Journal of Pharmacy and Biological Sciences*, 12(4), 63–65. <https://doi.org/10.9790/3008-1204046365>
- Kaushik, K.K., Sahu, P., & Nath, R. (2019). Length-weight relationship of *Notopterus notopterus* (Pallas, 1769), *Anabas testudineus* (Bloch, 1792) and *Clarias batrachus* (Linnaeus, 1758) from Pokoriya River, Morigaon, Assam, India. *International Journal of Scientific and Technology Research*, 8(10), 3087–3088. <https://doi.org/10.2307/1540>
- Kottelat, M. (2013). The fishes of the inland waters of Southeast Asia: a catalogue and core bibliography of the fishes known to occur in freshwaters, mangroves and estuaries. *The Raffles Bulletin of Zoology*, (27), 1–663.
- Kottelat, M., Whitten, A.J., Kartikasari, S., & Wirjoatmodjo, S. (1996). *Freshwater fishes of western Indonesia and Sulawesi*. Jakarta. Indonesia: Periplus Editions.
- Kumar, K.H., & Kiran, B.R. (2016). Relative condition factor of feather back, *Notopterus notopterus* (Pallas) from Jannapura pond, Bhadravathi Taluk, Karnataka. *International Journal of Scientific Research in Science, Engineering and Technology*, 2(1), 36–39.
- Kyaw, M.M., Kyaw, M.M., Lum, M.R., & Naing, S.S.S. (2020). Relationship of ecological factors and commercial bronze featherback fish, *Notopterus notopterus* (Pallas, 1769 ); Irrawaddy River, Myitkyina Segment, Kachin State, Myanmar. *University of Mandalay Research Journal*, 11, 98–111.
- Lavoué, S., Ghazali, S.Z., Jamaluddin, J.A.F., Nor, S.A.M.N., & Zain, K.M. (2020). Genetic evidence for the recognition of two allopatric species of Asian bronze featherback *Notopterus* (Teleostei, Osteoglossomorpha,

- Notopteridae). *Zoosyst*, 96(2), 449–454. <https://doi.org/10.3897/zse.96.51350>
- Mainero, A.H., Al-Jufaili, S.M., Jawad, L., & Reichenbacher, B. (2023). Sex dimorphism and evidence of sexually selected traits: A case study on the killifish *Aphaniops stoliczkanus* (Day, 1872). *Acta Zoologica*, 104, 473–487. <https://doi.org/10.1111/azo.12436>
- Mohanty, D., & Samanta, L. (2016). Chemosphere multivariate analysis of potential biomarkers of oxidative stress in *Notopterus notopterus* tissues from Mahanadi River as a function of concentration of heavy metals. *Chemosphere*, 155, 28–38. <https://doi.org/10.1016/j.chemosphere.2016.04.035>
- Muchlisin, Z. A. (2014). A general overview on some aspects of fish reproduction. *Aceh International Journal of Science and Technology*, 3(1), 43–52. <https://doi.org/10.13170/AIJST.0301.05>
- Muddasir, J., & Neelofar, J. (2017). Studies on the fecundity (F), gonadosomatic index (GSI) and hepatosomatic index (HSI) of *Salmo trutta fario* (Brown trout) at Kokernag trout fish farm, Anantnag, Jammu and Kashmir. *International Journal of Fisheries and Aquatic Studies*, 5(6), 170–173.
- Mulyani, I., & Budijono, B. (2020). Morphometric and meristic analysis of Asian knifefish (*Notopterus notopterus*) in Sail River, Pekanbaru Riau Province. *Jurnal Ilmiah Biologi Eksperimen dan Keanekaragaman Hayati*, 7(2), 59–64.
- Muslim, M., Heltonika, B., Sahusilawane, H., Wardani, W., & Rifai, R. (2020). *Ikan lokal perairan tawar Indonesia yang prospektif dibudidayakan* (Pertama). Purwokerto: Pena Persada.
- Muslim, M., & Syaifudin, M. (2022). Biodiversity of freshwater fish in Kelekar Floodplain Ogan Ilir Regency in Indonesia. *Journal of Tropical Biodiversity and Biotechnology*, 7(1), 67494. <https://doi.org/10.22146/jtbb.67494>
- Muslim, M., & Lestari, L.W. (2005). *Studi biodiversitas ikan di reservat perikanan Lebung Karang, Indralaya Ogan Ilir*. Indralaya.
- Mustafa, M.G., Singha, S., Islam, M., & Mallick, N. (2014). Population dynamics of *Notopterus notopterus* (Pallas, 1769) from the Kaptai Reservoir of Bangladesh. *SAARC Journal of Agriculture*, 12(2), 112–122. <https://doi.org/10.3329/sja.v12i2.21920>
- Naeem, B.M., Salam, A., Gillani, Q., & Ishtiaq, A. (2010). Length-weight relationships of *Notopterus notopterus* and introduced *Oreochromis niloticus* from the Indus River, southern Punjab, Pakistan. *Journal of Applied Ichthyology*, 26, 620. <https://doi.org/10.1111/j.1439-0426.2010.01480.x>
- Pandit, D.N., & Gupta, M.L. (2019). Hepato-somatic index, gonado-somatic index and condition factor of *Anabas testudineus* as bio-monitoring tools of nickel and chromium toxicity. *International Journal of Innovations in Engineering and Technology*, 12(3), 25–28.
- Parasmewaran, S., & Sinha, M. (1966). Observations on the biology of the feather-back, *Notopterus notopterus* (Pallas). *Indian Journal of Fisheries*, 13(1), 232–250.
- Parmaksiz, A., Oymak, S. A., Dogan, N., Naim, D. M., & Unlu, E. (2017). Reproductive characteristics of an invasive species *Carassius gibelio*

- (Bloch, 1782) in Ataturk Dam Lake, Turkey. *Indian Journal of Fisheries*, 64(4), 28–33. <https://doi.org/10.21077/ijf.2017.64.4.67478-04>
- Paul, M.G., Nath, P., & Dutta, A. (2022). Length-weight relationship, relative condition factor and fecundity of *Notopterus notopterus* (Pallas, 1769) from river Brahmaputra in Dhubri, Assam, India. *Journal of Fisheries*, 10(3), 103201. <https://doi.org/10.17017/j.fish.398>
- Purba, E.A., Efizon, D., & Putra, R.M. (2017). Study of morphometrics, meristics, and growth patterns of belida (*Notopterus notopterus*, Pallas 1769) in the Sail River of Pekanbaru City, Riau Province. *JOM Faprika*, 1, 1–8.
- Purushottama, G.B., Thakurdas, R.V., Dash, G., Akhilesh, K.V., Ramkumar, S., ... Zacharia, P.U. (2017). Reproductive biology and diet of the grey sharpnose shark *Rhizoprionodon oligoinx* Springer, 1964 (Chondrichthyes: Carcharhinidae) from the north-eastern Arabian Sea. *Indian Journal of Fisheries*, 64(4), 9–20. <https://doi.org/10.21077/ijf.2017.64.4.63379-02>
- Radhakrishnan, R.C., Kuttanelloor, R., & Balakrishna, M.K. (2020). Reproductive biology of the endemic cyprinid fish *Hypselobarbus thomassi* (Day, 1874) from Kallada River in the Western Ghats, India. *Journal of Applied Ichthyology*, 36(5), 604–612. <https://doi.org/10.1111/jai.14064>
- Rapita, R., Susiana, S., Kurniawan, D., Lestari, F., Sabriaty, D., & Rianti, U. (2021). Food habits of belida fish (*Notopterus notopterus*, Pallas 1769) in Sei Gesek Reservoir, Bintan Regency, Riau Island, Indonesia. *IOP Conference Series: Earth and Environmental Science* 919, 012003. <https://doi.org/10.1088/1755-1315/919/1/012003>
- Rianti, U., Susiana, S., & Kurniawan, D. (2021). Karakteristik morfometrik dan meristik ikan putak (*Notopterus notopterus*, Pallas 1769) di Waduk Sei Gesek Kabupaten Bintan. *BAWAL Widya Riset Perikanan Tangkap*, 13(3), 123–132.
- Singh, S., & Srivastava, A. K. (2015). Variations in hepatosomatic index (HSI) and gonadosomatic index (GSI) in fish *Heteropneustes fossilis* exposed to higher sub-lethal concentration to arsenic and copper. *Journal of Ecophysiology and Occupational Health*, 15(3–4), 89–93. <https://doi.org/10.15512/joeoh/2015/v15i3-4/121589>
- Srivastava, S., Rishnan, A., Singh, S., & Pandey, A. (2012). Embryo and larval development of threatened bronze featherback, *Notopterus notopterus* (Pallas). *Journal Experience Zoology India*, 15(2), 425–430.
- Srivastava, S.M., Singh, S.P., & Pandey, A.K. (2012). Food and feeding habits of threatened *Notopterus notopterus* in Gomti river, Lucknow (India). *Journal of Experimental Zoology India*, 15(2), 395–402.
- Sudarshan, S., & Kulkarni, R.S. (2013). Determination of condition factor (K), somatic condition factor (Ks), hepatic and gonado somatic indices in the fresh water fish *Notopterus notopterus*. *Zoology*, 2(11), 524–526.
- Sudarto, S. (2011). Ikan pipih yang potensial untuk ikan hias. *Media Akuakultur*, 6(1), 59–62.
- Sukendi, S., Thamrin, T., Putra, R., & Yulindra, A. (2020). Cultivation technology of bronze featherback (*Notopterus notopterus*, Pallas 1769) at different stocking densities and types of feed. *IOP Conf. Series: Earth and Environmental Science* 430, 012027. <https://doi.org/10.1088/1755-1315/430/1/012027>

- Sullivan, C.V., Yilmaz, O., Aquagyn, C., States, U., & Yilmaz, O. (2018). Vitellogenesis and yolk proteins, fish. *Reference Module in Life Sciences*, (January), 1–11. <https://doi.org/10.1016/B978-0-12-809633-8.20567-0>
- Uba, K.I.N. (2019). Sexual shape dimorphism in the monomorphic fish *Decapterus macrosoma* (Teleostei: Carangidae). *Computational Ecology and Software*, 9(4), 134–142.
- Ueng, Y., Lin, F., Chan, Y., Tsao, C., & Chen, M. (2022). Sexual dimorphism, length-weight relationships, fecundity, and diet of the striped eel catfish *Plotosus lineatus* (Plotosidae) on Taiwan's Southwest Coast. *Natural Resources*, 13, 77–90. <https://doi.org/10.4236/nr.2022.133006>
- Wibowo, A., Sunarno, M.T.D., Subagja, S., & Hidayah, T. (2009). Population characteristic of the putak (*Notopterus notopterus*) using fenotipic diversity analysis and 16sRNA region mitochondria DNA. *Jurnal Penelitian Perikanan Indonesia*, 15(1), 1–12.
- Winn, N.A., Sandi, P., Khaing, T., Nyunt, K.T., Kyaw, H.T., Sabai, M., & Aung, T.T.N. (2021). Length weight relationship of twelve freshwater fish species from Sunye Lake, Mandalay Region, Myanmar. *Greener Journal of Biological Sciences*, 11(2), 74–80.

## Cover Letter

### Dear Editor

I am pleased to submit an original research article entitled “**Sexual dimorphism and reproductive biology of bronze featherback (*Notopterus notopterus* Pallas 1769) from Kelekar River, Ogan Ilir, South Sumatra, Indonesia**” by Muslim Muslim, Mochamad Syaifudin, Ferdinand Hukama Taqwa, Muhammad iqbal Saputra.

For consideration for publication in the Baghdad Science Journal.

Our paper is very suitable with your journal because of the good results it contains. I agree for all journal conditions and I hope to get the acceptance.

This manuscript has not been published and is not under consideration for publication elsewhere.

Sexual dimorphism and reproductive biology are fundamental aspects of fish breeding studies. The aim of this research was to analyze the sexual dimorphism and reproductive biology of *N. notopterus*. A total of 74 *N. notopterus* collected from the Kelekar River in Ogan Ilir Regency, Indonesia, consisting 38 males (TL: 18–23.6 cm; BW: 35.1–92.1 g) and 36 females (TL: 19.6–26.3 cm; BW: 49.4–133.8 g). Seventeen morphometric characters, three meristic characters, and five reproductive biological parameters were analyzed. The results showed the differences in the morphometric characteristics of male and female *N. notopterus*. However, there was no difference in the meristic characters. The male gonadosomatic index ranged from 0.15 to 0.61%, and the female from 1.12 to 9.10%. The hepatosomatic index of males ranged from 0.19 to 1.38%, and that of females ranged from 0.15 to 1.23%, with a sex ratio of 1.056:1, fecundity of 1.007–3.901 eggs, ova diameters of 1.1–3.9 mm, and maturity levels of developed gonads to mature for males and mature to spawning for females. In conclusion, the differences between male and female *N. notopterus* can be identified morphologically.

### Best Regards

Corresponding Author

Corresponding Author Name: Muslim Muslim

Corresponding Author Address: Jl. Palembang-Prabumulih Km 32,  
Indralaya, South Sumatra, Indonesia

Corresponding Author E-mail: muslim\_bda@unsri.ac.id





Muslim &lt;muslim\_bda@unsri.ac.id&gt;

---

**[BSJ] Submission Acknowledgement**

1 message

---

**Prof. Dr. Fikrat M. Hassan** <fikrat@csw.uobaghdad.edu.iq>  
To: Muslim Muslim <muslim\_bda@unsri.ac.id>

Thu, Oct 19, 2023 at 5:26 AM

Dear Muslim Muslim:

Thank you for submitting the manuscript, "Sexual dimorphism and reproductive biology of bronze featherback (*Notopterus notopterus* Pallas 1769) from Kelekar River, Ogan Ilir, South Sumatra, Indonesia" to Baghdad Science Journal. With the online journal management system that we are using, you will be able to track its progress through the editorial process by logging in to the journal web site:

Submission URL: <https://bsj.uobaghdad.edu.iq/index.php/BSJ/authorDashboard/submission/9997>

Username: muslim\_unsri

If you have any questions, please contact me. Thank you for considering this journal as a venue for your work.

Prof. Dr. Fikrat M. Hassan

---

**Baghdad Science Journal,**

Al- Jadiryia, Baghdad P.O. BOX 4732. Baghdad-Iraq





E-mail : [bsj@csw.uobaghdad.edu.iq](mailto:bsj@csw.uobaghdad.edu.iq)

Mobile :+964 780 070 9148

## Participants

Salah Abdulla Hasoon (profsalah)

Muslim Muslim (muslim\_unsri)





Messages	
Note	From
<p>Dear Author</p> <p>Please correct your manuscript as requested by our journal, correction must be done on the same file attached, and fill out the attached form</p> <p>Best Regards</p> <p> <a href="#">Author certification.docx</a></p> <p> <a href="#">9997-Article Text-92567-102202-9-20240120.docx</a></p>	<p>profsalah</p> <p>2024-01-20 08:48 PM</p>
<p>▶ We resubmit the corrected manuscript and the completed form.</p> <p> <a href="#">9997-Certification of Proofreading-92593-102235-18-20240120.docx</a></p> <p> <a href="#">9997-Article Text-92592-102234-18-20240120.docx</a></p>	<p>muslim_unsri</p> <p>2024-01-21 06:01 PM</p>

[Add Message](#)

## Participants

Salah Abdulla Hasoon (profsalah)

Muslim Muslim (muslim\_unsri)

Messages	
Note	From
<p>Dear Author</p> <p>Please correct your manuscript as requested by our journal, correction must be done on the same file attached, and fill out the attached form</p> <p>Best Regards</p> <p> <a href="#">Author certification.docx</a></p> <p> <a href="#">9997-Article Text-92567-102202-9-20240120.docx</a></p>	<p>profsalah</p> <p>2024-01-20 08:48 PM</p>
<p>▶ We resubmit the corrected manuscript and the completed form.</p> <p> <a href="#">9997-Certification of Proofreading-92593-102235-18-20240120.docx</a></p> <p> <a href="#">9997-Article Text-92592-102234-18-20240120.docx</a></p>	<p>muslim_unsri</p> <p>2024-01-21 06:01 PM</p>

[Add Message](#)

## Sexual dimorphism and reproductive biology of bronze featherback (*Notopterus notopterus*, Pallas 1769) from Kelekar River, Ogan Ilir, South Sumatra, Indonesia

Muslim Muslim\*<sup>1</sup>  , Mochamad Syaifudin<sup>1</sup>  , Ferdinand Hukama Taqwa<sup>1</sup>  ,  
Muhammad Iqbal Saputra<sup>1</sup> 

Comment [N1]: ID?



\*Corresponding Author.

Comment [N2]: ???????  
Department? college? university?

Received dd/mm/yyyy, Revised dd/mm/yyyy, Accepted dd/mm/yyyy, Published Online First dd/mm/yyyy,  
Published dd/mm/yyyy



This work is licensed under a [Creative Commons Attribution 4.0 International License](https://creativecommons.org/licenses/by/4.0/).

### Abstract

Sexual dimorphism and reproductive biology are fundamental aspects of fish breeding studies. The aim of this research was to analyze the sexual dimorphism and reproductive biology of *N. notopterus*. A total of 74 *N. notopterus* were collected from the Kelekar River in Ogan Ilir Regency, Indonesia, consisting of 38 males (TL: 18–23.6 cm; BW: 35.1–92.1 g) and 36 females (TL: 19.6–26.3 cm; BW: 49.4–133.8 g). Seventeen morphometric characters, three meristic characters, and five reproductive biological parameters were analyzed. The results showed the differences in the morphometric characteristics of male and female *N. notopterus*. However, there was no difference in the meristic characters. The male gonadosomatic index ranged from 0.15 to 0.61%, and the female from 1.12 to 9.10%. The hepatosomatic index of males ranged from 0.19 to 1.38%, and that of females ranged from 0.15 to 1.23%, with a sex ratio of 1.056:1, fecundity of 1.007–3.901 eggs, ova diameters of 1.1–3.9 mm, and maturity levels of developed gonads to mature for males and mature to spawning for females. In conclusion, the differences between male and female *N. notopterus* can be identified morphologically.

**Keywords:** Broodstock selection; fish reproduction; gonochorism; secondary sexual characteristics

### Introduction

Bronze featherback (*Notopterus notopterus*, Pallas 1769) is a freshwater fish species native to Indonesia. *N. notopterus* is one of the fish species of the Notopteridae family<sup>1</sup>. The Notopteridae consist of approximately ten species in four genera: Chitala, Notopterus, Xenomystus, and Papyrocranus<sup>2</sup>. The only recognized species of Notopterus at present is *N. notopterus*<sup>3</sup>. The *N. notopterus* distribution in Asia includes Indonesia, India, Malaysia, Thailand, Bangladesh, Pakistan, and Vietnam<sup>1,2,4-7</sup>. In Indonesia, this species is distributed in Java, Sumatra, and Kalimantan<sup>1,8</sup>. *N. notopterus* was found in Kolong-Bendungan Simpur, Bangka Island<sup>9</sup>, Ogan River, Kelekar River (tributary of Musi River, South Sumatra), waters in Bangka Island as well as

Comment [N3]: Keywords: Required number of keywords is 5–8 words

Kota Bangun and Tanah Ulu (Mahakam River, East Kalimantan)<sup>10</sup>, Sail River, Pekanbaru City, Riau Province<sup>11</sup>, Sei Gesek Reservoir, Bintan, Riau Islands<sup>12</sup>, and Kelekar River<sup>13</sup>.


Several studies on *N. notopterus* have been conducted, among others, on aspects of length-weight relationship, relative condition factor and fecundity<sup>14-17</sup>, genetic diversity<sup>10</sup>, morphometrics and growth patterns<sup>11</sup>, food and feeding habits<sup>12</sup>, reproductive biology<sup>9</sup>, determination of condition factors, somatic condition, hepatosomatic index and gonadosomatic index<sup>18</sup>, behavior and reproduction<sup>19</sup>, potential biomarkers of oxidative stress<sup>20</sup>, embryo and larval development<sup>21</sup>, relative condition factors<sup>22,23</sup>, population dynamics<sup>5</sup>, relationship of ecological factors and commercial<sup>24</sup>, and its early domestication<sup>25,26</sup>.

Fishery management mainly depends on knowledge of fish reproductive biology<sup>27,28</sup>. In adult fish, sexual dimorphism is a frequent and well-known phenomenon<sup>29</sup>. The majority of fishes are bisexual and have various sexually dimorphic characteristics in their secondary sexual characteristics<sup>30</sup>. Distinguishing characteristics for the morphological qualities of many teleosts have been documented, including body color, ornamental patterns, breeding tubercles on the snout, pectoral fins and scales, fin hooks, body shape, fin size and shape<sup>31-35</sup>.

*N. notopterus* is one of the most valuable freshwater fish in South Sumatra, Indonesia. This fish is utilized by the community in South Sumatra, especially as a daily side dish and a raw material for the culinary industries of pempek (fish cake) and kerupuk-kemplang (fish crackers). Therefore, this species has the potential for cultivation. There is relatively little known about sexual dimorphism and reproductive biology in *N. notopterus*. Such information is crucial for initial data in the breeding of *N. notopterus*. The study aimed to analyze the sexual dimorphism and reproductive biology of *N. notopterus* from Sungai Kelekar, Ogan Ilir. The results of this study are beneficial in the selection of *N. notopterus* broodstock candidates for breeding programs.

## Materials and Methods

The fish collection was conducted in Kelekar River, Tanjung Baru Village, North Indralaya District, Ogan Ilir Regency, South Sumatra (Fig 1). Sampling was conducted in February–May 2023. The temperature, water acidity, and dissolved oxygen at the sampling site ranged from 24.4°C to 28.3°C, 5.5 to 6.4, and 4.9 to 6.4 mg.L<sup>-1</sup>, respectively. Fishing was conducted by fishermen using fish traps. A total of 74 *N. notopterus* samples were collected: 38 males (TL: 18–23.6 cm; BW: 35.1–92.1 g) and 36 females (TL: 19.6–26.3 cm; BW: 49.4–133.8 g). The *N. notopterus* samples were transported to the Fisheries Basic Laboratory, Aquaculture Study Program, Department of Fisheries, Faculty of Agriculture, Universitas Sriwijaya, for analysis of morphometric-meristic characters and also aspects of reproductive biology.



**Figure 1. Map of *Notopterus notopterus* (Pallas 1769) sample collection in Kelekar River, Ogan Ilir Regency, South Sumatra, Indonesia**

In the laboratory, fish were weighed using a digital scale (accuracy 0.01 gram). Then observe the fish morphologically and count the meristic characters (number of rays of the dorsal fin, number of rays of the pectoral fin, and number of rays of the anal fin). Furthermore, the samples were measured for morphometric characters using a caliper (accuracy 0.01 cm) and a ruler (accuracy 0.1 cm). The morphometric characters measured were seventeen characters (Fig 2). Next, the fish were dissected, and the gonads and liver were taken, separated from other organs. Gonads and liver were weighed for the calculation of gonadosomatic index (IGS) = [gonad weight (g) / fish weight (g) x 100] and hepatosomatic index (IHS) = [liver weight (g) / fish weight (g) x 100]<sup>36</sup>. The calculation of fecundity was done gravimetrically, namely, the fish gonads were taken partially, and then the number of eggs in the partial gonads was calculated. Fecundity = [number of sample gonad eggs x (total gonad weight/partial gonad weight)]<sup>37</sup>. Oocyte diameter was observed under a digital microscope. The maturity level of fish gonads was categorized based on the morphology and color of the gonads, IGS, and oocyte diameter. The sex ratio was calculated based on the number of male and female fish. The data obtained were grouped into two categories, namely male and female fish. Data tabulation using the Microsoft Excel program was then analyzed descriptively to obtain differences between male and female fish.




Figure 2. Morphometric character measurements of bronze featherback (*Notopterus notopterus*, Pallas 1769): TL (total length), SL (standard length), M-O (mouth to operculum distance), M-DF (mouth to dorsal fin distance), DF-C (dorsal fin to caudal distance), DFL (dorsal fin length), PFL (pectoral fin length), AF-CFL (anal fin to caudal fin length), BH (body height), HL (head length), HH (head height), ED (eye diameter), DTBH (Degree of Tilt of the Back of the Head), BT (body thickness), AT (abdominal thickness), HM (height of mouth), WM (width of mouth)

## Results and Discussion

A total of 74 *N. notopterus* samples, consisting of 38 males and 36 females, have been successfully measured for morphometric characters and calculated for meristic characters. The results of measurements and calculations of *N. notopterus* samples in this study are presented in Table 1. Morphological differences between male and female *N. notopterus* are presented in Fig 3.

**Table 1. Morphometric and meristic characters of male and female *Notopterus notopterus* (Pallas, 1770) from Kelekar River, Ogan Ilir Regency, South Sumatra, Indonesia**

Morphometric character	Male		Proportion (%)	Female		Proportion (%)
	Range (cm)	AVG±STD		Range (cm)	AVG±STD	
TL	18-23.6	20.39±1.39	-	19.6-26.3	22.46±1.56	-
SL	16-21.6	18.60±1.45	91.25 TL	17.3-24.2	20.45±1.59	91.07 TL
HL	1.3-3.5	2.07±0.51	10.18 TL	1.6-3.7	2.16±0.47	9.64 TL
HH	2.1-3.5	2.63±0.38	12.90 TL	2.4-4.3	2.85±0.39	12.70 TL
ED	0.5-0.7	0.64±0.06	30.91 HL	0.8-1.0	0.86±0.06	39.81 HL
BH	3.1-5.1	4.29±0.44	21.07 TL	4.8-6.4	5.62±0.50	25.05 TL
PFL	2.1-3.3	2.53±0.35	12.44 TL	2.2-3.6	2.73±0.37	12.16 TL
DFL	1.7-2.8	2.19±0.27	10.77 TL	1.9-3.2	2.33±0.30	10.38 TL
AF-CFL	13.2-18.4	15.42±1.22	75.62 TL	14.0-19.1	16.44±1.36	73.20 TL
M-O	3.0-4.7	3.65±0.40	17.91 TL	3.0-4.9	3.90±0.44	17.40 TL
M-DF	8.1-11.5	9.75±0.68	47.85 TL	9.5-12.3	10.58±0.80	47.14 TL
DF-C	8.0-11.6	9.85±1.02	48.32 TL	8.5-12.5	10.94±0.98	48.72 TL
BT	1.1-1.3	1.19±0.08	-	1.4-1.9	1.51±0.14	-
AT	0.6-1.0	0.89±0.11	-	1.1-1.6	1.26±0.13	-
MH	0.8-1.2	1.02±0.09	-	0.8-1.2	1.05±0.09	-
MW	0.8-1.1	0.96±0.10	-	0.8-1.2	1.00±0.11	-
DTBH (°)	32.0-50.0	42.36±3.51	-	47.0-63.0	52.6±3.06	-
<b>Meristic character</b>	<b>Number of rays of the male fish</b>		<b>Number of rays of the female fish</b>			
Pectoral (P)	10-14		11-14			
Dorsal (D)	6-9		6-9			
Anal+Caudal (A+C)	107-118		110-118			

The total length of *N. notopterus* used in this study ranges from 18 cm to 23.6 cm; body weight ranges from 35.1 g to 92.1 g for male fish; The length ranges from 19.6 cm to 26.3 cm; and body weight ranges from 49.4 g to 33.8 g for female fish. According to the study results of <sup>9</sup>, the total length of female fish ranged from 130 mm to 249 mm, males from 120 mm to 232 mm, and the body weight of males was 22.17–97.17 g and females 38.98–120.47 g. The total length of *N. notopterus* ranged from 30 mm to 249 mm. The total length of *N. notopterus* ranged from 30 mm to 249 mm. The total length of *N. notopterus* ranges from 401–950 mm <sup>10</sup>. The male *N. notopterus* measured between 19.1 cm and 24.1 cm in length and weighed between 48.7 g and 133.4 g, while the female measured between 17.2 cm and 25.6 cm in length and weighed between 69.7 g and 151.2 g <sup>16</sup>.

The dorsal fin of *N. notopterus* fused with the caudal fin and anal fin. Each fin contains only rays and no spine. Meristic characters have been successfully counted. The results showed that P.10-14, D.6-9, and A+C.107-118. The results of the study by <sup>38</sup> showed that P.11-14, D.5-7, and A.98-111. According to <sup>11</sup> study, the rays of the P. 11-14, D. 6-7, and A. 99-111. According to <sup>39</sup>, the number of rays on the P.13-14, D.7-9, and A.97-111. However, <sup>40</sup> stated that D.7-9, and A.97-111. According to <sup>8</sup>, P.13-14, and A. 99-111.

One of the vital physiological processes that is essential to the life cycle of any organism, including fish, is reproduction <sup>41</sup>. Planning improved conservation and management strategies for fishery resources requires a fundamental understanding of the reproductive cycle of fish <sup>28,42</sup>. Some aspects of *N. notopterus* reproductive biology observed in this study are presented in Table 2.



**Table 2. Reproductive aspects of male and female *Notopterus notopterus* (Pallas, 1770) from Kelekar River, Ogan Ilir Regency, South Sumatra, Indonesia**

Parameters	Sampling Period							
	February		March		April		May	
	Male	Female	Male	Female	Male	Female	Male	Female
GSI (%)	0.31±0.05	5.18±1.04	0.39±0.11	4.50±1.16	0.30±0.14	3.22±1.69	0.28±0.06	3.59±2.57
HSI (%)	0.65±0.22	0.53±0.10	0.52±0.19	0.51±0.22	0.79±0.23	0.80±0.36	0.79±0.27	0.73±0.19
Ova diameter (mm)	-	1.2-3.6	-	1.1-3.9	-	1.2-3.9	-	1.2-3.8
Fecundity	-	1361-3543	-	1455-3121	-	1060-1471	-	1007-3901
Sex ratio	1:1		1.14:1		1.5:1		0.86:1	

The GSI utilized to ascertain the fish's reproductive periodicity<sup>43</sup>. Table 2 shows that the GSI value of female fish was highest in February, and the male fish was highest in March. The GSI of female fish is higher than that of male fish. The GSI value is also related to the stage of gonad maturity. The GSI value increases with increasing gonad maturity. The increase in GSI indicates a developmental process in the gonads. The GSI value will reach a maximum value before spawning occurs. In addition, the GSI value is influenced by environmental factors related to the food availability as an energy source for somatic development and fish reproduction<sup>44</sup>.

HSI values for female fish were highest in April and for males in May. HSI is the percentage value of the ratio between liver weight and total body weight. One of the organs that plays a role in fish reproduction is the liver. The liver plays an important role in the vitellogenesis process. This process is triggered by the increasing concentration of estradiol in the blood. Estradiol is an important hormone produced by the ovaries in female fish that is beneficial in the process of vitellogenesis. The higher the level of estradiol concentration in the blood, the earlier the liver will carry out the vitellogenesis process, and gonad maturation will be faster<sup>45,46</sup>. The HSI indicated an energy reserve in the liver<sup>47</sup> and a bio-indicator of contaminant exposure<sup>48,49</sup>. The fecundity of *N. notopterus* in this study ranged from 1,007 to 3,901 eggs per individual.<sup>50</sup> showed that fecundity values ranged from 53 to 748 and from 175 to 4,494<sup>51</sup>. The value of fecundity is influenced by several factors, one of which is environmental. Variations in the number of fish fecundities are also caused by variations in the length and weight of fish.

The diameter of the ova of *N. notopterus* in this study ranged from 1.1 mm to 3.9 mm. Previous studies ranged from 1.05 to 2.2 mm<sup>9</sup>. The oocyte diameter is categorized as a mature ovary. The higher the oocyte diameter value, the higher the ovary's maturity. The results showed that the diameter of the oocyte in the ovary varied. This indicates that the development of oocytes in the ovary is not uniform. It is suspected that *N. notopterus* spawns gradually. This is by the opinion of<sup>9</sup> that *N. notopterus* releases mature oocytes gradually. *N. notopterus* spawns more than once during a spawning season. The sex ratio is an important component in the study of fish reproduction<sup>42,52</sup>. The balance of the number of male and female fish in a body of water affects the continued existence of a species. The results of this study show that the number of male fish is higher than the number of female fish, except in the May sampling period. This result indicates that the population of male fish is greater than that of female fish.

## Conclusion

There are differences in the morphometric characteristics of male and female *N. notopterus*. However, there are no differences in meristic characters. The gonadosomatic index of male fish ranged from 0.15% to 0.61%, and that of females ranged from 1.12% to 9.10%. The hepatosomatic index of male fish ranged from 0.19% to 1.38%, and females ranged from 0.15 to 1.23%; the sex ratio of the sample fish obtained was 1.056:1. The fecundity value of the sampled fish ranged from 1,007 to 3,901. The oocyte diameter of the

sample fish ranged from 1.1 to 3.9 mm. The gonadal maturity stage of the sample fish was at the developing to mature stage for males and mature to spawning for females.

### Acknowledgment

We appreciate the fisherman (Mr. Nur and Mr. Lukman) who assisted with the fish collection. We also extend thanks to Universitas Sriwijaya's Rector and Head of LPPM. The research of this article was funded by DIPA of the Public Service Agency of Universitas Sriwijaya 2023. Nomor SP DIPA-023.17.2.677515/2023, On November 30, 2022. Rector's Decree Number: 0188/UN9.3.1/SK/2023, On April 18, 2023.

### Authors' Declaration

- Conflicts of Interest: None.
- We hereby confirm that all the Figures and Tables in the manuscript are ours. Furthermore, any Figures and images, that are not ours, have been included with the necessary permission for re-publication, which is attached to the manuscript.
- Authors sign on ethical consideration's approval
- Ethical Clearance: The project was approved by the local ethical committee in Universitas Sriwijaya

### Authors' Contribution Statement

### References

1. Kottelat M. The fishes of the inland waters of Southeast Asia: a catalogue and core bibliography of the fishes known to occur in freshwaters, mangroves and estuaries. *Raffles Bull Zool.* 2013; (27): 1–663. [https://lknhm.nus.edu.sg/wp-content/uploads/sites/10/app/uploads/2017/04/rbz\\_S27.pdf](https://lknhm.nus.edu.sg/wp-content/uploads/sites/10/app/uploads/2017/04/rbz_S27.pdf)
2. Barby F, Ráb P, Lavoué S, Ezaz T, Bertollo L, Kilian A, et al. From chromosomes to genome: Insights into the evolutionary relationships and biogeography of old. *Genes.* 2018; 9(6): 306. <https://doi.org/10.3390/genes9060306>.
3. Lavoué S, Ghazali SZ, Jamaluddin JA, Nor SA, Zain KM. Genetic evidence for the recognition of two allopatric species of Asian bronze featherback *Notopterus notopterus* (Teleostei, Osteoglossomorpha, Notopteridae). *Zoosyst Evol.* 2020 Jan 7; 96(2): 449-54. <https://doi.org/10.3897/zse.96.51350>
4. Hilton EJ, Lavoué S. A review of the systematic biology of fossil and living bony-tongue fishes, Osteoglossomorpha (Actinopterygii: Teleostei). *Neotrop Ichthyol.* 2018 Oct 11; 16(3): e180031. <https://doi.org/10.1590/1982-0224-20180031>.
5. Mustafa MG, Singha S, Islam MR, Mallick N. Population dynamics of *Notopterus notopterus* (Pallas, 1769) from the Kaptai reservoir of Bangladesh. *SAARC J Agricult.* 2014; 12(2): 112-22. <https://doi.org/10.3329/sja.v12i2.21920>.
6. Naeem M, Salam A, Gillani Q, Ishtiaq A. Length- weight relationships of *Notopterus notopterus* and introduced *Oreochromis niloticus* from the Indus River, southern Punjab, Pakistan. *J Appl Ichthyol.* 2010 Aug; 26(4): 620-. <https://doi.org/10.1111/j.1439-0426.2010.01480.x>
7. Srivastava SM, Singh SP, Pandey AK. Food and feeding habits of threatened *Notopterus notopterus* in Gomti river, Lucknow (India). *J Exp Zoology India.* 2012; 15(2): 395–402.
8. Kottelat M, Whitten T. Freshwater fishes of Western Indonesia and Sulawesi: additions and corrections. Hong Kong: Periplus editions; 1996.
9. Gustomi A, Sulistiono S, Yonvitner Y. Reproductive biology featherback (*Notopterus notopterus* Pallas, 1769) in Simpur Reservoir, Bangka Island. *JIPi.* 2016; 21: 56-62. <https://doi.org/10.18343/jipi.21.1.56>
10. Wibowo A, Sunarno MT, Subagdja S, Hidayah T. Population characteristic of the putak (*Notopterus notopterus*) using fenotypic diversity analysis and 16sRNA region mitochondria DNA. *JIPi.* 2017 Feb 1; 15(1): 1-2. <https://dx.doi.org/10.15578/jipi.15.1.2009.1-12>
11. Purba EA, Efizon D, Putra RM. Study of morphometrics, meristics, and growth patterns of belida (*Notopterus notopterus*, Pallas 1769) in the Sail River of Pekanbaru City, Riau Province. *JOM Faprika.* 2017; 1: 1–8.

Comment [N4]: J Exp Zool India

Comment [N5]: Page , link?

Comment [N6]: ??????  
Indones. J. Agric. Sci

Comment [N7]: Issue?

Comment [N8]: ??????

- <https://jom.unri.ac.id/index.php/JOMFAPERIKA/article/view/16180/15711>
12. Rapita R, Susiana S, Kurniawan D, Lestari F, Sabriaty D, Rianti U. Food habits of belida fish (*Notopterus notopterus*, Pallas 1769) in Sei Gesek Reservoir, Bintan Regency, Riau Island, Indonesia. *IOP Conference Series: Earth and Environmental Science*. 2021 Nov 1; 919: 012003. <https://doi.org/10.1088/1755-1315/919/1/012003>
  13. Muslim M, Syaifudin M. Biodiversity of freshwater fish in Kelekar Floodplain Ogan Ilir Regency in Indonesia. *J Trop Biodivers Biotechnol*. 2022; 7(1): 67494. <https://doi.org/10.22146/jtbb.67494>.
  14. Kaur V, Rawal YK. Length-weight relationship (LWR) in *Notopterus notopterus* (Pallas) from Sukhna Lake, Chandigarh. *IOSR J Pharm Biol Sci*. 2017; 12(4): 63–65. <https://doi.org/10.9790/3008-1204046365>
  15. Kaushik KK, Sahu P, Nath R. Length-weight relationship of *Notopterus notopterus* (Pallas, 1769), *Anabas testudineus* (Bloch, 1792) and *Clarias batrachus* (Linnaeus, 1758) from Pokoriya River, Morigaon, Assam, India. *Int J Sci Technol Res*. 2019; 8(10): 3087–3088. <https://doi.org/10.2307/1540>
  16. Paul MG, Nath P, Dutta A. Length-weight relationship, relative condition factor and fecundity of *Notopterus notopterus* (Pallas, 1769) from river Brahmaputra in Dhubri, Assam, India *J Fish*. 2022; 10(3): 103201. <https://doi.org/10.17017/j.fish.398>
  17. Winn NA, Sandi P, Khaing T, Thi K. Length weight relationship of twelve freshwater fish species from Sunny Lake, Mandalay Region, Myanmar. *Greener J Biol Sci*. 2021; 11(2): 74–80. <https://gjournal.org/GJBS/Publication/2021/2/PDF/092221092%20Nandar%20et%20al.pdf>
  18. Sudarshan S, Kulkarni RS. Determination of condition factor (K) somatic condition factor (Ks) hepatic and gonado somatic indices in the fresh water fish *Notopterus Notopterus*. *Int J Sci Res*. 2013; 2(11): 524–6.
  19. Sukendi S, Thamrin T, Putra RM, Yulindra A. Cultivation technology of bronze featherback (*Notopterus notopterus*, Pallas 1769) at different stocking densities and types of feed. *IOP Conference Series: Earth and Environmental Science*. 2020 (Vol. 430, No. 1, p. 012027). IOP Publishing. <https://doi.org/10.1088/1755-1315/430/1/012027>
  20. Mohanty D, Samanta L. Multivariate analysis of potential biomarkers of oxidative stress in *Notopterus notopterus* tissues from Mahanadi River as a function of concentration of heavy metals. *Chemosphere*. 2016 Jul 1; 155: 28–38. <https://doi.org/10.1016/j.chemosphere.2016.04.035>
  21. Srivastava SM, Gopalakrishnan A, Singh SP, Pandey AK. Embryonic and larval development of threatened bronze featherback, *Notopterus notopterus* (Pallas). *J Exp Zool India*. 2012; 15(2): 425–30. <https://krishi.icar.gov.in/jspui/handle/123456789/4918>
  22. Achakzai WM, Saddozai S, Baloch WA, Soomro AN, Memon N. Length-weight relationship and condition factor of *Notopterus notopterus* (Pallas, 1769) from Manchar Lake Sindh, Pakistan. *SURJ*. 2015 Sep 15; 47(3): 515–18. <https://sujo.usindh.edu.pk/index.php/SURJ/article/view/5200>
  23. Kumar KH, Kiran BR. Relative condition factor of feather back, *Notopterus Notopterus* (Pallas) from Jannapura Pond, Bhadravathi Taluk, Karnataka. *Int J Sci Res Sci Eng Technol*. 2016; 2(1): 36–9. <https://ijsrset.com/IJSRSET162113>
  24. Kyaw MM, Kyaw MM, Lum MR, Naing SS. Relationship of ecological factors and commercial bronze featherback fish, *Notopterus notopterus* (Pallas, 1769 ); Irrawaddy River, Myitkyina Segment, Kachin State, Myanmar. *UMRJ*. 2020; 11: 98–111.
  25. Muslim M, Simanjuntak WJ. Growth and survival of bronze featherback (*Notopterus notopterus*, Pallas 1769) reared on bucket. *MSARR*. 2023; 9(1): 101–05. <https://doi.org/10.30574/msarr.2023.9.1.0133>.
  26. Muslim M, Pitriani E, Agustina H. Growth and survival of bronze featherback (*Notopterus notopterus*) adapted on box container and aquarium. *GSCARR*. 2023; 16(3): 133–37. <https://doi.org/10.30574/gscarr.2023.16.3.0368>
  27. Dinh QM, Tran NTT. Reproductive biological traits of the goby *Stigmatogobius pleurostigma* (Bleeker, 1849) from the Mekong Delta, Vietnam. *J Indian Fish*. 2018; 65(1): 20–25. <https://doi.org/10.21077/ijf.2018.65.1.68188-04>
  28. Parmaksız A, Oymak SA, Dogan N, Naim DM, Unlu E. Reproductive characteristics of an invasive species *Carassius gibelio* (Bloch, 1782) in Ataturk Dam Lake, Turkey. *J Indian Fish*. 2017 Jan 1; 64(4): 28–33. <https://doi.org/10.21077/ijf.2017.64.4.67478-04>
  29. Capasso LU, Del Re MC, Bravi SE, Taverne LP. Sexual dimorphism in a population of Pycnodontiformes of the Aptian (Lower Cretaceous) from Profeti (province of Caserta, Southern Apennines, Italy). *Boll Mus Civ Stor Nat. Verona*. 2021; 45: 31–41.
  30. Chacko JJ, Sekharan NM. Sexual dimorphism in structures, size and shape of the cyprinid Nilgiri melon barb. *Fish Aquat Life*. 2022 Sep 30; 30(3): 138–48. <https://doi.org/10.2478/aopf-2022-0013>.
  31. Ambus N, Moody EK. Size- based sexual dimorphism of nitrogen excretion in livebearing fishes. *Ecol Freshw Fish*. 2019 Apr; 28(2): 222–8. <https://doi.org/10.1111/eff.12445>
  32. Garcia EQ, Zuanon J. Sexual dimorphism in the electric knifefish, *Gymnorhamphichthys rondoni* (Rhamphichthyidae: Gymnotiformes). *Acta Amazonica*. 2019 Aug 12; 49: 213–20. <https://doi.org/http://dx.doi.org/10.1590/1809-4392201804392> original
  33. Mainero AH, Al-Jufaili SM, Jawad L, Reichenbacher B. Sex dimorphism and evidence of sexually selected traits: A

Comment [N9]: Abbreviation?

Comment [N10]: Abbreviation?

Comment [N11]: Wrong way?

Comment [N12]:  
Sindh Univ Res JourComment [N13]: ?????????/  
Abbreviation?

Comment [N14]: ????????

Comment [N15]: ???????

Comment [N16]: Link?

- case study on the killifish *Aphaniops stoliczkanus* (Day, 1872). *Acta Zoologica*. 2022 Aug 21; 104: 473–87. <https://doi.org/10.1111/azo.12436>
34. Uba KI. Sexual shape dimorphism in the monomorphic fish *Decapterus macrosoma* (Teleostei: Carangidae). *Comput Ecol Softw*. 2019 Dec 1; 9(4): 134-42. [http://www.iaees.org/publications/journals/ces/articles/2019-9\(4\)/sexual-shape-dimorphism-in-monomorphic-fish-Decapterus-macrosoma.pdf](http://www.iaees.org/publications/journals/ces/articles/2019-9(4)/sexual-shape-dimorphism-in-monomorphic-fish-Decapterus-macrosoma.pdf)
  35. Ueng YT, Lin FJ, Chan YS, Tsao CW, Chen MJ. Sexual dimorphism, length-weight relationships, fecundity, and diet of the striped eel catfish *Plotosus lineatus* (Plotosidae) on Taiwan's southwest coast. *Nat Resour*. 2022 Mar 22; 13(3): 77-90. <https://doi.org/10.4236/nr.2022.133006>
  36. Jan M, Jan N. Studies on the fecundity (F), gonadosomatic index (GSI) and hepatosomatic index (HSI) of *Salmo trutta fario* (Brown trout) at Kokernag trout fish farm, Anantnag, Jammu and Kashmir. *Int J Fish Aquat Stud*. 2017; 5(6): 170-3. <https://www.fisheriesjournal.com/archives/2017/vol5issue6/PartC/5-6-17-154.pdf>
  37. Bishai HM, Ishak MM, Labib W. Fecundity of the mirror carp *Cyprinus carpio* L. at the Serow Fish Farm (Egypt). *Aquaculture*. 1974 Jan 1; 4: 257-65. [https://doi.org/10.1016/0044-8486\(74\)90038-6](https://doi.org/10.1016/0044-8486(74)90038-6)
  38. Rianti U, Susiana S, Kurniawan D. Morphometric and meristic characteristics of bronze featherback (*Notopterus notopterus*, pallas 1769) in Sei Gesek Reservoir Bintan Regency. *BAWAL*. 2021 Dec 6; 13(3): 123-32. <https://dx.doi.org/10.15578/bawal.13.3.2021.123-132>
  39. Mulyani I, Budijono B. Morphometric and meristic analysis of Asian knifefish (*Notopterus notopterus*) in Sail River, Pekanbaru Riau Province. *J-BEKH*. 2020 Dec 1; 7(2): 59-64. <https://doi.org/10.23960/jbekh.v7i2.156>
  40. Sudarto S. Flatfish that have potential for ornamental fish. *Media Akuakultur*. 2011 Jun 30; 6(1): 59-62.
  41. Muchlisin ZA. A general overview on some aspects of fish reproduction. *AIJST*. 2014 Jan 1; 3(1):43-52. <https://doi.org/10.13170/AIJST.0301.05>
  42. Chelapurath RR, Kuttanelloor R, Balakrishna MK. Reproductive biology of the endemic cyprinid fish *Hypselobarbus thomassi* (Day, 1874) from Kallada River in the Western Ghats, India. *J Appl Ichthyol*. 2020 Oct; 36(5): 604-12. <https://doi.org/10.1111/jai.14064>
  43. Jewel MA, Haque MA, Ferdous MS, Khatun MS, Hasan J. Gonadosomatic index and fecundity of threatened reba carp, (Hamilton), in the Ganges River (northwest Bangladesh). *Fish Aquat Life*. 2019; 27(2): 80-5. <https://doi.org/10.2478/aopf-2019-0009>
  44. Kasmi M, Hadi S, Kantun W. Reproductive biology of Indian mackerel, *Rastreliger kanagurta* (Cuvier, 1816) in Takalar coastal waters, South Sulawesi. *J Iktiologi Indones*. 2017 Oct 23; 17(3): 259-71. <https://doi.org/10.32491/jii.v17i3.364>
  45. Garnayak SK, Mohanty J, Rao TV, Sahoo SK, Sahoo PK. Vitellogenin in Asian catfish, *Clarias batrachus*: Purification, partial characterization and quantification during the reproductive cycle by ELISA. *Aquaculture*. 2013 May 10; 392: 148-55. <https://doi.org/10.1016/j.aquaculture.2013.02.020>
  46. Sullivan CV, Yilmaz O. Vitellogenesis and yolk proteins, fish. *Encyclopedia of reproduction*. 2018 Jan 1; 6: 266-77. <https://doi.org/10.1016/B978-0-12-809633-8.20567-0>
  47. Pandit k, Gupta ML. Hepato-somatic index, gonado-somatic index and condition factor of *Anabas testudineus* as bio-monitoring tools of nickel and chromium toxicity. *Int J Innov Eng Technol*. 2019; 12(3): 25-8. <https://ijiet.com/wp-content/uploads/2019/03/5.pdf>
  48. Araújo FG, Morado CN, Parente TT, Paumgarten FJ, Gomes ID. Biomarkers and bioindicators of the environmental condition using a fish species (*Pimelodus maculatus* Lacepède, 1803) in a tropical reservoir in Southeastern Brazil. *Braz J Biol*. 2017 Aug 17; 78: 351-9. <https://doi.org/10.1590/1519-6984.167209>
  49. Singh S, Srivastava AK. Variations in hepatosomatic index (HSI) and gonadosomatic index (GSI) in fish *Heteropneustes fossilis* exposed to higher sub-lethal concentration to arsenic and copper. *J Ecophysiol Occup Health*. 2015 Dec 1; 15(3/4): 89-93. <https://doi.org/10.15512/joeoh/2015/v15i3-4/121589>
  50. Azadi MA, Islam MA, Nasiruddin M, Quader MF. Reproductive biology of *Notopterus notopterus* (Pallas) in Kaptai reservoir, Bangladesh. *Bangladesh J Zool*. 1995; 23: 215-20.
  51. Gupta S, Ray A. Review on biology and culture potential of *Notopterus notopterus* (Pallas, 1769). *Int J Aquat Biol*. 2022 Apr 1; 10(2):78-91. <https://ij-aquaticbiology.com/index.php/ijab/article/view/1511>.
  52. Purushottama GB, Ramasubramanian V, Dash G, Akhilesh KV, Ramkumar S, Kizhakudan SJ, et al. Reproductive biology and diet of the grey sharpnose shark *Rhizoprionodon oligolinx* Springer, 1964 (Chondrichthyes: Carcharhinidae) from the north-eastern Arabian Sea. *J Indian Fish*. 2017; 64(4): 9-20. <https://doi.org/10.21077/ijf.2017.64.4.63379-02>

Comment [N17]: ????

Comment [N18]: ????

Comment [N19]: Edition?, page?, link?

## إزدواج الشكل الجنسي وبيولوجيا الإنجاب للـ *Bronze featherback*

(*Notopterus notopterus* ، Pallas 1769) من نهر كيليكار، أوغان إير، جنوب سومطرة، إندونيسيا

### الخلاصة

يعد ازدواج الشكل الجنسي وعلم الأحياء الإنجابي من الجوانب الأساسية لدراسات تربية الأسماك. كان الهدف من هذا البحث هو تحليل ازدواج الشكل الجنسي والبيولوجيا الإنجابية لـ *N. notopterus*. تم جمع ما مجموعه 74 *N. notopterus* من نهر كيليكار في منطقة أوغان إير، إندونيسيا، وتتكون من 38 ذكرًا (طول الجسم: 18-23.6 سم؛ وزن الجسم: 35.1-92.1 جم) و36 أنثى (طول الجسم: 19.6-26.3 سم؛ وزن الجسم: 35.1-92.1 جم) و36 أنثى (طول الجسم: 19.6-26.3 سم؛ وزن الجسم: 49.4-133.8 جم). تم تحليل سبعة عشر شخصية مورفومترية وثلاثة صفات مرسية وخمسة معايير بيولوجية إنجابية. أظهرت النتائج وجود اختلافات في الخصائص المورفومترية للذكور والإناث *N. notopterus*. ومع ذلك، لم يكن هناك اختلاف في الشخصيات المرسية. ويتراوح مؤشر الغدد التناسلية عند الذكور من 0.15 إلى 0.61%، وعند الإناث من 1.12 إلى 9.10%. تراوح المؤشر الكبدى عند الذكور من 0.19 إلى 1.38%، وتراوح عند الإناث من 0.15 إلى 1.23%، مع نسبة جنس 1:1.056، وخصوبة 1.007-3.901 بيضة، وأقطار البويضات 1.1-3.9 ملم، ومستويات النضج. من الغدد التناسلية المتطورة لتتضح عند الذكور وتتضح لوضع البيض عند الإناث. في الختام، يمكن تحديد الاختلافات بين الذكور والإناث *N. notopterus* شكليا.




**الكلمات المفتاحية:** اختيار الأمهات؛ تكاثر الأسماك داء البنية. الخصائص الجنسية الثانوية

## Participants

Salah Abdulla Hasoon (profsalah)

Muslim Muslim (muslim\_unsri)

### Messages

Note	From
<p>Dear Sir</p> <p>1- Check again all references in your manuscript, correction must be done on the same file attached,</p> <p>2- the signature of your certification?</p> <p>Regards</p> <p> <a href="#">9997-Article Text-92767-102413-18-20240121.docx</a></p>	<p>profsalah</p> <p>2024-01-22 11:48 AM</p>
<p>▶ We send back the corrected manuscript and signed form.</p> <p> <a href="#">9997-Certification of Proofreading-92593-102235-18-20240120.docx</a></p> <p> <a href="#">9997-Article Text-92866-102519-18-20240122 (Rev).docx</a></p>	<p>muslim_unsri</p> <p>2024-01-22 05:25 PM</p>




[Add Message](#)

## Participants

Salah Abdulla Hasoon (profsalah)

Muslim Muslim (muslim\_unsri)

### Messages

Note	From
<p>Dear Sir</p> <p>1- Check again all references in your manuscript, correction must be done on the same file attached,</p> <p>2- the signature of your certification?</p> <p>Regards</p> <p> <a href="#">9997-Article Text-92767-102413-18-20240121.docx</a></p>	<p>profsalah</p> <p>2024-01-22 11:48 AM</p>
<p>▶ We send back the corrected manuscript and signed form.</p> <p> <a href="#">9997-Certification of Proofreading-92593-102235-18-20240120.docx</a></p> <p> <a href="#">9997-Article Text-92866-102519-18-20240122 (Rev).docx</a></p>	<p>muslim_unsri</p> <p>2024-01-22 05:25 PM</p>

[Add Message](#)

## Sexual dimorphism and reproductive biology of bronze featherback (*Notopterus notopterus*, Pallas 1769) from Kelekar River, Ogan Ilir, South Sumatra, Indonesia

Muslim Muslim\*<sup>1</sup>  , Mochamad Syaifudin<sup>1</sup>  , Ferdinand Hukama Taqwa<sup>1</sup>  ,  
Muhammad Iqbal Saputra<sup>1</sup>  

<sup>1</sup>Aquaculture Study Program, Department of Fisheries, Faculty of Agriculture, Universitas Sriwijaya, Indralaya, Indonesia

\*Corresponding Author.

Received dd/mm/yyyy, Revised dd/mm/yyyy, Accepted dd/mm/yyyy, Published Online First dd/mm/yyyy,  
Published dd/mm/yyyy



This work is licensed under a [Creative Commons Attribution 4.0 International License](https://creativecommons.org/licenses/by/4.0/).

### Abstract

Sexual dimorphism and reproductive biology are fundamental aspects of fish breeding studies. The aim of this research was to analyze the sexual dimorphism and reproductive biology of *N. notopterus*. A total of 74 *N. notopterus* were collected from the Kelekar River in Ogan Ilir Regency, Indonesia, consisting of 38 males (TL: 18–23.6 cm; BW: 35.1–92.1 g) and 36 females (TL: 19.6–26.3 cm; BW: 49.4–133.8 g). Seventeen morphometric characters, three meristic characters, and five reproductive biological parameters were analyzed. The results showed the differences in the morphometric characteristics of male and female *N. notopterus*. However, there was no difference in the meristic characters. The male gonadosomatic index ranged from 0.15 to 0.61%, and the female from 1.12 to 9.10%. The hepatosomatic index of males ranged from 0.19 to 1.38%, and that of females ranged from 0.15 to 1.23%, with a sex ratio of 1.056:1, fecundity of 1.007–3.901 eggs, ova diameters of 1.1–3.9 mm, and maturity levels of developed gonads to mature for males and mature to spawning for females. In conclusion, the differences between male and female *N. notopterus* can be identified morphologically.

**Keywords:** Broodstock selection; fish reproduction; gonochorism; Indonesian fish; knifefish; notopteridae; secondary sexual characteristics

### Introduction

Bronze featherback (*Notopterus notopterus*, Pallas 1769) is a freshwater fish species native to Indonesia. *N. notopterus* is one of the fish species of the Notopteridae family<sup>1</sup>. The Notopteridae consist of approximately ten species in four genera: Chitala, Notopterus, Xenomystus, and Papyrocranus<sup>2</sup>. The only recognized species of Notopterus at present is *N. notopterus*<sup>3</sup>. The *N. notopterus* distribution in Asia includes Indonesia, India, Malaysia, Thailand, Bangladesh, Pakistan, and Vietnam<sup>1,2,4-7</sup>. In Indonesia, this species is distributed in Java, Sumatra, and Kalimantan<sup>1,8</sup>. *N. notopterus* was found in Kolong-Bendungan Simpung, Bangka Island<sup>9</sup>, Ogan River, Kelekar River (tributary of Musi River, South Sumatra), waters in Bangka Island as well as Kota Bangun and Tanah Ulu (Mahakam River, East Kalimantan)<sup>10</sup>, Sail River, Pekanbaru City, Riau Province<sup>11</sup>, Sei Gesek Reservoir, Bintan, Riau Islands<sup>12</sup>, and Kelekar River<sup>13</sup>.




Several studies on *N. notopterus* have been conducted, among others, on aspects of length-weight relationship, relative condition factor and fecundity<sup>14-17</sup>, genetic diversity<sup>10</sup>, morphometrics and growth patterns<sup>11</sup>, food and feeding habits<sup>12</sup>, reproductive biology<sup>9</sup>, determination of condition factors, somatic condition, hepatosomatic index and gonadosomatic index<sup>18</sup>, behavior and reproduction<sup>19</sup>, potential biomarkers of oxidative stress<sup>20</sup>, embryo and larval development<sup>21</sup>, relative condition factors<sup>22,23</sup>, population dynamics<sup>5</sup>, relationship of ecological factors and commercial<sup>24</sup>, and its early domestication<sup>25,26</sup>.

Fishery management mainly depends on knowledge of fish reproductive biology<sup>27,28</sup>. In adult fish, sexual dimorphism is a frequent and well-known phenomenon<sup>29</sup>. The majority of fishes are bisexual and have various sexually dimorphic characteristics in their secondary sexual characteristics<sup>30</sup>. Distinguishing characteristics for the morphological qualities of many teleosts have been documented, including body color, ornamental patterns, breeding tubercles on the snout, pectoral fins and scales, fin hooks, body shape, fin size and shape<sup>31-35</sup>.

*N. notopterus* is one of the most valuable freshwater fish in South Sumatra, Indonesia. This fish is utilized by the community in South Sumatra, especially as a daily side dish and a raw material for the culinary industries of pempek (fish cake) and kerupuk-kemplang (fish crackers). Therefore, this species has the potential for cultivation. There is relatively little known about sexual dimorphism and reproductive biology in *N. notopterus*. Such information is crucial for initial data in the breeding of *N. notopterus*. The study aimed to analyze the sexual dimorphism and reproductive biology of *N. notopterus* from Sungai Kelekar, Ogan Ilir. The results of this study are beneficial in the selection of *N. notopterus* broodstock candidates for breeding programs.

## Materials and Methods

The fish collection was conducted in Kelekar River, Tanjung Baru Village, North Indralaya District, Ogan Ilir Regency, South Sumatra (Fig 1). Sampling was conducted in February–May 2023. The temperature, water acidity, and dissolved oxygen at the sampling site ranged from 24.4°C to 28.3°C, 5.5 to 6.4, and 4.9 to 6.4 mg.L<sup>-1</sup>, respectively. Fishing was conducted by fishermen using fish traps. A total of 74 *N. notopterus* samples were collected: 38 males (TL: 18–23.6 cm; BW: 35.1–92.1 g) and 36 females (TL: 19.6–26.3 cm; BW: 49.4–133.8 g). The *N. notopterus* samples were transported to the Fisheries Basic Laboratory, Aquaculture Study Program, Department of Fisheries, Faculty of Agriculture, Universitas Sriwijaya, for analysis of morphometric-meristic characters and also aspects of reproductive biology.



**Figure 1. Map of *Notopterus notopterus* (Pallas 1769) sample collection in Kelekar River, Ogan Ilir Regency, South Sumatra, Indonesia**

In the laboratory, fish were weighed using a digital scale (accuracy 0.01 gram). Then observe the fish morphologically and count the meristic characters (number of rays of the dorsal fin, number of rays of the pectoral fin, and number of rays of the anal fin). Furthermore, the samples were measured for morphometric characters using a caliper (accuracy 0.01 cm) and a ruler (accuracy 0.1 cm). The morphometric characters measured were seventeen characters (Fig 2). Next, the fish were dissected, and the gonads and liver were taken, separated from other organs. Gonads and liver were weighed for the calculation of gonadosomatic index (IGS) = [gonad weight (g) / fish weight (g) x 100] and hepatosomatic index (IHS) = [liver weight (g) / fish weight (g) x 100]<sup>36</sup>. The calculation of fecundity was done gravimetrically, namely, the fish gonads were taken partially, and then the number of eggs in the partial gonads was calculated. Fecundity = [number of sample gonad eggs x (total gonad weight/partial gonad weight)]<sup>37</sup>. Oocyte diameter was observed under a digital microscope. The maturity level of fish gonads was categorized based on the morphology and color of the gonads, IGS, and oocyte diameter. The sex ratio was calculated based on the number of male and female fish. The data obtained were grouped into two categories, namely male and female fish. Data tabulation using the Microsoft Excel program was then analyzed descriptively to obtain differences between male and female fish.




Figure 2. Morphometric character measurements of bronze featherback (*Notopterus notopterus*, Pallas 1769): TL (total length), SL (standard length), M-O (mouth to operculum distance), M-DF (mouth to dorsal fin distance), DF-C (dorsal fin to caudal distance), DFL (dorsal fin length), PFL (pectoral fin length), AF-CFL (anal fin to caudal fin length), BH (body height), HL (head length), HH (head height), ED (eye diameter), DTBH (Degree of Tilt of the Back of the Head), BT (body thickness), AT (abdominal thickness), HM (height of mouth), WM (width of mouth)

## Results and Discussion

A total of 74 *N. notopterus* samples, consisting of 38 males and 36 females, have been successfully measured for morphometric characters and calculated for meristic characters. The results of measurements and calculations of *N. notopterus* samples in this study are presented in Table 1. Morphological differences between male and female *N. notopterus* are presented in Fig 3.

**Table 1. Morphometric and meristic characters of male and female *Notopterus notopterus* (Pallas, 1770) from Kelekar River, Ogan Ilir Regency, South Sumatra, Indonesia**

Morphometric character	Male		Proportion (%)	Female		Proportion (%)
	Range (cm)	AVG±STD		Range (cm)	AVG±STD	
TL	18-23.6	20.39±1.39	-	19.6-26.3	22.46±1.56	-
SL	16-21.6	18.60±1.45	91.25 TL	17.3-24.2	20.45±1.59	91.07 TL
HL	1.3-3.5	2.07±0.51	10.18 TL	1.6-3.7	2.16±0.47	9.64 TL
HH	2.1-3.5	2.63±0.38	12.90 TL	2.4-4.3	2.85±0.39	12.70 TL
ED	0.5-0.7	0.64±0.06	30.91 HL	0.8-1.0	0.86±0.06	39.81 HL
BH	3.1-5.1	4.29±0.44	21.07 TL	4.8-6.4	5.62±0.50	25.05 TL
PFL	2.1-3.3	2.53±0.35	12.44 TL	2.2-3.6	2.73±0.37	12.16 TL
DFL	1.7-2.8	2.19±0.27	10.77 TL	1.9-3.2	2.33±0.30	10.38 TL
AF-CFL	13.2-18.4	15.42±1.22	75.62 TL	14.0-19.1	16.44±1.36	73.20 TL
M-O	3.0-4.7	3.65±0.40	17.91 TL	3.0-4.9	3.90±0.44	17.40 TL
M-DF	8.1-11.5	9.75±0.68	47.85 TL	9.5-12.3	10.58±0.80	47.14 TL
DF-C	8.0-11.6	9.85±1.02	48.32 TL	8.5-12.5	10.94±0.98	48.72 TL
BT	1.1-1.3	1.19±0.08	-	1.4-1.9	1.51±0.14	-
AT	0.6-1.0	0.89±0.11	-	1.1-1.6	1.26±0.13	-
MH	0.8-1.2	1.02±0.09	-	0.8-1.2	1.05±0.09	-
MW	0.8-1.1	0.96±0.10	-	0.8-1.2	1.00±0.11	-
DTBH (°)	32.0-50.0	42.36±3.51	-	47.0-63.0	52.6±3.06	-
<b>Meristic character</b>	<b>Number of rays of the male fish</b>		<b>Number of rays of the female fish</b>			
Pectoral (P)	10-14		11-14			
Dorsal (D)	6-9		6-9			
Anal+Caudal (A+C)	107-118		110-118			

The total length of *N. notopterus* used in this study ranges from 18 cm to 23.6 cm; body weight ranges from 35.1 g to 92.1 g for male fish; The length ranges from 19.6 cm to 26.3 cm; and body weight ranges from 49.4 g to 33.8 g for female fish. According to the study results of <sup>9</sup>, the total length of female fish ranged from 130 mm to 249 mm, males from 120 mm to 232 mm, and the body weight of males was 22.17–97.17 g and females 38.98–120.47 g. The total length of *N. notopterus* ranged from 30 mm to 249 mm. The total length of *N. notopterus* ranged from 30 mm to 249 mm. The total length of *N. notopterus* ranges from 401–950 mm <sup>10</sup>. The male *N. notopterus* measured between 19.1 cm and 24.1 cm in length and weighed between 48.7 g and 133.4 g, while the female measured between 17.2 cm and 25.6 cm in length and weighed between 69.7 g and 151.2 g <sup>16</sup>.

The dorsal fin of *N. notopterus* fused with the caudal fin and anal fin. Each fin contains only rays and no spine. Meristic characters have been successfully counted. The results showed that P.10-14, D.6-9, and A+C.107-118. The results of the study by <sup>38</sup> showed that P.11-14, D.5-7, and A.98-111. According to <sup>11</sup> study, the rays of the P. 11-14, D. 6-7, and A. 99-111. According to <sup>39</sup>, the number of rays on the P.13-14, D.7-9, and A.97-111. However, <sup>40</sup> stated that D.7-9, and A.97-111. According to <sup>8</sup>, P.13-14, and A. 99-111.

One of the vital physiological processes that is essential to the life cycle of any organism, including fish, is reproduction <sup>41</sup>. Planning improved conservation and management strategies for fishery resources requires a fundamental understanding of the reproductive cycle of fish <sup>28,42</sup>. Some aspects of *N. notopterus* reproductive biology observed in this study are presented in Table 2.

**Table 2. Reproductive aspects of male and female *Notopterus notopterus* (Pallas, 1770) from Kelekar River, Ogan Ilir Regency, South Sumatra, Indonesia**

Parameters	Sampling Period							
	February		March		April		May	
	Male	Female	Male	Female	Male	Female	Male	Female
GSI (%)	0.31±0.05	5.18±1.04	0.39±0.11	4.50±1.16	0.30±0.14	3.22±1.69	0.28±0.06	3.59±2.57
HSI (%)	0.65±0.22	0.53±0.10	0.52±0.19	0.51±0.22	0.79±0.23	0.80±0.36	0.79±0.27	0.73±0.19
Ova diameter (mm)	-	1.2-3.6	-	1.1-3.9	-	1.2-3.9	-	1.2-3.8
Fecundity	-	1361-3543	-	1455-3121	-	1060-1471	-	1007-3901
Sex ratio	1:1		1.14:1		1.5:1		0.86:1	

The GSI utilized to ascertain the fish's reproductive periodicity<sup>43</sup>. Table 2 shows that the GSI value of female fish was highest in February, and the male fish was highest in March. The GSI of female fish is higher than that of male fish. The GSI value is also related to the stage of gonad maturity. The GSI value increases with increasing gonad maturity. The increase in GSI indicates a developmental process in the gonads. The GSI value will reach a maximum value before spawning occurs. In addition, the GSI value is influenced by environmental factors related to the food availability as an energy source for somatic development and fish reproduction<sup>44</sup>.

HSI values for female fish were highest in April and for males in May. HSI is the percentage value of the ratio between liver weight and total body weight. One of the organs that plays a role in fish reproduction is the liver. The liver plays an important role in the vitellogenesis process. This process is triggered by the increasing concentration of estradiol in the blood. Estradiol is an important hormone produced by the ovaries in female fish that is beneficial in the process of vitellogenesis. The higher the level of estradiol concentration in the blood, the earlier the liver will carry out the vitellogenesis process, and gonad maturation will be faster<sup>45,46</sup>. The HSI indicated an energy reserve in the liver<sup>47</sup> and a bio-indicator of contaminant exposure<sup>48,49</sup>. The fecundity of *N. notopterus* in this study ranged from 1,007 to 3,901 eggs per individual.<sup>50</sup> showed that fecundity values ranged from 53 to 748 and from 175 to 4,494<sup>51</sup>. The value of fecundity is influenced by several factors, one of which is environmental. Variations in the number of fish fecundities are also caused by variations in the length and weight of fish.

The diameter of the ova of *N. notopterus* in this study ranged from 1.1 mm to 3.9 mm. Previous studies ranged from 1.05 to 2.2 mm<sup>9</sup>. The oocyte diameter is categorized as a mature ovary. The higher the oocyte diameter value, the higher the ovary's maturity. The results showed that the diameter of the oocyte in the ovary varied. This indicates that the development of oocytes in the ovary is not uniform. It is suspected that *N. notopterus* spawns gradually. This is by the opinion of<sup>9</sup> that *N. notopterus* releases mature oocytes gradually. *N. notopterus* spawns more than once during a spawning season. The sex ratio is an important component in the study of fish reproduction<sup>42,52</sup>. The balance of the number of male and female fish in a body of water affects the continued existence of a species. The results of this study show that the number of male fish is higher than the number of female fish, except in the May sampling period. This result indicates that the population of male fish is greater than that of female fish.

## Conclusion

There are differences in the morphometric characteristics of male and female *N. notopterus*. However, there are no differences in meristic characters. The gonadosomatic index of male fish ranged from 0.15% to 0.61%, and that of females ranged from 1.12% to 9.10%. The hepatosomatic index of male fish ranged from 0.19% to 1.38%, and females ranged from 0.15 to 1.23%; the sex ratio of the sample fish obtained was 1.056:1. The fecundity value of the sampled fish ranged from 1,007 to 3,901. The oocyte diameter of the

sample fish ranged from 1.1 to 3.9 mm. The gonadal maturity stage of the sample fish was at the developing to mature stage for males and mature to spawning for females.

### Acknowledgment

We appreciate the fisherman (Mr. Nur and Mr. Lukman) who assisted with the fish collection. We also extend thanks to Universitas Sriwijaya's Rector and Head of LPPM. The research of this article was funded by DIPA of the Public Service Agency of Universitas Sriwijaya 2023. Nomor SP DIPA-023.17.2.677515/2023, On November 30, 2022. Rector's Decree Number: 0188/UN9.3.1/SK/2023, On April 18, 2023.

### Authors' Declaration

- Conflicts of Interest: None.
- We hereby confirm that all the Figures and Tables in the manuscript are ours. Furthermore, any Figures and images, that are not ours, have been included with the necessary permission for re-publication, which is attached to the manuscript.
- Authors sign on ethical consideration's approval
- Ethical Clearance: The project was approved by the local ethical committee in Universitas Sriwijaya

### Authors' Contribution Statement

MM in conception, design, acquisition of data, analysis, interpretation, drafting the manuscript, revision and proofreading, MS in acquisition of data, analysis interpretation, drafting the manuscript, revision and proofreading, FHT in conception, design, acquisition of data, drafting the manuscript, revision, MIS in acquisition of data.

### References

1. Kottelat M. The fishes of the inland waters of Southeast Asia: a catalogue and core bibliography of the fishes known to occur in freshwaters, mangroves and estuaries. *Raffles Bull Zool.* 2013; (27): 1–663. [https://lknhm.nus.edu.sg/wp-content/uploads/sites/10/app/uploads/2017/04/rbz\\_S27.pdf](https://lknhm.nus.edu.sg/wp-content/uploads/sites/10/app/uploads/2017/04/rbz_S27.pdf)
2. Barby F, Ráb P, Lavoué S, Ezaz T, Bertollo L, Kilian A, et al. From chromosomes to genome: Insights into the evolutionary relationships and biogeography of old. *Genes.* 2018; 9(6): 306. <https://doi.org/10.3390/genes9060306>.
3. Lavoué S, Ghazali SZ, Jamaluddin JA, Nor SA, Zain KM. Genetic evidence for the recognition of two allopatric species of Asian bronze featherback *Notopterus notopterus* (Teleostei, Osteoglossomorpha, Notopteridae). *Zoosyst Evol.* 2020 Jan 7; 96(2): 449-54. <https://doi.org/10.3897/zse.96.51350>.
4. Hilton EJ, Lavoué S. A review of the systematic biology of fossil and living bony-tongue fishes, Osteoglossomorpha (Actinopterygii: Teleostei). *Neotrop Ichthyol.* 2018 Oct 11; 16(3): e180031. <https://doi.org/10.1590/1982-0224-20180031>.
5. Mustafa MG, Singha S, Islam MR, Mallick N. Population dynamics of *Notopterus notopterus* (Pallas, 1769) from the Kaptai reservoir of Bangladesh. *SAARC J Agricult.* 2014; 12(2): 112-22. <https://doi.org/10.3329/sja.v12i2.21920>.
6. Naeem M, Salam A, Gillani Q, Ishtiaq A. Length-weight relationships of *Notopterus notopterus* and introduced *Oreochromis niloticus* from the Indus River, southern Punjab, Pakistan. *J Appl Ichthyol.* 2010 Aug; 26(4): 620-. <https://doi.org/10.1111/j.1439-0426.2010.01480.x>.
7. Srivastava SM, Singh SP, Pandey AK. Food and feeding habits of threatened *Notopterus notopterus* in Gomti river, Lucknow (India). *J Exp Zool India.* 2012; 15(2): 395–402.
8. Kottelat M, Whitten T. Freshwater fishes of Western Indonesia and Sulawesi: additions and corrections. Hong Kong: Periplus editions; 1996. 8 pp. [https://www.researchgate.net/publication/270508914\\_Freshwater\\_fishes\\_of\\_western\\_Indonesia\\_and\\_Sulawesi\\_additions\\_and\\_corrections](https://www.researchgate.net/publication/270508914_Freshwater_fishes_of_western_Indonesia_and_Sulawesi_additions_and_corrections).
9. Gustomi A, Sulistiono S, Yonvitner Y. Reproductive biology featherback (*Notopterus notopterus* Pallas, 1769) in

- Simpur Reservoir, Bangka Island. Indones. J Agric Sci. 2016; 21(1): 56-62. <https://doi.org/10.18343/jipi.21.1.56>
10. Wibowo A, Sunarno MT, Subagdja S, Hidayah T. Population characteristic of the putak (*Notopterus notopterus*) using fenotypic diversity analysis and 16sRNA region mitochondria DNA. J Lit Perikan Ind. 2017; 15(1): 1-2. <https://dx.doi.org/10.15578/jppi.15.1.2009.1-12>
  11. Purba EA, Efizon D, Putra RM. Study of morphometrics, meristics, and growth patterns of belida (*Notopterus notopterus*, Pallas 1769) in the Sail River of Pekanbaru City, Riau Province. JOM Faprika. 2017; 1: 1–8. <https://jom.unri.ac.id/index.php/JOMFAPERIKA/article/view/16180/15711>
  12. Rapita R, Susiana S, Kurniawan D, Lestari F, Sabriaty D, Rianti U. Food habits of belida fish (*Notopterus notopterus*, Pallas 1769) in Sei Gesek Reservoir, Bintan Regency, Riau Island, Indonesia. IOP Conf Seri: Earth Environ Sci. 2021 Nov 1; 919: 012003. <https://doi.org/10.1088/1755-1315/919/1/012003>
  13. Muslim M, Syaifudin M. Biodiversity of freshwater fish in Kelekar Floodplain Ogan Ilir Regency in Indonesia. J Trop Biodivers Biotechnol. 2022; 7(1): 67494. <https://doi.org/10.22146/jtbb.67494>.
  14. Kaur V, Rawal YK. Length-weight relationship (LWR) in *Notopterus notopterus* (Pallas) from Sukhna Lake, Chandigarh. IOSR J Pharm Biol Sci. 2017; 12(4): 63–65. <https://doi.org/10.9790/3008-1204046365>
  15. Kaushik KK, Sahu P, Nath R. Length-weight relationship of *Notopterus notopterus* (Pallas, 1769), *Anabas testudineus* (Bloch,1792) and *Clarias batrachus* (Linnaeus, 1758) from Pokoriya River, Morigaon, Assam, India. Int J Sci Technol Res. 2019; 8(10): 3087–3088. <https://doi.org/10.2307/1540>
  16. Paul MG, Nath P, Dutta A. Length-weight relationship, relative condition factor and fecundity of *Notopterus notopterus* (Pallas, 1769) from river Brahmaputra in Dhubri, Assam, India J Fish. 2022; 10(3): 103201. <https://doi.org/10.17017/j.fish.398>
  17. Winn NA, Sandi P, Khaing T, Thi K. Length weight relationship of twelve freshwater fish species from Sunny Lake, Mandalay Region, Myanmar. Greener J Biol Sci. 2021; 11(2): 74-80. <https://gjournal.org/GJBS/Publication/2021/2/PDF/092221092%20Nandar%20et%20al.pdf>
  18. Sudarshan S, Kulkarni RS. Determination of condition factor (K) somatic condition factor (Ks) hepatic and gonado somatic indices in the fresh water fish *Notopterus Notopterus*. Int J Sci Res. 2013; 2(11): 524-6.
  19. Sukendi S, Thamrin T, Putra RM, Yulindra A. Cultivation technology of bronze featherback (*Notopterus notopterus*, Pallas 1769) at different stocking densities and types of feed. IOP Conf Seri: Earth Environ Sci.. 2020; 430(1): 012027. <https://doi.org/10.1088/1755-1315/430/1/012027>
  20. Mohanty D, Samanta L. Multivariate analysis of potential biomarkers of oxidative stress in *Notopterus notopterus* tissues from Mahanadi River as a function of concentration of heavy metals. Chemosphere. 2016 Jul 1; 155: 28-38. <https://doi.org/10.1016/j.chemosphere.2016.04.035>
  21. Srivastava SM, Gopalakrishnan A, Singh SP, Pandey AK. Embryonic and larval development of threatened bronze featherback, *Notopterus notopterus* (Pallas). J Exp Zool India. 2012; 15(2): 425–30. <https://krishi.icar.gov.in/jspui/handle/123456789/4918>
  22. Achakzai WM, Saddozai S, Baloch WA, Soomro AN, Memon N. Length-weight relationship and condition factor of *Notopterus notopterus* (Pallas, 1769) from Manchar Lake Sindh, Pakistan. Sindh Univ Res Jour. 2015 Sep 15; 47(3): 515–18. <https://sujo.usindh.edu.pk/index.php/SURJ/article/view/5200>
  23. Kumar KH, Kiran BR. Relative condition factor of feather back, *Notopterus Notopterus* (Pallas) from Jannapura Pond, Bhadravathi Taluk, Karnataka. Int J Sci Res Sci Eng Technol. 2016; 2(1): 36-9. <https://ijsrset.com/IJSRSET162113>
  24. Kyaw MM, Kyaw MM, Lum MR, Naing SS. Relationship of ecological factors and commercial bronze featherback fish, *Notopterus notopterus* (Pallas, 1769); Irrawaddy River, Myitkyina Segment, Kachin State, Myanmar. Mandalay University, Research Journal. 2020; 11: 98–111.
  25. Muslim M, Simanjuntak WJ. Growth and survival of bronze featherback (*Notopterus notopterus*, Pallas 1769) reared on bucket. Magna Scientia Advanced Research and Reviews. 2023; 9(1): 101–05. <https://doi.org/10.30574/msarr.2023.9.1.0133>.
  26. Muslim M, Pitriani E, Agustina H. Growth and survival of bronze featherback (*Notopterus notopterus*) adapted on box container and aquarium. GSC Advanced Research and Reviews. 2023; 16(3): 133-37. <https://doi.org/10.30574/gscarr.2023.16.3.0368>
  27. Dinh QM, Tran NTT. Reproductive biological traits of the goby *Stigmatogobius pleurostigma* (Bleeker, 1849) from the Mekong Delta, Vietnam. J Indian Fish. 2018; 65(1): 20–25. <https://doi.org/10.21077/ijf.2018.65.1.68188-04>
  28. Parmaksız A, Oymak SA, Dogan N, Naim DM, Unlu E. Reproductive characteristics of an invasive species *Carassius gibelio* (Bloch, 1782) in Ataturk Dam Lake, Turkey. J Indian Fish. 2017 Jan 1; 64(4): 28-33. <https://doi.org/10.21077/ijf.2017.64.4.67478-04>
  29. Capasso LU, Del Re MC, Bravi SE, Taverne LP. Sexual dimorphism in a population of Pycnodontiformes of the Aptian (Lower Cretaceous) from Profeti (province of Caserta, Southern Apennines, Italy). Boll Mus Civ Stor Nat. Verona. 2021; 45: 31-41.
  30. Chacko JJ, Sekharan NM. Sexual dimorphism in structures, size and shape of the cyprinid Nilgiri melon barb. Fish

Comment [N1]: Not found Check again

Comment [N2]: Magna Scientia Adv Res Rev

Comment [N3]: GSC Adv Res Rev

- Aquat Life. 2022 Sep 30; 30(3): 138-48. <https://doi.org/10.2478/aopf-2022-0013>.
31. Ambus N, Moody EK. Size- based sexual dimorphism of nitrogen excretion in livebearing fishes. *Ecol Freshw Fish*. 2019 Apr; 28(2): 222-8. <https://doi.org/10.1111/eff.12445>
  32. Garcia EQ, Zuanon J. Sexual dimorphism in the electric knifefish, *Gymnorhamphichthys rondoni* (Rhamphichthyidae: Gymnotiformes). *Acta Amazonica*. 2019 Aug 12; 49: 213-20. <https://doi.org/http://dx.doi.org/10.1590/1809-4392201804392> original
  33. Mainero AH, Al-Jufaili SM, Jawad L, Reichenbacher B. Sex dimorphism and evidence of sexually selected traits: A case study on the killifish *Aphaniops stoliczkanus* (Day, 1872). *Acta Zoologica*. 2022 Aug 21; 104: 473–87. <https://doi.org/10.1111/azo.12436>
  34. Uba KI. Sexual shape dimorphism in the monomorphic fish *Decapterus macrosoma* (Teleostei: Carangidae). *Comput Ecol Softw*. 2019 Dec 1; 9(4): 134-42. [http://www.iaees.org/publications/journals/ces/articles/2019-9\(4\)/sexual-shape-dimorphism-in-monomorphic-fish-Decapterus-macrosoma.pdf](http://www.iaees.org/publications/journals/ces/articles/2019-9(4)/sexual-shape-dimorphism-in-monomorphic-fish-Decapterus-macrosoma.pdf)
  35. Ueng YT, Lin FJ, Chan YS, Tsao CW, Chen MJ. Sexual dimorphism, length-weight relationships, fecundity, and diet of the striped eel catfish *Plotosus lineatus* (Plotosidae) on Taiwan's southwest coast. *Nat Resour*. 2022 Mar 22; 13(3): 77-90. <https://doi.org/10.4236/nr.2022.133006>
  36. Jan M, Jan N. Studies on the fecundity (F), gonadosomatic index (GSI) and hepatosomatic index (HSI) of *Salmo trutta fario* (Brown trout) at Kokernag trout fish farm, Anantnag, Jammu and Kashmir. *Int J Fish Aquat Stud*. 2017; 5(6): 170-3. <https://www.fisheriesjournal.com/archives/2017/vol5issue6/PartC/5-6-17-154.pdf>
  37. Bishai HM, Ishak MM, Labib W. Fecundity of the mirror carp *Cyprinus carpio* L. at the Serow Fish Farm (Egypt). *Aquaculture*. 1974 Jan 1; 4: 257-65. [https://doi.org/10.1016/0044-8486\(74\)90038-6](https://doi.org/10.1016/0044-8486(74)90038-6)
  38. Rianti U, Susiana S, Kurniawan D. Morphometric and meristic characteristics of bronze featherback (*Notopterus notopterus*, pallas 1769) in Sei Gesek Reservoir Bintan Regency. *Bawal: Widya Riset Perikanan Tangkap*. 2021 Dec 6; 13(3): 123-32. <https://dx.doi.org/10.15578/bawal.13.3.2021.123-132>
  39. Mulyani I, Budijono B. Morphometric and meristic analysis of Asian knifefish (*Notopterus notopterus*) in Sail River, Pekanbaru Riau Province. *J-BEKH*. 2020 Dec 1; 7(2): 59-64. <https://doi.org/10.23960/jbekh.v7i2.156>
  40. Sudarto S. Flatfish that have potential for ornamental fish. *Media Akuakultur*. 2011 Jun 30; 6(1): 59-62.
  41. Muchlisin ZA. A general overview on some aspects of fish reproduction. *Aceh International Journal of Science and Technology*. 2014 Jan 1; 3(1):43-52. <https://doi.org/10.13170/AIJST.0301.05>
  42. Chelapurath RR, Kuttanelloor R, Balakrishna MK. Reproductive biology of the endemic cyprinid fish *Hypselobarbus thomassi* (Day, 1874) from Kallada River in the Western Ghats, India. *J Appl Ichthyol*. 2020 Oct; 36(5): 604-12. <https://doi.org/10.1111/jai.14064>
  43. Jewel MA, Haque MA, Ferdous MS, Khatun MS, Hasan J. Gonadosomatic index and fecundity of threatened reba carp, (Hamilton), in the Ganges River (northwest Bangladesh). *Fish Aquat Life*. 2019; 27(2): 80-5. <https://doi.org/10.2478/aopf-2019-0009>
  44. Kasmi M, Hadi S, Kantun W. Reproductive biology of Indian mackerel, *Rastreliger kanagurta* (Cuvier, 1816) in Takalar coastal waters, South Sulawesi. *J Iktiologi Indones*. 2017 Oct 23; 17(3): 259-71. <https://doi.org/10.32491/jii.v17i3.364>
  45. Garnayak SK, Mohanty J, Rao TV, Sahoo SK, Sahoo PK. Vitellogenin in Asian catfish, *Clarias batrachus*: Purification, partial characterization and quantification during the reproductive cycle by ELISA. *Aquaculture*. 2013 May 10; 392: 148-55. <https://doi.org/10.1016/j.aquaculture.2013.02.020>
  46. Sullivan CV, Yilmaz O. Vitellogenesis and yolk proteins, fish. *Encyclopedia of reproduction*. 2018 Jan 1; 6: 266-77. <https://doi.org/10.1016/B978-0-12-809633-8.20567-0>
  47. Pandit k, Gupta ML. Hepato-somatic index, gonado-somatic index and condition factor of *Anabas testudineus* as bio-monitoring tools of nickel and chromium toxicity. *Int J Innov Eng Technol*. 2019; 12(3): 25-8. <https://ijiet.com/wp-content/uploads/2019/03/5.pdf>
  48. Araújo FG, Morado CN, Parente TT, Paumgartten FJ, Gomes ID. Biomarkers and bioindicators of the environmental condition using a fish species (*Pimelodus maculatus* Lacepède, 1803) in a tropical reservoir in Southeastern Brazil. *Braz J Biol*. 2017 Aug 17; 78: 351-9. <https://doi.org/10.1590/1519-6984.167209>
  49. Singh S, Srivastava AK. Variations in hepatosomatic index (HSI) and gonadosomatic index (GSI) in fish *Heteropneustes fossilis* exposed to higher sub-lethal concentration to arsenic and copper. *J Ecophysiol Occup Health*. 2015 Dec 1; 15(3/4): 89-93. <https://doi.org/10.15512/joeoh/2015/v15i3-4/121589>
  50. Azadi MA, Islam MA, Nasiruddin M, Quader MF. Reproductive biology of *Notopterus notopterus* (Pallas) in Kaptai reservoir, Bangladesh. *Bangladesh J Zool*. 1995; 23: 215-20.
  51. Gupta S, Ray A. Review on biology and culture potential of *Notopterus notopterus* (Pallas, 1769). *Int J Aquat Biol*. 2022 Apr 1; 10(2):78-91. <https://ij-aquaticbiology.com/index.php/ijab/article/view/1511>.
  52. Purushottama GB, Ramasubramanian V, Dash G, Akhilesh KV, Ramkumar S, Kizhakudan SJ, et al. Reproductive biology and diet of the grey sharpnose shark *Rhizoprionodon oligolinx* Springer, 1964 (Chondrichthyes: Carcharhinidae) from the north-eastern Arabian Sea. *J Indian Fish*. 2017; 64(4): 9-20.



## إزدواج الشكل الجنسي وبيولوجيا الإنجاب لل *Bronze featherback*

(*Notopterus notopterus* ، Pallas 1769) من نهر كيليكار، أوغان إير، جنوب سومطرة، إندونيسيا

### الخلاصة



يعد ازدواج الشكل الجنسي وعلم الأحياء الإنجابي من الجوانب الأساسية لدراسات تربية الأسماك. كان الهدف من هذا البحث هو تحليل ازدواج الشكل الجنسي والبيولوجيا الإنجابية لـ *N. notopterus*. تم جمع ما مجموعه 74 *N. notopterus* من نهر كيليكار في منطقة أوغان إير، إندونيسيا، وتتكون من 38 ذكرًا (طول الجسم: 18-23.6 سم؛ وزن الجسم: 1.35-92.1 جم) و36 أنثى (طول الجسم: 19.6-26.3 سم؛ وزن الجسم: 1.35-92.1 جم) و36 أنثى (طول الجسم: 19.6-26.3 سم؛ وزن الجسم: 4.4-133.8 جم). تم تحليل سبعة عشر شخصية مورفومترية وثلاثة صفات مرسية وخمسة معايير بيولوجية إنجابية. أظهرت النتائج وجود اختلافات في الخصائص المورفومترية للذكور والإناث *N. notopterus*. ومع ذلك، لم يكن هناك اختلاف في الشخصيات المرسية. ويتراوح مؤشر الغدد التناسلية عند الذكور من 0.15 إلى 0.61%، وعند الإناث من 1.12 إلى 9.10%. تراوح المؤشر الكبدي عند الذكور من 0.19 إلى 1.38%، وتراوح عند الإناث من 0.15 إلى 1.23%، مع نسبة جنس 1:1.056، وخصوبة 1.007-3.901 بيضة، وأقطار البويضات 1.1-3.9 ملم، ومستويات النضج. من الغدد التناسلية المتطورة لتتضح عند الذكور وتتضح لوضع البيض عند الإناث. في الختام، يمكن تحديد الاختلافات بين الذكور والإناث *N. notopterus* شكلياً.

**الكلمات المفتاحية:** اختيار الأمهات؛ تكاثر الأسماك داء البنية. الخصائص الجنسية الثانوية

## Participants

Salah Abdulla Hasoon (profsalah)



Muslim Muslim (muslim\_unsri)

Messages	
Note	From
<p>Dear Sir</p> <p>Please correct or change reference (15) in your manuscript, correction must be done on the same file attached</p> <p>Regards</p> <p> <a href="#">9997-Article Text-92907-102560-18-20240122.docx</a></p>	<p>profsalah</p> <p>2024-01-22 08:40 PM</p>
<p>▶ Dear Sir</p> <p>We resubmitted the corrected manuscript, and reference (10) has been changed.</p> <p>Best Regards</p> <p></p>	<p>muslim_unsri</p> <p>2024-01-23 03:42 AM</p>

## Participants

Salah Abdulla Hasoon (profsalah)

Muslim Muslim (muslim\_unsri)

Messages	
Note	From
<p>Dear Sir</p> <p>Please correct or change reference (15) in your manuscript, correction must be done on the same file attached</p> <p>Regards</p> <p> <a href="#">9997-Article Text-92907-102560-18-20240122.docx</a></p>	<p>profsalah</p> <p>2024-01-22 08:40 PM</p>
<p>▶ Dear Sir</p> <p>We resubmitted the corrected manuscript, and reference (10) has been changed.</p> <p>Best Regards</p> <p></p>	<p>muslim_unsri</p> <p>2024-01-23 03:42 AM</p>



Muslim &lt;muslim\_bda@unsri.ac.id&gt;

---

**[BSJ] Primarily acceptance Letter**

2 messages

---

**Prof. Dr. Fikrat M. Hassan** <bsj-info@csw.uobaghdad.edu.iq>  
Reply-To: "Prof. Dr. Fikrat M. Hassan" <fikrat@csw.uobaghdad.edu.iq>  
To: Muslim Muslim <muslim\_bda@unsri.ac.id>

Tue, Jan 23, 2024 at 10:32 AM

Dear Muslim Muslim,

We have reached a decision regarding your submission to Baghdad Science Journal, "Sexual dimorphism and reproductive biology of bronze featherback (Notopterus notopterus Pallas 1769) from Kelekar River, Ogan Ilir, South Sumatra, Indonesia".

Our decision is to: a primarily accept the submission.

Please, complete the fee publication, for more information on [Fees of Publication](#)

The publication fee is \$160

Prof. Dr. Fikrat M. Hassan  
[fikrat@csw.uobaghdad.edu.iq](mailto:fikrat@csw.uobaghdad.edu.iq)

---

**Baghdad Science Journal,**

Al- Jadirya, Baghdad P.O. BOX 4732. Baghdad-Iraq

E-mail : [bsj@csw.uobaghdad.edu.iq](mailto:bsj@csw.uobaghdad.edu.iq)

Mobile :+964 780 070 9148

---




**Muslim Muslim** <muslim\_bda@unsri.ac.id>  
To: "Prof. Dr. Fikrat M. Hassan" <fikrat@csw.uobaghdad.edu.iq>

Wed, Jan 31, 2024 at 4:51 PM




We will pay the publication fee via paypal method. Please make an invoice. Thank you

[Quoted text hidden]

**Messages**

Note	From
<p>Dear Author,</p> <p>Hope this email finds you well. BSJ would like you to carefully check on the format of your paper according to the journal template. This includes the content of the tables and figures, as well as the corrections in the body of the paper itself.</p> <p>You are, also, kindly required to correct all the comments written in the manuscript.</p> <p>Please take note that your approval after the proofreading cannot be changed. Moreover, it is to be noticed that publishing the manuscript will be delayed into later issues in case the correction is not sent within <b>2 days</b> from the date of our sending this message.</p> <p>Kind regards.</p> <p> <a href="#">animal welfare statement.docx</a></p> <p> <a href="#">Certification of Proof after layout edit.docx</a></p> <p> <a href="#">10-9997-Article Text-93075-102730-11-20240123 (1).docx</a></p>	<p>bsj 2024-03-12 11:03 AM</p>
<p>▶ Dear Chief in Editor,</p>	<p>muslim_unsri</p>


**Messages**

Note	From
<p>Dear Author,</p> <p>Hope this email finds you well. BSJ would like you to carefully check on the format of your paper according to the journal template. This includes the content of the tables and figures, as well as the corrections in the body of the paper itself.</p> <p>You are, also, kindly required to correct all the comments written in the manuscript.</p> <p>Please take note that your approval after the proofreading cannot be changed. Moreover, it is to be noticed that publishing the manuscript will be delayed into later issues in case the correction is not sent within <b>2 days</b> from the date of our sending this message.</p> <p>Kind regards.</p> <p> <a href="#">animal welfare statement.docx</a></p> <p> <a href="#">Certification of Proof after layout edit.docx</a></p> <p> <a href="#">10-9997-Article Text-93075-102730-11-20240123 (1).docx</a></p>	<p>bsj 2024-03-12 11:03 AM</p>
<p>▶ Dear Chief in Editor,</p>	<p>muslim_unsri</p>

## Participants

Baghdad Science Journal (bsj)

Muslim Muslim (muslim\_unsri)


Messages	
Note	From
Dear Author, Herewith the attached file we send the acceptance letter. Best regards   <a href="#">22 9997.pdf</a>	bsj 2024-05-02 12:52 PM

[Add Message](#)

## Participants

Baghdad Science Journal (bsj)

Muslim Muslim (muslim\_unsri)

Messages	
Note	From
Dear Author, Herewith the attached file we send the acceptance letter. Best regards	bsj 2024-05-02 12:52 PM
 <a href="#">22 9997.pdf</a>	

[Add Message](#)