

6. Growth and Survival of Bronze Featherback (*Notopterus notopterus*) Adapted On Box Container and Aquarium *by Muslim Muslim*

Submission date: 16-Jun-2024 05:37AM (UTC+0700)

Submission ID: 2402902068

File name: Notopterus_notopterus_Adapted_On_Box_Container_and_Aquarium.pdf (1.06M)

Word count: 2729

Character count: 14159



(RESEARCH ARTICLE)



5

Growth and survival of bronze featherback (*Notopterus notopterus*) adapted on box container and aquarium

Muslim MUSLIM *, Elika PITRIANI and Heni AGUSTINA

¹⁵ *Aquaculture Study Program, Department of Fisheries, Faculty of Agriculture, Universitas Sriwijaya, Indralaya, South Sumatra, Indonesia.*

GSC Advanced Research and Reviews, 2023, 16(03), 133–137

Publication history: Received on 30 July 2023; revised on 12 September 2023; accepted on 14 September 2023

Article DOI: <https://doi.org/10.30574/gscarr.2023.16.3.0368>

Abstract

The adaptation of wild fish to the cultured environment is basic information in fish farming. The aim of this study was to adapt *N. notopterus* to box containers and aquaria. This study consisted of two experiments. Experiment 1 adapted *N. notopterus* to a box container with different stocking densities of 1 fish and 2 fish per box container. Experiment 2 adapted *N. notopterus* to an aquarium with different feeding rates of 1% and 5%. The results showed that *N. notopterus* successfully adapted to the controlled environment of a box container and aquarium. *N. notopterus* survived and grew in box containers and aquaria. The results of this study provide basic information for the cultivation of *N. notopterus*.

Keywords: Aquaculture system; Aquarium fish; Fish adaptation; Notopteridae

1 Introduction

16

Aquaculture is the breeding, raising, and harvesting of fish, shellfish, and aquatic plants. Basically, it's farming with water. According to [1], aquaculture is the process of raising, reproducing, and harvesting aquatic species, including animals and plants, in regulated aquatic habitats such as oceans, lakes, rivers, ponds, and streams. Some aquaculture production systems are suitable for freshwater environments, such as land-based freshwater ponds, tarpaulin ponds, running-water ponds, waring cages, floating cages, pen-culture, cement ponds, aquaria, box containers, buckets, and others, while others are suitable for marine environments. Production systems differ in their setup and the species they are suitable for.

18

The bronze featherback (*Notopterus notopterus*, Pallas 1770) is the only fish species of the genus *Notopterus* [2]. This species belongs to the family *Notopteridae*, a species of fish with fan-shaped fins (Class *Actinopterygii*). The distinctive feature of this species is its knife-shaped back. This species lives in fresh or brackish water. This species has a short dorsal fin, a long caudal fin, and small pelvic fins [3]. South and south-east Asian nations, including Bangladesh, Malaysia, Thailand, Myanmar, Indonesia, India, and Viet Nam, are home to *N. notopterus* [4-8]. Due to overexploitation, pollution, and habitat degradation, the wild population of *N. notopterus* has decreased recently [9,10]. The Indonesian government has listed *N. notopterus* as an endangered species because of its declining population in the wild [11].

In Indonesia, fish is still a staple diet and a major source of protein, which is easily digestible and has high biological value. *N. notopterus* is one of Indonesia's freshwater fish species that has high economic value [12]. This species is used by Indonesian people as a daily side dish and is processed into culinary specialties, namely pempek (fish cake) and kerupuk-kemplang (fish cracker). The increasing demand for this fish has led to increased fishing in the wild. As a result, its population in the wild is declining. In order to preserve this species, it needs to be domesticated. Early domestication is aimed at improving fish survival and growth in aquaculture environments. The adaptation of wild fish to the

¹¹ Corresponding author: Muslim MUSLIM

Copyright © 2023 Author(s) retain the copyright of this article. This article is published under the terms of the Creative Commons Attribution License 4.0.

aquaculture environment can cause physiological disorders of growth and development. In the media, cultivation can cause fish not to survive and not to grow. This study aimed to adapt *N. notopterus* in container boxes and aquaria. The results of this study provide basic information for the aquaculture of *N. notopterus*.

2. Material and methods

2.1. Experiment 1 Bronze featherback (*N. notopterus*) adapted on box container

2.1.1. Preparation of the box containers

The rearing container uses 12 box containers (50x32x30 cm). The box containers are black in color. The box containers were cleaned using dug-well water to remove embedded dirt. After cleaning, the box containers were dried, and after drying, they were filled with water as high as 20 cm. The water in the tubs was deposited for 24 hours. The box containers are placed in an open space in the backyard. The box containers are protected with a net to prevent fish-eating animals from entering.

2.1.2. Fish rearing

N. notopterus is put into the box containers in the morning when the weather conditions are not yet hot, so that the fish placed in the box containers do not experience stress. Before being put into the box containers, the fish were weighed using a digital scale (accuracy 0.01 g), and the total length was measured using a ruler (accuracy 0.1 cm). The measurement data of body weight and total length were used as initial data for the experiment. Next, acclimatization was carried out for 15 minutes. Fish rearing is carried out for 30 days. The stocking density of fish in box containers, namely 1 fish per box container and 2 fish per box container, was repeated five times. The feed given is in the form of small swamp shrimp (*Caridina gracillirostris*; Atyidae). Ad libitum feeding. The frequency of feeding is done twice a day at 08.00 and 17.00 WIB. Sampling was carried out at the beginning and end of the experiment to determine fish growth.

2.1.3. Measured parameters

Total length growth

The total length growth was calculated using the formula: $L = L_t - L_0$ (Formula 1). where L is the growth in total length of the fish (cm), L_t is the length of the fish at the end of rearing (cm), and L_0 is the length of the fish at the beginning of rearing (cm).

Absolute weight growth

Absolute weight growth was calculated using the formula: $W = W_t - W_0$ (Formula 2). where W is the absolute weight growth of the experimental fish (g), W_t is the weight of the experimental fish at the end of the rearing period (g), and W_0 is the weight of the fish at the beginning of the rearing period (g).

Specific growth rate

A specific growth rate was calculated using the formula: $SGR = \frac{[\ln W_t - \ln W_0]}{t} \times 100$ (Formula 3). where SGR is the specific growth rate (%), W_t is the average weight of fish at the end of the rearing period (g), W_0 is the average weight of fish at the beginning of the rearing period (g), and t is the number of days of rearing (days).

Survival rate

Survival rate was calculated using the formula: $SR = \frac{N_t}{N_0} \times 100$ (Formula 4). Where, SR is the survival rate of fish reared (%), N_t is the number of fish at the end of the rearing period (fish), and N_0 is the number of fish at the beginning of the rearing period (fish).

Water quality

The water quality parameters measured were water temperature, pH, and ammonia. The measurements are carried out periodically. Temperature and pH measurements are carried out daily, while ammonia is measured weekly.

2.2. Experiment 2 Bronze featherback (*N. notopterus*) adapted on aquarium

2.2.1. Preparation of the aquaria

N. notopterus were reared in an aquaria (34x39x25 cm). A total of 12 aquaria were used in this study. The aquarium used is made of glass with a thickness of 5 mm. The aquarium is cleaned of dirt and dust by washing it with clean water. After cleaning, the aquarium is drained and then filled with 20 liters of clean water. Aeration installations were installed in each aquarium. Each aquarium is covered with black plastic on the outside.

2.2.2. Fish rearing

Before *N. notopterus* was put into the aquarium, the fish were weighed using a digital scale (accuracy 0.01 g), and the total length was measured using a ruler (accuracy 0.1 cm). The measurement data of body weight and total length were used as initial data for the experiment. Next, acclimatization was carried out for 15 minutes. Fish rearing is carried out for 30 days. Sampling was carried out at the beginning and end of the experiment to determine fish growth.

2.2.3. Feeding

During the rearing period, *N. notopterus* is given natural food, namely small swamp shrimp (*Caridina gracillirostris*). The feeding rate was between 1% and 5% of the total fish biomass that was maintained. Every week, the body weight of the test fish is measured as a basis for calculating the amount of food that will be given in the next week. The frequency of feeding is twice a day, namely at 08.00 and 16.00 WIB.

2.2.4. Measured parameters

Parameters measured included the total length growth of the experimental fish, absolute body weight growth, specific growth rate, feed efficiency, and survival rate of the experimental fish. The calculation of these parameters refers to the formula used in experiment 1.

3. Results and discussion

3.1. Experiment 1 Bronze featherback (*N. notopterus*) adapted on box container

The results showed that *N. notopterus* reared in box containers successfully survived and grew. The growth of trial fish is reflected in the increase in total length and body weight. The experimental results of rearing *N. notopterus* in box containers are presented in Table 1.

Table 1 Experimental results of rearing *N. notopterus* in box containers

Parameters	Stocking density	
	1 fish/box	2 fish/box
Initial total length (cm)	11.50	9.20
Final total length (cm)	12.94	11.50
Growth in length (cm)	1.44	2.30
Initial body weight (g)	12.00	16.00
Final body weight (g)	13.50	20.00
Growth in body weight (g)	1.50	4.00
Specific growth rate (%)	5.00	13.3
Survival rate (%)	100	100
Water quality		
Temperature (°C)	27-29.7	27-29.7
pH (unit)	6.0-7.2	6.85-7.92
Dissolved oxygen (mg/L)	4.5-6.9	5.3-7.8
Ammonia (mg/L)	0.041-0.055	0.043-0.055

3.2. Experiment 2. Bronze featherback (*N. notopterus*) reared on aquaria

The results showed that *N. notopterus* successfully survived and grew in the aquarium (Table 2).

Table 2 Experimental results of rearing *N. notopterus* in aquaria

Parameters	Feeding rate	
	1%	5%
Initial total length (cm)	3.92	4.68
Final total length (cm)	4.14	5.17
Growth in length (cm)	0.22	0.49
Initial body weight (g)	7.86	10.81
Final body weight (g)	8.55	11.96
Growth in body weight(g)	0.69	1.15
Specific growth rate (%)	2.3	3.8
Survival rate (%)	100	100
Water quality		
Temperature (°C)	26–28.9	26–28.2
pH (unit)	6.6–8.0	6.5–7.9
Dissolved oxygen (mg/L)	4.9–7.5	6.0–7.5
Ammonia (mg/L)	0.041–0.050	0.041–0.055

Growth is a physiological process that involves both internal and external physiological factors, such as health, stress, and reproductive status, as well as environmental factors like food quality and quantity, temperature, and water quality. The survival of young fish is frequently correlated with their growth. Growth is the steady growth in average fish weight, which for many species can be depicted by an asymptotic sigmoid curve [13]. Thus, the observed increase can be approximated by exponential, linear, or asymptotic functions depending on the start and end times of an experiment. Aquaculturists often report growth using one of three metrics: absolute growth rate, relative growth rate, and specific growth rate. Aquaculture fish grow through a complicated mechanism that transforms the energy they consume into biomass. The organism's capacity for growth, its trophic level, and a number of abiotic factors, including food availability, temperature, and unfavorable environmental effects brought on by the conditions under which the fish are raised, all influence how effectively this conversion occurs [14].

Several ecological variables, including temperature, dissolved oxygen, salinity, and photoperiod, have an impact on how quickly something grows. Water temperature, food availability, and environmental stressors like decreased oxygen or increased sedimentation all affect the rates at which fish and their scales grow [15]. One of the physical parameters that has the greatest impact on fish production and growth is water temperature [16,17]. The health and subsequent growth of fish are directly related to the quality of water in which the fish are raised. In general, factors affecting fish growth and production in freshwater aquatic systems can be classified as physical, chemical/biochemical, or a combination thereof. The physical properties of water that are important to fish production and growth include temperature and the concentrations of suspended and settleable solids; important chemical parameters include pH, alkalinity, hardness, and metals [17].

4. Conclusion

The adaptation of *N. notopterus* to container boxes and aquaria has been successful. *N. notopterus* was adaptive to the cultural environment so that it could successfully survive and grow. The results of this study are useful as basic information for *N. notopterus* aquaculture.

Compliance with ethical standards

Acknowledgments

We would like to thank Mr. Lukman, a fisherman in Tanjung Baru Village, Ogan Ilir Regency for providing the test fish for this study. We also express our gratitude to the students who have helped us in this study.

Disclosure of conflict of interest

We declare that there is no conflict of interest in this study.

References

- [1] Subasinghe R, Soto D, and Jia J. Global aquaculture and its role in sustainable development. *Reviews in aquaculture*. 2009;1(1): 2-9.
- [2] Ng, H.H. *Notopterus notopterus*. 2020; e.T166433A60584003. doi:10.2305/IUCN.UK.2020-1.RLTS.T166433A60584003.en
- [3] Froese, Rainer, and Pauly. Species of *Notopterus* [Internet]. Fishbase, @ 2023 [cited 2023 Sep 05] Available from <https://www.fishbase.se/summary/Notopterus-notopterus.html>
- [4] Azadi M, Islam M, Nasiruddin M, and Quader M.. Reproductive biology of *Notopterus notopterus* (Pallas) in Kaptai reservoir, Bangladesh. *Bangladesh Journal of Zoology*. 1995; 23: 215–220.
- [5] Mustafa MG, Singha S, Islam M, and Mallick N. Population dynamics of *Notopterus notopterus* (Pallas, 1769) from the Kaptai Reservoir of Bangladesh. *SAARC Journal of Agriculture*. 2014. 12(2): 112–122.
- [6] Paul MG, Nath P, and Dutta A. Length-weight relationship, relative condition factor and fecundity of *Notopterus notopterus* (Pallas, 1769) from river Brahmaputra in Dhubri, Assam, India. *Journal of Fisheries*. 2022; 10(3): 103201. <https://doi.org/10.17017/j.fish.398>
- [7] Wibowo A, Sunarno MTD, Subagja S, and Hidayah T. Population characteristic of the putak (*Notopterus notopterus*) using fenotypic diversity analysis and 16sRNA region mitochondria DNA. *Jurnal Penelitian Perikanan Indonesia*. 2009; 15(1): 1–12
- [8] Muslim M, and Syaifudin M. Biodiversity of freshwater fish in Kelekar Floodplain Ogan Ilir Regency in Indonesia. *Journal of Tropical Biodiversity and Biotechnology*. 2022; 7(1): 67494.
- [9] Memon KH, Memon AM, Ursani TJ, Shaikh AM, and Memon SW. Population parameters and exploitation status of clown knifefish *Notopterus chitala* (Hamilton, 1822) from Indus River, Pakistan. *Pakistan Journal of Zoology*. 2022; 54(2): 889–899.
- [10] Yulindra A, Lumbantoruan RP, Zulkifli Z, and Sukendi S. Effect of granting ovaprim with different dosage to ovulation and eggs quality of knife fish (*Notopterus notopterus*). *International Journal of Oceans and Oceanography*. 2017; 11(2): 189–199
- [11] Iskandar A, Muslim M, Hendriana A, and Wiyoto W. Jenis-jenis ikan Indonesia yang kritis dan terancam punah. *Jurnal Sains Terapan*. 2020;10(1): 53-59.
- [12] Sukendi S, Thamrin T, Putra R, and Yulindra A. Cultivation technology of bronze featherback (*Notopterus notopterus*, Pallas 1769) at different stocking densities and types of feed. *IOP Conf. Series: Earth and Environmental Science* 430. 2020: 012027. <https://doi.org/10.1088/1755-1315/430/1/012027>
- [13] Ricker WE. Growth rates and models. Pages 677-743 in Hoar WS, Randall DJ and Brett JR, editors. *Fish physiology*, volume VIII. Bioenergetics and growth. Academic Press, New York, USA. 1979.
- [14] Soderberg RW. Factors affecting fish growth and production. In *Dynamics of pond aquaculture* (pp. 199-213). CRC Press. 2017.
- [15] Dutta H. Growth in fishes. *Gerontology*. 1994; 40(2-4): 97-112.
- [16] Burel C, Person-Le Ruyet J, Gaumet F, Le Roux A, Severe A, and Boeuf G. Effects of temperature on growth and metabolism in juvenile turbot. *Journal of Fish Biology*. 1996; 49(4): 678-692.
- [17] Viadero R C. Factors affecting fish growth and production. *Water Encyclopedia*. 2005; 3: 129-133.

6. Growth and Survival of Bronze Featherback (Notopterus notopterus) Adapted On Box Container and Aquarium

ORIGINALITY REPORT

20%

SIMILARITY INDEX

13%

INTERNET SOURCES

11%

PUBLICATIONS

%

STUDENT PAPERS

PRIMARY SOURCES

1	ojs.unud.ac.id Internet Source	1%
2	pubs.sciepub.com Internet Source	1%
3	D F Putra, F Nur, S A E Rahimi, N Othman. " Effects of different live feed on growth and survival rate of clown loach, ", IOP Conference Series: Earth and Environmental Science, 2019 Publication	1%
4	Auste Asadauskas, Markus M. Luedi, Richard D. Urman, Lukas Anderegg. "Modern Approaches to the Treatment of Acute Facial Pain", Current Pain and Headache Reports, 2024 Publication	1%
5	mail.ijaers.com Internet Source	1%

6	<p>Wahyu Afrilasari, Widanarni, Anja Meryandini. "Effect of Probiotic Bacillus megaterium PTB 1.4 on the Population of Intestinal Microflora, Digestive Enzyme Activity and the Growth of Catfish (Clarias sp.)", HAYATI Journal of Biosciences, 2016</p>	1 %
Publication		
7	<p>Kevin D. Hopkins. "Reporting Fish Growth: A Review of the Basics", Journal of the World Aquaculture Society, 9/1992</p>	1 %
Publication		
8	<p>digitalarchive.worldfishcenter.org</p>	1 %
Internet Source		
9	<p>garuda.kemdikbud.go.id</p>	1 %
Internet Source		
10	<p>Carl F. Mazur, Donald Tillapaugh, John R. Brett, George K. Iwama. "The effect of feeding level and rearing density on growth, feed conversion and survival in chinook salmon (Oncorhynchus tshawytscha) reared in salt water", Aquaculture, 1993</p>	1 %
Publication		
11	<p>academicjournals.org</p>	1 %
Internet Source		
12	<p>link.springer.com</p>	1 %
Internet Source		

13	G N Susanto, Sutyarso, H Busman, N R Kurniawan, S M Hasanah. "Genital reversal of betta fish by immersion using steroid extract of sea urchins", IOP Conference Series: Earth and Environmental Science, 2021 Publication	1 %
14	bmcecolevol.biomedcentral.com Internet Source	1 %
15	jurnal.ugm.ac.id Internet Source	1 %
16	www.coursehero.com Internet Source	1 %
17	pjms.com.pk Internet Source	1 %
18	Arif Wibowo, Haryono Haryono, Kurniawan Kurniawan, Vitas Atmadi Prakoso et al. "Genetic and morphological evidence of a single species of bronze featherback (Notopterus notopterus) in Sundaland", Global Ecology and Conservation, 2023 Publication	1 %
19	Puspahati Kurnia Ghani, Dicky Harwanto, Rosa Amalia, Seto Windarto et al. "The Effect of Combination of Coconut Water (Cocos nucifera) and Shallot Crude Extract (Allium cepa L.) on the Growth of Caulerpa	1 %

racemosa", IOP Conference Series: Earth and Environmental Science, 2023

Publication

Exclude quotes On

Exclude matches < 1%

Exclude bibliography On