

Chemical Constituents from Stem Bark of *Flacourtia rukam* Zoll. & Mor. and Their Antioxidant Activities

(Kandungan Kimia daripada Kulit Batang *Flacourtia rukam* Zoll. & Mor. dan Aktiviti Antioksidannya)

MUHARNI, ELFITA*, HENI YOHANDINI, JULINAR, YASRINA & MIRANTI

ABSTRACT

Exploration of chemical compounds and biological activities of extract stem bark of Flacourtia rukam have been successfully. We have isolated and characterized three compounds friedelin (1), poliothryoside (2), and β -sitosteryl-3 β -glucopyranoside (3). The structures were identified used UV, IR and NMR spectrum data and comparison of their NMR data with literature. The antioxidant activities of each of them were tested used DPPH (1,1-diphenyl-2-picryl hydrazyl) method. All compounds given IC_{50} value $>100 \mu\text{g/mL}$ and were categorized weaker antioxidant. Compounds 2 and 3 have been reported from other species from genus Flacourtia but this is the first report of the isolation from F. rukam.

Keywords: Antioxidant; β -sitosteryl-3 β -glucopyranoside; Flacourtia rukam; friedelin; poliothryoside

ABSTRAK

Penerokaan sebatian kimia dan aktiviti biologi ekstrak kulit batang Flacourtia rukam telah berjaya. Kami telah dapat mengasingkan dan mencirikan tiga sebatian friedelin (1), poliotryosida (2), dan β -sitosteril-3 β -glukopyranosida (3). Struktur telah dikenal pasti menggunakan data spektrum UV, IR dan NMR dan perbandingan data NMR mereka dengan data kepustakaan. Kegiatan antioksidan masing-masing diuji menggunakan kaedah DPPH (1,1-difenil-2-pikril hidrazil). Semua sebatian mengandungi nilai $IC_{50} >100 \mu\text{g/mL}$ dan dikategorikan sebagai antioksidan yang lemah. Sebatian 2 dan 3 telah dilaporkan daripada spesies lain daripada genus Flacourtia tetapi ini adalah laporan pertama pengasingan daripada F. rukam.

Kata kunci: Antioksidan; Flacourtia rukam; friedelin; poliotrisosida; β -sitosteril-3 β -glukopiranosida

INTRODUCTION

Flacourtia rukam Zoll. & Mor. (syn. *F. edulis* Griff., *F. euphlebica* Merr., *F. megaphylla* Ridl.), is a species of genus *Flacourtia*, locally known as rukem or ganda rukem. This plant is native to Indonesia and is widely found in China, India and most of Southeast Asia (Lim 2013). In Indonesia, the stem bark of *Flacourtia rukam* traditionally used as high blood pressure (Yustian et al. 2012), the leaves is used to inflamed eye-lids and the fruits as medicine against diarrhoea and dysentery (Susilo & Denny 2016; Wiart et al. 2006). In the Philippines, a decoction of the roots is taken by women after childbirth (Ragasa et al. 2016). In Thailand, the wood used as herbal components the traditional for skin infectious treatments (Chusri et al. 2011). The fruit of the *F. rukam* is edible and is used for making pickle or mixed sugar made into jams, meanwhile the wood is hard and strong can be used for house construction and furniture and young leaves *F. rukam* can be eaten directly (Ragasa et al. 2016).

Chemical content information from *F. rukam* very still limited. Ragasa et al. (2016) have been reported 5 compound from the dichloromethane extract the fruit of *F. rukam* that is monogalactosyl diacylglycerols, β -sitosteryl-3 β -glucopyranoside-6 β -O-fatty acid esters,

β -sitosterol, triacylglycerols, and chlorophyll a. Compound monogalactosyl diacylglycerols is active as anti-inflammatory agent (Imbs et al. 2013) and also reported to be cytotoxic (Nguyen et al. 2004; Tsai et al. 2012). Ikram et al. (2009) reported that the fruits contain phenolics and demonstrate the activation of antioxidants with an inhibitory value of 70.9%. Saree (1998) isolated two chemical constituents from the twigs of *Flacourtia rukam* Zol. and Mor. and were identified as steroid, stigmastan-3, 6-dione, and triterpene friedelin. Phytochemical screening of extract ethanol of stem bark *F. rukam* showed the chemical constituents terpenoid, steroid, flavonoid, and fenol (Muharni et al. 2016).

The investigation chemical constituents *F. rukam* of Indonesia, three compounds have been isolation i.e. friedelin (1), poliothryoside (2), and β -sitosteryl-3 β -glucopyranoside from ethyl acetate extract of the stem bark. In this paper, we report the isolation, structure elucidation and antioxidant activity assay of these compounds. The structure of these compounds was determined to base on UV, IR and NMR data spectroscopy and compared with the reported data (Man et al. 2011; Peshin & Kar 2017). Antioxidant activity of these compound was determined with the DPPH method.

MATERIALS AND METHODS

GENERAL EXPERIMENTAL PROCEDURES

Melting point was determined by Gallen Camp melting point apparatus. UV spectrum was recorded on spectrophotometer ultraviolet Beck DU-7500. The IR (KBr) spectrum was recorded on Perkin Elmer-FTIR spectrometer, Shimadzu UV-168A. The $^1\text{H-NMR}$ and $^{13}\text{C-NMR}$ spectra were recorded on Agilent DD2 (500 MHz for $^1\text{H-NMR}$ and 125 MHz for $^{13}\text{C-NMR}$) with TMS as an internal standard (chemical shifts in δ , ppm) in CDCl_3 or $(\text{CD}_3)_2\text{CO}$ or DMSO. Coupling constants (J) are reported in Hz. UV-visible spectrophotometer using a single beam provided by Shimadzu-UV mini 1240 instrument. ultra-violet lamps used CAMAG 254 nm, radiation ($\lambda = 254$ and 365 nm) Thin layer chromatography (TLC) was performed with silica gel 60G F₂₅₄ using Merck (Art.5554) and spots were visualized by cerium sulfate vapors and ultraviolet light. All solvents were analytical reagent grade.

EXTRACTION AND ISOLATION

The fresh stem bark was collected in the month of January 2018 from Regency of Musi Banyu Asin South Sumatera Indonesia and identified by Dr. Laila Hanum (number specimen VIC 2702), Head of the Botany Laboratory, University of Sriwijaya. After the collection, the sample is minimized and dried at room temperature for 15 days and was crushed to obtain coarse powder which could pass through sieve number 60. About 1500 g of dried powder was extraction by maceration method with step gradient polarity solvent (n-hexane, ethyl acetate, and methanol) each for 24 h. This extraction process was repeated three times. The combined of each extract was evaporated to dryness under high vacuum at about 60°C to obtained crude n-hexane extract (0.92 g), ethyl acetate extract (13 g) and methanol (52 g). Each extracts were determined antioxidant activity by DPPH method.

The dry extract of ethyl acetate (13 g) was separated by using a vacuum column chromatography (KCV) method with silica gel F₂₄₅ 230 - 400 mesh particle size stationary phase. The column was prepared by introducing 65 g of silica gel 60 as stationary phase and compressed by pressing press, then eluting with n-hexane to compress the column. Dried ethyl acetate extract (13 g) was dissolved in the minimum quantity of acetone and adsorbed on silica gel 60 70-230 mesh particle size (13 g). The slurry formed was allowed to dry. The adsorbed extract was charged into the column. The column was first eluted with n-hexane and then with the solvent by gradually increasing the percentage of ethyl acetate in n-hexane. Eluen accommodated with Bottle. Furthermore, each bottle was analyzed TLC with eluent n-hexane: ethyl acetate 7: 3. Eluent in bottle with showed same TLC pattern were pooled and concentrated to minimum volume resulted and obtained F1 (2.4 g), F2 (2.2 g), F3 (1.5 g), and F4 (3.25 g). Fraction F1 yielded a residue, which was

recrystallized in acetone obtained white crystals needles 23 mg (1). Analyzes TLC at various eluent with spots were visualized by serum sulfate vapors showed a single spot. Based on this TLC analysis, it is assumed that the isolation has been pure compound.

The F4 fraction (3.25 g) was separated and purified by open column chromatography techniques. Fraction F4 (3.25 g) was taken and dissolved in the minimum quantity of acetone and adsorbed on silica gel (3.25 g) of 70-230 mesh particle size. The slurry formed was allowed to dry material and then crushed so that the sample and silica are homogeneous. A dried column (60 × 2 cm) was taken a cotton plug was put at the base of the column and the column is filled with an n-hexane solvent and while the tap is opened n-hexane with silica gel (60 g) was poured into the column gradually. Silica in the column was compacted by flowing solvent repeatedly. The adsorbed extract was charged into the column. The column was first eluted with n-hexane and then with the solvent by gradually increasing the percentage of ethyl acetate in n-hexane, n-hexane:ethyl acetate 9:1 (100 mL), 8:2 (100 mL), 6:4 (100 mL), 5:5 (150 mL), 3:7 (100 mL), and ethyl acetate 100% (200 mL). Eluate were collected in bottle volume of 10 mL and each bottle was analyzed by thin layer chromatography (TLC). The bottle showed same TLC with the pattern were pooled and concentrated to minimum volume resulted and obtained 4 subfractions F 4.1 (0.43 g), F 4.2 (0.50 g), F4.3 (0.22 g), and F4.4 (0.95 g). In F4.3 fraction is formed crystals and after purified by ethyl acetate obtained white amorphous solid 20 mg (2).

A part of methanol extract (10 g) also was separated by using vacuum column chromatography (KCV) in the same way as the separation of ethyl acetate extract with the solvent by gradually increasing the percentage of ethyl acetate in n-hexane and obtained 4 fractions F1 (0.75), F2 (3.25), F3 (2.1 g) dan F4 (1.25 g). The F4 fraction (1.25 g) was separated and purified by open column chromatography techniques with the solvent by gradually increasing the percentage of ethyl acetate in n-hexane: n-hexane: ethyl acetate 9:1 – 3:7 (each 100 mL), n-hexane: ethyl acetate 2: 8 (150 mL), EtOAc (100 mL) and EtOAc:MeOH 9:1 (200 mL) and obtained 3 fractions : F4.1 (0.35 g), F4.2 (0.25 g), and F4.3 (0.45 g). The fraction F4.3 (0.45 g) re-purified by open column chromatography techniques and obtained 4 subfractions: F4.3.1 (95 mg), F4.3.2 (vial 87 mg), F4.3.3 (40 mg) and F4.3.4 (125 mg). Fraction F4.3.3 after concentrated to minimum volume resulted in the crystal. and after recrystallized in ethyl acetate obtained amorphous solid 17 mg (3). Each isolated compounds also were determined antioxidant activity by DPPH method.

ANALYSIS OF ANTIOXIDANT ACTIVITY BY DPPH METHOD

The experiment should use a freshly prepared DPPH solution. DPPH (1.98 mg) was dissolved in 100 mL methanol (Shyu et al. 2009). The test conditions were as follows: 200 μL of each extract (concentration 1000, 500, 250, 125, and 62.5

$\mu\text{g/mL}$) and isolated compounds (concentration 100, 50, 25, 12.5, and 6.25 $\mu\text{g/mL}$) and 3800 μL of DPPH solution (0.05 mM). This mixture was shaken and allowed to react at room temperature during 30 min. Absorbance was recorded in a UV-visible spectrophotometer at 517 nm. The results were recorded as the percentage of inhibition according to the following equation: % inhibition = [(absorbance control – absorbance sample)/absorbance control] \times 100 (Molyneux 2004; Nagai et al. 2003; Selvi et al. 2003). The % inhibition value was plotted to concentration to obtain the IC_{50} against DPPH by linear regression.

RESULTS

Antioxidant activity of extracts

IC_{50} value of n-hexane extract 2239.80 $\mu\text{g/mL}$, ethyl acetate extracts 183.92 $\mu\text{g/mL}$, methanol extract 83.40 $\mu\text{g/mL}$.

Antioxidant activity of isolated compound

IC_{50} value of isolated compounds 1: 162.9 $\mu\text{g/mL}$, compound 2: 154.6 $\mu\text{g/mL}$, and compound 3: 109.0 $\mu\text{g/mL}$.

Compound 1

The structure of compound 1 according to spectroscopy data were found to be as follows:

UV spectra no absorption above 210 nm. IR peak at 2927 and 2866 (aliphatic C-H stretching), 1712 (C=O absorption peak ketone) other absorption peaks; 1456 cm^{-1} (CH₂) 1384 (gem dimethyl), 1047 (cycloalkane). NMR spectrum data compound 1 showed in Table 3.

Compound 2

UV spectra absorption at λ_{max} nm: 225 and 282 and addition of NaOH showed. The bathochromic shift at 282 nm to 299. IR peak at 3406 (OH), 2922; 2860 (CH aliphatic), 1701. (C=O ester), 1614, 1506, 1464 (C=C aromatic), and 1068 (C-O). NMR spectrum data compound 2 showed at Table 4.

Compound 3

UV spectrum no patterns above 210. IR peak at 3427 cm^{-1} a hydroxyl group (-OH), 2935-2866 cm^{-1} (aliphatic C-H stretching of -CH₃, -CH₂- and C-H), 1462 cm^{-1} and 1377 (CH₂ and CH₃ bending), 1068 cm^{-1} (C-O-C linkage). NMR spectrum data compound 3 showed at Table 5.

DISCUSSION

This research based on the used of the stem bark of *F. rukam* as a traditional medicine in the Musi Banyuasin for the treatment of hypertension (Yustian et al. 2012). Based on literature studies, researchers have not found any information on chemical content or biological activity for the stem bark part of this plant. To scientifically information what compounds provide antihypertensive activity from this plant extract, the researchers were interested in identifying secondary metabolites contained in these plant extracts. It is known that hypertension is a

degenerative disease that is related to oxidants. Therefore, the beginning of this study was to test the antioxidant activity of each extract by DPPH method, and the separation and purification of the compound from the active extract. Compounds that were successfully isolated were tested for antioxidant activity.

Extraction (1500 g) of dried powder of *F. rukam* with step gradient polarity solvent (n-hexane, ethyl acetate, and methanol) after evaporated to dryness under high vacuum at about 60°C to obtained crude n-hexane extract (0.92 g), ethyl acetate extract (13 g) and methanol (52 g). Antioxidant activity of the extracts were determined used DPPH method base on absorption DPPH at λ maks 517 nm and calculated by percent inhibition (% I). The IC_{50} value each extracts determined by linear regression. The result show in Table 1.

TABLE 1. The antioxidant activity of each extract

Extract	IC_{50} ($\mu\text{g/mL}$)
n-Hexane	2239.80
Ethyl acetate	183.92
Methanol	83.40

The results showed that methanol extract have a better antioxidant activity compared to ethyl acetate and n-hexane extracts. The extract is categorized as active as an antioxidant if it has an IC_{50} of less than 200 $\mu\text{g/mL}$. If the IC_{50} value is found at around 200 to 1000 $\mu\text{g/mL}$ in category weak antioxidant and above 1000 $\mu\text{g/mL}$ is categorized as inactive (Molyneux 2004). Base on this steatment, methanol and ethyl acetate extract is categorized as active antioxidant, and n-hexane extract in category in active antioxidant. Separation and purification of ethyl acetate extract to obtained 2 compound: White crystals needles 23 mg (1) and white amorphous solid 20 mg (2). While from methanol extract obtained one compound form amorphous solid 17 mg (3).

Antioxidant activity of the compounds was determined by DPPH method and the results are shown in Table 2. According to Selvi et al. (2003) the pure compound categorized has potent antioxidant activity value to IC_{50} <10 $\mu\text{g/mL}$. IC_{50} 10-100 $\mu\text{g/mL}$ categorized as active antioxidants and when have IC_{50} > 100 $\mu\text{g/mL}$ categorized as weaker antioxidant. The data proved that all compound were categorized as weaker antioxidant or in active antioxidant. Meanwhile ascorbic acid used as a standard gives an IC_{50} value IC_{50} 9.05 $\mu\text{g/mL}$.

TABLE 2. The antioxidant activity of the isolated compounds

Compound	IC_{50} ($\mu\text{g/mL}$)
1	162.9
2	154.6
3	109.0
Ascorbit acid	9.05

The antioxidant activity test of the methanol and ethyl acetate extract in category antioxidant activity (Table 1), but all the compounds that were isolated from extract (compound **1**, **2**, **3**) only show weak antioxidant categories ($IC_{50} > 100$). Base on this data, it was concluded that researchers had not succeeded in identifying compounds antioxidants in the stem bark extract of *F. rukam*. Therefore, it has not been shown scientifically what compounds act as antioxidants from bark extracts of rukam plants. For isolated compounds, spectroscopic analysis is performed to determine the molecular structure.

The compound **1** is substance form white crystalline needles with a melting point of 144-146°C. UV spectrum showed no patterns above 210. IR spectroscopic analysis showed the absorption bands at 2727 and 2866 cm^{-1} that is characteristic of stretching aliphatic C-H and Absorption at 1712 cm^{-1} is due to in ketone carbonyl. Other absorption frequencies at 1456 cm^{-1} is a bending frequency for cyclic (CH₂)_n and 1384 cm^{-1} for -C-CH₃)₂. The absorption frequency at 1047 cm^{-1} signifies cycloalkane. The IR spectrum data resemble the absorption frequencies observed for friedelin. The structures of the

isolated compound were determined by ¹H and ¹³C NMR spectroscopy and by comparison of the spectral data with published data, which showed the presence of 30 carbons and suggestive of a triterpenoid typical of friedelane skeleton (Man et al. 2011) (Table 3). Based on the data, isolated compound was identified as friedelin (**1**). The above IR, ¹H-NMR, and ¹³C NMR spectral data and their comparison with those described in the literature showed the structure of compound **1** to be the friedelin (Figure 1). All data are in good agreement to that of friedelane skeleton (Man et al. 2011).

Compound **2** was obtained as white amorphous solid. The UV spectra of **2** exhibited absorption at λ_{max} nm: 225 and 282 and addition of NaOH showed. The bathochromic shift at 282 nm to 299, it can conclude that there was a phenolic group. IR spectrum showed the functional group such as 3406 (OH), 2922; 2860 (CH aliphatic), 1701 (C=O ester), 1614, 1506.41, 1464. (C=C aromatic), and 1068 (C-O). The ¹H-NMR spectrum showed the chemical shift at δ 6.54 (1H, *dd*, *J*=3.05; 8.50), 6.82 (1H, *d*, *J*=3.05), and 7.11 (1H, *d*, *J*=8.5) were the three aromatic protons which is ortho meta coupled from the aromatic three substitutions.

TABLE 3. ¹H-NMR (500 MHz), ¹³C-NMR (125 MHz), and DEPT data compound 1^b (CDCl₃, in ppm) and friedelin^a (in CDCl₃) (Man et al. 2011)

Carbon No	C-NMR ^a	H-NMR ^a	C-NMR ^b	H NMR ^b	DEPT
1	22.3	1.96 ; 1.68	22.3	1.96; 1.89 (2 H, <i>m</i>)	CH ₂
2	41.5	2.39 ; 2.28	41.5	2.39 (2H, <i>m</i>)	CH ₂
3	213.2	-	213.2	-	C
4	58.2	2.25	58.2	2.28 (1H, <i>m</i>)	CH
5	42.1	-	42.1	-	C
6	41.3	1.75 ; 1.28	41.3	1.28; 1.70 (2H, <i>m</i>)	CH ₂
7	18.2	1.49 ; 1.37	18.6	1.41; 1.43	CH ₂
8	53.1	1.39	53.1	1.37	CH
9	37.4	-	37.4	-	C
10	59.4	1.53	59.5	1.73 (1H, <i>m</i>)	CH
11	35.6	1.46 ; 1.26	35.6	1.26 ; 1.38	CH ₂
12	30.4	1.34 ; 1.34	30.4	1.32 (2H, <i>m</i>)	CH ₂
13	39.7	-	39.7	-	C
14	38.3	-	38.3	-	C
15	32.4	1.46; 1.27	32.4	1.31; 1.48(2H, <i>m</i>)	CH ₂
16	36.0	1.57; 1.36	36.0	1.37; 1.52	CH ₂
17	30.0	-	31.8	-	C
18	42.8	1.56	42.8	1.54 (1H, <i>m</i>)	CH
19	35.3	1.38; 1.20	35.3	1.26 ;1.41	CH ₂
20	2811	-	28.2	-	C
21	32.7	1.51; 1.30	32.8	1.26;1.44	CH ₂
22	39.2	1.50 ; 0.94	39.2	0.98 ;1.51 (2H, <i>m</i>)	CH ₂
23	6,8	0.87	6.8	0.89(3H, <i>s</i>)	CH ₃
24	14.6	0.71	14.6	0.73 (3H, <i>s</i>)	CH ₃
25	17.9	0.86	17.9	0.87 (3H, <i>s</i>)	CH ₃
26	20.2	1.0	20.3	0.96 (3H, <i>s</i>)	CH ₃
27	18.6	1.05	18.2	1.00 (3H, <i>s</i>)	CH ₃
28	32.1	1.17	32.1	1.06(3H, <i>s</i>)	CH ₃
29	35.0	1.00	35.0	0.94(3H, <i>s</i>)	CH ₃
30	31.8	0.95	32.3	1.01(3H, <i>s</i>)	CH ₃




FIGURE 1. Structure of compound 1-3

There are also three other groups of proton aromatics on 7.56 (2H, *m*), 7.68 (1H, *m*) and 8.09 (2H, *m*) which were protons in mono substitution aromatic. The proton NMR spectrum also showed sinyal chemical shifts region δ 3.55 (3H, *s*), 3.83 (1H, *m*), 4.43 (1H), 4.45 (1H), 4.7 - 4.8 (3H) showed the presence of protons of the sugar moiety. The ^{13}C -NMR spectrum (in acetone) showed twenty carbon signals. The presence of sugar was supported by ^{13}C -NMR data which showed a group of sugar signals at δC 104.52 (C-1''), 77.9 (C-3''), δC 75.1 (C-5''), δC 74.9 (C-2''), δC 71.6 (C-4'') and δC 6.1 (C-6'') and fourteen other signals to the aglycone moiety. From fourteen carbon signals for aglycone were attributed to two methylene signals, eight methine signals and four quaternary carbons by DEPT and HSQC spectra.

The HMBC correlation H-1'' at δ 4.77 to C-7 at δ 166.6 is shown and correlation H-6'' at 4.75 at δ 150.2 that the linkage position with the glucose unit is at C-7 and C-1'. The correlations of one methylene protons from methyl alcohol at δ 4.43 to C-2' at δ 134.3 in HMBC spectrum indicated that one methylene group is connected to C-2'. NMR Data of ^1H , ^{13}C , and HMBC NMR of compound 2 (Table 4), key correlations at HMBC the structure of compound 2 (Figure 2). By analyzing the ^1H - and ^{13}C -NMR data and compared with the reported data (Hussain et al. 2016), the compound was elucidated as β -D-glucopyranoside, 4-hydroxy-2-(hydroxymethyl)phenyl, 6-benzoate (poliothryoside).

Compound 3 was obtained as an amorphous solid. m.p: 290°-292°C, UV spectrum showed no patterns above 210. IR spectrum showed an absorption peak in the region




FIGURE 2. Key correlation HMBC from proton to carbon compound 2

(3427) cm^{-1} indicating the presence of a hydroxyl group (-OH) of glucoside linkage and the absorption bands at 2935-2866 cm^{-1} indicated the presence of -CH aliphatic asymmetric and symmetric stretching of -CH₃, -CH₂- and C-H. Besides these, another two band at 1462 cm^{-1} and 1377 is a bending frequency for CH₂ and CH₃ and a band observed at 1068 cm^{-1} was assigned to the C-O-C linkage.

^1H -NMR (DMSO-*d*₆) spectrum showed the chemical shift at δ 5.33 (1H, *m*) for one olefinic double bond proton. The proton NMR spectrum also showed the six signals at δ 0.65 (3H), 0.95 (3H), for H-18 and H-19, respectively, and sinyal at δ 0.78 - 0.84 (6H) and 0.89 - 0.93 (6H) for 4 other methyl groups. The peak at chemical shifts region δ 2.85-3.65 as a multiplet was showed the presence of protons of

TABLE 4. ¹H-NMR (500 MHz), ¹³C-NMR (125 MHz), HSQC and HMBC data of **2** (CD₃)₂CO, in ppm)

Carbon No	C-NMR	H-NMR	HMBC H C	DEPT
1	131.2	-	-	C
2,6	130.3	8.09(2H, m) J = 7.45; 4.05	129.4 (C3,5); 134.0 (C4); 166.6 (C7)	CH
3,5	129.4	7.56 (2H, dd, J = 8.0, 7.45)	129.4 (C3,5), 130.3 (C2,6); 134.0 (C4)	CH
4	134.0	7.68 (1H, t) 8.0	129.4; 130.3 (C2,6)	CH
7	166.6	-	-	C
1'	150.2	-	-	C
2'	134.3	-	-	C
3'	116.0	6.82 (1H, d, J = 3.05)	115.0 (C5'); 150.2 (C1')	CH
4'	153.2	-	-	C
5'	115.0	6.54 (1H, dd J = 3.05; 8.5)	116.0 (C3'); 150.2 (C1')	CH
6'	119.2	7.11 (1H, d, J = 8.5)	134.2 (C2'); 150.2 (C1'); 153.2 (C4')	CH
7'	61.1	4.75; 4.43	150.2 (C1'), 116.0 (C3'), 134.3 (C2')	CH ₂
7' OH		4.30	61.1	
1''	104.5	4.77(1H, d, J = 4.85)	150.2 (C1')	CH
2''	74.9	3.55 (1H, m)	104.5; 77.9	CH
2'' OH		4.60	104.5 (C1'')	
3''	77.9	3.55 (1H, m)	71.5	CH
		4.90		
4''	71.6	3.55 (1H, m)	77.9 (C3')	CH
5''	75.1	3.83 (1H, dt, J = 9.55; 9.35)	77.9 (C3')	CH
6''	65.1	4.45 ; 4.76 (2H, m)	75.11; 166.62	CH ₂

the sugar moiety and the very downfield chemical shift at δ 4.21 was assigned for the proton of C-H group of glycoside.

The ¹³C NMR spectrum of the compound **3** showed 35 carbon signals, of which six were for the sugar moiety and 29 others signal to the aglycone moiety. The ¹³C NMR spectrum of the compound **3** showed 35 carbon signals, of which six were for the sugar moiety and 29 others signal to the aglycone moiety. The chemical shift at δ 11.6 and 19.0 ppm was signal to two methyl group at position C-18, C-19. The signals observed at δ 11.7, 18.6, and 18.9 and 19.7 is signal for 4 methyl group others. The carbon signals of the sugar moiety observed at δ 100.7 (C-1'), 73.4 (C-2'), 76.6 (C-3'), 70.0 (C-4'), 76.7 (C-5'), and δ 61.0 (C-6'), according with signal-signal of glucose moiety. The aglycone moiety signals were observed at δ 140.4 and 121.2 ppm were assigned for the olefinic carbon.

The ¹H and ¹³C NMR values for all the protons and carbons were the determined basis of heteronuclear single quantum coherence spectroscopy (HSQC and (Distortionless Enhancement by Polarization Transfer (DEPT), which are given in Table 5. The ¹³C NMR spectra data were a comparison with those described in the literature (Peshin & Car 2017) and these data confirmed that compound **3** β -sitosterol-3-O- β -D-glucoside (Figure 1). Friedelin (**1**) has been reported from twigs of *F. rukam* (Saree 1998). Poliothryoside (**2**) has been reported from *F. indica* (Sashidhara et al. 2013) and β -sitosterol-3-O-

β -D-glucoside (**3**) also has been obtained of *Flacourtia ramontchi* (Satyanarayana et al. 1991). Compound **3** also have been reported from different genus i.e. *Phyllanthus* species (Mahbuka et al. 2012) and *Salvadora persica* LINN (Arora et al. 2013) and no reports of the presence of compounds **2** and **3** of *F. rukam*. Based on literature study, this is the first report on the isolation compounds **2** and **3** from *F. rukam*.

CONCLUSION

In the work, three compounds were isolated from extracts active antioxidant which is friedelin (**1**), poliothryoside (**2**) and β -sitosterol-D-glucoside (**3**). Compounds **2** and **3** have been reported from other species from genus *Flacourtia* but the first report on the isolation from *F. rukam*, but all compounds showed weaker antioxidant activity. This study has not succeeded in identifying antioxidant compounds from the stem bark extract *F. rukam*.

ACKNOWLEDGEMENTS

The authors are thankful to the Ministry of Technology Research, Directorate of Research and Community Service, Directorate General Strengthening Research and Development of Ministry of Research, Technology and High Education Republic of Indonesia for Basic Research of College 2018 research grant and the Department of

TABLE 5. ¹H-NMR (500 MHz), ¹³C-NMR (125 MHz), data of compound **3**^b (DMSO-*d*₆, in ppm) and β-sitosterol-3-O-β-glucoside^a (CDCl₃) (Peshin & Car 2017)

Carbon No	C-NMR ^a	C-NMR ^b	H-NMR ^a	H-NMR ^b	DEPT
1	36.8	36.8	1.25 (<i>m</i> , 2H)	0.99 (1H, <i>m</i>) ; 1.80 (1H, <i>m</i>)	CH ₂
2	29.2	29.2	1.33 (<i>m</i> , 2H)	1.65 (2H, <i>m</i>)	CH ₂
3	78.6	76.8	3.13 (<i>m</i> , 1H)	3.14 (1H, <i>m</i>)	CH
4	39.3		2.14 (<i>m</i> , 2H)	1.15 (1H, <i>m</i>); 1.95 (1H, <i>m</i>)	CH ₂
5	140.4	140.4	-	-	C
6	121.2	121.2	5.09 (<i>bs</i> , 1H)	5.33 (1H, <i>bs</i>)	CH
7	31.4	31.3	1.73 (<i>m</i> , 2H)	1.51; 1.94 (1H, <i>m</i>)	CH ₂
8	31.3	31.4	1.22 (<i>m</i> , 1H)	1.40 (1H, <i>m</i>)	CH
9	49.6	49.5	1.22 (<i>m</i> , 1H)	0.89 (1H, <i>m</i>)	CH
10	36.2	36.2	-	-	C
11	20.6	20.6	1.33 (<i>m</i> , 2H)	1.40 (1H, <i>m</i>); 1.44 (1H, <i>m</i>)	CH ₂
12	38.3	38.2	1.33 (<i>m</i> , 2H)	2.12 (1H, <i>m</i>); 2.37 (1H, <i>m</i>)	CH ₂
13	41.8	41.8	-	-	C
14	56.1	56.1	1.22 (1H, <i>m</i>)	0.99 (1H, <i>m</i>)	CH
15	23.8	23.8	1.73 (2H, <i>m</i>)	1.24 (2H, <i>m</i>)	CH ₂
16	27.8	27.8	1.73 (1H, <i>m</i>)	1.22 (2H, <i>m</i>)	CH ₂
17	55.4	55.4	1.73 (2H, <i>m</i>)	1.09 (1H, <i>m</i>)	CH
18	11.6	11.6	0.62 (3H, <i>s</i>)	0.65 (3H, <i>s</i>)	CH ₃
19	19.1	19.0	0.94 (3H, <i>s</i>)	0.95 (3H, <i>s</i>)	CH ₃
20	35.4	35.4	1.32 (1H, <i>m</i>)	1.3 (1H, <i>m</i>)	CH
21	18.6	18.6	0.84 (3H, <i>d</i>)	0.90 (3H, <i>s</i>)	CH ₃
22	33.3	33.3	1.73 (2H, <i>m</i>)	0.98 (2H, <i>m</i>)	CH ₂
23	25.4	25.4	1.73 (2H, <i>m</i>)	-	CH ₂
24	45.1	45.1	1.12 (1H, <i>m</i>)	0.93 (<i>m</i> , 1H)	CH
25	28.6	28.7	2.14 (1H, <i>m</i>)	-	CH
26	19.7	19.7	0.75 (3H, <i>d</i>)	0.82 (<i>d</i> , 3H)	CH ₃
27	18.9	18.9	0.73 (3H, <i>d</i>)	0.80 (<i>d</i> , 3H)	CH ₃
28	22.1	22.6	1.33 (2H, <i>m</i>)	-	CH ₂
29	11.7	11.7	0.77(3H, <i>t</i>)	0.82 (<i>s</i> , 3H)	CH ₃
1'	100.7	100.7	4.11 (1H, <i>d</i>)	4.21 (<i>d</i> , 1H)	CH
2'	73.4	73.4	3.14 (1H, <i>m</i>)	2.89 (<i>m</i> , 1H)	CH
3'	76.9	76.6	3.14 (1H, <i>m</i>)	3.07 (<i>m</i> , 1H)	CH
4'	70.0	70.0	3.4 (1H, <i>m</i>)	3.01 (<i>m</i> , 1H)	CH
5'	76.7	76.7	3.06 (1H, <i>m</i>)	3.11 (1H, <i>m</i>)	CH
6'	62.8	61.0	2.94 (2H, <i>m</i>)	3.40 (1H, <i>m</i>), 3.64 (1H, <i>m</i>)	CH ₂

Chemistry, University of Sriwijaya for the facilities provided.

REFERENCES

- Arora, M. & Kalia, A.N. 2013. Isolation and characterization of stigmasterol and β-sitosterol-D-glycoside from the ethanolic extract of stems of *Salvadora persica* LINN. *International Journal of Pharmacy and Pharmaceutical Sciences* 5(1): 245-249.
- Chusri, S., Chaicoch, N., Thongza-ard, W., Limsuwan, S. & Voravuthikunchai, S.P. 2012. *In vitro* antibacterial activity of ethanol extracts of nine herbal formulas and its plant components used for skin infections in Southern Thailand. *Journal of Medicinal Plants Research* 6(44): 5616-5623.
- Hussain, S.M., Mohammed, S.H., Arbaaz, A.L. & Nahida, A. 2016. Characterization of isolated bioactive phytoconstituents from *Flacourtia indica* as potential phytopharmaceuticals-An *in silico* perspective. *Journal of Pharmacognosy and Phytochemistry* 5(6): 323-331.

- Ikrum, E.H.K., Eng, K.H., Jalil, A.M.M., Islail, M., Idris, S., Azlan, A., Nasri, H.S.M., Diatom, N.K.M. & Mokhtar, R.A.M. 2009. Antioxidant capacity and total phenolic content of Malaysian underutilized fruits. *Journal of Food Composition and Analysis* 22: 388-393.
- Imbs, T.I., Ermakova, S.P., Fedoreyev, S.A., Anastuyuk, S.D. & Zvyagintseva, T.N. 2013. Isolation of fucoxanthin and highly unsaturated monogalactosyldiacylglycerol from brown alga *Fucus evanescens* C Agardh and *in vitro* investigation of their antitumor activity. *Mar. Biotechnol.* 15(5): 606-612.
- Lim, T.K. 2013. *Flacourtia rukam*. *Edible Medicinal and Non-Medicinal Plants*. Volume 5. Springer Science Business Media. pp. 776-779.
- Mahbuka, K., Mirajum, B. & Abdul, Q. 2012. Sterols and sterol glucoside from *Phyllanthus* species. *Dhaka University Journal of Science* 60(1): 5-10.
- Muharni, F. & Nurmaliana, R. 2016. Phytochemical screening antioxidant and antibacterial activity of traditional medicine ethnic Musi Palembang. Ministry of Health Republic of Indonesia.
- Man, A., Ibrahim, K., Oyewale, A.O., Amupitan, J.O., Fatope, M.O. & Okogun, J.I. 2011. Antimycobacterial friedelanterpenoid from the root bark of *Terminalia Avicennioides*. *American Journal of Chemistry* 1(2): 52-55.
- Molyneux, P. 2004. The use of the stable free radicals diphenylpicrylhydrazyl (DPPH) for estimating antioxidant activity. *Songklanakarinn J. Sci. Technol.* 26(2): 211-219.
- Nagai, T., Inoue, R., Inoue, H. & Suzuki, N. 2003. Preparations and antioxidant properties of water extract of propolis. *Food Chemistry* 80: 29-33.
- Nguyen, A.T., Malonne, H., Duez, P., Vanhaelen, F.R., Vanhaelen, M. & Fontaine, J. 2004. Cytotoxic constituents from *Plumbago zeylanica*. *Fitoterapia* 75(5): 500-504.
- Peshin, T. & Kar, H.K. 2017. Isolation and characterization of β -sitosterol-D-glycoside from the extract of the flowers of *Viola odorata*. *British Journal of Pharmaceutical Research* 16(4): 1-8.
- Ragasa, C.Y., Jo, M.A.Y., Theresa, J.T., Maria, C.S., Irving, D., Robert, B. & Sylvia, U. 2016. Chemical constituents of *Flacourtia rukam* Zoli. & Moritzi fruit. *International Journal of Pharmaceutical and Clinical Research* 8(12): 1625-1628.
- Sashidhara, K.V., Suriya, P.S., Shiv, V.S., Rajeev, K.S., Kumkum, S., Saxena, J.K. & Sunil, K.P. 2013. Isolation and identification of β -hematin inhibitors from *Flacourtia indica* as promising antiplasmodial agents. *European Journal of Medicinal Chemistry* 60: 497-502.
- Saree, O. 1998. Chemical constituents from *Flacourtia rukam* Zol. and Mor. Faculty of Education. Demonstration School Prince of Songkla Univ. Pattani, Thailand (Unpublished).
- Selvi, A.T., Joseph, G. & Jayaprakasha, G. 2003. Inhibition of growth and aflatoxin production in *Aspergillus flavus* by *Garcinia indica* extract and its antioxidant activity. *Food Microbiol.* 20: 455-460.
- Shyu, Y.S., Lin, J.T., Chang, Y.T., Chiang, C.J. & Yang, D.J. 2009. Evaluation of antioxidant ability of ethanolic extract from dill (*Anethum graveolens* L.) flower. *Food Chemistry* 115: 515-521.
- Satyanarayana, V., Kurupadanam, G.L. & Srimanaraya, G. 1991. A butyrolactone lignin disaccharide from *Flacourtia ramontchi*. *Phytochem.* 30(3): 1026-1029.
- Susilo, A. & Denny, S. 2016. Diversity and potential use of plants in the secondary natural forest in RPH Cisujen KPH Sukabumi, West Java. *Pros. Semnas Masy. Biodiv. Indon* 2(2): 256-262.
- Tsai, P.W., de Castro-Cruz, K., Shen, C.C. & Ragasa, C.Y. 2012. Chemical constituents of *Ficus odorata*. *Pharmaceutical Chemistry Journal* 46(4): 225-227.
- Wiant, C. 2006. *Medical plants of the Asia-Pacific: Drugs for the Future*. World Scientific Publishing Co. Pte. Ltd.
- Yustian, I., Muharni, S., Zulaicha, S. & Arbi, M. 2012. *Special Research on the Exploration of Ethnomedicine and Local Community Medicinal Plants in Indonesia (Ethnic Musi II)*. Palembang. Ministry of Health Republic of Indonesia.
- Muharni, Elfita*, Heni Yohandini & Julinar
Department of Chemistry
Faculty of Mathematics and Natural Sciences
University of Sriwijaya
Indralaya, Ogan Ilir
South Sumatra, 30662
Indonesia
- Yasrina & Miranti
Department of Pharmacy
Faculty of Mathematics and Natural Sciences
University of Sriwijaya
South Sumatra, 30662
Indonesia

*Corresponding author; email: elfita69@gmail.com

Received: 6 August 2018

Accepted: 9 July 2019