

Association of Vitamin D Deficiency and Diabetic Foot Ulcer Patients in Dr Mohammad Hoesin General Hospital Palembang

by Krisna Murti

Submission date: 25-Nov-2024 09:24AM (UTC+0700)

Submission ID: 2400944810

File name: ion_of_Vitamin_D_Deficiency_and_Diabetic_Foot_Ulcer_Patients.pdf (285.13K)

Word count: 5043

Character count: 26760



Association of Vitamin D Deficiency and Diabetic Foot Ulcer Patients in Dr Mohammad Hoesin General Hospital Palembang

Kemas Dahlan^{1*}, Fahmi Jaka Yusuf², Irfannuddin³, Krisna Murti⁴,
 Akhmadu Muradi⁵, Iche Andriyani Liberty⁶
 RSUP Dr. Mohammad Hoesin Palembang, Indonesia^{1,6}
 Universitas Sriwijaya, Palembang, Indonesia^{2,3,5}
 Universitas Indonesia, Depok, Indonesia⁴
 Email: dokterdahlabsb@gmail.com

ABSTRACT

Keywords:
 Vitamin D; Diabetes
 Mellitus; Diabetic Foot
 Ulcers

Vitamin D plays a role in the healing process of diabetic foot ulcers (DFU); it can improve the immune system, regulate inflammation function, and regulate growth factors in diabetes mellitus (DM). The immunomodulatory effects of vitamin D include reduced production of IL-2, IL-6, and TNF-alpha and increased production of IL-10. It also increases the activity of antimicrobial peptides (AMP) such as cathelicidin, and it promotes the differentiation of monocytes into macrophages, phagocytosis, and the production of lysosomal enzymes. Furthermore, vitamin D stimulates growth factors including vascular endothelial growth factor (VEGF) and transforming growth factor beta (TGF-β). In this study, the incidence of DM and DFU in DM patients at RSUP Dr. Mohammad Hoesin Palembang will be compared to serum vitamin D levels. This research uses quantitative methods with an analytical observational design. The Mean vitamin D serum level was 11,727 ng/mL, and there was a significant association between serum vitamin D deficiency and DFU (p-value; 0.036 < 0.05). There was a significant association between serum vitamin D deficiency and DFU occurrence (p-value; 0.036 < 0.05) in Mohammad Hoesin General Hospital Palembang. Based on the results of this study, we proposed that patients with diabetic foot ulcers should have their serum vitamin D levels checked, and that vitamin D should be given to them as part of their therapy.

Corresponden Author: Kemas Dahlan

Email: dokterdahlabsb@gmail.com

Artikel dengan akses terbuka dibawah lisensi



Introduction

One chronic consequence of diabetes mellitus (DM) is diabetic foot ulcers (DFU). The prevalence rate of DFU in the DM population is quite high, about 4-15% and (Harrisa et al., 2020; Ji et al., 2021; Yunir et al., 2022). The immune system and growth factors, which are crucial to the healing of wounds, perform less well in people with diabetes. In hyperglycemic conditions, proinflammatory mediators increase, leading to chronic inflammation and delayed wound healing time in DM patients (Cimmino et al., 2013). As an inflammatory marker, fibrinogen promotes the expression of some protease enzymes such as matrix metalloproteinases-9 (MMP-9) (Giatsidis et al., 2018).

DFU patients had higher blood sedimentation rate (ESR) and C-reactive protein (CRP) levels. Elevation of CRP and ESR in DFU can be used as indicators of wound severity (Muller et al., 2008). The higher the CRP or ESR of DFU patients, the worse the wound. Defense systems are unable to create an initial response to damage as a result of decreased immune function, growth factor, and inflammatory system activity. Formation of granulation tissue will be delayed in DFU (Lobman et al., 2002).

MMP-9 plays a role in the degradation of the extracellular matrix in the wound-healing process. TIMP-1 is an inhibitor of several protease enzymes including MMP-9. High levels of MMP-9 and low TIMP-1 are predictors of obstacles to DFU healing in DM patients (Caley et al., 2015; López-López et al., 2014; Shin et al., 2019). Based on research conducted by Lopez et al. (2014), vitamin D3 could regulate MMP gene expression from keratinocyte cells in DFU patients (Ross et al., 2011).

In addition to controlling calcium metabolism to preserve bone health, vitamin D can help diabetic foot ulcers (DFU) heal more quickly because it strengthens the immune system, controls inflammatory cytokines, and modifies the activity of growth factors, proteolytic enzymes like MMPs, and their inhibitors in diabetic mellitus (DM) (Dai et al., 2019; Ghezal et al., 2019; Kamble & Swarnkar, 2019; Kinesya et al., 2023; Ross et al., 2011; Yamine et al., 2020).

By boosting the action of antimicrobial peptides (AMP) like cathelicidin, stimulating monocyte differentiation into macrophages, enhancing phagocytosis function, and elevating lysosomal enzyme synthesis, vitamin D modulates the immune system. Vitamin D also affects the action of the inflammation system for several interleukins and growth factors by reducing the production of IL-2, IL-6, and TNF- α and increasing the production of IL-10. TGF- β and VEGF are two examples of growth factors that are activated by vitamin D (Kamble & Swarnkar, 2019).

Numerous MMP genes that are generated by keratinocyte cells are influenced by vitamin D. Furthermore, increased blood vitamin D levels have been demonstrated in recent research to upregulate MMP-1 expression, which is considered a trustworthy marker of DFU wound healing (8, 10).

Sunlight's ultraviolet B rays cause the skin to spontaneously create the prohormone vitamin D. The activation of vitamin D's effect in the skin is attributed to vitamin D receptors (VDR) in the skin. The relationship between vitamin D and VDR affects the control of basal cell keratinocyte growth, which acts as a physical barrier. Vitamin D also helps to regulate cell differentiation in the skin, which in turn helps to synthesise keratin, involucrin, transglutaminase, loricrin, and filaggrin (Tiwari et al., 2014).

By affecting the function of the Matrix Metalloproteinase (MMPs) enzymes and the activity of the Tissue Inhibitor Matrix Metalloproteinase (TIMP), vitamin D also plays a part in the proteolysis of the extracellular matrix (ECM), which is generated during the remodeling phase of DFU wound healing. In DM patients, there are high glucose levels in blood serum; this condition increases macrophage infiltration, increasing TNF- α in the wound. Increased levels of TNF- α cause more expression of MMP produced by keratinocyte cells and delay wound healing in DFU patients, as in invitro research conducted by Huang et al. (2019). Elevated glucose levels trigger M-1 macrophage polarization, which hinders keratinocyte migration through TNF- α . This is a crucial mechanism that postpones the healing of diabetic wounds. For DFU among other disorders, vitamin D supplementation treatment is an option (Dadaei et al., 2015; Holick et al., 2011).

A level of vitamin D less than 50 nmol/mL (<20 ng/mL) is considered a deficiency of vitamin D. Serum Vitamin D level more than 50 nmol/L (20 ng/mL) is recommended. In healthy

people, daily requirement for vitamin D is 15 mcg (600 IU) (Masood et al., 2015; Rastogi et al., 2022). The Endocrine Society states that people need a daily vitamin D supplement of at least 37.5 to 50 mcg (1,500–2,000 IU) in order to maintain blood 25(OH)D levels above 75 nmol/L (30 ng/mL). High-dose vitamin D is more effective in promoting wound healing in DFU (Marcinowska-Suchowierska et al., 2018; Rastogi et al., 2022).

Based on other studies, it has also been proven that low levels of Vitamin D3 have a close relationship with the occurrence of LKD due to prolonged inflammation and infection processes 28–30. Studies such as the Randomized Double-blinded Clinical Trial by Halschou-Jensen et al. in Denmark in 2021 and the study by Razzaghi et al. in Iran in 2017 have demonstrated that administering additional high-dose vitamin D3 therapy to patients with LKD improves the wound healing process more than administering low doses.

Further research is required to determine the relationship between serum vitamin D levels and the incidence of diabetes mellitus and DFU, there is still a dearth of information in Indonesia, particularly in South Sumatra, regarding vitamin D levels and the degree of DFU and the high risk of disability due to DFU.

Research Methods

This study used quantitative methods with an analytical observational design to determine the relationship between serum vitamin D levels in DM patients with DFU. The research was conducted at the Mohammad Hoesin Palembang General Hospital (RSMH) from January to September 2023. The study population included all patients at RSMH who underwent blood sampling to examine serum vitamin D levels. Retrieving, inputting, and collecting data were carried out using medical record data stored in e-Medical Records (EMR). The data collection period that satisfied the inclusion and exclusion criteria was used for data collection. Data recorded included the subject's identity, serum vitamin D levels, current blood sugar values, and whether the patient had diabetes mellitus with diabetic foot ulcers (DFU) or not.

Before analyzing research data, determine the data type to be processed based on each research variable. In each study, the data obtained were converted into ordinal form before bivariate analysis. All calculations of the results of this research were done using SPSS software. This research was conducted with approval from the Dr. Mohammad Hoesin General Hospital Health Research Ethics Committee No. DP.04.03/ D.XVIII.6.11/ ETIK/ 99/ 2023.

Results and Discussion

Result

The connection between blood vitamin D levels and DFU patients at Mohammad Hoesin General Hospital Palembang was investigated using a retrospective analytical observational design from January to September 2023. 658 patients satisfied the inclusion criteria.

Table 1 Patient Demographic Characteristics and Health Status

Variable	n	%
Gender		
Man	166	25.2
Woman	492	74.8
Age		
≥ 60 Years	302	45.9
50 – 59 Years	275	41.8
≤ 49 Years	81	12.3

Vitamin D levels		
Deficiency	555	84.3
Insufficiency	84	12.8
Sufficient	19	2.9
Diabetes Mellitus Status		
DM + DFU	30	4.6
DM	111	16.9
Non-DM	517	78.6

In this study, 658 samples of patients were treated at RSUP Dr. Mohammad Hoesin Palembang. 166 (25.2%) samples were male and 492 (74.8%) were female. The highest age group of the sample was in the group aged more than 60 years with 302 (45.9%) samples. In this study, there were 555 (84.3%) samples who had Vitamin D deficiency, there were 111 (16.9%) samples suffering from DM, and there were 30 (4.6%) samples with DFU (Table 1).

Table 2 Serum Vitamin D Levels of Patients at RSMH (ng/mL) (n=658)

	Mean	Min	Max
DM + DFU	11.727	5.5	23.3
DM	24.736	4.9	156
Non-DM	22.849	5.1	151.3
All Patients	22.660	4.9	156

DM= Diabetes Mellitus, DFU= Diabetic Foot Ulcer

The average vitamin D level across all patients in this research is 22,66 ng/mL; in the DM group, it is 24,736 ng/mL; and in patients with DFU (30), the average blood vitamin D level is around 11,727 ng/mL (Table 2).

Table 3 Association Between Serum Vitamin D Concentration and Diabetes Mellitus Patients in Mohammad Hoesin Hospital Palembang (n=658)

Variable	Deficiency		Insufficiency		Sufficient		Total		P-Value
	N	%	n	%	n	%	n	%	
DM + DFU	30	100%	0	0%	0	0%	30	100%	0,026
DM	90	81.1%	14	12.6%	7	6.3%	111	100%	
Non-DM	435	84.1%	70	13.5%	12	2.3%	517	100%	
Total	555	84,3%	84	12,8%	19	2,9%	658	100%	

DM= Diabetes Mellitus, DFU= Diabetic Foot Ulcer

The association between each hospital patient's vitamin D level and the Chi-Square test was examined. The study demonstrated a strong connection (p-value of 0.026) between the occurrence of diabetes mellitus (DM) in patients at Mohammad Hoesin General Hospital in Palembang and serum vitamin D deficiency (Table 3).

Table 4 Association Between Serum Vitamin D Concentration and DFU Patients in Mohammad Hoesin Hospital Palembang (n=120)

Variable	Deficiency		Insufficiency		Sufficient		Total		P-Value
	n	%	n	%	n	%	n	%	
DM + DFU	30	100%	0	0%	0	0%	30	100%	0,036
DM	90	81.1%	14	12.6%	7	6.3%	111	100%	
Total	120	85.1%	14	0.9%	7	0.5%	141	100%	

DM= Diabetes Mellitus, DFU= Diabetic Foot Ulcer

The chi-square test was used to look at the connection between vitamin D levels and diabetes mellitus with DFU. The Chi-Square test produced a p-value of 0.036 <0.05, indicating a possible correlation between the frequency of DFU patients in RSMH and the blood vitamin D level (Table 4).

Discussion

In this study, we analyzed 658 samples. The samples were divided into two groups DM and Non-DM. The samples consisted of 492 (74.8%) females and 166 (25.2%) males; in this study, we found females more than males. The largest sample age in the age group is over 60 years, with a total of 302 (45.9%) samples. A complication of DM is more frequently found in elderly patients because in the elderly group, there is more atherosclerosis of the vessels, especially in the lower extremity, and the immune system's capacity is declining, along with the body's resistance to infection and the inflammatory response.

From all of the 658 samples in this study, there were 517 (78.6%) samples without DM, 111 (16.9%) samples with DM, and 30 (4.6%) samples DM with DFU. Out of all the samples, we discovered that 555 (84.3%) had inadequate levels of vitamin D, 84 (12.8%) had inadequate levels, and 19 (2.9%) had appropriate levels. Many samples had vitamin D deficiency, both from patients. Diabetes Mellitus patients and patients who do not suffer from Diabetes Mellitus. It is well recognized that type 2 diabetic mellitus (T2DM) and vitamin D insufficiency have combined to become a contemporary epidemic (Vijay et al., 2023). A growing amount of evidence suggests that vitamin D deficiency may contribute to the onset and consequences of diabetes. Numerous studies have shown that a drop in 25(OH)D (25-hydroxy vitamin D) levels is linked to a forty percent higher risk of developing diabetes, while an increase in 25(OH)D concentration is linked to a twenty-four percent lower risk of developing diabetes (Bleizgys, 2021).

In this study, all DM patients with DFU (30; 4.6%) have vitamin D deficiency, with an average serum level of vitamin D 11.727 ng/mL. Samples from DM without DFU group have an average serum vitamin D level of 24.736 ng/mL. The average value of serum vitamin D levels in patients who do not suffer from diabetes mellitus is 22.849 ng/mL.

We used the Pearson chi-square test to examine the relationship between blood vitamin D levels and the prevalence of DM. A p-value of 0.026 indicated a significant relationship between vitamin D levels and the occurrence of diabetes mellitus. According to research by Arafat et al. (2020), vitamin D deficiency has been associated to poor control and outcomes in T2DM patients, and it is commonly observed in DFU patients. Furthermore, vitamin D has been shown by Mousa et al. (2018) to lessen persistent low-grade inflammation in people with type 2 diabetes (Arafat et al., 2020).

The relationship between blood vitamin D levels and DFU's Diabetes Mellitus status was examined using the chi-square test. With a p-value of 0.036 (<0.05), the findings indicated a

significant relationship between blood vitamin D levels and DM with DFU. Serum **vitamin D levels and DFU with infection have been linked in** several studies. For instance, vitamin D administration has been shown in the research by **8** Karonova et al. (2020) to improve clinical symptoms and decrease indicators of inflammation **in individuals with type 2 diabetes.**

Vitamin D plays a role in immune system regulation as well as in the generation of genes that code for two antimicrobial peptides (AMP) that fight infections and keep the right balance of skin flora: cathelicidin and β -defensin (Benson et al., 2023).

Vitamin D is associated with the control of infections and immune responses because it is a critical regulator of the innate immune response to the threat of bacterial infection. This has been stated in a number of papers. Decreased immune ability against microbial infections and excessive inflammatory response are the causes of delayed wound **12** healing in DFU patients (Furtado et al., 2020). In DFU patients, there is a decrease in levels of **Antimicrobial Peptides (AMP) such as cathelicidin (LL-37)**, an increase in inflammatory cytokines, and excessive biofilm formation (Wang et al., 2022).

According to Irma Gonzalez-Curiel et al., patients with DFU showed reduced antimicrobial peptides (AMP) levels in the lesion region, which may have impeded the healing process of their wounds. It showed the potential for vitamin D as supplementation therapy in treating DFU (Afonso et al., 2021; Gonzalez-Curiel et al., 2014; Mendoza-Marí et al., 2022; Miranda et al., 2023).

Several other studies found that patients with DM and DFU tended to have low serum vitamin D levels (Mozaffari-Khosravi et al., 2016; Wang et al., 2022). Thus, it is postulated that in individuals with diabetes mellitus and DFU, persistent inflammation and infections that are hard to cure are linked to vitamin D insufficiency (Lin et al., 2023; Y. Tang et al., 2023). The use of vitamin D can be considered as an additional therapy for DFU to reduce infection (Macido, 2018; W. Tang et al., 2022).

According to Dai et al. (2019), there is a strong **13** correlation between severe **vitamin D deficiency and a high incidence of diabetic foot ulcers**, as well **6** a large drop in vitamin D levels in diabetic foot ulcers. According to Zubair et al. (2013), low **level 5** of 25(OH)D are correlated with the ulcer's grade and have a major influence in the etiology **of foot ulcers**. According to **1** **wari et al.** (2014), patients with diabetes who also had ulcer infection had substantially lower 25(OH)D levels.

Conclusion

The average **vitamin D level in** 658 samples was 11.727 ng/mL. Of them, 492 (74.8%) were female, 302 (45.9%) were over 60, 555 (84.3%) had vitamin D insufficiency, 120 (85.1%) had DM, and sample DM had DFU 30 (100%). Every patient in the study had significant associations between **2** their serum vitamin D levels and diabetes mellitus (p-value of $0.026 < 0.05$) and between their **serum vitamin D levels and DFU** (p-value of $0.036 < 0.05$). This suggests that individuals with Diabetes Mellitus, both those with and those without DFU, have low blood levels of Vitamin D.

Our research led us to conclude that, in order to effectively treat individuals with diabetic foot ulcers, vitamin D must be given and blood vitamin D levels must be monitored.

Bibliography

- Afonso, A. C., Oliveira, D., Saavedra, M. J., Borges, A., & Simões, M. (2021). Biofilms in Diabetic Foot Ulcers: Impact, Risk Factors and Control Strategies. *International Journal of Molecular Sciences*, 22(15), 8278. <https://doi.org/10.3390/ijms22158278>
- Arafat, E. S., Taha, I. M., Kattan, S. W., Babteen, N. A., & Fawzy, I. (2020). Associations between Vitamin D and Type 2 Diabetes Mellitus: The Role of Vitamin D Receptor and Binding Protein. *Journal of Diabetes Mellitus*, 10(04), 222–235. <https://doi.org/10.4236/jdm.2020.104018>
- Benson, R., Unnikrishnan, M. K., Kurian, S. J., Velladath, S. U., Rodrigues, G. S., Chandrashekar Hariharapura, R., Muraleedharan, A., Bangalore Venkateshiah, D., Banerjee, B., Mukhopadhyay, C., Johnson, A. S., Munisamy, M., Rao, M., Kochikuzhyil, B. M., & Sekhar Miraj, S. (2023). Vitamin D attenuates biofilm-associated infections via immunomodulation and cathelicidin expression: a narrative review. *Expert Review of Anti-Infective Therapy*, 21(1), 15–27. <https://doi.org/10.1080/14787210.2023.2151439>
- Bleizgys, A. (2021). Vitamin D Dosing: Basic Principles and a Brief Algorithm (2021 Update). *Nutrients*, 13(12), 4415. <https://doi.org/10.3390/nu13124415>
- Caley, M. P., Martins, V. L. C., & O'Toole, E. A. (2015). Metalloproteinases and Wound Healing. *Advances in Wound Care*, 4(4), 225–234. <https://doi.org/10.1089/wound.2014.0581>
- Cimmino, G., Ragni, M., Cirillo, P., Petrillo, G., Loffredo, F., Chiariello, M., Gresele, P., Falcinelli, E., & Golino, P. (2013). C-reactive protein induces expression of matrix metalloproteinase-9: A possible link between inflammation and plaque rupture. *International Journal of Cardiology*, 168(2), 981–986. <https://doi.org/10.1016/j.ijcard.2012.10.040>
- Dadaei, T., Safapoor, M. H., Asadzadeh Aghdaei, H., Balaii, H., Pourhoseingholi, M. A., Naderi, N., Zojaji, H., Azimzadeh, P., Mohammadi, P., & Zali, M. R. (2015). Effect of vitamin D3 supplementation on TNF- α serum level and disease activity index in Iranian IBD patients. *Gastroenterology and Hepatology from Bed to Bench*, 8(1), 49–55.
- Dai, J., Jiang, C., Chen, H., & Chai, Y. (2019). Vitamin D and diabetic foot ulcer: a systematic review and meta-analysis. *Nutrition & Diabetes*, 9(1), 8. <https://doi.org/10.1038/s41387-019-0078-9>
- Furtado, S. C., Srinivasan, B., & Abraham, S. (2020). Wound healing concepts: contemporary practices and future perspectives. *Int J App Pharm*, 12(5), 7–15.
- Ghezel, A., Salekzamani, S., Mehralizadeh, H., Jafarabadi, M. A., & Gargari, B. P. (2019). Vitamin D supplementation has no effect on matrix metalloproteinases-2, -9, and tissue inhibitor matrix metalloproteinase-1 in subjects with metabolic syndrome: A pilot study. *International Journal for Vitamin and Nutrition Research*, 89(5–6), 227–237. <https://doi.org/10.1024/0300-9831/a000559>
- Giatsidis, G., Orgill, D. P., & Yannas, I. V. (2018). *Dermal Regeneration and Induction of Wound Closure in Diabetic Wounds* (pp. 155–172). https://doi.org/10.1007/978-3-319-89869-8_9
- Gonzalez-Curiel, I., Trujillo, V., Montoya-Rosales, A., Rincon, K., Rivas-Calderon, B., deHaro-Acosta, J., Marin-Luevano, P., Lozano-Lopez, D., Enciso-Moreno, J. A., & Rivas-Santiago, B. (2014). 1,25-Dihydroxyvitamin D3 Induces LL-37 and HBD-2 Production in Keratinocytes from Diabetic Foot Ulcers Promoting Wound Healing: An In Vitro Model. *PLoS ONE*, 9(10), e111355. <https://doi.org/10.1371/journal.pone.0111355>
- Halschou-Jensen, P. M., Sauer, J., Bouchelouche, P., Fabrin, J., Brorson, S., & Ohrt-Nissen, S. (2023). Improved Healing of Diabetic Foot Ulcers After High-dose Vitamin D: A Randomized Double-blinded Clinical Trial. *The International Journal of Lower Extremity Wounds*, 22(3), 466–474. <https://doi.org/10.1177/15347346211020268>
- Harrisa, S., Suhartono, R., & Kekalih, A. (2020). Correlation between Serum Level of Vitamin D and Severity Degree of Diabetic Foot Ulcer in Patients with Normal Ankle Brachial Index. *Journal of Indonesian Society for Vascular and Endovascular Surgery*, 1(1), 34–36.

- Holick, M. F., Binkley, N. C., Bischoff-Ferrari, H. A., Gordon, C. M., Hanley, D. A., Heaney, R. P., Murad, M. H., & Weaver, C. M. (2011). Evaluation, Treatment, and Prevention of Vitamin D Deficiency: an Endocrine Society Clinical Practice Guideline. *The Journal of Clinical Endocrinology & Metabolism*, *96*(7), 1911–1930. <https://doi.org/10.1210/jc.2011-0385>
- Huang, S.-M., Wu, C.-S., Chiu, M.-H., Wu, C.-H., Chang, Y.-T., Chen, G.-S., & Lan, C.-C. E. (2019). High glucose environment induces M1 macrophage polarization that impairs keratinocyte migration via TNF- α : An important mechanism to delay the diabetic wound healing. *Journal of Dermatological Science*, *96*(3), 159–167. <https://doi.org/10.1016/j.jdermsci.2019.11.004>
- Ji, S., Liu, X., Huang, J., Bao, J., Chen, Z., Han, C., Hao, D., Hong, J., Hu, D., Jiang, Y., Ju, S., Li, H., Li, Z., Liang, G., Liu, Y., Luo, G., Lv, G., Ran, X., Shi, Z., ... Xia, Z. (2021). Consensus on the application of negative pressure wound therapy of diabetic foot wounds. *Burns & Trauma*, *9*. <https://doi.org/10.1093/burnst/tkab018>
- Kamble, A., & Swamkar, M. (2019). A study of the prevalence and severity of vitamin D deficiency in patient with diabetic foot and its association with vascular calcification and effect on healing. *International Surgery Journal*, *6*(5), 1654. <https://doi.org/10.18203/2349-2902.isj20191886>
- Karonova, T., Stepanova, A., Bystrova, A., & Jude, E. B. (2020). High-Dose Vitamin D Supplementation Improves Microcirculation and Reduces Inflammation in Diabetic Neuropathy Patients. *Nutrients*, *12*(9), 2518. <https://doi.org/10.3390/nu12092518>
- Kinesya, E., Santoso, D., Gde Arya, N., Putri Cintya, E., Seriari Ambarini, P., Kinesya, B., Stephanie Kartjito, M., & Mannagalli, Y. (2023). Vitamin D as adjuvant therapy for diabetic foot ulcers: Systematic review and meta-analysis approach. *Clinical Nutrition ESPEN*, *54*, 137–143. <https://doi.org/10.1016/j.clnesp.2023.01.011>
- Lin, J., Mo, X., Yang, Y., Tang, C., & Chen, J. (2023). Association between vitamin D deficiency and diabetic foot ulcer wound in diabetic subjects: A meta-analysis. *International Wound Journal*, *20*(1), 55–62. <https://doi.org/10.1111/iwj.13836>
- Lobmann, R., Ambrosch, A., Schultz, G., Waldmann, K., Schiweck, S., & Lehnert, H. (2002). Expression of matrix-metalloproteinases and their inhibitors in the wounds of diabetic and non-diabetic patients. *Diabetologia*, *45*(7), 1011–1016. <https://doi.org/10.1007/s00125-002-0868-8>
- López-López, N., González-Curiel, I., Treviño-Santa Cruz, M. B., Rivas-Santiago, B., Trujillo-Paez, V., Enciso-Moreno, J. A., & Serrano, C. J. (2014). Expression and vitamin D-mediated regulation of matrix metalloproteinases (MMPs) and tissue inhibitors of metalloproteinases (TIMPs) in healthy skin and in diabetic foot ulcers. *Archives of Dermatological Research*, *306*(9), 809–821. <https://doi.org/10.1007/s00403-014-1494-2>
- Macido, A. (2018). Diabetic Foot Ulcers and Vitamin D Status: A Literature Review. *SAGE Open Nursing*, *4*, 237796081878902. <https://doi.org/10.1177/2377960818789027>
- Marcinowska-Suchowierska, E., Kupisz-Urbańska, M., Łukaszkiwicz, J., Płudowski, P., & Jones, G. (2018). Vitamin D Toxicity—A Clinical Perspective. *Frontiers in Endocrinology*, *9*. <https://doi.org/10.3389/fendo.2018.00550>
- Masood, M. Q., Khan, A., Awan, S., Dar, F., Naz, S., Naureen, G., Saghir, S., & Jabbar, A. (2015). Comparison of Vitamin D Replacement Strategies with High-Dose Intramuscular or Oral Cholecalciferol: A Prospective Intervention Study. *Endocrine Practice*, *21*(10), 1125–1133. <https://doi.org/10.4158/EP15680.OR>
- Mendoza-Marí, Y., García-Ojalvo, A., Fernández-Mayola, M., Rodríguez-Rodríguez, N., Martínez-Jimenez, I., & Berlanga-Acosta, J. (2022). Epidermal growth factor effect on lipopolysaccharide-induced inflammation in fibroblasts derived from diabetic foot ulcer. *Scars, Burns & Healing*, *8*, 20595131211067380. <https://doi.org/10.1177/20595131211067380>
- Miranda, E., Bramono, K., Yunir, E., Reksodiputro, M. H., Suwarsa, O., Rengganis, I., Harahap, A. R., Subekti, D., Suwanto, S., Hayun, H., Bardosono, S., & Baskoro, J. C. (2023). Efficacy

- of LL-37 cream in enhancing healing of diabetic foot ulcer: a randomized double-blind controlled trial. *Archives of Dermatological Research*, 315(9), 2623–2633. <https://doi.org/10.1007/s00403-023-02657-8>
- Mousa, A., Naderpoor, N., Teede, H., Scragg, R., & de Courten, B. (2018). Vitamin D supplementation for improvement of chronic low-grade inflammation in patients with type 2 diabetes: a systematic review and meta-analysis of randomized controlled trials. *Nutrition Reviews*, 76(5), 380–394. <https://doi.org/10.1093/nutrit/nux077>
- Mozaffari-Khosravi, H., Haratian-Arab, M., Tavakkoli, H. M., & Nadjarzadeh, A. (2016). Comparative effect of two different doses of vitamin D on diabetic foot ulcer and inflammatory indices among the type 2 diabetic patients: a randomized clinical trial. *Iranian Journal of Diabetes and Obesity*, 8(4), 164–171.
- Muller, M., Trocme, C., Lardy, B., Morel, F., Halimi, S., & Benhamou, P. Y. (2008). Matrix metalloproteinases and diabetic foot ulcers: the ratio of MMP-1 to TIMP-1 is a predictor of wound healing. *Diabetic Medicine*, 25(4), 419–426. <https://doi.org/10.1111/j.1464-5491.2008.02414.x>
- Rastogi, A., Bhansali, A., Khare, N., Suri, V., Yaddanapudi, N., Sachdeva, N., Puri, G. D., & Malhotra, P. (2022). Short term, high-dose vitamin D supplementation for COVID-19 disease: a randomised, placebo-controlled, study (SHADE study). *Postgraduate Medical Journal*, 98(1156), 87–90. <https://doi.org/10.1136/postgradmedj-2020-139065>
- Razzaghi, R., Pourbagheri, H., Momen-Heravi, M., Bahmani, F., Shadi, J., Soleimani, Z., & Asemi, Z. (2017). The effects of vitamin D supplementation on wound healing and metabolic status in patients with diabetic foot ulcer: A randomized, double-blind, placebo-controlled trial. *Journal of Diabetes and Its Complications*, 31(4), 766–772. <https://doi.org/10.1016/j.jdiacomp.2016.06.017>
- Ross, A. C., Taylor, C. L., Yaktine, A. L., & Del Valle, H. B. (2011). Tolerable upper intake levels: Calcium and Vitamin D. In *Dietary reference intakes for calcium and vitamin D*. National Academies Press (US).
- Shin, M. H., Lee, Y., Kim, M.-K., Lee, D. H., & Chung, J. H. (2019). UV increases skin-derived 1 α ,25-dihydroxyvitamin D3 production, leading to MMP-1 expression by altering the balance of vitamin D and cholesterol synthesis from 7-dehydrocholesterol. *The Journal of Steroid Biochemistry and Molecular Biology*, 195, 105449. <https://doi.org/10.1016/j.jsbmb.2019.105449>
- Tang, W., Chen, L., Ma, W., Chen, D., Wang, C., Gao, Y., & Ran, X. (2022). Association between vitamin D status and diabetic foot in patients with type 2 diabetes mellitus. *Journal of Diabetes Investigation*, 13(7), 1213–1221. <https://doi.org/10.1111/jdi.13776>
- Tang, Y., Huang, Y., Luo, L., Xu, M., Deng, D., Fang, Z., Zhao, X., & Chen, M. (2023). Level of 25-hydroxyvitamin D and vitamin D receptor in diabetic foot ulcer and factor associated with diabetic foot ulcers. *Diabetology & Metabolic Syndrome*, 15(1), 30. <https://doi.org/10.1186/s13098-023-01002-3>
- Tiwari, S., Pratyush, D. D., Gupta, S. K., & Singh, S. K. (2014). Vitamin D deficiency is associated with inflammatory cytokine concentrations in patients with diabetic foot infection. *British Journal of Nutrition*, 112(12), 1938–1943. <https://doi.org/10.1017/S0007114514003018>
- Vijay, G. S., Ghonge, S., Vajjala, S. M., & Palal, D. (2023). Prevalence of Vitamin D Deficiency in Type 2 Diabetes Mellitus Patients: A Cross-Sectional Study. *Cureus*. <https://doi.org/10.7759/cureus.38952>
- Wang, F., Zhou, L., Zhu, D., & Yang, C. (2022). A Retrospective Analysis of the Relationship Between 25-OH-Vitamin D and Diabetic Foot Ulcer. *Diabetes, Metabolic Syndrome and Obesity: Targets and Therapy*, Volume 15, 1347–1355. <https://doi.org/10.2147/DMSO.S358170>
- Yamine, K., Hayek, F., & Assi, C. (2020). Is there an association between vitamin D and diabetic foot disease? A meta-analysis. *Wound Repair and Regeneration*, 28(1), 90–96. <https://doi.org/10.1111/wrr.12762>

- Yunir, E., Tarigan, T. J. E., Iswati, E., Sarumpaet, A., Christabel, E. V., Widiyanti, D., Wisnu, W., Purnamasari, D., Kurniawan, F., Rosana, M., Anestherita, F., Muradi, A., & Tahapary, D. L. (2022). Characteristics of Diabetic Foot Ulcer Patients Pre- and During COVID-19 Pandemic: Lessons Learnt From a National Referral Hospital in Indonesia. *Journal of Primary Care & Community Health*, 13, 215013192210897. <https://doi.org/10.1177/21501319221089767>
- Zubair, M., Malik, A., Meerza, D., & Ahmad, J. (2013). 25-Hydroxyvitamin D [25(OH)D] levels and diabetic foot ulcer: Is there any relationship? *Diabetes & Metabolic Syndrome: Clinical Research & Reviews*, 7(3), 148–153. <https://doi.org/10.1016/j.dsx.2013.06.008>

Association of Vitamin D Deficiency and Diabetic Foot Ulcer Patients in Dr Mohammad Hoesin General Hospital Palembang

ORIGINALITY REPORT

12%

SIMILARITY INDEX

11%

INTERNET SOURCES

11%

PUBLICATIONS

%

STUDENT PAPERS

PRIMARY SOURCES

1	mdpi-res.com Internet Source	3%
2	"Handbook of vitamin D in human health", Brill, 2013 Publication	1%
3	jusindo.publikasiindonesia.id Internet Source	1%
4	Muhammad Holid Fajar, Anita Fira Waluyo. "Sistem Monitoring dan Penyiraman Otomatis Pada Bibit Sawit Berbasis Android", Infotek: Jurnal Informatika dan Teknologi, 2024 Publication	1%
5	"Extraskeletal Effects of Vitamin D", Springer Science and Business Media LLC, 2018 Publication	1%
6	mts.intechopen.com Internet Source	1%
7	mro.massey.ac.nz Internet Source	

1 %

8

"Handbook of nutrition in heart health", Brill, 2017

Publication

1 %

9

link.springer.com

Internet Source

1 %

10

www.frontiersin.org

Internet Source

1 %

11

www.researchgate.net

Internet Source

1 %

12

www.wjgnet.com

Internet Source

1 %

13

Ying Tang, Yixuan Huang, Li Luo, Murong Xu, Datong Deng, Zhaohui Fang, Xiaotong Zhao, Mingwei Chen. "Decreased expression of 25-hydroxyvitamin D in peripheral plasma and vitamin D receptor in wound margin tissue associated with diabetic foot ulcers in patients with type 2 diabetes mellitus", Research Square Platform LLC, 2022

Publication

1 %

Exclude quotes On

Exclude matches < 1%

Exclude bibliography On

