

BUKTI KORESPONDENSI
ARTIKEL DI JURNAL TERINDEX SINTA 2


Judul artikel : An Aeration Pressure, Plant Spacing, And Dissolved Oxygen For
Growth Of Lettuce (Lactuca Sativa L.) Plants In Floating
Hydroponics

Jurnal : Jurnal Penelitian Pertanian Terapan (JPPT)

Penulis : **Puspitahati**, Fidel Harmanda Prima, Nurul Izzah, Nikita Prettisha

No.	Perihal	Tanggal
1.	Bukti konfirmasi submit artikel	17 September 2024
2.	Bukti konfirmasi artikel masuk ke editorial	17 September 2024
3.	Bukti konfirmasi submission accepted	27 September 2024
4.	Bukti proses review artikel	27 september 2024
5.	Bukti sudah mengirimkan artikel yang sudah direvisi	4 Januari 2025
6.	Bukti konfirmasi Accepted dan publish online	5 januari 2025

1. Bukti Konfirmasi Submit artikel



Comments for the Editor


Participants [Edit](#)

puspitahtati puspitahtati (pita)


Messages


Note	From
Kepada Yth. Editor Jurnal Penelitian Pertanian Terapan Dengan hormat, Kami mengajukan artikel kami yang berjudul : "Aeration Pressure, Plant Spacing, And Dissolved Oxygen For Growth Of Lettuce (Lactuca Sativa L.) Plants In Floating Hydroponics" untuk dipertimbangkan dalam publikasi di jurnal ini. Kami juga ingin menginformasikan bahwa artikel ini belum pernah dipublikasikan. Kami juga menyediakan Link untuk Referensi pada artikel ini (https://drive.google.com/drive/folders/11w2h2FTRMNFRX6c-MQ3RhGLhXKVbeK7B?usp=sharing) Kami sangat berharap artikel ini dapat memberikan kontribusi yang signifikan dalam pengembangan teknik hidroponik dan pertanian terapan. Semoga artikel ini dapat diterima di Jurnal ini. Terima kasih atas perhatian dan kesempatannya. [Dr. Puspitahtati, S.T.P., M.P]	pita2024-09-16 05:40 PM
▶ Kepada Yth Editor Jurnal Penelitian Pertanian Terapan MOhon maaf sebelumnya, saya mau bertanya, apakah artikel kami ini dapat diterima dan dilanjutkan untuk di review? mohon informasinya Bapak/IBu, Terimakasih Banyak	pita2024-10-04 09:22 PM

Add Message



2. Bukti konfirmasi artikel masuk ke editorial





3. Bukti konfirmasi submission accepted

Notifications

[JPPT] Editor Decision

2025-01-09 06:40 AM

puspitahati puspitahati: The editing of your submission, "A Aeration Pressure, Plant Spacing, And Dissolved Oxygen For Growth Of Lettuce (Lactuca Sativa L.) Plants In Floating Hydroponics," is complete. We are now sending it to production. Submission URL:

<https://jurnal.polinela.ac.id/jppt/authorDashboard/submission/3722>

Jurnal Penelitian Pertanian Terapan <http://jurnal.polinela.ac.id/index.php/JPPT>

Round 1

Round 1 Status

Submission accepted.

Notifications

[\[JPPT\] Editor Decision](#)

2025-01-05 01:06 PM

[\[JPPT\] Editor Decision](#)

2025-01-09 06:40 AM

Reviewer's Attachments

14926-1

, 3722-Article Text-14757-1-4-20241128.doc

December 27, 2024

Search

Revisions

Search

Upload File

4. bukti proses review/perbaikan (27 september 2024)

Dokumen review :

jppt@polinela.ac.id

Aktif

Google Apps

Perbaikan JPPT

Eksternal

Kotak Masuk x

JPPT UPPM

<jppt@polinela.ac.id>

kepada saya

27 Des 2024, 10.11

Yth. Author JPPT

Mohon untuk melakukan perbaikan , sesuai saran dari reviewer.

Terima Kasih

Satu lampiran • Dipindai dengan Gmail

3722-Article Text...

puspitahti fp

<puspitahti@fp.unsri.ac.id>

kepada JPPT

29 Des 2024, 08.08

Baik bapak /ibu-. terimakasih atas informasinya

ff 3722-14926-1-5-20241227 (1) - Compatibility Mode • Saved

sert Draw Design Layout References Mailings Review View Zotero Help Grammarly

Font Paragraph Styles

ABSTR INDO ABSTRACT AFILIASI

Find Replace Select Add-ins Open Grammar

eration pressure correlates with increased dissolved oxygen levels, suggesting that higher aeration pressure correlates with increased dissolved oxygen levels. The Suggestion was Further research should be conducted on the appropriate size of the nutrient solution reservoir concerning the aeration pressure applied.

Keywords: Lettuce (*Lactuca Sativa L.*), Dissolved Oxygen, Aeration Pressure, Plant Spacing, Floating Hydroponics

Submitted: September **Received:** **Approved:**

INTRODUCTION

The number of Indonesia's population is increasing, and the need for horticultural commodities, especially vegetables, every year is always increasing in line with the increasing demand for lettuce (*Lactuca sativa L.*) (Pujiiswanto and Pangaribuan, 2010). Lettuce (*Lactuca sativa L.*) is a plant that is very popular among urban communities. This is because, in addition to being nutritious, it is also good for the body's vitamin needs (Cahyani, Hasibuan and Ch, 2019). Lettuce (*Lactuca sativa L.*) is one of the vegetable plants that are consumed by the public in fresh form (Warganegara *et al.*, 2015). Furthermore, lettuce is very effective when grown using hydroponics (Hamyana, Budianto and Bukori, 2023). Lettuce is cultivated hydroponically without using soil (Mardina *et al.*, 2019) and utilizes a nutrient solution as a source of essential elements (Zenita, 2019); (Umarie, 2020); (Sipin Solis and U. Gabutan, 2023). One of the nutrients commonly used in hydroponics is AB Mix fertilizer (Nazara *et al.*, 2023); (Lestari, Rahayu and Mulyaningsih, 2022); (Romalasari and Sobari, 2019) which is capable of increasing plant height, leaf count, and leaf width. (Zahririma, Sutariati and Rakian, 2019). In order to grow well, plants cultivated hydroponically need to get complete nutrients, which consist of macro (N, P, K, Ca, Mg, S) and micro (Cl, Mn, Fe, Cu, Zn, B, and Mo) elements (Warjoto *et al.*, 2020).

SKD 2020
Please add at least one paragraph regarding the novelty of this research and its benefits from both economic and non-economic perspectives.

English (United States)

ere to search

1 The research was conducted from May 2021 to September 2021 at the plant house on Jalan Raoang Selasa, Bukit Besar, Palembang, and at the Basic Fisheries Laboratory, Department of Fisheries, Agriculture, Faculty Sriwijaya University. The tools used in this research were 1) Aerator Set, 2) Punching Tool, 3) Writing Instruments, 4) Floating Raft System Tank, 5) DO Meter, 6) Camera, 7) Seedling Tray, 8) Netpot, 9) Ruler, 10) pH Meter, 11) Stirring Spoon, 12) Styrofoam, 13) TDS EC Meter, 14) Thermohygrometer, 15) Analytical Scale. The materials used in the implementation of this research were: 1) Water, 2) Lettuce seeds, 3) AB mix nutrients, and 4) Rockwool.

This research uses a Factorial Randomized Block Design (FRBD) with two factors, as follows: The first factor is planting distance (D) which consists of two levels, namely: D1 = 20 cm, D2 = 30 cm, and the second factor is aeration pressure (P) which consists of three levels, namely: P1 = 12 kPa, P2 = 14 kPa, B3 = 16 kPa. Each treatment is replicated three times. The research data are examined using analysis of variance (ANOVA) with a 5% significance threshold, and the results are further analyzed with the Honest Significant Difference (HSD) test at the same 5% level. A tabulation method is employed to observe the effects of dissolved oxygen, pH, and EC during each week of plant growth. The parameters observed in this study are as follows: Plant Height (cm), Number of Leaves (pieces), Canopy Width (cm), Root Length (cm), Total Fresh Weight Per Plant (g), and dissolved oxygen (ml/l).

RESULTS AND DISCUSSION

The environmental conditions during the research were as follows: the air temperature was 28.1°C, the air humidity was 72.5%, the pH of the nutrient solution ranged from 6.5 to 6.6, and the electrical conductivity (EC) of the nutrient solution was between 1070 and 1103 mS/cm. The results of the significance

SKD 2020
references in the discussion are still few, please add and sharpen them, if possible there is a discussion from the economic side related to the application of hydroponics

Volume X Nomor X, September 2024 Hal 2

5. Bukti sudah mengirimkan artikel yang sudah direvisi (4 januari 2025)

REVISION

Participants [Edit](#)

Edy Humaidi (edyhumaidi145)

puspitahtati puspitahtati (pita)

Messages


Note	From
Yang terhormat dewan reviewer dan editor jurnal berikut saya kirimkan revisi artikel yang sudah diperbaiki terimakasih atas perhatiannya mohon diproses lebih lanjut wasalam	pita2025-01-04 10:06 AM

[pita, 3722-Article Text-14757-puspitahtati_Revisi.docx](#)

Add Message

jppt@polinela.ac.id

4 dari 7


**puspitahtati fp** <puspitahtati@fp.unsri.ac.id>
kepada JPPT

4 Jan 2025, 16:59

Yth Dewan Editor dan Reviewer Jurnal JPT UPPM (POLinela)
berikut saya kirimkan revisi artikel yang berjudul :An Aeration Pressure, Plant Spacing, And Dissolved Oxygen For Growth Of Lettuce (Lactuca Sativa L.) Plants In Floating Hydroponics
mohon kiranya dapat diterima dan diproses, terimakasih atas perhatiannya. wassalam

Pada Jum, 27 Des 2024 pukul 10.11 JPPT UPPM <jppt@polinela.ac.id> menulis:
Yth . Author JPPT
Mohon untuk melakukan perbaikan , sesuai saran dari reviewer.
Terima Kasih

Satu lampiran • Dipindai dengan Gmail


3722-Article Text...

INTRODUCTION

The number of Indonesia's population is increasing, and the need for horticultural commodities, especially vegetables, every year is always increasing in line with the increasing demand for lettuce (*Lactuca sativa* L.) (Pujisiswanto and Pangaribuan, 2010). Lettuce (*Lactuca sativa* L.) is a plant that is very popular among urban communities. This is because, in addition to being nutritious, it is also good for the body's vitamin needs (Cahyani, Hasibuan and Ch, 2019). Lettuce (*Lactuca sativa* L.) is one of the vegetable plants that are consumed by the public in fresh form (Warganegara *et al.*, 2015). Furthermore, lettuce is very effective when grown using hydroponics (Hamyana, Budianto and Bukori, 2023). Lettuce is cultivated hydroponically without using soil (Mardina *et al.*, 2019) and utilizes a nutrient solution as a source of essential elements (Zenita, 2019); (Umarie, 2020); (Sipin Solis and U. Gabutan, 2023). One of the nutrients commonly used in hydroponics is AB Mix fertilizer (Nazara *et al.*, 2023);(Lestari, Rahayu and Mulyaningsih, 2022); (Romalasari and Sobari, 2019) which is capable of increasing plant height, leaf count, and leaf width. (Zahrima, Sutariati and Rakian, 2019)

RESULTS AND DISCUSSION

The environmental conditions during the research were as follows: the air temperature was 28.1°C, the air humidity was 72.5%, the pH of the nutrient solution ranged from 6.5 to 6.6, and the electrical conductivity (EC) of the nutrient solution was between 1070 and 1103 mS/cm. The results of the significance test indicate that planting distance does not significantly impact plant height, whereas aeration pressure does have a significant effect on plant height, as detailed in Table 1.

Table 1. BNJ 5% test table of the effect of planting distance on the height of lettuce plants (cm) from 1 MST to 4 WAP.

Plant Age	Factor	Average	BNJ 0,05
1 WAP	D1	0,828 a	0,131
	D2	0,882 a	0,131
2 WAP	D1	2,453 a	0,164
	D2	2,476 a	0,164
3 WAP	D1	7,435 a	0,288
	D2	7,515 a	0,288
4 WAP	D1	11,608 a	0,248
	D2	11,785 a	0,248

Note: The average values followed by different letters in the same column indicate a significant effect.

The results of the significance test indicate that planting distance does not have a significant effect on plant height, suggesting that other factors, such as environmental conditions or genetic factors, might play a more important role in determining plant height. On the other hand, aeration pressure significantly impacts plant height, as supported by research showing that soil aeration can affect plant growth by enhancing root oxygen availability, which is crucial for root respiration and nutrient absorption (Ben-Noah and Friedman, 2018);(Li *et al.*, 2019).

According to the BNJ test results at a 5% significance level, planting distances ranging from 1 to 4 WAP do not significantly impact plant height. However, the standard planting distance for lettuce, which is 14 cm at 35 HST, does influence leaf growth and plant height (Vu Phong *et al.*, 2019). On the other hand, aeration pressure does have a noticeable effect on plant height. At 4 MST, the tallest plant was observed in treatment P0 at 11.909 cm, while the shortest was in treatment P1 at 11.401 cm. The results of the 5% got the effect of aeration pressure on plant height (in cm) from 1 to 4 WAP are presented in Table 2.

Table 2. The Results of the 5% BNJ test on the effect of aeration pressure on plant height (cm) from 1 to 4 WAP.

Plant Age	Factor	Average	BNJ 0,05
1 WAP	P3	0,728 a	0,252
	P2	0,773 ab	0,252
	P1	0,914 ab	0,252
	P0	1,006 b	0,252
2 WAP	P3	2,217 a	0,315
	P2	2,274 a	0,315
	P0	2,651 b	0,315
	P1	2,717 b	0,315
3 WAP	P3	7,125 a	0,552
	P2	7,299 ab	0,552
	P1	7,664 ab	0,552
	P0	7,813 b	0,552
4 WAP	P1	11,401 a	0,476
	P2	11,626 ab	0,476
	P3	11,850 ab	0,476
	P0	11,909 b	0,476

Note: The average values followed by different letters in the same column indicate a significant effect.

The application of aeration pressure on plants can assist in root growth and obtain dissolved oxygen in water, thereby enhancing the absorption of nutrients (Virha and Bayfurqon, 2020), specifically, the nutrients N, P, and K found in the AB MIX nutrient content that can promote plant growth. (Soleha, Suroso and Wijaya, 2021). The impact of planting distance on the number of lettuce leaves, as determined by the BNJ 5% significance level, is detailed in Table 3.

Table 3. BNJ 5% Test of the effect of planting distance on the number of lettuce leaves from 1 MST to 4 WAP.

Plant Age	Factor	Average	BNJ 0,05
1 WAP	D1	5,5 a	0,390
	D2	5,7 a	0,390
2 WAP	D1	7,8 a	0,475
	D2	8,2 a	0,475
3 WAP	D1	10,9 a	0,468
	D2	11,5 b	0,468
4 WAP	D1	13,8 a	0,468
	D2	14,3 b	0,468

Note: The average values followed by different letters in the same column indicate a significant effect.

The BNJ test at the 5% level showed that the planting distance treatment significantly affected the number of plant leaves between 3 and 4 WAP, a greater planting distance can enhance the number of leaves because the wider the planting distance, the lower the competition among roots in absorbing nutrients (Kurniawati, 2020a). The AB Mix solution can stimulate leaf growth if its absorption by the roots is done optimally (Sudewi *et al.*, 2022), allowing each plant to grow well. The results of the BNJ 5% test on aeration pressure against the number of lettuce leaves can be seen in Table 4.

Table 4. BNJ 5% effect of aeration pressure on the number of leaves of lettuce aged 1 to 4 WAP

Plant Age	Factor	Average	BNJ 0,05
1 WAP	P0	5,2 a	0,748
	P1	5,3 ab	0,748
	P2	5,8 ab	0,748
	P3	6,0 b	0,748
2 WAP	P0	7,4 a	0,911
	P1	7,7 ab	0,911
	P2	8,5 b	0,911
	P3	8,5 b	0,911
3 WAP	P1	10,5 a	0,897
	P2	10,8 ab	0,897
	P3	11,7 b	0,897
	P4	11,8 b	0,897
4 WAP	P1	13,3 a	0,897
	P2	13,8 a	0,897
	P3	14,2 ab	0,897
	P4	14,8 b	0,897

Note: Average values followed by different letters in the same column indicate a significant effect.

In Table 4 of the BNJ 5% test results, it is evident that aeration pressure significantly affected the number of lettuce leaves from 1 to 4 WAP. This is because higher levels of dissolved oxygen in the water improve nutrient absorption for plants and support the respiration process, leading to an increased number of plant leaves (Dharmayanti, Sumiyati and Yulianti, 2021).

Aerating nutrient solutions generally enhances plant growth, including leaf production, by improving nutrient and water uptake. Specifically, aeration boosts the absorption of essential nutrients such as potassium (K), phosphorus (P), and magnesium (Mg) (Camarillo *et al.*, 2020). Aeration pressure also influences plant respiration and photosynthesis processes. It has been observed that as total pressure increases in high humidity, the respiration rate decreases linearly, while the net photosynthesis rate can increase up to 1.6 times due to higher total pressure and CO₂ partial pressure (Takeishi *et al.*, 2013). At low pressures, gas diffusivity increases, impacting the rate of CO₂ absorption and water transpiration through the stomata. Stomatal conductance, which regulates gas exchange, is inversely related to stomatal opening, thereby affecting plant respiration (Gohil, Correll and Sinclair, 2011).

Table 5. BNJ 5% test table for the effect of planting distance on the canopy width (cm) of lettuce plants aged 1 to 4 WAP.

Plant Age	Factor	Average	BNJ 0,05
1 WAP	D1	3,990 a	0,276
	D2	4,239 a	0,276
2 WAP	D1	16,260 a	0,677
	D2	16,986 b	0,677
3 WAP	D1	28,037 a	0,407
	D2	30,000 b	0,407

4 WAP	D1	29,372 a	0,191
	D2	31,200 b	0,191

Note: Average values in the same column that are marked with different letters indicate a significant difference.

The BNJ test results at a 5% significance level reveal that planting distance significantly affects canopy width and plant growth at ages 2 to 4 WAP. It is suggested that plants will perform better if spaced appropriately, allowing them to absorb nutrients more effectively and grow to their full potential. Consequently, increasing the planting distance will benefit both the canopy width and overall plant growth. At 4 WAP, treatment P3 had an average canopy width of 30.721 cm, which was not different from treatment B2 but differed from treatments P0 and P1. Treatment P0 had an average canopy width of 30.232 cm, which was not different from P3 but differed from treatments P1 and P2. According to (Puspitaningrum et al., 2012), the dissolved oxygen content an important role in the plant respiration process, which causes the plant canopy to become wider. DO levels are affected by photosynthesis and respiration. Plants produce oxygen during photosynthesis, which can increase DO levels in the surrounding water. On the other hand, respiration consumes oxygen, so DO levels decrease (Sunardi et al., 2020); (Akomeah and Lindenschmidt, 2017) An increase in DO can lead to a larger leaf area index (LAI), which is a measure of leaf area per unit area of soil and is directly related to the width of the canopy. (Ouyang et al., 2021).

Table 6. BNJ 5% test results for the impact of aeration pressure on lettuce canopy width (cm) at ages 1 to 4 WAP.

Plant Age	Factor	Average	BNJ 0,05
1 WAP	P0	3,958 a	0,530
	P3	4,039 a	0,530
	P1	4,167 a	0,530
	P2	4,293 a	0,530
2 WAP	P0	16,060 a	1,299
	P3	16,429 a	1,299
	P1	16,935 a	1,299
	P2	17,069 a	1,299
3 WAP	P0	28,457 a	0,780
	P3	28,925 ab	0,780
	P1	29,258 b	0,780
	P2	29,436 b	0,780
4 WAP	P0	30,232 a	0,366
	P3	30,364 a	0,366
	P1	30,566 ab	0,366
	P2	30,721 b	0,366

Note: Average values in the same column with differing letters denote a significant difference.

The results of the 5% BNJ test on the effect of planting distance on the root length (cm) of lettuce plants can be seen in Table 7.

Table 7. BNJ 5% test results for the influence of planting distance and aeration pressure on the root length (cm) of lettuce plants.

Factor	Average	BNJ 0,05	Factor	Average	BNJ 0,05
--------	---------	----------	--------	---------	----------

			1,455		
0,759					
D1	19,275	a	P0	19,868	A
D2	22,144	b	P1	20,178	Ab
			P2	21,294	B
			P3	21,681	B

Note: Average values within the same column that have different letters indicate a significant difference.

The BNJ test results at a 5% significance level show that increasing the planting distance generally leads to longer root lengths in lettuce plants. When the planting distance is wider, there is more space for growth, which means the plants do not have to compete for nutrients from the AB Mix. Conversely, the closer the planting distance, the less air is available for the plants to absorb (Rajagukguk, Aswandi and M. Si., 2023). Meanwhile, the influence of aeration pressure on the root length of lettuce plants showed that an increase in aeration pressure and oxygen content leads to an increase in root length. The increase in temperature causes a decrease in DO levels (Puspitahati and Andica, 2023). The results of the BNJ 5% test on the effect of planting distance on the fresh weight total (g) of lettuce plants can be seen in Table 8.

Table 8. BNJ 5% test table on the effect of planting distance on the fresh weight total (g) of lettuce plants

Factor	Average	BNJ 0,05	Factor	Average	BNJ 0,05
3,653			7,007		
D1	50,354	a	P0	59,777	a
D2	78,250	b	P1	62,332	ab
			P2	67,223	b
			P3	67,875	b

Note: Average values in the same column marked with different letters reflect a significant difference.

The BNJ test results at a 5% significance level show that treatment A1 differs significantly from treatment D2. Increasing the planting distance leads to a higher total fresh weight of the plants. Furthermore, the BNJ test indicates that treatment P0 is comparable to treatment P1 in terms of significance, while treatments P2 and P3 exhibit significant differences. According to (Kurniawati, 2020b), adequate dissolved oxygen in the root zone is crucial for optimal plant growth.

From the observations, the highest dissolved oxygen content was provided by the aeration pressure treatment of 16 kPa, which ranged from 7.17 to 7.22 mg/l, as shown in Figure 1




Figure 1. The Average Dissolved Oxygen Value for each treatment (mg/l)

In Figure 1, the trend of the dissolved oxygen (DO) graph against the aeration pressure applied is linear. It is evident that as the aeration pressure increases, the dissolved oxygen value tends to rise as well. According to (Pudjiwati and Asmina, 2020), The higher the dissolved oxygen content in hydroponic water, the better the respiration process of the plants. Aeration can help plant roots obtain dissolved oxygen in water and enhance the nutrient content that can be absorbed by the plants (Virha and Bayfurqon, 2020).

Aeration affects the dissolved oxygen (DO) content in hydroponics for plant growth. This process involves transferring oxygen from the air into the water, which can be done through surface or subsurface aeration, using air bubbles (Baiyin et al., 2021). The aeration efficiency is affected by the size of the bubbles, the surface area, the flow rate, and the diameter of the nozzle, all of which contribute to increased oxygen transfer. In addition, aerators and venturi are very effective in increasing DO levels, as they can improve the displacement of water and air masses, in hydroponic systems (Zhang et al., 2024). Correlation between Dissolved Oxygen (mg/l) and Aeration Pressure shown on Figure 2.





Figure 2. Correlation between Dissolved Oxygen (mg/l) and Aeration Pressure

Figure 2 showed that the relationship between dissolved oxygen (DO) levels and aeration pressure has a linear relationship shown by an R of 0.9993, which means that as the aeration pressure increases, the concentration of dissolved oxygen in the water also increases. This happens because the higher aeration pressure increases the amount of oxygen transferred from the air into the water, thereby increasing the availability of oxygen for the plant's roots .

The energy needed for root growth and ion absorption comes from respiration which requires oxygen. Without sufficient oxygen for respiration, the absorption of water and ions stops, and the plant roots die (Krisna et al., 2017). Plants that receive sufficient oxygen will have better growth, and the movement of nutrients will circulate, allowing the plants to absorb nutrients effectively (Puspitahati et al., 2022). According to Paramitha et al, (2014), Dissolved oxygen levels are affected by factors such as atmospheric pressure, temperature, salinity, water movement, as well as photosynthesis and respiration processes.

4. Bukti konfirmasi Accepted dan publish online (6 Januari 2025)




Notifications

[JPPT] Editor Decision

2025-01-05 01:06 PM

puspatahati puspatahati: We have reached a decision regarding your submission to Jurnal Penelitian Pertanian Terapan, "A Aeration Pressure, Plant Spacing, And Dissolved Oxygen For Growth Of Lettuce (Lactuca Sativa L.) Plants In Floating Hydroponics". Our decision is to: Accept Submission_____ Jurnal Penelitian Pertanian Terapan <http://jurnal.polinela.ac.id/index.php/JPPT>

Pengiriman LOA



Jurnal Penelitian Pertanian Terapan | ISSN: 1410-5020 E-ISSN: 2407-1781
Unit Penelitian dan Pengabdian Kepada Masyarakat, Politeknik Negeri Lampung
Jl. Soekarno-Hatta No. 10 Rajabasa, Bandar Lampung
Email : jppt@polinela.ac.id

No : 24.4.4/JPPT/J242/2024
Prihal : **Surat Pemberitahuan Naskah Diterima
untuk Publikasi**

Yth Author
Di tempat

Dengan hormat,
Redaksi Jurnal Penelitian Pertanian Terapan dengan ini menginformasikan bahwa naskah saudara :


ID Artikel : 3722
Email : puspitahati@fp.unsri.ac.id
Judul : Aeration Pressure, Plant Spacing, And Dissolved Oxygen For Growth Of Lettuce (Lactuca Sativa L.) Plants In Floating Hydroponics
Penulis : Puspitahati, Fidel Harmanda Prima, Nurul Izzah Aulia and Nikita Prettisha

Dinyatakan **diterima (accepted)** untuk dipublikasikan pada Jurnal Penelitian Pertanian Terapan Volume 24 (4) Edisi Desember 2024".

Demikian pemberitahuan ini disampaikan agar dapat dipergunakan sebagaimana mestinya. Atas partisipasi dan perhatiannya diucapkan terimakasih.



Bandar Lampung, 06 Januari 2025
Editor in Chief


Dr. Ir. Yana Sukaryana, M.P
NIP 196203241989031003

Workflow

Publication

Status: Published

This version has been published and can not be edited.

Title & Abstract

Bahasa Indonesia

English

Contributors

Metadata

References

Prefix

Examples: A, The

Title

An Aeration Pressure, Plant Spacing, And Dissolved Oxygen For Growth Of Lettuce (Lactuca Sativa L.) Plants In Floating H

Subtitle

Abstract

B

I

\times^*

\times_2

ϕ

20 cm independent lettuce plants were grown in terms of canopy width, root length, and total fresh weight. When the aeration pressure increases, the concentration of dissolved oxygen in the water also increases, the canopy width, root length, and total fresh weight of lettuce. The average dissolved oxygen ranged from 7.17 to 7.22 mg/l, so the aeration pressure correlates with increased dissolved oxygen levels, suggesting that higher aeration pressure correlates with increased dissolved oxygen levels. The Suggestion was Further research should be conducted on the appropriate size of the nutrient solution reservoir concerning the aeration pressure applied.