

Semarang, 24-25 September 2024

| No         | : MNP03/ICTCRED 2024/VIII/2024 | Semarang, August 8 <sup>th</sup> , 2024 |
|------------|--------------------------------|---|
| Attachment | : 1 exp                        |   |
| Subject    | : Acceptance letter            |   |

Dear Authors,

Congratulations! After careful reviews, we are delighted to inform you that the abstract with the details:

| : | Rozirwan   |
|---|--|
|   | Nopriani   |
|   | Fauziyah   |
|   | Wike Ayu Eka Putri                                     |
|   | Melki  |
|   | Iskhaq Iskandar  |
|   | Ela Novianti   |
|   | Apon Zaenal Mustopa                                    |
|   | Fatimah  |
|   | Redho Yoga Nugroho                                     |
| : | THE NATURAL CALCIUM AND PHOSPORUS CONTENT IN           |
|   | TWO FISHBONE OF COMMERCIAL SPECIES FROM ENGGANO WATERS |
|   | :  |

has been **accepted** for <del>Oral</del>/**Poster** presentation at "The 8<sup>th</sup> International Conference on Tropical and Coastal Region Eco Development 2024" on September 24-25th, 2024 in Aruss Hotel Semarang, Indonesia.

Please e-mail us to ictcred@live.undip.ac.id if you require any further information regarding with your participation. We look forward to welcoming you to our conference and wish you a fruitful and pleasant stay in Semarang.

Sincerely Yours,

A.H. Condro Haditomo, S.Pi., M.Si., Ph.D Chair Person ICTCRED 2024

Conference Secretariat: Faculty of Fisheries and Marine Science, Diponegoro University Jl. Prof. Jacub Rais, Tembalang, Kota Semarang, 50275, Indonesia Email: ictcred@live.undip.ac.id Website: http://ictcred.undip.ac.id



The 8<sup>th</sup> International Conference on Tropical and Coastal Region Eco Development

2024

Semarang, 24-25 September 2024

#### **Tentative Schedule**

| Time                      | Program  |  |  |  |
|---------------------------|--|--|--|--|
| Day 1 (24 September 2024) |  |  |  |  |
| 07.30-08.30               | Registration                                   |  |  |  |
| 07.30-08.30               | Opening speech                                 |  |  |  |
|                           | Plenary session                                |  |  |  |
| 08.45-09.30               | Keynote Speaker I (Prof. Slamet Budi Prayitno) |  |  |  |
| 09.30-10.15               | Keynote Speaker II (Dr. Noor Faizul H. Nordin) |  |  |  |
| 10.15-10.30               | Q & A  |  |  |  |
| 10.30-11.15               | Keynote Speaker III (Dr. Hansan Park)          |  |  |  |
| 11.15-12.00               | Keynote Speaker IV (Prof. Toshio Suga)         |  |  |  |
| 12.00-12.15               | Q & A  |  |  |  |
| 12.15-13.15               | Lunch and praying break                        |  |  |  |
| 13.30-14.30               | Scientific session I                           |  |  |  |
| 14.30-15.30               | Scientific session II                          |  |  |  |
| 15.30-16.00               | Coffee break while continuing presentation     |  |  |  |
| 15.30-16.30               | Scientific Session III                         |  |  |  |
| 16.30-17.30               | Scientific Session IV                          |  |  |  |
| 17:30-17:45               | Clossing                                       |  |  |  |
| Day 2 (25 September 2024) |  |  |  |  |
| City tour                 |  |  |  |  |

Chair Person ICTCRED 2024,

A.H. Condro Haditomo, S.Pi., M.Si, Ph.D



Redho Yoga Nugroho <redhoyn.29@gmail.com>

### Fwd: ICTCRED 2024: ABSTRACT AND EARLY BIRD PAYMENT NOTIFICATION

2 pesan

Rozirwan ROZIRWAN <rozirwan@unsri.ac.id> Kepada: Redho Yoga Nugroho <redhoyn.29@gmail.com> 22 September 2024 pukul 21.45

------ Forwarded message ------Dari: ICTCRED Undip <ictcred@live.undip.ac.id> Date: Sab, 10 Agu 2024 pukul 13.44 Subject: ICTCRED 2024: ABSTRACT AND EARLY BIRD PAYMENT NOTIFICATION To: <rozirwan@unsri.ac.id>

#### Dear Rozirwan,

Thank you for submitting your abstract to "The 8th International Conference on Tropical and Coastal Region Eco-Development 2024" to be held in Semarang, Indonesia.

After careful reviews, on behalf of the Program Committee, we are delighted to inform you that the abstract with the title: THE NATURAL CALCIUM AND PHOSPORUS CONTENT IN TWO FISHBONE OF COMMERCIAL SPECIES FROM ENGGANO WATERS has been

**accepted for Poster** presentation at "The 8th International Conference on Tropical and Coastal Region Eco-Development 2024" on September 24-25<sup>th</sup>, 2024 in Aruss Hotel Semarang, Indonesia.

Attached are your Letter of Acceptance and instruction for payment, please complete your registration fee by August 15<sup>th</sup>, 2024 04.00 PM GMT+7 to get the early bird payment.

The details of your payment are as follows : Virtual Account Number: 8788220400100412 Remittance Amount: 850,000 IDR Note:

- 1. Please complete your payment according to the amount mentioned above.
- 2. If you wish to book Aruss Hotel with the special rate (limited room), please contact Ms. Annisa Aulia Lukman (+6285205227373).

Failure to pay by the deadline will lead to cancellation of early bird payment, and your Virtual Account Number will expire. You will get the new virtual account number for regular payment.

Through this email, we also would like to express our deep regrets for notification delay due to an error on our system, and we are still trying to fix it. As long as our system still cannot be resolved, we will send conference notification via email.

https://mail.google.com/mail/u/0/?ik=419f7db4a5&view=pt&search=all&permthid=thread-f:1810907885794094190&simpl=msg-f:1810907885794094190&simpl=msg-a:r8855708803915434607

We look forward to welcoming you in Semarang, Indonesia this year. Should you require any further assistance, please do not hesitate to contact us.

For further information please check our website at https://ictcred.fpik.undip.ac.id/.

Thank you for your understanding and cooperation,

Kind Regards,

A.H. Condro Haditomo, S.Pi., M.Si., Ph.D Chairperson of ICTCRED 2024 +628158281374

**Dr. Rozirwan** Head of Marine Bioecology Laboratory Department of Marine Science Faculty of Mathematics and Natural Sciences Sriwijaya University Jalan Raya Palembang-Prabumulih KM 32, Indralaya Ogan Ilir, Sumatera Selatan, Indonesia, Pos Code: 30862 Email: rozirwan@unsri.ac.id, rozirwan@gmail.com

#### 2 lampiran

| <b>1</b> | Payment | Instruction_ | VA BNI.pdf |
|----------|---------|--------------|------------|
|          | 135K    |              |            |



Redho Yoga Nugroho <redhoyn.29@gmail.com> Kepada: rolazyoga96@gmail.com

[Kutipan teks disembunyikan]

#### 2 lampiran

Payment Instruction\_VA BNI.pdf 135K

| 7- | LoA_ | MNP03.pdf |
|----|------|-----------|
|    | 149K |           |

22 September 2024 pukul 21.47

#### THE NATURAL CALCIUM AND PHOSPORUS CONTENT IN TWO FISHBONE OF COMMERCIAL SPECIES FROM ENGGANO WATERS

Rozirwan<sup>1\*</sup>, Nopriani<sup>1</sup>, Fauziyah<sup>1</sup>, Wike Ayu Eka Putri<sup>1</sup>, Melki<sup>1</sup>, Iskhaq Iskandar<sup>2</sup>, Ela Novianti<sup>3</sup>, Apon Zaenal Mustopa<sup>3</sup>, Fatimah<sup>3</sup>, Redho Yoga Nugroho<sup>1</sup>

<sup>1</sup>Department of Marine Science, Faculty of Mathematics and Natural Science, Sriwijaya University, Indralaya, South Sumatra 30862, Indonesia

<sup>2</sup>Department of Physics, Faculty of Mathematics and Natural Science, Sriwijaya University, Indralaya, South Sumatra, 30862, Indonesia

<sup>3</sup>Research Center for Genetic Engineering, National Research and Innovation Agency, Bogor, West Java 16911, Indonesia

\*: rozirwan@unsri.ac.id

#### Abstract

Enggano waters, located at the outermost part of the Indian Ocean, host a diverse range of local commercial fish species with significant economic and nutritional potential. This study focuses on two such species, Epinephelus fuscoguttatus and Lethrinus lentjan, known for their high calcium and phosphorus content. The aim of this research was to determine the natural calcium and phosphorus content in the fishbones of these two commercial species from Enggano waters. The study was conducted from September to November 2023, with samples collected from local waters in Banjarsari Village, Enggano Island. The fishbones were cleaned, dried, and ground into powder. Calcium content was measured using Atomic Absorption Spectrophotometry (AAS), while phosphorus content was measured using UV-Vis Spectrophotometry. The results revealed that the calcium content in the fishbones of the E. fuscoguttatus was 8.84%, while in the L. lentjan was 7.97%. The phosphorus content was 1.4% in the E. fuscoguttatus and 1.43% in the L. lentjan. These findings indicated that the calcium content in the *E. fuscoguttatus* bones was higher compared to the *L. lentjan* bones, whereas the phosphorus content in the two species was relatively similar, with a slight edge for the L. lentjan. This study highlighted the valuable natural source of calcium and phosphorus found in the fishbones of E. fuscoguttatus and L. lentjan from Enggano waters. The high calcium content in E. fuscoguttatus bones made them a potential candidate for applications in the food and health industries. Meanwhile, the phosphorus content in both species as their nutritional value, supports the comprehensive utilization of these marine resources. In conclusion, the fishbones of E. fuscoguttatus and L. lentjan from Enggano waters are potential natural sources of calcium and phosphorus, offering benefits for various industrial applications and nutritional enhancements.

Keywords: calcium, Enggano waters, fishbones, phosphorus

THE 8THINTERNATIONAL CONFERENCE ON TROPICAL AND COASTAL REGION ECO DEVELOPMENT (ICTORED) 2024 & E-ASIA MHWMEETING 2024



THE NATURAL CALCIUM AND PHOSPORUS CONTENT IN TWO FISHBONE OF COMMERCIAL SPECIES FROM ENGGANO WATERS

# Rozirwanı, Noprianiı, Fauziyahı, Wike Ayu Eka Putriı, Melkiı, Iskhaq Iskandarz, Ela Noviantia, Apon Zaenal Mustopaa, Fatimaha, Redho Yoga Nugroho

<sup>1</sup> Department of Marine Science, Sriwijaya University, Indralaya, South Sumatra 30862, Indonesia <sup>2</sup> Department of Physics, Sriwijaya University, Indralaya, South Sumatra, 30862, Indonesia <sup>3</sup> Research Center for Genetic Engineering, National Research and Innovation Agency, West Java 16911, Indonesia



September 25<sup>th</sup>, 2024

As presenter "Redho Yoga Nugroho, S.Kel. M.Si" Department of Marine Science Sriwijaya University





No: 246/UN7.F10/HK/IX/2024

This certificate is presented to

Redho Yoga Nugroho, S.Kel., M.Si

Title: The natural Calcium and Phosporus Content in Two Fishbone of Commercial Species From Enggano Waters

has successfully participated in

# "The 8th International Conference on Tropical and Coastal Region Eco Development (ICTCRED) 2024"

as

# Presenter

Organized by Faculty of Fisheries and Marine Science, Diponegoro University

Semarang, September 24th, 2024

Chair, Organizing Committee

A. H. Condro Haditomo, S.Pi., M.Si., Ph.D. NIP. 198309082006041001

NIP. 196903231995121001

an of Faculty



8 Oktober 2024 pukul 13.38

### Fwd: ICTCRED 2024: Final Paper Submission Guidelines and Deadlines

2 pesan

Rozirwan ROZIRWAN <rozirwan@unsri.ac.id> Kepada: Redho Yoga Nugroho <redhoyn.29@gmail.com>

----- Forwarded message ------

Dari: ICTCRED Undip <ictcred@live.undip.ac.id>

Date: Sel, 8 Okt 2024 pukul 09.49

Subject: ICTCRED 2024: Final Paper Submission Guidelines and Deadlines

To: <khoirunnisalaina@gmail.com>, <fronthea.swastawati@live.undip.ac.id>, <sunardi2000@mail.ugm.ac.id>, Mey Dian <dianlapoliwa@gmail.com>, <t.ersti@lecturer.unri.ac.id>, <baigrienhs@unram.ac.id>, <nurdewisatsmoko@gmail.com>, <shobrinasilmi@untidar.ac.id>, <lailalatifah@students.undip.ac.id>, <santimariaa30@gmail.com>, <adityabasir88@gmail.com>, Subhan FPIK UHO <subhan ikl@uho.ac.id>, <yosepinecinta@gmail.com>, <sigitbayhuiryanthony@gmail.com>, <bobychristiansinaga@students.undip.ac.id>, <wahyuadi@ubb.ac.id>, <pascamsdp@live.undip.ac.id>, <kakartikadewi@gmail.com>, <dafina.shaquilla@gmail.com>, Al Fajar Alam <alfajaralamstrpi@mail.ugm.ac.id>, Miguel Enrigue Ma. A. AZCUNA <miguel.azcuna@g.batstate-u.edu.ph>, <dwiharyanti@lecturer.undip.ac.id>, <alifalahcahyadi23@gmail.com>, <febritrynto1@gmail.com>, <rfrancisco@msi.upd.edu.ph>, pattyana95ina <Pattyana95ina@yahoo.co.id>, <ibnumaulana1988@mail.ugm.ac. id>, <mimie.saputri@univ-lr.fr>, <vinanurnadiro@gmail.com>, <agilla.aditiara@gmail.com>, <dplongitm@gmail.com>, Nenik Kholilah <nenikkholil@gmail.com>, <adiguna.rn@gmail.com>, <siswanti@staff.uns.ac.id>, Aprilia Da Cruz Tita <apriliacruz20@gmail.com>, <diahayuningrum21@lecturer.undip.ac.id>, <rascaningrat@gmail.com>, <tvani455@gmail.com>, <sekarariane.sam@gmail.com>, <anikkusmiatun@gmail.com>, <finanovia34@gmail.com>, <jasiel.j.ks@gmail.com>, <putrantorizkita@gmail.com>, <rizalattagwa@student.undip.ac.id>, <primadintarahmasyam@mail.ugm.ac.id>, ira handayani <irahndyn@gmail.com>, <majariana.krisanti@apps.ipb.ac.id>, <dinda.chrisnawati@gmail.com>, <cbaran@msi.upd.edu.ph>, <rbaskara@staff.uns.ac.id>, <dewiks2903@gmail.com>, <muh.azril@untidar.ac.id>, <ekonofridiansyah@unib.ac.id>, <regina.melnawati@yahoo.com>, <a-husni@ugm.ac.id>, Nining Nursalim <nursalime1i015052@gmail.com>, <usman.ngc225@gmail.com>, <ade.pugara@live.undip.ac.id>, <pranatacandra981@gmail.com>, <soniaputrim@student.ub.ac.id>, <yanida@student.ub.ac.id>, <hanani.adiwira.c1@tohoku.ac.jp>, <candrawahyuningsih657@gmail.com>, <msiadmin@msi.upd.edu.ph>, <lptupaz@msi.upd.edu.ph>, <kismawardhani@dc.tohoku.ac.jp>, <fadlyirmawan911@gmail.com>, <nmdumato@up.edu.ph>, Reni Pertiwi <tyasasrining29@gmail.com>, <m.rais@itb.ac.id>, <adiningsih.sekar.s1@tohoku.ac.ip>, <yasunaka@tohoku.ac.jp>, <saqfikri@gmail.com>, <idiocton@msi.upd.edu.ph>, <crepollo@msi.upd.edu.ph>, <fpik@iunpad.ac.id>, <hanif.prasetyo@gmail.com>, <rozirwan@unsri.ac.id>, <suga@tohoku.ac.jp>, <sekarahma25@gmail.com>, <marellakerenbanget@gmail.com>, <hefni.effendi@apps.ipb.ac.id>, <medestiartono@lecturer.undip.ac.id>, <anjani.puspa07@gmail.com>, <aprizonputra@gmail.com>, <ima.wijayanti@live.undip.ac.id>, <daenkhimawa8@gmail.com>, <pdhiyaaflah05@gmail.com>, Retno Hartati <retnohartati.undip@yahoo.com>, <endrawijaya333@univpancasila.ac.id>, <ririzaiskandar@gmail.com>, rockie supit <supitrockie@gmail.com>, <ahma122@brin.go.id>, <wmuskita@gmail.com>, <wahyuendrak@ub.ac.id>, MUNTI SARIDA <munti.sarida@fp.unila.ac.id>, <suzanneundap@unsrat.ac.id>, <herunur.krisno@gmail.com>, <munasik@live.undip.ac.id>, <iwansetiabudi@undiksha.ac.id>, <asni@unismuh.ac.id>, <wilson tisera@yahoo.com>, <parichatw@g.swu.ac.th>, <bessiedonny.25@gmail.com>, <oktosupratman@gmail.com>, <kasealfred@yahoo.com>, <nandakhoirunisa@fmipa.unmul.ac.id>, <yeftaoliviashn@gmail.com>, <ristiawan1976@gmail.com>, <harmesa@brin.go.id>, <eko.susanto@live.undip.ac.id>, <yppertiwi@gmail.com>, <ikamaha45@gmail.com>, <novydwistifar@gmail.com>, <raflyadipratama77@gmail.com>, <puryajati.ardiansyah.desmont.s8@dc.tohoku.ac.jp>, Elvi Mandiangan <elvi.mandiangan@gmail.com>, <agielhasyim@yahoo.co.id>, <satriasatriaginanjar@gmail.com>, <sitisiwiwulandari@gmail.com>, <annisanabilaramdini@mail.ugm.ac.id>, <berlaniamahardikaputri@mail.ugm.ac.id>, <dpaskal11@gmail.com>, <suhaeli.fahmi@live.undip.ac.id>, <widodo.pranowo@gmail.com>, <rizalattagwa@students.undip.ac.id>, <rita.rostika@unpad.ac.id>

Dear ICTCRED 2024 Participants,

We are pleased to inform you about the final steps for paper submission following the **8th International Conference on Tropical and Coastal Region Eco-Development (ICTCRED) 2024.** Please carefully review the submission guidelines and journal options provided below.

Journal Options and APCs (Article Processing Charges):

- IOP (SCOPUS Proceeding): USD 95
- International Journal of Marine Science (IJMS) (Q3):
  - USD 175 (International)
  - Rp 2.500.000 (Local)
- Journal of Coastal and Tropical Studies (JKT) (SINTA 2):
  - USD 100 (International)
  - Rp 1.000.000 (Local)
- Buletin Oseanografi Marina (Buloma) (SINTA 2): Rp 1.000.000

#### **Important Announcement**:

1. For participants intending to submit to the **Food Research Journal**, we regret to inform you that submissions to this journal have been redirected to other journals due to insufficient participant quota.

- 2. The rule of **IOP** states that there should be **no more than 2 papers per author** (including as a **co-author**).
- 3. All Papers submitted to intended journals should be in English including for JKT and Buloma.

#### **Article Template:**

Before submitting your paper, please ensure it follows the required format. You can download the article template here:

Article Template: https://bit.ly/FullArticleTemplateICTCRED2024

#### **Submission Process:**

Participants are kindly requested to submit their papers via the links provided below:

- IOP: https://www.morressier.com/call-for-papers/664730b0843840bbdb0fcd6b
- IJMS, JKT, Buloma: https://bit.ly/FullPaperSubmissionICTCRED2024

The deadline for submission is **31 October 2024**. Once submitted, the ICTCRED review team will conduct an initial review of your paper. Following the review, a recommendation letter for journal submission will be provided. Final journal submission, including payment of the APC, will be the responsibility of each participant.

#### 6/1/25, 10:33 PM

#### Gmail - Fwd: ICTCRED 2024: Final Paper Submission Guidelines and Deadlines

We appreciate your attention to these guidelines and timely submission. Should you have any questions or require further clarification, we are at your disposal for assistance.

Thank you for your continued support, and we look forward to your contributions.

Warmest Regards,

Committee of ICTCRED 2022 ictcred.fpik.undip.ac.id ictcred@live.undip.ac.id

**Dr. Rozirwan** Head of Marine Bioecology Laboratory Department of Marine Science Faculty of Mathematics and Natural Sciences Sriwijaya University Jalan Raya Palembang-Prabumulih KM 32, Indralaya Ogan Ilir, Sumatera Selatan, Indonesia, Pos Code: 30862 Email: rozirwan@unsri.ac.id, rozirwan@gmail.com

Redho Yoga Nugroho <redhoyn.29@gmail.com> Kepada: rolazyoga96@gmail.com

[Kutipan teks disembunyikan]



Rozirwan ROZIRWAN <rozirwan@unsri.ac.id>

Your Paper for ICTCRED-2024

1 pesan

**Morressier Team** <discover@morressier.com> Kepada: rozirwan@unsri.ac.id 20 Januari 2025 pukul 22.17



Hi there,

Thank you for your Paper Submission to 'ICTCRED-2024'.

The Editor has requested that you resubmit your Paper 'The Natural Calcium and Phosporus Content in Two Fishbone of Commercial Species from Enggano Waters' with some changes. Go to My Submissions to see any comments from the Reviewers or Editor and submit an updated version of your Paper. The Deadline for resubmitting is January 26, 2025.

Please note that your revised Paper must be a camera-ready manuscript without any highlighted changes. A summary of the changes you have made to your Paper can be included in your Response to Reviewers, which you may upload as a separate document.

Go to My Submissions

# **IOP** Publishing

### **Response to Reviewers**

| Editor  | Response to Reviewer       |
|---|----------------------------|
| Based on the reviewer's evaluation, your manuscript is minor    | We have revise the last    |
| revision. Please revise your manuscript following reviewer's    | updated article, the       |
| comments by 8 January 2025. The similarity of your manuscript   | similarity was 12%         |
| is 29%. Please reduce it to less than 25%.                      |                            |
|   |                            |
| Reviewer 1  |                            |
| Did you repeat the test for each sample? If you did it, you can | Unfortunately, we did not  |
| present the average of the results in table or diagram          | perform a repeat on each   |
|   | sample, so we simply       |
|   | present the results as     |
|   | before, but we updated and |
|   | added references to some   |
|   | sentences and secondary    |
|   | supporting data            |

# The Natural Calcium and Phosporus Content in Two Fishbone of Commercial Species from Enggano Waters

Rozirwan<sup>1\*</sup>, Nopriani<sup>1</sup>, Fauziyah<sup>1</sup>, Wike Ayu Eka Putri<sup>1</sup>, Melki<sup>1</sup>, Iskhaq Iskandar<sup>2</sup>, Ela Novianti<sup>3</sup>, Apon Zaenal Mustopa<sup>3</sup>, Fatimah<sup>3</sup>, Redho Yoga Nugroho<sup>1</sup>

<sup>1</sup>Department of Marine Science, Faculty of Mathematics and Natural Sciences, Sriwijaya University, Jalan Palembang-Prabumulih KM. 32, Indralaya, South Sumatra 30662, Indonesia

<sup>2</sup>Department of Physics, Faculty of Mathematics and Natural Sciences, Sriwijaya University, Jalan Palembang-Prabumulih KM. 32, Indralaya, South Sumatra, 30862, Indonesia

<sup>3</sup>Research Center for Genetic Engineering, National Research and Innovation Agency, West Java 16911, Indonesia

\*Corresponding author: rozirwan@unsri.ac.id

**Abstract.** Calcium and phosphorus are important elements needed for the growth and development of the body both in humans and other living things. The bones of Tiger Grouper Fish (*Epinephelus fuscoguttatus*) and Lencam Fish (*Lethrinus lentjan*) have a fairly high content of calcium and phosphorus. The purpose of this study was to determine the calcium and phosphorus content in the bones of Tiger Grouper Fish and Lencam Fish. This research was carried out in September-November 2023 with direct sampling of collectors in Banjarsari Village, Enggano Island. The calcium content test was performed using an Atomic Absorption Spectrophotometer (AAS) and the phosphorus content test was performed using a UV-Vis spectrophotometer. The results of this study showed that the calcium content produced in both samples was 1.4% for Tiger Grouper Fish of 1.4% and 1.43% for Lencam Fish. Based on the results of the study, the calcium content in Tiger Grouper is greater than that of Lencam, while the phosphorus content in Tiger Grouper is smaller than Lencam Fish.

Keywords: Calcium, Enggano waters, Fishbone, Phosphorus

#### 1. Introduction

The production of catches produced on Enggano island can reach 27.89% of the total production of capture fisheries in North Bengkulu [1,2]. Snapper, Grouper and Pomfret are the three types of marine capture fisheries that produce the highest production from sea waters compared to coastal areas [3]. The high development of the industry in the field of processing fishery products will have the potential to increase organic waste in the area [4–6]. According to [7], The most common types of fishery waste are fish bones, fish heads, fish guts, and even shells, while this waste cannot be recycled due to the many limitations of its advanced processing [8]. Actually, it is estimated to be a material that can be used again, especially the use of fish bones which have natural calcium and phosphorus content.

Calcium and phosphorus are very important for bone and tooth growth and calcium intake must be considered early to meet the needs of bone growth in humans [9]. Several health problems can arise

due to calcium and phosphorus deficiency, namely the risk of osteoporosis. In Indonesia, osteoporosis is caused by the low average calcium intake of the Indonesian population which is only 254 mg per day. The average daily calcium requirement is 300-400 mg for infants, 500 mg for children, 600-700 mg for adolescents, and 500-800 mg for adults [10].

Fish bones are a type of fishery waste that contains a number of important benefits for humans. The main components of fish bones are calcium, phosphorus, and carbonate, so that fish bones themselves have the highest calcium content compared to other parts of the fish's body [11]. Calcium derived from fish bones is a waste product and is currently not used on a large scale for human needs. Therefore, further research is needed on the benefits of fish bone waste to be used as processed products that are beneficial for fisheries and marine problems [12]. One of the uses of fish bones is to process them into fish bone meal.

According to [13], calcium and phosphorus are macrominerals that are directly related to the development and maintenance of the skeleton system and participate in various physiological processes of the organism's body. The nutritional content of fish bones in 100 g of fish bone meal is calcium 735 mg, protein 9.2 g, fat 44 mg, phosphorus 345 mg, iron 78 mg, ash 24.5 g, and carbohydrates 0.1 mg [14]. Calcium has a role as an important micronutrient in addition to phosphorus for the growth and development of children and calcium and phosphorus are very important in the bone mineralization process [15]. Calcium plays an important role in physiological processes, such as cell metabolism, muscle contraction, and bone growth [16]. Phosphorus has metabolic functions to build bones, scales, adenosine triphosphate, cell membranes, and nucleic acids [4]. Chemical analysis revealed that fish bones are a valuable source of calcium phosphate as an economical source for synthetic hydrocytopathy. Bone is a biological composition consisting of an inorganic phase (calcium phosphate with a structure like carbonated hydroxyapati) [17].

The fisheries from the waters of Enggano that are consumed are Tiger Grouper Fish and Lencam Fish, is one type of fish waste that has the potential to be researched because it is included in the target fish species and is also the most dominant economic fish found [18]. Morphologically and physiologically the two samples are different, but the living habits or habitats of the two samples are almost the same, so this test was carried out to determine the difference in calcium and phosphorus content of Tiger Grouper and Lencam fish.

One of the indicators of the existence of Tiger Grouper fish is the presence of coral reefs in a water area to be able to shelter and as a place to forage [19]. Adult Tiger Grouper Fish can reach depths of more than 2 m. This type of fish often lives in clear waters with many coral reefs and depths of up to 60 m. The habitat of this fish is in the shallow waters of coral reefs, rocky seabeds, lagoon tops, coral channels, and the outermost part of steep coral reefs. This Grouper also likes waters with a sandy, rocky bottom and seaweed growing at the bottom [20]. In general, groupers are solitary, but when they want to spawn the fish in groups together. In general, this fish lives at a depth of 5-20 m in all types of corals in good condition. Most Tiger Grouper fish use burrows/cavities/holes in coral reefs as shelter and usually move around.

Lencam fish usually live in coastal areas, coral waters and belong to the group of underwater fish. The habitat of the Lencam fish is coral reefs, seagrass meadows, mangrove areas and shallow coastal waters with a sandy bottom at a depth of 30 m [21]. Lencam fish consists of 29 species spread around the world and about 81% of them are found in Indonesian waters. This fish continues to be threatened both directly and indirectly due to exploitation caused by various human activities [22].

#### 2. Materials and Methods

#### 2.1 Study area

This research was carried out in September – December 2023 with sampling obtained from fish collectors in Banjarsari Village, Enggano Island, Enggano District, North Bengkulu Regency, Bengkulu Province (Figure 1). Further sample analysis was carried out at the Marine Bioecology Laboratory and the Oceanography and Marine Instrumentation Laboratory at Sriwijaya University.



Figure 1. Sampling location of tiger grouper and lencam fish in Enggano Waters

#### 2.2 Sample Identification and Preparation

The samples used were taken from fish collectors in Banjarsari Village, Enggano Island. The fish samples amounted to 2 Tiger Grouper Fish and 2 Lencam Fish. The fish was still fresh, seen from its red color and not slimy. The sample preparation process was carried out at the Marine Bioecology Laboratory, starting with dissecting the fish using a knife. Then the dissected sample was washed and then boiled using a hot plate with a temperature of 60-80 °C for 60 min. After boiling, the fish was cleaned again using a brush to remove the meat that was still attached to the fish bones, then NaOH alkaline extraction was carried out which aimed to remove fat and blood levels in the sample. Samples that have been extracted for 60 min were cleaned again using aquades and then baked for 48 h at a temperature of 60°C. After the sample dried, it was crushed with mortal until it became fish bone meal powder which will later be used to make calcium and phosphorus test solutions. Fish bone meal powder was taken as much as 1 g (calcium) and 2 g (phosphorus) weighed using an analytical scale. Each sample taken was put into the erlenmeyer to make a test solution. The preparation of this test solution was carried out in the fume hood with the addition of NaOH 1.5 N, HNO<sub>3</sub>, HCI, H<sub>2</sub>O<sub>2</sub>, NaHCO<sub>3</sub>, (NH<sub>4</sub>)<sub>6</sub>Mo<sub>7</sub>O<sub>24</sub>.4H<sub>2</sub>O, H<sub>2</sub>O(SbO)C<sub>4</sub>H<sub>4</sub>O<sub>6</sub> 0.5K, H<sub>2</sub>SO<sub>4</sub> and PO<sub>4</sub>. After the fish bone sample becomes powder, a destructive process is carried out to determine the calcium and phosphorus content. The calcium content test itself was carried out using an Atomic Absorption Spectrophotometer (AAS) with a wavelength of 420 nm, while the phosphorus content testing was carried out using a tool called the Uv-Vis Spectrophotometer with a wavelength of 889 nm.

#### 2.3 Sample Destruction

Common destructions to determine the mineral components present in foodstuffs were known to be of two types, namely dry and wet [23]. Sample destruction in this study will use the wet destruction method referring to the study [24]. Their results showed that wet destruction gave better results than dry destruction because the temperature used could not exceed the boiling point of the solution and in general the carbon was destroyed faster than using dry ashing.

#### 2.4 Calcium level testing

The calcium content was determined by measuring the sample by wet destruction using an Atomic Absorption Spectrophotometer (AAS) with a wavelength of 420 nm, and the calcium content was checked using a modified method [25]. To analyze the calcium content, it can be done by weighing as much as 0.1 g of fine samples and transferring them to a 100 mL measuring flask. The destruction process was carried out by adding 15 mL of hydrochloric acid (HCl) to digest the sample. The solution was decomposed until it was clear and then cooled. Filtration volumes up to 100 mL can be measured with atomic absorption spectrophotometer (AAS) [26].

#### 2.5 Phosphorus level testing

The determination of phosphorus content in the solution of fish bone samples aimed to obtain the concentration of UV-Vis spectrophotometer (ppm) readings. The determination of the maximum wavelength P was carried out by analyzing the sample of phosphorus sample using a UV-Vis spectrophotometer at a wavelength of 889 nm. With this determination of wavelength, the maximum result of the P content in the fishbone sample [27]. The dissolved concentration obtained from measurements using a UV-Vis spectrophotometer was then recorded so that a graph of the wavelength and absorbance relationship in each fishbone sample used in testing the content of each sample was obtained.

2.6 Data analysis

2.6.1 Moisture content analysis

%Moisture content was calculated based on SNI-19-1705-2000 and AOAC (1995). %Moisture content =  $\frac{Wo-Wt}{Wt} \times 100\%$   $fk = \frac{100}{100-\% Moisture content}$ Information: Wt : Final weight of fish bone meal (g) Wo : Initial weight of the fish bone (g)

Fk : 100 / (100 - % of air)

#### 2.6.2 Calcium concentration data analysis

The concentration obtained from the curve was a concentration in mg/L. While the sample used in this study can be in the form of a solid so that it will be converted into mg/kg. To calculate the level of calcium in fish bone samples in ppm by sample weight, which was formulated as follows:

$$\label{eq:mineral} \begin{split} \text{Mineral concentration (mg/kg)} = & \frac{\text{Sample concentartion } \left(\frac{\text{mg}}{\text{L}}\right) \text{x sample volume (L)}}{\text{Sample weight (kg)}} \text{ x } D_{f} \\ \text{Information :} \end{split}$$

D<sub>f</sub> : Dilution factor

#### 2.6.3 Phosphorus concentration data analysis

The processing of phosphorus concentration data that was previously in the form of ppm units is converted into % which is formulated as follows:

 $P_2O_5$  levels available (ppm) = ppm curve x ml extract/1,000 mL x 1,000 g (g example) x  $D_f$  x 142/190 x  $M_f$ 

Information :

| ppm curve        | : | the sample level obtained from the relationship curve between the standard series and its reading after blank correction |
|------------------|---|--|
| $D_{\mathrm{f}}$ | : | dilution factor (if any)   |
| 142/190          | : | PO <sub>4</sub> to P <sub>2</sub> O form conversion factor   |
| $M_{\mathrm{f}}$ | : | moisture content correction factor   |

#### 3. Result and Discussion

3.1 Description of Tiger Grouper Fish

Based on the identification [28,29] the fish obtained were classified as *Epinephelus fuscoguttatus species*. Observations of the morphology were size, body shape of the fish, the position of the mouth, the type of fins and also the type of scales.



Figure 2. (a) Fish body parts, (b) Mouth type, (c) Scale type, (d) Fin type

Observation of the morphology of the Tiger Grouper fish consists of parts of the fish's body consisting of the head, body, fins and tail. The body shape of this fish in figure 3 (a) is compressed. In Figure (b) the mouth type is a terminal type whose upper and lower jaws can be seen to be the same length (parallel), and is often used to eat prey floating in front of it. Figure (c) body type covered by scales with a brownish body color and filled with reddish spots. While Figure (d) is a rounded fin type, meaning that the dorsal fin is elongated and totals 8-10 pieces and the caudal fin totals 15-17 pieces.

#### 3.2 Description of Lencam Fish

The determination of the morphological characteristics of Lencam fish includes body shape, mouth position, caudal fin shape, and scale type.



Figure 3. (a) Fish body parts, (b) Mouth type, (c) Scale type, (d) Fin type

Lencam fish are characterized by a flattened body that is covered with many scales, smooth cheeks, at the base of the ventral fin there are large scales (axillary scale) that are well developed. The mouth is located at the front end of the fish's head (terminal type) with a relatively thick lip, the front of the jaw has strong canines, on the side of the jaw there are conical teeth and large molars, there are no teeth at the top of the oral cavity. The first dorsal fin is fused with the second fin, consisting of 9 hard fingers and 9 or 10 soft fingers, the anal fin with 3 hard fingers, 8-10 soft fingers. (a) is a part of the

body of a fish consisting of 3 parts, namely the head, body and tail, in Figure (b) the mouth type of the Lencam fish is a terminal mouth type, meaning it has the same length of upper and lower jaws, and often eats prey floating in front of it. Figure (c) the shape of the body is rather tall and flattened, and the body is covered by scales. While Figure (d) is the type of Homocercal fin, the shape of the pinna caudalis is curved or not and supported by the spokes of the tail fin. In this type, the length between the two ends of the tail is the same length and also forms like the letter V.

#### 3.3 Yield Value of Tiger Grouper Fish and Lencam Fish

The yield value in this study is the result of a comparison between the initial weight of the fish bone and the amount produced from the final weight of the fish bone powder. According to [30], the level of yield value produced shows that the number of products produced is also high, indicating that the process used is more effective.

#### Table 1. Yield Value

| Types of fish      | Bone weight (g) | Fish bone powder (g) | Yield value (%) |
|--------------------|-----------------|----------------------|-----------------|
| Tiger Grouper Fish | 17.98           | 6.7                  | 2.6836          |
| Lencam Fish        | 13.04           | 4.05                 | 3.2198          |

The yield value obtained in this study can be an indicator of the quality of bone powder in both types of fish samples. It can be assumed that a high yield value indicates good flour quality. The magnitude of the yield obtained is influenced by the effectiveness in the extraction process. According to [31], a high yield will not necessarily produce the highest calcium levels, but it is also determined by other factors such as the low protein content in the material. The low yield is also suspected to be due to the influence of drying, where drying is the process of removing or disposing of liquid materials from a material.

#### 3.4 Calcium Content of Tiger Grouper Fish and Lencam Fish Bones

The analysis of calcium content in this study was obtained from measurements or readings using an atomic absorption spectrophotometer (AAS).

| Table 2. Calcium Content |            |        |                          |
|--------------------------|------------|--------|--------------------------|
| Types of Fish Bones      | Conc (ppm) | Ca (%) | Quality Standards<br>(%) |
| Tiger Grouper Fish       | 8.479      | 8.48   | 20.20                    |
| Lencam Fish              | 7.967      | 7.97   | 20-30                    |

# The results of the measurement of calcium content in the bone samples of Tiger Grouper and

Lencam Fish are known that the calcium content in Tiger Grouper is 8.84%, while in Lencam Fish it is 7.97%.



Figure 4. Measurement of calcium content

The difference in calcium content in the two samples can be influenced by the difference in the samples used, because basically different types of fish affect the value of the calcium levels produced. The results of measuring the calcium content of both Tiger Grouper and Lencam fish bones with different sizes are assumed that the sample size affects the mineral content in the fish's body, one of which is the bones. This is in line with the opinion [32,33], that minerals play an important role in the maintenance of every body function at the level of cells, tissues, organs, and overall body functions, and the presence of macro minerals in the body of fish is influenced by several factors, such as the size of the fish.

#### 3.5 Phosphorus Content of Tiger Grouper Fish and Lencam Fish Bones

The analysis of phosphorus content in this study was obtained by reading the results of the UV-Vis Spectrophotometer. The phosphorus content obtained in the bones of Tiger Grouper Fish and Lencam Fish with the same weight category in this study is seen in Table 3.

| Types of Fish Bones | Conc (ppm) | P (%) | Quality Standards<br>(%) |
|---------------------|------------|-------|--------------------------|
| Tiger Grouper Fish  | 1,401.49   | 1.4   | 20-30 %                  |
| Lencam Fish         | 1,432.58   | 1.43  |                          |

Table 3. Phosphorus Content of Fish Bones

Results of testing phosphorus content in the bones of Tiger Grouper and Lencam fish in the same weight category in mg/kg and percentage (%). Based on the quality standards of fish bone meal according to SNI 01-3158-1992, it is understood that the phosphorus content in the two samples does not meet the quality requirements. Furthermore, the results that have been obtained are visualized in the form of a graph to show a comparison of the results between the two fish, which can be seen in the following graph.



Figure 5. Measurement of phosphorus content

The phosphorus content in Lencam fish bone samples was higher than the phosphorus content in Tiger Grouper fish bone samples. The phosphorus content in Tiger Grouper can be seen at 1.4%, while in Lencam fish the content is slightly higher at 1.43%. The comparison of these two samples is around 0.03%. The type of sample, the material and the procedure used to measure the calcium and phosphorus content can also affect the results obtained. Different types of fish used in the study can affect the calcium and phosphorus content obtained, because the mineral content in each fish is different. Another factor that can cause this difference is also due to the sample size and the process in handling the sample.

The type of sample, materials and procedures in the study used to measure calcium and phosphorus levels can also affect the results obtained. This was revealed by [12] in the results of the study, the analysis of various types of frequencies in the boiling process affected the calcium content produced in the bones of Belida fish (*Chitala lopis*) ranging from 28.25% and the phosphorus content varied between 3.98% - 4.04%. The difference in NaOH concentration during the extraction process

affects the calcium and phosphorus content obtained, namely the concentration of 1.5 N provides the best content compared to the concentration of 1 N and 2 N. The selection of the test method in measuring the calcium and phosphorus content can affect the results obtained in each sample used.

#### 3.6 Calcium and phosphorus content in several types of fish bones

Based on Table 4, it is the result of a comparison of calcium and phosphorus content from various sources and also from various other types of marine fish. The different calcium and phosphorus content in the table below is caused by various factors, especially from the type of fish, the size of the fish and also influenced by the sample handling process before testing the content level in each sample. The method used in each source is also different, so it can be stated that the content is different because it is also due to the method used, because each method uses a different solution and also the processing process before being made into a solution is also different.

| No | Types of Fish Bones  | Calcium %   | Phosphorus % | References   |
|----|----------------------|-------------|--------------|--------------|
| 1  | Epinephelus sp       | 8.48        | 1.4          | This results |
| 2  | Lethrinus sp         | 7.97        | 1.43         | This results |
| 3  | Chitala lopis        | 28.25-31.31 | 3.98-4.06    | [12]         |
| 4  | P, albilabris        | 48.1        | 35.4         | [34]         |
| 5  | Thunnus sp.          | 39.24       | 13.66        | [35]         |
| 6  | Trichiurus savala    | 19.33       | 7.21         | [36]         |
| 7  | Sardinella longiceps | 26.39       | 16.57        | [36]         |
| 8  | Scomber scombrus     | 14.3        | 8.6          | [37]         |
| 9  | Salmo trutta         | 14.7        | 8.7          | [37]         |
| 10 | Salmo salar          | 13.5        | 8.1          | [37]         |
| 11 | Micromesistius       | 17.0        | 87           | [37]         |
| 11 | poutassou            | 17.0        | 0.7          | [57]         |
| 12 | Clupea harengus      | 16.1        | 9.4          | [37]         |

| Table 4 | . Calc | ium and | Phosp | horus | Content | in Several | l Types | of Fish Bones |
|---------|--------|---------|-------|-------|---------|------------|---------|---------------|
|         |        |         |       |       |         |            |         |               |

Based on the comparison table of calcium content in the table above, it can be stated that the calcium content in fish bones is lower than the calcium content in crab shell shells. This was also explained by [38,39], it was explained that the calcium carbonate content reached 53.70%-78.40% contained in the shell of the shell. As for the phosphorus content in fish bones from various sources presented in the table above, it is known that the phosphorus content is higher than the phosphorus content in seaweed. According to [40], his research explained that the phosphorus content in seaweed is >0.05, it can also be concluded that the phosphorus content is indeed higher than the content of other species. The difference in calcium content is also caused by the difference in the species analyzed, because every difference in a living creature can cause the results of the analysis related to the nutrient content in that type or species to be different. The difference between this study and the previous study lies in the category of fish sample size used and the chosen destruction method. The sample size was made in 3 categories to see the difference between them, while the selection of H<sub>2</sub>PO<sub>4</sub><sup>-</sup> and PO<sub>4</sub><sup>3-</sup>. In line with the results of a study [41], which tested soil samples with two different methods.

Calcium is an important mineral that can be relied on in the human body in a variety of several important processes [42]. Most of the calcium in the body is in the bones. Your circulatory system also has a certain amount of calcium which is essential for blood clotting as well as the function of blood vessels, nerves, muscles and heart. Calcium and phosphorus are one of the substances that have an important role in various bodily functions, such as bone mineralization, functioning as cofactors in many enzyme systems, maintaining muscle and nerve excitation, and in the case of iron, maintaining oxygen transport and blood capacity [43]. Many female athletes consume less calcium than the recommended nutritional adequacy rate. It is also a concern due to the need to reach peak bone mass during adolescence and the possible relationship of poor calcium intake with stress-induced fractures. Calcium and

phosphorus are available in various supplement forms, both can be taken together as individual supplements, and there are calcium phosphate supplements that are a combination of the two minerals. The role of phosphorus in the human body functions in the growth and development of energy in the body. The high content of calcium and phosphorus in fish bones makes fish bones can be processed into natural ingredients to replace bones or hydroxyapathic materials. According to [26], hydroxyapatite  $(Ca_{10}(PO_4)_6(OH)_2)$  is a bioceramic material that is biocompatible, biodegradable, and non-toxic. Hydroxyapatite can be synthesized from natural materials that contain relatively high levels of calcium which will later be used as calcium precursors because hydrosapathic compounds can be obtained by mixing calcium precursors with phosphate precursors. The functions of calcium in the body, especially in fish, include as the main component of the formation of bones, teeth, skin and scales, as well as other functions can maintain the strength of the body's skeleton, thicken the blood, become an intracellular regulator which can also help regulate the activity of skeletal muscles, heart and other tissues, muscle contraction and relaxation, help the absorption of vitamin B12, maintain osmotic balance [9][44]. This is supported by research [13], which states that calcium plays an important role in skeletal mineralization and bone formation, not only that calcium is also needed for bone growth and strength.

#### Conclusion

Grouper Fish Bones have a higher calcium content of 8.48% compared to Lencam Fish bones of 7.97%. Meanwhile, Lencam fish bones have a phosphorus content 1.43% higher than Tiger Grouper 1.4%. Based on the quality requirements of fish bone meal according to SNI 01-3158-1992, the calcium and phosphorus content of the bones of Tiger Grouper Fish and Lencam Fish do not meet the quality requirements.

#### References

- [1] Limi M A, Ningsih S R, Fyka S A, Syarni P, Dewi H S and Attamimi U 2022 Analysis of crab fisherman's household structure and livelihood strategy after the Kendari Bay revitalization *IOP Conference Series: Earth and Environmental Science* vol 1118 (IOP Publishing) p 12075
- [2] Rozirwan, Adefta A, Apri R, Fauziyah, Putri W A E, Melki, Iskandar I, Novianti E, Mustopa A Z, Fatimah and Nugroho R Y 2024 The Ecological Health of Coral Reefs In The Waters of Enggano Island, The Outermost Island In The Eastern Indian Ocean *Int. J. Conserv. Sci.* 15 1947–58
- [3] Rozirwan, Nugroho R Y, Fauziyah, Putri W A E, Melki, Ulqodry T Z, Isnaini, Absori A and Iskandar I 2023 Mollusks diversity in the protected coastline of Berbak-Sembilang National Park Indonesia *Int. J. Conserv. Sci.* **14** 1627–40
- [4] Fraser T W K, Witten P E, Albrektsen S, Breck O, Fontanillas R, Nankervis L, Thomsen T H, Koppe W, Sambraus F and Fjelldal P G 2019 Phosphorus nutrition in farmed Atlantic salmon (Salmo salar): Life stage and temperature effects on bone pathologies *Aquaculture* 511 734246
- [5] Khotimah N N, Rozirwan, Putri W A E, Fauziyah, Aryawati R, Isnaini and Nugroho R Y 2024 Bioaccumulation and Ecological Risk Assessment of Heavy Metal Contamination (Lead and Copper) Build Up in the Roots of Avicennia alba and Excoecaria agallocha J. Ecol. Eng. 25 101– 13
- [6] Rozirwan, Saputri A P, Nugroho R Y, Khotimah N N, Putri W A E, Fauziyah and Purwiyanto A I S 2023 An Assessment of Pb and Cu in Waters, Sediments, and Mud Crabs (Scylla serrata) from Mangrove Ecosystem Near Tanjung Api-Api Port Area, South Sumatra, Indonesia Sci. Technol. Indones. 8 675–83
- [7] Venugopal V 2022 Green processing of seafood waste biomass towards blue economy *Curr. Res. Environ. Sustain.* **4** 100164
- [8] Rozirwan, Nanda, Nugroho R Y, Diansyah G, Muhtadi, Fauziyah, Putri W A E and Agussalim A 2023 Phytochemical composition, total phenolic content and antioxidant activity of Anadara granosa (Linnaeus, 1758) collected from the east coast of South Sumatra, Indonesia *Baghdad Sci. J.* 20 1258–65
- [9] Raskh S 2020 The importance and role of calcium on the growth and development of children and its complications *Int. J. Res. Appl. Sci. Biotechnol.* **7** 162–7

- [10] Stoutjesdijk E 2020 *Brain-selective nutrients in pregnancy and lactation* (Groningen: University of Groningen)
- [11] Meiyasa F and Tarigan N 2020 Utilization of tuna fish bone waste (Thunnus sp.) as a source of calcium in making seaweed sticks *J. Teknol. Pertan. Andalas* **24** 66–75
- [12] Kondolele S L, Asikin A N, Kusumaningrum I, Diachanty S and Zuraida I 2022 Effect of Boiling Temperature on Physicochemical Characteristics of Mackerel Fish Bone Flour (Scomberomorus commerson) *Media Teknol. Has. Perikan.* 10 177–84
- [13] Ciosek Ż, Kot K, Kosik-Bogacka D, Łanocha-Arendarczyk N and Rotter I 2021 The effects of calcium, magnesium, phosphorus, fluoride, and lead on bone tissue *Biomolecules* **11** 506
- [14] Kaliky N, Lessy A, Rieuwpassa F, Kaya A O, Mailoa M N and Sormin R B D 2024 Physicochemical Characteristics of Red Snapper Fish Bone Flour (Lutjanus sp.) BIOSEL (Biology Sci. Educ. J. Penelit. Sci. dan Pendidik. 13 129–40
- Bhadada S K and Rao S D 2021 Role of phosphate in biomineralization Calcif. Tissue Int. 108 32–40
- [16] Battafarano G, Rossi M, Marampon F, Minisola S and Del Fattore A 2020 Bone control of muscle function Int. J. Mol. Sci. 21 1178
- [17] Kanis J A, Norton N, Harvey N C, Jacobson T, Johansson H, Lorentzon M, McCloskey E V, Willers C and Borgström F 2021 SCOPE 2021: a new scorecard for osteoporosis in Europe Arch. Osteoporos. 16 82
- [18] Scabra A R, Marzuki M and Yarni B M 2023 The Effect of Calcium Hydroxide (CAOH2) and Phosphorus (P) on the Growth of Vannamei Shrimp (Litopenaeus Vannamei) in Freshwater Media J. Ruaya J. Penelit. Dan Kaji. Ilmu Perikan. Dan Kelaut. 11 39–51
- [19] Paxton A B, Harter S L, Ross S W, Schobernd C M, Runde B J, Rudershausen P J, Johnson K H, Shertzer K W, Bacheler N M and Buckel J A 2021 Four decades of reef observations illuminate deep-water grouper hotspots *Fish Fish.* 22 749–61
- [20] Craig M T, de Mitcheson Y J S and Heemstra P C 2024 *Groupers of the world: a field and market guide* (Honolulu: CRC Press)
- [21] Prihatiningsih P, Pane A R P, Herlisman H and Hartati S T 2021 Reproduction and Growth of Lencam Fish (Lethrinus atkinsoni Seale, 1910) in Wakatobi Waters, Southeast Sulawesi Bawal Widya Ris. Perikan. Tangkap 13 111–22
- [22] Saini V P, Singh A K, Paul T, Thakur A and Singh M 2024 Advances in Environment Management for Sustainable Fisheries & livestock production (Patna: Bihar Animal Sciences University)
- [23] El Hosry L, Sok N, Richa R, Al Mashtoub L, Cayot P and Bou-Maroun E 2023 Sample preparation and analytical techniques in the determination of trace elements in food: A review *Foods* 12 895
- [24] Rozirwan, Ramadani S, Ayu W, Putri E, Nur N and Nugroho R Y 2023 Evaluation of Calcium and Phosphorus content in Scallop Shells (Placuna placenta) and Blood Cockle Shells (Anadara granosa) from Banyuasin Waters, South Sumatra Egypt. J. Aquat. Biol. Fish. Zool. Dep. Fac. Sci. 27 1053–68
- [25] Gómez-Nieto B, Motyzhov V, Gismera M J, Procopio J R and Sevilla M T 2020 Fast-sequential determination of cadmium and copper in milk powder and infant formula by direct solid sampling high-resolution continuum source graphite furnace atomic absorption spectrometry *Microchem. J.* 159 105335
- [26] Sareen A, Gupta R C, Bansal G and Singh V 2020 Comparison of key mineral elements in wild edible fruits of Ziziphus Mauritiana and Z. nummularia using atomic absorption spectrophotometer (AAS) and flame photometer *Int. J. Fruit Sci.* **20** S987–94
- [27] Pratiwi R A, Bayu A and Nandiyanto D 2022 How to Read and Interpret UV-VIS Spectrophotometric Results in Determining the Structure of Chemical Compounds Indones. J. Educ. Res. Technol. 2 1–20
- [28] Webb P W and Weihs D 1986 Functional locomotor morphology of early life history stages of fishes *Trans. Am. Fish. Soc.* **115** 115–27
- [29] Strauss R E and Bond C E 1990 Taxonomic methods: morphology Methods fish Biol. 109–40

- [30] Rosida R, Handayani L and Apriliani D 2018 Utilization of goat fish bone waste (Abalistes stellaris) as gelatin using variations in CH3COOH concentration Acta Aquat. Aquat. Sci. J. 5 93– 9
- [31] Henchion M, Hayes M, Mullen A M, Fenelon M and Tiwari B 2017 Future protein supply and demand: strategies and factors influencing a sustainable equilibrium *Foods* **6** 53
- [32] Ali A A H 2023 Overview of the vital roles of macro minerals in the human body *J. Trace Elem. Miner.* **4** 100076
- [33] Lall S P and Kaushik S J 2021 Nutrition and Metabolism of Minerals in Fish *Animals* 11 2711
- [34] Sumarto, Desmelati, Sari N I, Angraini R M and Arieska L 2021 Characteristic of Nano-Calcium Bone from a Different Species of Catfish (Pangasius hypophthalmus, Clarias batrachus, Hemibagrus nemurus and Paraplotosus albilabris) *IOP Conf. Ser. Earth Environ. Sci.* **695**
- [35] Trilaksani W, Salamah E and Nabil M 2006 Utilization of tuna fish bone waste (Thunnus sp.) as a source of calcium using the protein hydrolysis method *Bul. Tekonologi Has. Perikan.* **9** 34–45
- [36] Logesh A R, Pravinkumar M, Raffi S M and Kalaiselvam M 2012 Calcium and phosphorus determination in bones of low value fishes, Sardinella longiceps (Valenciennes) and Trichiurus savala (Cuvier), from Parangipettai, Southeast Coast of India Asian Pacific J. Trop. Dis. 2 S254– 6
- [37] Toppe J, Albrektsen S, Hope B and Aksnes A 2007 Chemical composition, mineral content and amino acid and lipid profiles in bones from various fish species *Comp. Biochem. Physiol. - B Biochem. Mol. Biol.* 146 395–401
- [38] Jun J-Y, Jung M J, Jeong I-H, Kim G, Sim J, Nam S and Kim B 2019 Effects of crab shell extract as a coagulant on the textural and sensorial properties of tofu (soybean curd) *Food Sci. Nutr.* 7
- [39] Yihun F, Egusa M, Kaminaka H, Izawa H, Morimoto M, Saimoto H and Ifuku S 2016 Protein/CaCO3/Chitin Nanofiber Complex Prepared from Crab Shells by Simple Mechanical Treatment and Its Effect on Plant Growth Int. J. Mol. Sci. 17 1600
- [40] Budi W S, Winarko W, Rokhmalia F, Darjati D and Poerwati S 2023 Analysis of Nitrogen, Phosphorus, Potassium Content in Humus in Soil at Temporary Shelters J. Penelit. Kesehat. Suara Forikes 14 62–6
- [41] Umaternate G R, Abidjulu J and Wuntu A D 2014 Olsen and Bray Method Test in Analyzing Available Phosphate Content in Paddy Field Soil in West Konarom Village, North Dumoga District J. MIPA 3 6–10
- [42] Soetan K O, Olaiya C O and Oyewole O E 2010 The importance of mineral elements for humans, domestic animals and plants: A review *African J. food Sci.* **4** 200–22
- [43] Malik D, Narayanasamy N, Pratyusha V A, Thakur J and Sinha N 2023 Inorganic Nutrients: Macrominerals *Textbook of Nutritional Biochemistry* (Springer) pp 391–446
- [44] Bouillon R and Suda T 2014 Vitamin D: calcium and bone homeostasis during evolution *Bonekey Rep.* **3**

#### Acknowledgements

This research was supported by the RIIM LPDP Grant and BRIN, (Grant Number: 12/II.7/HK/2023; Decree Number: 76/IV/KS/05/2023; 0236/UN9.3.1/PL/2023). We also thanks to the Sriwijaya University and Research Center for Genetic Engineering, BRIN.



Rozirwan ROZIRWAN <rozirwan@unsri.ac.id>

Your Paper for ICTCRED-2024

1 pesan

**Morressier Team** <discover@morressier.com> Kepada: rozirwan@unsri.ac.id 10 Februari 2025 pukul 16.06



Hi there,

Thank you for your Paper Submission to 'ICTCRED-2024'.

The Editor has requested that you resubmit your Paper 'The Natural Calcium and Phosporus Content in Two Fishbone of Commercial Species from Enggano Waters' with some changes. Go to My Submissions to see any comments from the Reviewers or Editor and submit an updated version of your Paper. The Deadline for resubmitting is February 13, 2025.

Please note that your revised Paper must be a camera-ready manuscript without any highlighted changes. A summary of the changes you have made to your Paper can be included in your Response to Reviewers, which you may upload as a separate document.

Go to My Submissions

# **IOP** Publishing

### **Response to Reviewers**

| Editor  | Response to Reviewer |
|---|----------------------|
| Please follow reviewer's comments by 26 January 2025          | Done                 |
|   |                      |
| Reviewer 1  |                      |
| In Figure 5, decimal still use coma not dot. Please revise it | We have revised the  |
|   | Figure 5.            |

# The Natural Calcium and Phosporus Content in Two Fishbone of Commercial Species from Enggano Waters

Rozirwan<sup>1\*</sup>, Nopriani<sup>1</sup>, Fauziyah<sup>1</sup>, Wike Ayu Eka Putri<sup>1</sup>, Melki<sup>1</sup>, Iskhaq Iskandar<sup>2</sup>, Ela Novianti<sup>3</sup>, Apon Zaenal Mustopa<sup>3</sup>, Fatimah<sup>3</sup>, Redho Yoga Nugroho<sup>1</sup>

<sup>1</sup>Department of Marine Science, Faculty of Mathematics and Natural Sciences, Sriwijaya University, Jalan Palembang-Prabumulih KM. 32, Indralaya, South Sumatra 30662, Indonesia

<sup>2</sup>Department of Physics, Faculty of Mathematics and Natural Sciences, Sriwijaya University, Jalan Palembang-Prabumulih KM. 32, Indralaya, South Sumatra, 30862, Indonesia

<sup>3</sup>Research Center for Genetic Engineering, National Research and Innovation Agency, West Java 16911, Indonesia

\*Corresponding author: rozirwan@unsri.ac.id

**Abstract.** Calcium and phosphorus are important elements needed for the growth and development of the body both in humans and other living things. The bones of Tiger Grouper Fish (*Epinephelus fuscoguttatus*) and Lencam Fish (*Lethrinus lentjan*) have a fairly high content of calcium and phosphorus. The purpose of this study was to determine the calcium and phosphorus content in the bones of Tiger Grouper Fish and Lencam Fish. This research was carried out in September-November 2023 with direct sampling of collectors in Banjarsari Village, Enggano Island. The calcium content test was performed using an Atomic Absorption Spectrophotometer (AAS) and the phosphorus content test was performed using a UV-Vis spectrophotometer. The results of this study showed that the calcium content produced in both samples was 1.4% for Tiger Grouper Fish of 1.4% and 1.43% for Lencam Fish. Based on the results of the study, the calcium content in Tiger Grouper is greater than that of Lencam, while the phosphorus content in Tiger Grouper is smaller than Lencam Fish.

Keywords: Calcium, Enggano waters, Fishbone, Phosphorus

#### **1. Introduction**

The production of catches produced on Enggano island can reach 27.89% of the total production of capture fisheries in North Bengkulu [1,2]. Snapper, Grouper and Pomfret are the three types of marine capture fisheries that produce the highest production from sea waters compared to coastal areas [3]. The high development of the industry in the field of processing fishery products will have the potential to increase organic waste in the area [4–6]. According to [7], The most common types of fishery waste are fish bones, fish heads, fish guts, and even shells, while this waste cannot be recycled due to the many limitations of its advanced processing [8]. Actually, it is estimated to be a material that can be used again, especially the use of fish bones which have natural calcium and phosphorus content.

Calcium and phosphorus are very important for bone and tooth growth and calcium intake must be considered early to meet the needs of bone growth in humans [9]. Several health problems can arise

due to calcium and phosphorus deficiency, namely the risk of osteoporosis. In Indonesia, osteoporosis is caused by the low average calcium intake of the Indonesian population which is only 254 mg per day. The average daily calcium requirement is 300-400 mg for infants, 500 mg for children, 600-700 mg for adolescents, and 500-800 mg for adults [10].

Fish bones are a type of fishery waste that contains a number of important benefits for humans. The main components of fish bones are calcium, phosphorus, and carbonate, so that fish bones themselves have the highest calcium content compared to other parts of the fish's body [11]. Calcium derived from fish bones is a waste product and is currently not used on a large scale for human needs. Therefore, further research is needed on the benefits of fish bone waste to be used as processed products that are beneficial for fisheries and marine problems [12]. One of the uses of fish bones is to process them into fish bone meal.

According to [13], calcium and phosphorus are macrominerals that are directly related to the development and maintenance of the skeleton system and participate in various physiological processes of the organism's body. The nutritional content of fish bones in 100 g of fish bone meal is calcium 735 mg, protein 9.2 g, fat 44 mg, phosphorus 345 mg, iron 78 mg, ash 24.5 g, and carbohydrates 0.1 mg [14]. Calcium has a role as an important micronutrient in addition to phosphorus for the growth and development of children and calcium and phosphorus are very important in the bone mineralization process [15]. Calcium plays an important role in physiological processes, such as cell metabolism, muscle contraction, and bone growth [16]. Phosphorus has metabolic functions to build bones, scales, adenosine triphosphate, cell membranes, and nucleic acids [4]. Chemical analysis revealed that fish bones are a valuable source of calcium phosphate as an economical source for synthetic hydrocytopathy. Bone is a biological composition consisting of an inorganic phase (calcium phosphate with a structure like carbonated hydroxyapati) [17].

The fisheries from the waters of Enggano that are consumed are Tiger Grouper Fish and Lencam Fish, is one type of fish waste that has the potential to be researched because it is included in the target fish species and is also the most dominant economic fish found [18]. Morphologically and physiologically the two samples are different, but the living habits or habitats of the two samples are almost the same, so this test was carried out to determine the difference in calcium and phosphorus content of Tiger Grouper and Lencam fish.

One of the indicators of the existence of Tiger Grouper fish is the presence of coral reefs in a water area to be able to shelter and as a place to forage [19]. Adult Tiger Grouper Fish can reach depths of more than 2 m. This type of fish often lives in clear waters with many coral reefs and depths of up to 60 m. The habitat of this fish is in the shallow waters of coral reefs, rocky seabeds, lagoon tops, coral channels, and the outermost part of steep coral reefs. This Grouper also likes waters with a sandy, rocky bottom and seaweed growing at the bottom [20]. In general, groupers are solitary, but when they want to spawn the fish in groups together. In general, this fish lives at a depth of 5-20 m in all types of corals in good condition. Most Tiger Grouper fish use burrows/cavities/holes in coral reefs as shelter and usually move around.

Lencam fish usually live in coastal areas, coral waters and belong to the group of underwater fish. The habitat of the Lencam fish is coral reefs, seagrass meadows, mangrove areas and shallow coastal waters with a sandy bottom at a depth of 30 m [21]. Lencam fish consists of 29 species spread around the world and about 81% of them are found in Indonesian waters. This fish continues to be threatened both directly and indirectly due to exploitation caused by various human activities [22].

#### 2. Materials and Methods

#### 2.1 Study area

This research was carried out in September – December 2023 with sampling obtained from fish collectors in Banjarsari Village, Enggano Island, Enggano District, North Bengkulu Regency, Bengkulu Province (Figure 1). Further sample analysis was carried out at the Marine Bioecology Laboratory and the Oceanography and Marine Instrumentation Laboratory at Sriwijaya University.



Figure 1. Sampling location of tiger grouper and lencam fish in Enggano Waters

#### 2.2 Sample Identification and Preparation

The samples used were taken from fish collectors in Banjarsari Village, Enggano Island. The fish samples amounted to 2 Tiger Grouper Fish and 2 Lencam Fish. The fish was still fresh, seen from its red color and not slimy. The sample preparation process was carried out at the Marine Bioecology Laboratory, starting with dissecting the fish using a knife. Then the dissected sample was washed and then boiled using a hot plate with a temperature of 60-80 °C for 60 min. After boiling, the fish was cleaned again using a brush to remove the meat that was still attached to the fish bones, then NaOH alkaline extraction was carried out which aimed to remove fat and blood levels in the sample. Samples that have been extracted for 60 min were cleaned again using aquades and then baked for 48 h at a temperature of 60°C. After the sample dried, it was crushed with mortal until it became fish bone meal powder which will later be used to make calcium and phosphorus test solutions. Fish bone meal powder was taken as much as 1 g (calcium) and 2 g (phosphorus) weighed using an analytical scale. Each sample taken was put into the erlenmeyer to make a test solution. The preparation of this test solution was carried out in the fume hood with the addition of NaOH 1.5 N, HNO<sub>3</sub>, HCI, H<sub>2</sub>O<sub>2</sub>, NaHCO<sub>3</sub>, (NH<sub>4</sub>)<sub>6</sub>Mo<sub>7</sub>O<sub>24</sub>.4H<sub>2</sub>O, H<sub>2</sub>O(SbO)C<sub>4</sub>H<sub>4</sub>O<sub>6</sub> 0.5K, H<sub>2</sub>SO<sub>4</sub> and PO<sub>4</sub>. After the fish bone sample becomes powder, a destructive process is carried out to determine the calcium and phosphorus content. The calcium content test itself was carried out using an Atomic Absorption Spectrophotometer (AAS) with a wavelength of 420 nm, while the phosphorus content testing was carried out using a tool called the Uv-Vis Spectrophotometer with a wavelength of 889 nm.

#### 2.3 Sample Destruction

Common destructions to determine the mineral components present in foodstuffs were known to be of two types, namely dry and wet [23]. Sample destruction in this study will use the wet destruction method referring to the study [24]. Their results showed that wet destruction gave better results than dry destruction because the temperature used could not exceed the boiling point of the solution and in general the carbon was destroyed faster than using dry ashing.

#### 2.4 Calcium level testing

The calcium content was determined by measuring the sample by wet destruction using an Atomic Absorption Spectrophotometer (AAS) with a wavelength of 420 nm, and the calcium content was checked using a modified method [25]. To analyze the calcium content, it can be done by weighing as much as 0.1 g of fine samples and transferring them to a 100 mL measuring flask. The destruction process was carried out by adding 15 mL of hydrochloric acid (HCl) to digest the sample. The solution was decomposed until it was clear and then cooled. Filtration volumes up to 100 mL can be measured with atomic absorption spectrophotometer (AAS) [26].

#### 2.5 Phosphorus level testing

The determination of phosphorus content in the solution of fish bone samples aimed to obtain the concentration of UV-Vis spectrophotometer (ppm) readings. The determination of the maximum wavelength P was carried out by analyzing the sample of phosphorus sample using a UV-Vis spectrophotometer at a wavelength of 889 nm. With this determination of wavelength, the maximum result of the P content in the fishbone sample [27]. The dissolved concentration obtained from measurements using a UV-Vis spectrophotometer was then recorded so that a graph of the wavelength and absorbance relationship in each fishbone sample used in testing the content of each sample was obtained.

2.6 Data analysis

2.6.1 Moisture content analysis

%Moisture content was calculated based on SNI-19-1705-2000 and AOAC (1995). %Moisture content =  $\frac{Wo-Wt}{Wt} \times 100\%$   $fk = \frac{100}{100-\% Moisture content}$ Information: Wt : Final weight of fish bone meal (g)

Wo : Initial weight of the fish bone (g)

Fk : 100 / (100 - % of air)

## 2.6.2 Calcium concentration data analysis

The concentration obtained from the curve was a concentration in mg/L. While the sample used in this study can be in the form of a solid so that it will be converted into mg/kg. To calculate the level of calcium in fish bone samples in ppm by sample weight, which was formulated as follows:

$$\label{eq:mineral} \begin{split} \text{Mineral concentration } (mg/kg) = & \frac{\text{Sample concentartion } \left(\frac{mg}{L}\right) x \text{ sample volume } (L)}{\text{Sample weight } (kg)} \ x \ D_f \end{split}$$
 Information :

 $D_{\rm f}$  : Dilution factor

#### 2.6.3 Phosphorus concentration data analysis

The processing of phosphorus concentration data that was previously in the form of ppm units is converted into % which is formulated as follows:

 $P_2O_5$  levels available (ppm) = ppm curve x ml extract/1,000 mL x 1,000 g (g example) x  $D_f$  x 142/190 x  $M_f$ 

Information :

| ppm curve        | : | the sample level obtained from the relationship curve between the standard series and its reading after blank correction |
|------------------|---|--|
| $D_{\mathrm{f}}$ | : | dilution factor (if any)   |
| 142/190          | : | PO <sub>4</sub> to P <sub>2</sub> O form conversion factor   |
| $M_{\mathrm{f}}$ | : | moisture content correction factor   |

#### 3. Result and Discussion

3.1 Description of Tiger Grouper Fish

Based on the identification [28,29] the fish obtained were classified as *Epinephelus fuscoguttatus species*. Observations of the morphology were size, body shape of the fish, the position of the mouth, the type of fins and also the type of scales.



Figure 2. (a) Fish body parts, (b) Mouth type, (c) Scale type, (d) Fin type

Observation of the morphology of the Tiger Grouper fish consists of parts of the fish's body consisting of the head, body, fins and tail. The body shape of this fish in figure 3 (a) is compressed. In Figure (b) the mouth type is a terminal type whose upper and lower jaws can be seen to be the same length (parallel), and is often used to eat prey floating in front of it. Figure (c) body type covered by scales with a brownish body color and filled with reddish spots. While Figure (d) is a rounded fin type, meaning that the dorsal fin is elongated and totals 8-10 pieces and the caudal fin totals 15-17 pieces.

#### 3.2 Description of Lencam Fish

The determination of the morphological characteristics of Lencam fish includes body shape, mouth position, caudal fin shape, and scale type.



Figure 3. (a) Fish body parts, (b) Mouth type, (c) Scale type, (d) Fin type

Lencam fish are characterized by a flattened body that is covered with many scales, smooth cheeks, at the base of the ventral fin there are large scales (axillary scale) that are well developed. The mouth is located at the front end of the fish's head (terminal type) with a relatively thick lip, the front of the jaw has strong canines, on the side of the jaw there are conical teeth and large molars, there are no teeth at the top of the oral cavity. The first dorsal fin is fused with the second fin, consisting of 9 hard fingers and 9 or 10 soft fingers, the anal fin with 3 hard fingers, 8-10 soft fingers. (a) is a part of the

body of a fish consisting of 3 parts, namely the head, body and tail, in Figure (b) the mouth type of the Lencam fish is a terminal mouth type, meaning it has the same length of upper and lower jaws, and often eats prey floating in front of it. Figure (c) the shape of the body is rather tall and flattened, and the body is covered by scales. While Figure (d) is the type of *Homocercal fin*, the shape of the pinna caudalis is curved or not and supported by the spokes of the tail fin. In this type, the length between the two ends of the tail is the same length and also forms like the letter V.

#### 3.3 Yield Value of Tiger Grouper Fish and Lencam Fish

The yield value in this study is the result of a comparison between the initial weight of the fish bone and the amount produced from the final weight of the fish bone powder. According to [30], the level of yield value produced shows that the number of products produced is also high, indicating that the process used is more effective.

#### Table 1. Yield Value

| Types of fish      | Bone weight (g) | Fish bone powder (g) | Yield value (%) |
|--------------------|-----------------|----------------------|-----------------|
| Tiger Grouper Fish | 17.98           | 6.7                  | 2.6836          |
| Lencam Fish        | 13.04           | 4.05                 | 3.2198          |

The yield value obtained in this study can be an indicator of the quality of bone powder in both types of fish samples. It can be assumed that a high yield value indicates good flour quality. The magnitude of the yield obtained is influenced by the effectiveness in the extraction process. According to [31], a high yield will not necessarily produce the highest calcium levels, but it is also determined by other factors such as the low protein content in the material. The low yield is also suspected to be due to the influence of drying, where drying is the process of removing or disposing of liquid materials from a material.

#### 3.4 Calcium Content of Tiger Grouper Fish and Lencam Fish Bones

The analysis of calcium content in this study was obtained from measurements or readings using an atomic absorption spectrophotometer (AAS).

| Table 2. Calcium Content |            |        |                          |
|--------------------------|------------|--------|--------------------------|
| Types of Fish Bones      | Conc (ppm) | Ca (%) | Quality Standards<br>(%) |
| Tiger Grouper Fish       | 8.479      | 8.48   | 20.20                    |
| Lencam Fish              | 7.967      | 7.97   | 20-30                    |

The results of the measurement of calcium content in the bone samples of Tiger Grouper and Lencam Fish are known that the calcium content in Tiger Grouper is 8.84%, while in Lencam Fish it is 7.97%.



The difference in calcium content in the two samples can be influenced by the difference in the samples used, because basically different types of fish affect the value of the calcium levels produced. The results of measuring the calcium content of both Tiger Grouper and Lencam fish bones with different sizes are assumed that the sample size affects the mineral content in the fish's body, one of which is the bones. This is in line with the opinion [32,33], that minerals play an important role in the maintenance of every body function at the level of cells, tissues, organs, and overall body functions, and the presence of macro minerals in the body of fish is influenced by several factors, such as the size of the fish.

#### 3.5 Phosphorus Content of Tiger Grouper Fish and Lencam Fish Bones

The analysis of phosphorus content in this study was obtained by reading the results of the UV-Vis Spectrophotometer. The phosphorus content obtained in the bones of Tiger Grouper Fish and Lencam Fish with the same weight category in this study is seen in Table 3.

| Types of Fish Bones | Conc (ppm) | P (%) | Quality Standards<br>(%) |
|---------------------|------------|-------|--------------------------|
| Tiger Grouper Fish  | 1,401.49   | 1.40  | 20-30 %                  |
| Lencam Fish         | 1,432.58   | 1.43  | 20 30 70                 |

Table 3. Phosphorus Content of Fish Bone

Results of testing phosphorus content in the bones of Tiger Grouper and Lencam fish in the same weight category in mg/kg and percentage (%). Based on the quality standards of fish bone meal according to SNI 01-3158-1992, it is understood that the phosphorus content in the two samples does not meet the quality requirements. Furthermore, the results that have been obtained are visualized in the form of a graph to show a comparison of the results between the two fish, which can be seen in the following graph.



Figure 5. Measurement of phosphorus content

The phosphorus content in Lencam fish bone samples was higher than the phosphorus content in Tiger Grouper fish bone samples. The phosphorus content in Tiger Grouper can be seen at 1.4%, while in Lencam fish the content is slightly higher at 1.43%. The comparison of these two samples is around 0.03%. The type of sample, the material and the procedure used to measure the calcium and phosphorus content can also affect the results obtained. Different types of fish used in the study can affect the calcium and phosphorus content obtained, because the mineral content in each fish is different. Another factor that can cause this difference is also due to the sample size and the process in handling the sample.

The type of sample, materials and procedures in the study used to measure calcium and phosphorus levels can also affect the results obtained. This was revealed by [12] in the results of the study, the analysis of various types of frequencies in the boiling process affected the calcium content produced in the bones of Belida fish (*Chitala lopis*) ranging from 28.25% and the phosphorus content varied between 3.98% - 4.04%. The difference in NaOH concentration during the extraction process

affects the calcium and phosphorus content obtained, namely the concentration of 1.5 N provides the best content compared to the concentration of 1 N and 2 N. The selection of the test method in measuring the calcium and phosphorus content can affect the results obtained in each sample used.

#### 3.6 Calcium and phosphorus content in several types of fish bones

Based on Table 4, it is the result of a comparison of calcium and phosphorus content from various sources and also from various other types of marine fish. The different calcium and phosphorus content in the table below is caused by various factors, especially from the type of fish, the size of the fish and also influenced by the sample handling process before testing the content level in each sample. The method used in each source is also different, so it can be stated that the content is different because it is also due to the method used, because each method uses a different solution and also the processing process before being made into a solution is also different.

| No | Types of Fish Bones         | Calcium %   | Phosphorus % | References   |
|----|-----------------------------|-------------|--------------|--------------|
| 1  | Epinephelus sp              | 8.48        | 1.40         | This results |
| 2  | Lethrinus sp                | 7.97        | 1.43         | This results |
| 3  | Chitala lopis               | 28.25-31.31 | 3.98-4.06    | [12]         |
| 4  | P, albilabris               | 48.10       | 35.40        | [34]         |
| 5  | Thunnus sp.                 | 39.24       | 13.66        | [35]         |
| 6  | Trichiurus savala           | 19.33       | 7.21         | [36]         |
| 7  | Sardinella longiceps        | 26.39       | 16.57        | [36]         |
| 8  | Scomber scombrus            | 14.3        | 8.6          | [37]         |
| 9  | Salmo trutta                | 14.7        | 8.7          | [37]         |
| 10 | Salmo salar                 | 13.5        | 8.1          | [37]         |
| 11 | Micromesistius<br>poutassou | 17.0        | 8.7          | [37]         |
| 12 | Clupea harengus             | 16.1        | 9.4          | [37]         |

Table 4. Calcium and Phosphorus Content in Several Types of Fish Bones

Based on the comparison table of calcium content in the table above, it can be stated that the calcium content in fish bones is lower than the calcium content in crab shell shells. This was also explained by [38,39], it was explained that the calcium carbonate content reached 53.70%-78.40% contained in the shell of the shell. As for the phosphorus content in fish bones from various sources presented in the table above, it is known that the phosphorus content is higher than the phosphorus content in seaweed. According to [40], his research explained that the phosphorus content in seaweed is >0.05, it can also be concluded that the phosphorus content is indeed higher than the content of other species. The difference in calcium content is also caused by the difference in the species analyzed, because every difference in a living creature can cause the results of the analysis related to the nutrient content in that type or species to be different. The difference between this study and the previous study lies in the category of fish sample size used and the chosen destruction method. The sample size was made in 3 categories to see the difference between them, while the selection of the method for phosphorus testing with P Bray I where the phosphorus read was only in the form of H<sub>2</sub>PO<sub>4</sub><sup>-</sup> and PO<sub>4</sub><sup>3-</sup>. In line with the results of a study [41], which tested soil samples with two different methods.

Calcium is an important mineral that can be relied on in the human body in a variety of several important processes [42]. Most of the calcium in the body is in the bones. Your circulatory system also has a certain amount of calcium which is essential for blood clotting as well as the function of blood vessels, nerves, muscles and heart. Calcium and phosphorus are one of the substances that have an important role in various bodily functions, such as bone mineralization, functioning as cofactors in many enzyme systems, maintaining muscle and nerve excitation, and in the case of iron, maintaining oxygen transport and blood capacity [43]. Many female athletes consume less calcium than the recommended nutritional adequacy rate. It is also a concern due to the need to reach peak bone mass during adolescence and the possible relationship of poor calcium intake with stress-induced fractures. Calcium and

phosphorus are available in various supplement forms, both can be taken together as individual supplements, and there are calcium phosphate supplements that are a combination of the two minerals. The role of phosphorus in the human body functions in the growth and development of energy in the body. The high content of calcium and phosphorus in fish bones makes fish bones can be processed into natural ingredients to replace bones or hydroxyapathic materials. According to [26], hydroxyapatite  $(Ca_{10}(PO_4)_6(OH)_2)$  is a bioceramic material that is biocompatible, biodegradable, and non-toxic. Hydroxyapatite can be synthesized from natural materials that contain relatively high levels of calcium which will later be used as calcium precursors because hydrosapathic compounds can be obtained by mixing calcium precursors with phosphate precursors. The functions of calcium in the body, especially in fish, include as the main component of the formation of bones, teeth, skin and scales, as well as other functions can maintain the strength of the body's skeleton, thicken the blood, become an intracellular regulator which can also help regulate the activity of skeletal muscles, heart and other tissues, muscle contraction and relaxation, help the absorption of vitamin B12, maintain osmotic balance [9][44]. This is supported by research [13], which states that calcium plays an important role in skeletal mineralization and bone formation, not only that calcium is also needed for bone growth and strength.

#### Conclusion

Grouper Fish Bones have a higher calcium content of 8.48% compared to Lencam Fish bones of 7.97%. Meanwhile, Lencam fish bones have a phosphorus content 1.43% higher than Tiger Grouper 1.4%. Based on the quality requirements of fish bone meal according to SNI 01-3158-1992, the calcium and phosphorus content of the bones of Tiger Grouper Fish and Lencam Fish do not meet the quality requirements.

#### References

- [1] Limi M A, Ningsih S R, Fyka S A, Syarni P, Dewi H S and Attamimi U 2022 Analysis of crab fisherman's household structure and livelihood strategy after the Kendari Bay revitalization *IOP Conference Series: Earth and Environmental Science* vol 1118 (IOP Publishing) p 12075
- [2] Rozirwan, Adefta A, Apri R, Fauziyah, Putri W A E, Melki, Iskandar I, Novianti E, Mustopa A Z, Fatimah and Nugroho R Y 2024 The Ecological Health of Coral Reefs In The Waters of Enggano Island, The Outermost Island In The Eastern Indian Ocean Int. J. Conserv. Sci. 15 1947–58
- [3] Rozirwan, Nugroho R Y, Fauziyah, Putri W A E, Melki, Ulqodry T Z, Isnaini, Absori A and Iskandar I 2023 Mollusks diversity in the protected coastline of Berbak-Sembilang National Park Indonesia Int. J. Conserv. Sci. 14 1627–40
- [4] Fraser T W K, Witten P E, Albrektsen S, Breck O, Fontanillas R, Nankervis L, Thomsen T H, Koppe W, Sambraus F and Fjelldal P G 2019 Phosphorus nutrition in farmed Atlantic salmon (Salmo salar): Life stage and temperature effects on bone pathologies *Aquaculture* 511 734246
- [5] Khotimah N N, Rozirwan, Putri W A E, Fauziyah, Aryawati R, Isnaini and Nugroho R Y 2024 Bioaccumulation and Ecological Risk Assessment of Heavy Metal Contamination (Lead and Copper) Build Up in the Roots of Avicennia alba and Excoecaria agallocha J. Ecol. Eng. 25 101– 13
- [6] Rozirwan, Saputri A P, Nugroho R Y, Khotimah N N, Putri W A E, Fauziyah and Purwiyanto A I S 2023 An Assessment of Pb and Cu in Waters, Sediments, and Mud Crabs (Scylla serrata) from Mangrove Ecosystem Near Tanjung Api-Api Port Area, South Sumatra, Indonesia Sci. Technol. Indones. 8 675–83
- [7] Venugopal V 2022 Green processing of seafood waste biomass towards blue economy *Curr. Res. Environ. Sustain.* **4** 100164
- [8] Rozirwan, Nanda, Nugroho R Y, Diansyah G, Muhtadi, Fauziyah, Putri W A E and Agussalim A 2023 Phytochemical composition, total phenolic content and antioxidant activity of Anadara granosa (Linnaeus, 1758) collected from the east coast of South Sumatra, Indonesia *Baghdad Sci. J.* 20 1258–65
- [9] Raskh S 2020 The importance and role of calcium on the growth and development of children and its complications *Int. J. Res. Appl. Sci. Biotechnol.* **7** 162–7

- [10] Stoutjesdijk E 2020 Brain-selective nutrients in pregnancy and lactation (Groningen: University of Groningen)
- [11] Meiyasa F and Tarigan N 2020 Utilization of tuna fish bone waste (Thunnus sp.) as a source of calcium in making seaweed sticks *J. Teknol. Pertan. Andalas* **24** 66–75
- [12] Kondolele S L, Asikin A N, Kusumaningrum I, Diachanty S and Zuraida I 2022 Effect of Boiling Temperature on Physicochemical Characteristics of Mackerel Fish Bone Flour (Scomberomorus commerson) *Media Teknol. Has. Perikan.* 10 177–84
- [13] Ciosek Ż, Kot K, Kosik-Bogacka D, Łanocha-Arendarczyk N and Rotter I 2021 The effects of calcium, magnesium, phosphorus, fluoride, and lead on bone tissue *Biomolecules* **11** 506
- [14] Kaliky N, Lessy A, Rieuwpassa F, Kaya A O, Mailoa M N and Sormin R B D 2024 Physicochemical Characteristics of Red Snapper Fish Bone Flour (Lutjanus sp.) BIOSEL (Biology Sci. Educ. J. Penelit. Sci. dan Pendidik. 13 129–40
- [15] Bhadada S K and Rao S D 2021 Role of phosphate in biomineralization Calcif. Tissue Int. 108 32–40
- [16] Battafarano G, Rossi M, Marampon F, Minisola S and Del Fattore A 2020 Bone control of muscle function *Int. J. Mol. Sci.* **21** 1178
- [17] Kanis J A, Norton N, Harvey N C, Jacobson T, Johansson H, Lorentzon M, McCloskey E V, Willers C and Borgström F 2021 SCOPE 2021: a new scorecard for osteoporosis in Europe Arch. Osteoporos. 16 82
- [18] Scabra A R, Marzuki M and Yarni B M 2023 The Effect of Calcium Hydroxide (CAOH2) and Phosphorus (P) on the Growth of Vannamei Shrimp (Litopenaeus Vannamei) in Freshwater Media J. Ruaya J. Penelit. Dan Kaji. Ilmu Perikan. Dan Kelaut. 11 39–51
- [19] Paxton A B, Harter S L, Ross S W, Schobernd C M, Runde B J, Rudershausen P J, Johnson K H, Shertzer K W, Bacheler N M and Buckel J A 2021 Four decades of reef observations illuminate deep-water grouper hotspots *Fish Fish.* 22 749–61
- [20] Craig M T, de Mitcheson Y J S and Heemstra P C 2024 *Groupers of the world: a field and market guide* (Honolulu: CRC Press)
- [21] Prihatiningsih P, Pane A R P, Herlisman H and Hartati S T 2021 Reproduction and Growth of Lencam Fish (Lethrinus atkinsoni Seale, 1910) in Wakatobi Waters, Southeast Sulawesi Bawal Widya Ris. Perikan. Tangkap 13 111–22
- [22] Saini V P, Singh A K, Paul T, Thakur A and Singh M 2024 Advances in Environment Management for Sustainable Fisheries & livestock production (Patna: Bihar Animal Sciences University)
- [23] El Hosry L, Sok N, Richa R, Al Mashtoub L, Cayot P and Bou-Maroun E 2023 Sample preparation and analytical techniques in the determination of trace elements in food: A review *Foods* 12 895
- [24] Rozirwan, Ramadani S, Ayu W, Putri E, Nur N and Nugroho R Y 2023 Evaluation of Calcium and Phosphorus content in Scallop Shells (Placuna placenta) and Blood Cockle Shells (Anadara granosa) from Banyuasin Waters, South Sumatra Egypt. J. Aquat. Biol. Fish. Zool. Dep. Fac. Sci. 27 1053–68
- [25] Gómez-Nieto B, Motyzhov V, Gismera M J, Procopio J R and Sevilla M T 2020 Fast-sequential determination of cadmium and copper in milk powder and infant formula by direct solid sampling high-resolution continuum source graphite furnace atomic absorption spectrometry *Microchem. J.* 159 105335
- [26] Sareen A, Gupta R C, Bansal G and Singh V 2020 Comparison of key mineral elements in wild edible fruits of Ziziphus Mauritiana and Z. nummularia using atomic absorption spectrophotometer (AAS) and flame photometer *Int. J. Fruit Sci.* **20** S987–94
- [27] Pratiwi R A, Bayu A and Nandiyanto D 2022 How to Read and Interpret UV-VIS Spectrophotometric Results in Determining the Structure of Chemical Compounds Indones. J. Educ. Res. Technol. 2 1–20
- [28] Webb P W and Weihs D 1986 Functional locomotor morphology of early life history stages of fishes *Trans. Am. Fish. Soc.* **115** 115–27
- [29] Strauss R E and Bond C E 1990 Taxonomic methods: morphology Methods fish Biol. 109-40

- [30] Rosida R, Handayani L and Apriliani D 2018 Utilization of goat fish bone waste (Abalistes stellaris) as gelatin using variations in CH3COOH concentration Acta Aquat. Aquat. Sci. J. 5 93– 9
- [31] Henchion M, Hayes M, Mullen A M, Fenelon M and Tiwari B 2017 Future protein supply and demand: strategies and factors influencing a sustainable equilibrium *Foods* **6** 53
- [32] Ali A A H 2023 Overview of the vital roles of macro minerals in the human body *J. Trace Elem. Miner.* **4** 100076
- [33] Lall S P and Kaushik S J 2021 Nutrition and Metabolism of Minerals in Fish Animals 11 2711
- [34] Sumarto, Desmelati, Sari N I, Angraini R M and Arieska L 2021 Characteristic of Nano-Calcium Bone from a Different Species of Catfish (Pangasius hypophthalmus, Clarias batrachus, Hemibagrus nemurus and Paraplotosus albilabris) *IOP Conf. Ser. Earth Environ. Sci.* **695**
- [35] Trilaksani W, Salamah E and Nabil M 2006 Utilization of tuna fish bone waste (Thunnus sp.) as a source of calcium using the protein hydrolysis method *Bul. Tekonologi Has. Perikan.* **9** 34–45
- [36] Logesh A R, Pravinkumar M, Raffi S M and Kalaiselvam M 2012 Calcium and phosphorus determination in bones of low value fishes, Sardinella longiceps (Valenciennes) and Trichiurus savala (Cuvier), from Parangipettai, Southeast Coast of India Asian Pacific J. Trop. Dis. 2 S254– 6
- [37] Toppe J, Albrektsen S, Hope B and Aksnes A 2007 Chemical composition, mineral content and amino acid and lipid profiles in bones from various fish species *Comp. Biochem. Physiol. - B Biochem. Mol. Biol.* 146 395–401
- [38] Jun J-Y, Jung M J, Jeong I-H, Kim G, Sim J, Nam S and Kim B 2019 Effects of crab shell extract as a coagulant on the textural and sensorial properties of tofu (soybean curd) *Food Sci. Nutr.* 7
- [39] Yihun F, Egusa M, Kaminaka H, Izawa H, Morimoto M, Saimoto H and Ifuku S 2016 Protein/CaCO3/Chitin Nanofiber Complex Prepared from Crab Shells by Simple Mechanical Treatment and Its Effect on Plant Growth Int. J. Mol. Sci. 17 1600
- [40] Budi W S, Winarko W, Rokhmalia F, Darjati D and Poerwati S 2023 Analysis of Nitrogen, Phosphorus, Potassium Content in Humus in Soil at Temporary Shelters J. Penelit. Kesehat. Suara Forikes 14 62–6
- [41] Umaternate G R, Abidjulu J and Wuntu A D 2014 Olsen and Bray Method Test in Analyzing Available Phosphate Content in Paddy Field Soil in West Konarom Village, North Dumoga District J. MIPA 3 6–10
- [42] Soetan K O, Olaiya C O and Oyewole O E 2010 The importance of mineral elements for humans, domestic animals and plants: A review *African J. food Sci.* **4** 200–22
- [43] Malik D, Narayanasamy N, Pratyusha V A, Thakur J and Sinha N 2023 Inorganic Nutrients: Macrominerals *Textbook of Nutritional Biochemistry* (Springer) pp 391–446
- [44] Bouillon R and Suda T 2014 Vitamin D: calcium and bone homeostasis during evolution *Bonekey Rep.* **3**

#### Acknowledgements

This research was supported by the RIIM LPDP Grant and BRIN, (Grant Number: 12/II.7/HK/2023; Decree Number: 76/IV/KS/05/2023; 0236/UN9.3.1/PL/2023). We also thanks to the Sriwijaya University and Research Center for Genetic Engineering, BRIN.



Rozirwan ROZIRWAN <rozirwan@unsri.ac.id>

### Paper(s) for ICTCRED-2024 entering Production

1 pesan

**Morressier Team** <discover@morressier.com> Kepada: rozirwan@unsri.ac.id 23 April 2025 pukul 14.31



# **IOP** Publishing

Dear Rozirwan,

We are pleased to inform you that the following Papers have passed the Publisher's checks and are being finalized for publication:

• The Natural Calcium and Phosporus Content in Two Fishbone of Commercial Species from Enggano Waters

These are now entering the production process to prepare them for publication on the IOPscience platform. An overview of the publication procedure is available here.

To view the paper log in to the My Submissions section via the Morressier platform.

Go to My submissions

Please contact your conference organizer for questions regarding:

- publication timeline.
- payments and refunds.
- certificates and letters of approval.
- conference details.

Please contact support@morressier.com or use the Get help option in your personal menu if you need any assistance using the platform.

Fancy catching up on other event proceedings? Take a look here.

Best wishes, IOP Publishing

Check out the FAQ page

# **IOP** Publishing

#### **IOP** Publishing

No.2, The Distillery Glassfields, Avon St, Bristol BS2 0GR, United Kingdom

Powered by Morresster



Rozirwan ROZIRWAN <rozirwan@unsri.ac.id>

### [Action Required] Proofreading Confirmation for IOP Conference Proceeding Article

2 pesan

ICTCRED Undip <ictcred@live.undip.ac.id> Kepada: rozirwan@unsri.ac.id 19 Mei 2025 pukul 10.19

Dear Author,

Attached file is the proof read article of your IOP conference proceeding before publication.

We kindly ask you to carefully review the document. Please note that the IOP Publisher will only accept corrections that affect the article's discoverability, including: title, author list, affiliations, display issues (e.g., corrupted or unreadable figures)

Content-related changes or revised papers will not be accepted. If you identify any issues related to the title, author list, affiliation, or figures, please send your corrections in both PDF and Word formats by **21 May 2025 at 08:00 AM WIB**.

Should we not receive any response by the deadline, we will proceed with submitting the document to the IOP Publisher in its current form.

Thank you for your attention and cooperation. Warmest Regards,

Committee of ICTCRED 2024 ictcred.fpik.undip.ac.id ictcred@live.undip.ac.id

#### Rozirwan\_2025\_IOP\_Conf.\_Ser.\_\_Earth\_Environ.\_Sci.\_1496\_012013.pdf 1241K

Rozirwan ROZIRWAN <rozirwan@unsri.ac.id> Kepada: ICTCRED Undip <ictcred@live.undip.ac.id> 19 Mei 2025 pukul 14.21

Yes, everything is correct. Thank you [Kutipan teks disembunyikan]

Prof. Dr. Rozirwan, S.Pi., M.Sc Head of Marine Bioecology Laboratory Department of Marine Science Faculty of Mathematics and Natural Sciences Sriwijaya University Jalan Raya Palembang-Prabumulih KM 32, Indralaya Ogan Ilir, Sumatera Selatan, Indonesia, Pos Code: 30862 Email: rozirwan@unsri.ac.id, rozirwan@gmail.com



Rozirwan ROZIRWAN <rozirwan@unsri.ac.id>

### Payment Information for IOP Publication – ICTCRED 2024

2 pesan

ICTCRED Undip <ictcred@live.undip.ac.id> Kepada: rozirwan@unsri.ac.id 29 Mei 2025 pukul 07.44

Dear Author,

Thank you for your participation in the International Conference on Tropical and Coastal Region Eco-Development (ICTCRED 2024). We are pleased to inform you that your article has been accepted for publication in the IOP Conference Series: Earth and Environmental Science (EES).

To proceed with the publication process, we kindly request you to complete the publication fee payment as outlined below:

Payment Method : Bank Transfer Bank Name : Bank Mandiri Account Number : 136-00-0604107-0 Account Holder : Anindya Wira Satriya **Payment Deadline : Monday, 02 June 2025** 

#### Important:

- Please include the note: "ICTCRED [First Author's Name]" in the bank transfer description/reference.
- After completing the transfer, kindly send the proof of payment to: ictcred@live.undip.ac.id

This invoice and payment process serve as an official requirement for the submission of your article to IOP. Failure to complete the payment by the deadline may result in the exclusion of your paper from the publication process.

Should you have any questions, please don't hesitate to contact us. Thank you for your attention and cooperation.

Warmest regards, ICTCRED 2024 Committee Email: ictcred@live.undip.ac.id Website: https://ictcred.undip.ac.id

1496\_012013\_Invoice\_Rozirwan.pdf 255K Rozirwan ROZIRWAN <rozirwan@unsri.ac.id> Kepada: ICTCRED Undip <ictcred@live.undip.ac.id>

Dear ICTCRED Committee,

Thank you very much for the notification and for accepting our article for publication in the *IOP Conference Series: Earth and Environmental Science (EES).* 

We are pleased to inform you that we have completed the Article Processing Charge (APC) payment via bank transfer. Please find the proof of payment attached to this email for your reference.

Thank you for your support and cooperation.

#### Best regards,

[Kutipan teks disembunyikan]

**Prof. Dr. Rozirwan, S.Pi., M.Sc** Head of Marine Bioecology Laboratory Department of Marine Science Faculty of Mathematics and Natural Sciences Sriwijaya University Jalan Raya Palembang-Prabumulih KM 32, Indralaya Ogan Ilir, Sumatera Selatan, Indonesia, Pos Code: 30862 Email: rozirwan@unsri.ac.id, rozirwan@gmail.com



# International Conference on Tropical and Coastal Region Eco-development (ICTCRED) 2024



 26/04/2025

 VOL. SERIES
 1496

 NUMBER
 012013

 DUE DATE
 02/06/2025

Jl. Prof. Jacub Rais, Kampus Universitas Diponegoro, Tembalang, Semarang, Pos Code 50275

Tel/fax. 024 - 7474698, email: ictcred@live.undip.ac.id / fpik@undip.ac.id

| BILL TO   | FOR                    |
|---|------------------------|
| ROZIRWAN  | Manuscript Publication |
| Department of Marine Science, Faculty of Mathematics and Natural Sciences, Sriwijaya University, Jalan Palembang-Prabumulih KM. 32, Indralaya, South Sumatra 30662, Indonesia |                        |
| Tel.fax.: +62 82134788194   |                        |
| Email: rozirwan@unsri.ac.id   |                        |
|   |                        |
| Title   |                        |
| The Natural Calcium and Phosporus Content in Two Fishbone of Commercial Species from Enggano<br>Waters  |                        |

| ITEM DESCRIPTION  | TAXED | AMOUNT (IDR)     | AMOUNT (USD)             |
|---|-------|------------------|--------------------------|
| Payment for manuscript publication to IOP Conference Series:<br>Earth and Environmental Science 1496 through ICTCRED 2024 |       | IDR 1.600.000,00 | USD 95                   |
|   |       |                  |                          |
|   |       |                  |                          |
|   |       |                  |                          |
|   |       | -                |                          |
|   |       | Subtotal         | IDR 1.600.00,00 / USD 95 |
|   |       | Tay Data         |                          |

| Total Cost  | IDR 1.600.000,00 / USD 95 |
|-------------|---------------------------|
| Other Costs |                           |
| Tax Rate    |                           |
| Subiolal    | IDK 1.000.00,00 / 03D 95  |

| : Bank Transfer        |
|------------------------|
| : Bank Mandiri         |
| : 136-00-0604107-0     |
| : Anindya Wira Satriya |
|                        |

Notes:

- Please complete the payment before [Monday, 02 June 2025].
- Kindly send the payment proof to email: <u>ictcred@live.undip.ac.id</u>.
- Please include the note: "ICTCRED [First Author's Name]" in your bank transfer description/reference.
- This invoice serves as an official billing document for reimbursement or institutional purposes.



# Transfer berhasil

# Rp1.600.000

# 29 Mei 2025 • 13:43:15 WIB • Ref ID: 20250529134305851728

# Penerima

ANINDYA WIRA SATRIYA MANDIRI • 1360006041070

# Sumber dana

ROZIRWAN TAPLUS • \*\*\*\*\*764

# Detail transfer

Nominal Biaya transaksi Metode transfer BIZ ID Tujuan transaksi

Catatan

20250529B Publication fe

BIN

Rp1.600.000 Rp2.500 BI-FAST

20250529BNINIDJA0100 0240608959

Lainnya

Publication fee Rozirwan et al the natural calcium

# Rp1.602.500

Total



#### **PAPER • OPEN ACCESS**

# The Natural Calcium and Phosporus Content in Two Fishbone of Commercial Species from Enggano Waters

To cite this article: Rozirwan et al 2025 IOP Conf. Ser.: Earth Environ. Sci. 1496 012013

View the article online for updates and enhancements.

#### You may also like

- <u>Tiger: Concept Study for a New Frontiers</u> Enceladus Habitability Mission Elizabeth M. Spiers, Jessica M. Weber, Chandrakanth Venigalla et al.
- Abundance, diversity, and trophic state status of estuary based on Bacillariophyta using eDNA method in Jakarta Bay, Indonesia
   Majariana Krisanti, Ali Mashar, Hefni Effendi et al
- Impact of Mining Activities: Concentration of Heavy Metals (As, Cd, Cu, Ni) in Three Filter Feeder Species in Southeast Sulawesi, Indonesia Mimie Saputri, Muhammad Zainuri, Suryanti Suryanti et al.

# Join the Society Led by Scientists, for Scientists Like You!



This content was downloaded from IP address 114.125.249.206 on 01/06/2025 at 16:06

## The Natural Calcium and Phosporus Content in Two Fishbone of Commercial Species from Enggano Waters

Rozirwan<sup>1\*</sup>, Nopriani<sup>1</sup>, Fauziyah<sup>1</sup>, Wike Ayu Eka Putri<sup>1</sup>, Melki<sup>1</sup>, Iskhaq Iskandar<sup>2</sup>, Ela Novianti<sup>3</sup>, Apon Zaenal Mustopa<sup>3</sup>, Fatimah<sup>3</sup>, Redho Yoga Nugroho<sup>1</sup>

<sup>1</sup>Department of Marine Science, Faculty of Mathematics and Natural Sciences, Sriwijaya University, Jalan Palembang-Prabumulih KM. 32, Indralaya, South Sumatra 30662, Indonesia

<sup>2</sup>Department of Physics, Faculty of Mathematics and Natural Sciences, Sriwijaya University, Jalan Palembang-Prabumulih KM. 32, Indralaya, South Sumatra, 30862, Indonesia

<sup>3</sup>Research Center for Genetic Engineering, National Research and Innovation Agency, West Java 16911, Indonesia

\*Corresponding author: rozirwan@unsri.ac.id

Abstract. Calcium and phosphorus are important elements needed for the growth and development of the body both in humans and other living things. The bones of Tiger Grouper Fish (Epinephelus fuscoguttatus) and Lencam Fish (Lethrinus lentjan) have a fairly high content of calcium and phosphorus. The purpose of this study was to determine the calcium and phosphorus content in the bones of Tiger Grouper Fish and Lencam Fish. This research was carried out in September-November 2023 with direct sampling of collectors in Banjarsari Village, Enggano Island. The calcium content test was performed using an Atomic Absorption Spectrophotometer (AAS) and the phosphorus content test was performed using a UV-Vis spectrophotometer. The results of this study showed that the calcium content in the bones of Tiger Grouper was 8.84% and for Lencam Fish was 7.97%, and the phosphorus content produced in both samples was 1.4% for Tiger Grouper Fish of 1.4% and 1.43% for Lencam Fish. Based on the results of the study, the calcium content in Tiger Grouper is greater than that of Lencam, while the phosphorus content in Tiger Grouper is smaller than Lencam Fish.

#### 1. Introduction

The production of catches produced on Enggano island can reach 27.89% of the total production of capture fisheries in North Bengkulu [1,2]. Snapper, Grouper and Pomfret are the three types of marine capture fisheries that produce the highest production from sea waters compared to coastal areas [3]. The high development of the industry in the field of processing fishery products will have the potential to increase organic waste in the area [4–6]. According to [7], The most common types of fishery waste are fish bones, fish heads, fish guts, and even shells, while this waste cannot be recycled due to the many limitations of its advanced processing [8]. Actually, it is estimated to be a material that can be used again, especially the use of fish bones which have natural calcium and phosphorus content.

Calcium and phosphorus are very important for bone and tooth growth and calcium intake must be considered early to meet the needs of bone growth in humans [9]. Several health problems can arise due to calcium and phosphorus deficiency, namely the risk of osteoporosis. In Indonesia, osteoporosis is caused by the low average calcium intake of the Indonesian population which is only 254 mg per day.

Content from this work may be used under the terms of the Creative Commons Attribution 4.0 licence. Any further distribution of this work must maintain attribution to the author(s) and the title of the work, journal citation and DOI. Published under licence by IOP Publishing Ltd 1

The average daily calcium requirement is 300-400 mg for infants, 500 mg for children, 600-700 mg for adolescents, and 500-800 mg for adults [10].

Fish bones are a type of fishery waste that contains a number of important benefits for humans. The main components of fish bones are calcium, phosphorus, and carbonate, so that fish bones themselves have the highest calcium content compared to other parts of the fish's body [11]. Calcium derived from fish bones is a waste product and is currently not used on a large scale for human needs. Therefore, further research is needed on the benefits of fish bone waste to be used as processed products that are beneficial for fisheries and marine problems [12]. One of the uses of fish bones is to process them into fish bone meal.

According to [13], calcium and phosphorus are macrominerals that are directly related to the development and maintenance of the skeleton system and participate in various physiological processes of the organism's body. The nutritional content of fish bones in 100 g of fish bone meal is calcium 735 mg, protein 9.2 g, fat 44 mg, phosphorus 345 mg, iron 78 mg, ash 24.5 g, and carbohydrates 0.1 mg [14]. Calcium has a role as an important micronutrient in addition to phosphorus for the growth and development of children and calcium and phosphorus are very important in the bone mineralization process [15]. Calcium plays an important role in physiological processes, such as cell metabolism, muscle contraction, and bone growth [16]. Phosphorus has metabolic functions to build bones, scales, adenosine triphosphate, cell membranes, and nucleic acids [4]. Chemical analysis revealed that fish bones are a valuable source of calcium phosphate as an economical source for synthetic hydrocytopathy. Bone is a biological composition consisting of an inorganic phase (calcium phosphate with a structure like carbonated hydroxyapati) [17].

The fisheries from the waters of Enggano that are consumed are Tiger Grouper Fish and Lencam Fish, is one type of fish waste that has the potential to be researched because it is included in the target fish species and is also the most dominant economic fish found [18]. Morphologically and physiologically the two samples are different, but the living habits or habitats of the two samples are almost the same, so this test was carried out to determine the difference in calcium and phosphorus content of Tiger Grouper and Lencam fish.

One of the indicators of the existence of Tiger Grouper fish is the presence of coral reefs in a water area to be able to shelter and as a place to forage [19]. Adult Tiger Grouper Fish can reach depths of more than 2 m. This type of fish often lives in clear waters with many coral reefs and depths of up to 60 m. The habitat of this fish is in the shallow waters of coral reefs, rocky seabeds, lagoon tops, coral channels, and the outermost part of steep coral reefs. This Grouper also likes waters with a sandy, rocky bottom and seaweed growing at the bottom [20]. In general, groupers are solitary, but when they want to spawn the fish in groups together. In general, this fish lives at a depth of 5-20 m in all types of corals in good condition. Most Tiger Grouper fish use burrows/cavities/holes in coral reefs as shelter and usually move around.

Lencam fish usually live in coastal areas, coral waters and belong to the group of underwater fish. The habitat of the Lencam fish is coral reefs, seagrass meadows, mangrove areas and shallow coastal waters with a sandy bottom at a depth of 30 m [21]. Lencam fish consists of 29 species spread around the world and about 81% of them are found in Indonesian waters. This fish continues to be threatened both directly and indirectly due to exploitation caused by various human activities [22].

#### 2. Materials and Methods

#### 2.1 Study area

This research was carried out in September – December 2023 with sampling obtained from fish collectors in Banjarsari Village, Enggano Island, Enggano District, North Bengkulu Regency, Bengkulu Province (Figure 1). Further sample analysis was carried out at the Marine Bioecology Laboratory and the Oceanography and Marine Instrumentation Laboratory at Sriwijaya University.

#### doi:10.1088/1755-1315/1496/1/012013



Figure 1. Sampling location of tiger grouper and lencam fish in Enggano Waters

#### 2.2 Sample Identification and Preparation

The samples used were taken from fish collectors in Banjarsari Village, Enggano Island. The fish samples amounted to 2 Tiger Grouper Fish and 2 Lencam Fish. The fish was still fresh, seen from its red color and not slimy. The sample preparation process was carried out at the Marine Bioecology Laboratory, starting with dissecting the fish using a knife. Then the dissected sample was washed and then boiled using a hot plate with a temperature of 60-80 °C for 60 min. After boiling, the fish was cleaned again using a brush to remove the meat that was still attached to the fish bones, then NaOH alkaline extraction was carried out which aimed to remove fat and blood levels in the sample. Samples that have been extracted for 60 min were cleaned again using aquades and then baked for 48 h at a temperature of 60°C. After the sample dried, it was crushed with mortal until it became fish bone meal powder which will later be used to make calcium and phosphorus test solutions. Fish bone meal powder was taken as much as 1 g (calcium) and 2 g (phosphorus) weighed using an analytical scale. Each sample taken was put into the erlenmeyer to make a test solution. The preparation of this test solution was carried out in the fume hood with the addition of NaOH 1.5 N, HNO<sub>3</sub>, HCI, H<sub>2</sub>O<sub>2</sub>, NaHCO<sub>3</sub>, (NH<sub>4</sub>)<sub>6</sub>Mo<sub>7</sub>O<sub>24</sub>.4H<sub>2</sub>O, H<sub>2</sub>O(SbO)C<sub>4</sub>H<sub>4</sub>O<sub>6</sub> 0.5K, H<sub>2</sub>SO<sub>4</sub> and PO<sub>4</sub>. After the fish bone sample becomes powder, a destructive process is carried out to determine the calcium and phosphorus content. The calcium content test itself was carried out using an Atomic Absorption Spectrophotometer (AAS) with a wavelength of 420 nm, while the phosphorus content testing was carried out using a tool called the Uv-Vis Spectrophotometer with a wavelength of 889 nm.

#### 2.3 Sample Destruction

Common destructions to determine the mineral components present in foodstuffs were known to be of two types, namely dry and wet [23]. Sample destruction in this study will use the wet destruction method referring to the study [24]. Their results showed that wet destruction gave better results than dry destruction because the temperature used could not exceed the boiling point of the solution and in general the carbon was destroyed faster than using dry ashing.

#### 2.4 Calcium level testing

The calcium content was determined by measuring the sample by wet destruction using an Atomic Absorption Spectrophotometer (AAS) with a wavelength of 420 nm, and the calcium content was checked using a modified method [25]. To analyze the calcium content, it can be done by weighing as much as 0.1 g of fine samples and transferring them to a 100 mL measuring flask. The destruction process was carried out by adding 15 mL of hydrochloric acid (HCl) to digest the sample. The solution was decomposed until it was clear and then cooled. Filtration volumes up to 100 mL can be measured with atomic absorption spectrophotometer (AAS) [26].

#### 2.5 Phosphorus level testing

The determination of phosphorus content in the solution of fish bone samples aimed to obtain the concentration of UV-Vis spectrophotometer (ppm) readings. The determination of the maximum wavelength P was carried out by analyzing the sample of phosphorus sample using a UV-Vis spectrophotometer at a wavelength of 889 nm. With this determination of wavelength, the maximum result of the P content in the fishbone sample [27]. The dissolved concentration obtained from measurements using a UV-Vis spectrophotometer was then recorded so that a graph of the wavelength and absorbance relationship in each fishbone sample used in testing the content of each sample was obtained.

2.6 Data analysis

2.6.1 Moisture content analysis

%Moisture content was calculated based on SNI-19-1705-2000 and AOAC (1995). %Moisture content =  $\frac{Wo-Wt}{Wt} \times 100\%$  fk =  $\frac{100}{100-\% Moisture content}$ 

Information:

Wt : Final weight of fish bone meal (g)

Wo : Initial weight of the fish bone (g)

: 100 / (100 - % of air) Fk

#### 2.6.2 Calcium concentration data analysis

The concentration obtained from the curve was a concentration in mg/L. While the sample used in this study can be in the form of a solid so that it will be converted into mg/kg. To calculate the level of calcium in fish bone samples in ppm by sample weight, which was formulated as follows:

Mineral concentration (mg/kg) =  $\frac{\text{Sample concentartion}\left(\frac{\text{mg}}{\text{L}}\right) \text{x sample volume (L)}}{\text{Sample weight (kg)}} \text{x } D_{\text{f}}$ Information :

D<sub>f</sub> : Dilution factor

#### 2.6.3 Phosphorus concentration data analysis

The processing of phosphorus concentration data that was previously in the form of ppm units is converted into % which is formulated as follows:

 $P_2O_5$  levels available (ppm) = ppm curve x ml extract/1,000 mL x 1,000 g (g example) x  $D_f x 142/190$  $x M_{f}$ 

Information :

| ppm curve        | : | the sample level obtained from the relationship curve between the standard series |
|------------------|---|---|
|                  |   | and its reading after blank correction  |
| $D_{\mathrm{f}}$ | : | dilution factor (if any)  |
| 142/190          | : | PO <sub>4</sub> to P <sub>2</sub> O form conversion factor                        |
| $M_{\mathrm{f}}$ | : | moisture content correction factor  |

#### 3. Result and Discussion

3.1 Description of Tiger Grouper Fish

Based on the identification [28,29] the fish obtained were classified as *Epinephelus fuscoguttatus species*. Observations of the morphology were size, body shape of the fish, the position of the mouth, the type of fins and also the type of scales.



Figure 2. (a) Fish body parts, (b) Mouth type, (c) Scale type, (d) Fin type

Observation of the morphology of the Tiger Grouper fish consists of parts of the fish's body consisting of the head, body, fins and tail. The body shape of this fish in figure 3 (a) is compressed. In Figure (b) the mouth type is a terminal type whose upper and lower jaws can be seen to be the same length (parallel), and is often used to eat prey floating in front of it. Figure (c) body type covered by scales with a brownish body color and filled with reddish spots. While Figure (d) is a rounded fin type, meaning that the dorsal fin is elongated and totals 8-10 pieces and the caudal fin totals 15-17 pieces.

#### 3.2 Description of Lencam Fish

The determination of the morphological characteristics of Lencam fish includes body shape, mouth position, caudal fin shape, and scale type.



Figure 3. (a) Fish body parts, (b) Mouth type, (c) Scale type, (d) Fin type

Lencam fish are characterized by a flattened body that is covered with many scales, smooth cheeks, at the base of the ventral fin there are large scales (axillary scale) that are well developed. The mouth is located at the front end of the fish's head (terminal type) with a relatively thick lip, the front of the jaw has strong canines, on the side of the jaw there are conical teeth and large molars, there are no teeth at the top of the oral cavity. The first dorsal fin is fused with the second fin, consisting of 9 hard fingers and 9 or 10 soft fingers, the anal fin with 3 hard fingers, 8-10 soft fingers. (a) is a part of the

body of a fish consisting of 3 parts, namely the head, body and tail, in Figure (b) the mouth type of the Lencam fish is a terminal mouth type, meaning it has the same length of upper and lower jaws, and often eats prey floating in front of it. Figure (c) the shape of the body is rather tall and flattened, and the body is covered by scales. While Figure (d) is the type of *Homocercal fin*, the shape of the pinna caudalis is curved or not and supported by the spokes of the tail fin. In this type, the length between the two ends of the tail is the same length and also forms like the letter V.

#### 3.3 Yield Value of Tiger Grouper Fish and Lencam Fish

The yield value in this study is the result of a comparison between the initial weight of the fish bone and the amount produced from the final weight of the fish bone powder. According to [30], the level of yield value produced shows that the number of products produced is also high, indicating that the process used is more effective.

#### Table 1. Yield Value

| Types of fish      | Bone weight (g) | Fish bone powder (g) | Yield value (%) |
|--------------------|-----------------|----------------------|-----------------|
| Tiger Grouper Fish | 17.98           | 6.7                  | 2.6836          |
| Lencam Fish        | 13.04           | 4.05                 | 3.2198          |

The yield value obtained in this study can be an indicator of the quality of bone powder in both types of fish samples. It can be assumed that a high yield value indicates good flour quality. The magnitude of the yield obtained is influenced by the effectiveness in the extraction process. According to [31], a high yield will not necessarily produce the highest calcium levels, but it is also determined by other factors such as the low protein content in the material. The low yield is also suspected to be due to the influence of drying, where drying is the process of removing or disposing of liquid materials from a material.

#### 3.4 Calcium Content of Tiger Grouper Fish and Lencam Fish Bones

The analysis of calcium content in this study was obtained from measurements or readings using an atomic absorption spectrophotometer (AAS).

| Table 2. Calcium Content |            |        |                   |
|--------------------------|------------|--------|-------------------|
| Types of Fish Bones      | Conc (ppm) | Ca (%) | Quality Standards |
|                          | 0.470      | 0.40   | (70)              |
| Liger Grouper Fish       | 8.4/9      | 8.48   | 20-30             |
| Lencam Fish              | 7.967      | 7.97   | 20 50             |

The results of the measurement of calcium content in the bone samples of Tiger Grouper and Lencam Fish are known that the calcium content in Tiger Grouper is 8.84%, while in Lencam Fish it is 7.97%.



The difference in calcium content in the two samples can be influenced by the difference in the samples used, because basically different types of fish affect the value of the calcium levels produced. The results of measuring the calcium content of both Tiger Grouper and Lencam fish bones with different sizes are assumed that the sample size affects the mineral content in the fish's body, one of which is the bones. This is in line with the opinion [32,33], that minerals play an important role in the maintenance of every body function at the level of cells, tissues, organs, and overall body functions, and the presence of macro minerals in the body of fish is influenced by several factors, such as the size of the fish.

#### 3.5 Phosphorus Content of Tiger Grouper Fish and Lencam Fish Bones

The analysis of phosphorus content in this study was obtained by reading the results of the UV-Vis Spectrophotometer. The phosphorus content obtained in the bones of Tiger Grouper Fish and Lencam Fish with the same weight category in this study is seen in Table 3.

| Types of Fish Bones | Conc (ppm) | P (%) | Quality Standards<br>(%) |
|---------------------|------------|-------|--------------------------|
| Tiger Grouper Fish  | 1,401.49   | 1.40  | 20-30 %                  |
| Lencam Fish         | 1,432.58   | 1.43  |                          |

Table 3. Phosphorus Content of Fish Bones

Results of testing phosphorus content in the bones of Tiger Grouper and Lencam fish in the same weight category in mg/kg and percentage (%). Based on the quality standards of fish bone meal according to SNI 01-3158-1992, it is understood that the phosphorus content in the two samples does not meet the quality requirements. Furthermore, the results that have been obtained are visualized in the form of a graph to show a comparison of the results between the two fish, which can be seen in the following graph.



Figure 5. Measurement of phosphorus content

The phosphorus content in Lencam fish bone samples was higher than the phosphorus content in Tiger Grouper fish bone samples. The phosphorus content in Tiger Grouper can be seen at 1.4%, while in Lencam fish the content is slightly higher at 1.43%. The comparison of these two samples is around 0.03%. The type of sample, the material and the procedure used to measure the calcium and phosphorus content can also affect the results obtained. Different types of fish used in the study can affect the calcium and phosphorus content obtained, because the mineral content in each fish is different. Another factor that can cause this difference is also due to the sample size and the process in handling the sample.

The type of sample, materials and procedures in the study used to measure calcium and phosphorus levels can also affect the results obtained. This was revealed by [12] in the results of the study, the analysis of various types of frequencies in the boiling process affected the calcium content produced in the bones of Belida fish (*Chitala lopis*) ranging from 28.25% and the phosphorus content varied between 3.98% - 4.04%. The difference in NaOH concentration during the extraction process

affects the calcium and phosphorus content obtained, namely the concentration of 1.5 N provides the best content compared to the concentration of 1 N and 2 N. The selection of the test method in measuring the calcium and phosphorus content can affect the results obtained in each sample used.

#### 3.6 Calcium and phosphorus content in several types of fish bones

Based on Table 4, it is the result of a comparison of calcium and phosphorus content from various sources and also from various other types of marine fish. The different calcium and phosphorus content in the table below is caused by various factors, especially from the type of fish, the size of the fish and also influenced by the sample handling process before testing the content level in each sample. The method used in each source is also different, so it can be stated that the content is different because it is also due to the method used, because each method uses a different solution and also the processing process before being made into a solution is also different.

| No | Types of Fish Bones  | Calcium %   | Phosphorus % | References   |
|----|----------------------|-------------|--------------|--------------|
| 1  | Epinephelus sp       | 8.48        | 1.40         | This results |
| 2  | Lethrinus sp         | 7.97        | 1.43         | This results |
| 3  | Chitala lopis        | 28.25-31.31 | 3.98-4.06    | [12]         |
| 4  | P, albilabris        | 48.10       | 35.40        | [34]         |
| 5  | Thunnus sp.          | 39.24       | 13.66        | [35]         |
| 6  | Trichiurus savala    | 19.33       | 7.21         | [36]         |
| 7  | Sardinella longiceps | 26.39       | 16.57        | [36]         |
| 8  | Scomber scombrus     | 14.3        | 8.6          | [37]         |
| 9  | Salmo trutta         | 14.7        | 8.7          | [37]         |
| 10 | Salmo salar          | 13.5        | 8.1          | [37]         |
| 11 | Micromesistius       | 17.0        | 87           | [37]         |
| 11 | poutassou            | 17.0        | 0.7          | [57]         |
| 12 | Clupea harengus      | 16.1        | 9.4          | [37]         |

| Table 7. Calcium and Thosphorus Content in Several Types of This Dones |
|--|
|--|

Based on the comparison table of calcium content in the table above, it can be stated that the calcium content in fish bones is lower than the calcium content in crab shell shells. This was also explained by [38,39], it was explained that the calcium carbonate content reached 53.70%-78.40% contained in the shell of the shell. As for the phosphorus content in fish bones from various sources presented in the table above, it is known that the phosphorus content is higher than the phosphorus content in seaweed. According to [40], his research explained that the phosphorus content in seaweed is >0.05, it can also be concluded that the phosphorus content is indeed higher than the content of other species. The difference in calcium content is also caused by the difference in the species analyzed, because every difference in a living creature can cause the results of the analysis related to the nutrient content in that type or species to be different. The difference between this study and the previous study lies in the category of fish sample size used and the chosen destruction method. The sample size was made in 3 categories to see the difference between them, while the selection of the method for phosphorus testing with P Bray I where the phosphorus read was only in the form of H<sub>2</sub>PO<sub>4</sub><sup>-</sup> and PO<sub>4</sub><sup>3-</sup>. In line with the results of a study [41], which tested soil samples with two different methods.

Calcium is an important mineral that can be relied on in the human body in a variety of several important processes [42]. Most of the calcium in the body is in the bones. Your circulatory system also has a certain amount of calcium which is essential for blood clotting as well as the function of blood vessels, nerves, muscles and heart. Calcium and phosphorus are one of the substances that have an important role in various bodily functions, such as bone mineralization, functioning as cofactors in many enzyme systems, maintaining muscle and nerve excitation, and in the case of iron, maintaining oxygen transport and blood capacity [43]. Many female athletes consume less calcium than the recommended nutritional adequacy rate. It is also a concern due to the need to reach peak bone mass during adolescence and the possible relationship of poor calcium intake with stress-induced fractures. Calcium and

phosphorus are available in various supplement forms, both can be taken together as individual supplements, and there are calcium phosphate supplements that are a combination of the two minerals. The role of phosphorus in the human body functions in the growth and development of energy in the body. The high content of calcium and phosphorus in fish bones makes fish bones can be processed into natural ingredients to replace bones or hydroxyapathic materials. According to [26], hydroxyapatite  $(Ca_{10}(PO_4)_6(OH)_2)$  is a bioceramic material that is biocompatible, biodegradable, and non-toxic. Hydroxyapatite can be synthesized from natural materials that contain relatively high levels of calcium which will later be used as calcium precursors because hydrosapathic compounds can be obtained by mixing calcium precursors with phosphate precursors. The functions of calcium in the body, especially in fish, include as the main component of the formation of bones, teeth, skin and scales, as well as other functions can maintain the strength of the body's skeleton, thicken the blood, become an intracellular regulator which can also help regulate the activity of skeletal muscles, heart and other tissues, muscle contraction and relaxation, help the absorption of vitamin B12, maintain osmotic balance [9][44]. This is supported by research [13], which states that calcium plays an important role in skeletal mineralization and bone formation, not only that calcium is also needed for bone growth and strength.

#### Conclusion

Grouper Fish Bones have a higher calcium content of 8.48% compared to Lencam Fish bones of 7.97%. Meanwhile, Lencam fish bones have a phosphorus content 1.43% higher than Tiger Grouper 1.4%. Based on the quality requirements of fish bone meal according to SNI 01-3158-1992, the calcium and phosphorus content of the bones of Tiger Grouper Fish and Lencam Fish do not meet the quality requirements.

#### References

- [1] Limi M A, Ningsih S R, Fyka S A, Syarni P, Dewi H S and Attamimi U 2022 Analysis of crab fisherman's household structure and livelihood strategy after the Kendari Bay revitalization *IOP Conference Series: Earth and Environmental Science* vol 1118 (IOP Publishing) p 12075
- [2] Rozirwan, Adefta A, Apri R, Fauziyah, Putri W A E, Melki, Iskandar I, Novianti E, Mustopa A Z, Fatimah and Nugroho R Y 2024 The Ecological Health of Coral Reefs In The Waters of Enggano Island, The Outermost Island In The Eastern Indian Ocean Int. J. Conserv. Sci. 15 1947–58
- [3] Rozirwan, Nugroho R Y, Fauziyah, Putri W A E, Melki, Ulqodry T Z, Isnaini, Absori A and Iskandar I 2023 Mollusks diversity in the protected coastline of Berbak-Sembilang National Park Indonesia *Int. J. Conserv. Sci.* **14** 1627–40
- [4] Fraser T W K, Witten P E, Albrektsen S, Breck O, Fontanillas R, Nankervis L, Thomsen T H, Koppe W, Sambraus F and Fjelldal P G 2019 Phosphorus nutrition in farmed Atlantic salmon (Salmo salar): Life stage and temperature effects on bone pathologies *Aquaculture* 511 734246
- [5] Khotimah N N, Rozirwan, Putri W A E, Fauziyah, Aryawati R, Isnaini and Nugroho R Y 2024 Bioaccumulation and Ecological Risk Assessment of Heavy Metal Contamination (Lead and Copper) Build Up in the Roots of Avicennia alba and Excoecaria agallocha J. Ecol. Eng. 25 101– 13
- [6] Rozirwan, Saputri A P, Nugroho R Y, Khotimah N N, Putri W A E, Fauziyah and Purwiyanto A I S 2023 An Assessment of Pb and Cu in Waters, Sediments, and Mud Crabs (Scylla serrata) from Mangrove Ecosystem Near Tanjung Api-Api Port Area, South Sumatra, Indonesia Sci. Technol. Indones. 8 675–83
- [7] Venugopal V 2022 Green processing of seafood waste biomass towards blue economy *Curr. Res. Environ. Sustain.* **4** 100164
- [8] Rozirwan, Nanda, Nugroho R Y, Diansyah G, Muhtadi, Fauziyah, Putri W A E and Agussalim A 2023 Phytochemical composition, total phenolic content and antioxidant activity of Anadara granosa (Linnaeus, 1758) collected from the east coast of South Sumatra, Indonesia *Baghdad Sci. J.* 20 1258–65
- [9] Raskh S 2020 The importance and role of calcium on the growth and development of children and its complications *Int. J. Res. Appl. Sci. Biotechnol.* **7** 162–7

- doi:10.1088/1755-1315/1496/1/012013
- [10] Stoutjesdijk E 2020 *Brain-selective nutrients in pregnancy and lactation* (Groningen: University of Groningen)
- [11] Meiyasa F and Tarigan N 2020 Utilization of tuna fish bone waste (Thunnus sp.) as a source of calcium in making seaweed sticks *J. Teknol. Pertan. Andalas* **24** 66–75
- [12] Kondolele S L, Asikin A N, Kusumaningrum I, Diachanty S and Zuraida I 2022 Effect of Boiling Temperature on Physicochemical Characteristics of Mackerel Fish Bone Flour (Scomberomorus commerson) *Media Teknol. Has. Perikan.* 10 177–84
- [13] Ciosek Ż, Kot K, Kosik-Bogacka D, Łanocha-Arendarczyk N and Rotter I 2021 The effects of calcium, magnesium, phosphorus, fluoride, and lead on bone tissue *Biomolecules* **11** 506
- [14] Kaliky N, Lessy A, Rieuwpassa F, Kaya A O, Mailoa M N and Sormin R B D 2024 Physicochemical Characteristics of Red Snapper Fish Bone Flour (Lutjanus sp.) BIOSEL (Biology Sci. Educ. J. Penelit. Sci. dan Pendidik. 13 129–40
- Bhadada S K and Rao S D 2021 Role of phosphate in biomineralization Calcif. Tissue Int. 108 32–40
- [16] Battafarano G, Rossi M, Marampon F, Minisola S and Del Fattore A 2020 Bone control of muscle function *Int. J. Mol. Sci.* 21 1178
- [17] Kanis J A, Norton N, Harvey N C, Jacobson T, Johansson H, Lorentzon M, McCloskey E V, Willers C and Borgström F 2021 SCOPE 2021: a new scorecard for osteoporosis in Europe Arch. Osteoporos. 16 82
- [18] Scabra A R, Marzuki M and Yarni B M 2023 The Effect of Calcium Hydroxide (CAOH2) and Phosphorus (P) on the Growth of Vannamei Shrimp (Litopenaeus Vannamei) in Freshwater Media J. Ruaya J. Penelit. Dan Kaji. Ilmu Perikan. Dan Kelaut. 11 39–51
- [19] Paxton A B, Harter S L, Ross S W, Schobernd C M, Runde B J, Rudershausen P J, Johnson K H, Shertzer K W, Bacheler N M and Buckel J A 2021 Four decades of reef observations illuminate deep-water grouper hotspots *Fish Fish.* 22 749–61
- [20] Craig M T, de Mitcheson Y J S and Heemstra P C 2024 *Groupers of the world: a field and market guide* (Honolulu: CRC Press)
- [21] Prihatiningsih P, Pane A R P, Herlisman H and Hartati S T 2021 Reproduction and Growth of Lencam Fish (Lethrinus atkinsoni Seale, 1910) in Wakatobi Waters, Southeast Sulawesi Bawal Widya Ris. Perikan. Tangkap 13 111–22
- [22] Saini V P, Singh A K, Paul T, Thakur A and Singh M 2024 Advances in Environment Management for Sustainable Fisheries & livestock production (Patna: Bihar Animal Sciences University)
- [23] El Hosry L, Sok N, Richa R, Al Mashtoub L, Cayot P and Bou-Maroun E 2023 Sample preparation and analytical techniques in the determination of trace elements in food: A review *Foods* 12 895
- [24] Rozirwan, Ramadani S, Ayu W, Putri E, Nur N and Nugroho R Y 2023 Evaluation of Calcium and Phosphorus content in Scallop Shells (Placuna placenta) and Blood Cockle Shells (Anadara granosa) from Banyuasin Waters, South Sumatra Egypt. J. Aquat. Biol. Fish. Zool. Dep. Fac. Sci. 27 1053–68
- [25] Gómez-Nieto B, Motyzhov V, Gismera M J, Procopio J R and Sevilla M T 2020 Fast-sequential determination of cadmium and copper in milk powder and infant formula by direct solid sampling high-resolution continuum source graphite furnace atomic absorption spectrometry *Microchem. J.* 159 105335
- [26] Sareen A, Gupta R C, Bansal G and Singh V 2020 Comparison of key mineral elements in wild edible fruits of Ziziphus Mauritiana and Z. nummularia using atomic absorption spectrophotometer (AAS) and flame photometer *Int. J. Fruit Sci.* **20** S987–94
- [27] Pratiwi R A, Bayu A and Nandiyanto D 2022 How to Read and Interpret UV-VIS Spectrophotometric Results in Determining the Structure of Chemical Compounds Indones. J. Educ. Res. Technol. 2 1–20
- [28] Webb P W and Weihs D 1986 Functional locomotor morphology of early life history stages of fishes *Trans. Am. Fish. Soc.* **115** 115–27
- [29] Strauss R E and Bond C E 1990 Taxonomic methods: morphology Methods fish Biol. 109-40

- [30] Rosida R, Handayani L and Apriliani D 2018 Utilization of goat fish bone waste (Abalistes stellaris) as gelatin using variations in CH3COOH concentration Acta Aquat. Aquat. Sci. J. 5 93–9
- [31] Henchion M, Hayes M, Mullen A M, Fenelon M and Tiwari B 2017 Future protein supply and demand: strategies and factors influencing a sustainable equilibrium *Foods* **6** 53
- [32] Ali A A H 2023 Overview of the vital roles of macro minerals in the human body *J. Trace Elem. Miner.* **4** 100076
- [33] Lall S P and Kaushik S J 2021 Nutrition and Metabolism of Minerals in Fish *Animals* 11 2711
- [34] Sumarto, Desmelati, Sari N I, Angraini R M and Arieska L 2021 Characteristic of Nano-Calcium Bone from a Different Species of Catfish (Pangasius hypophthalmus, Clarias batrachus, Hemibagrus nemurus and Paraplotosus albilabris) *IOP Conf. Ser. Earth Environ. Sci.* **695**
- [35] Trilaksani W, Salamah E and Nabil M 2006 Utilization of tuna fish bone waste (Thunnus sp.) as a source of calcium using the protein hydrolysis method *Bul. Tekonologi Has. Perikan.* **9** 34–45
- [36] Logesh A R, Pravinkumar M, Raffi S M and Kalaiselvam M 2012 Calcium and phosphorus determination in bones of low value fishes, Sardinella longiceps (Valenciennes) and Trichiurus savala (Cuvier), from Parangipettai, Southeast Coast of India Asian Pacific J. Trop. Dis. 2 S254– 6
- [37] Toppe J, Albrektsen S, Hope B and Aksnes A 2007 Chemical composition, mineral content and amino acid and lipid profiles in bones from various fish species *Comp. Biochem. Physiol. - B Biochem. Mol. Biol.* 146 395–401
- [38] Jun J-Y, Jung M J, Jeong I-H, Kim G, Sim J, Nam S and Kim B 2019 Effects of crab shell extract as a coagulant on the textural and sensorial properties of tofu (soybean curd) *Food Sci. Nutr.* 7
- [39] Yihun F, Egusa M, Kaminaka H, Izawa H, Morimoto M, Saimoto H and Ifuku S 2016 Protein/CaCO3/Chitin Nanofiber Complex Prepared from Crab Shells by Simple Mechanical Treatment and Its Effect on Plant Growth Int. J. Mol. Sci. 17 1600
- [40] Budi W S, Winarko W, Rokhmalia F, Darjati D and Poerwati S 2023 Analysis of Nitrogen, Phosphorus, Potassium Content in Humus in Soil at Temporary Shelters J. Penelit. Kesehat. Suara Forikes 14 62–6
- [41] Umaternate G R, Abidjulu J and Wuntu A D 2014 Olsen and Bray Method Test in Analyzing Available Phosphate Content in Paddy Field Soil in West Konarom Village, North Dumoga District J. MIPA 3 6–10
- [42] Soetan K O, Olaiya C O and Oyewole O E 2010 The importance of mineral elements for humans, domestic animals and plants: A review *African J. food Sci.* **4** 200–22
- [43] Malik D, Narayanasamy N, Pratyusha V A, Thakur J and Sinha N 2023 Inorganic Nutrients: Macrominerals *Textbook of Nutritional Biochemistry* (Springer) pp 391–446
- [44] Bouillon R and Suda T 2014 Vitamin D: calcium and bone homeostasis during evolution *Bonekey Rep.* **3**

#### Acknowledgements

This research was supported by the RIIM LPDP Grant and BRIN, (Grant Number: 12/II.7/HK/2023; Decree Number: 76/IV/KS/05/2023; 0236/UN9.3.1/PL/2023). We also thanks to the Sriwijaya University and Research Center for Genetic Engineering, BRIN.