

Heavy Metal Accumulation and Ecological Risk on Seagrass *Cymodocea* and *Thalassia* in Pahawang Island, Indonesia

Ariqoh Athallah Gusri¹, Rozirwan^{2*}, Wike Ayu Eka Putri², Melki², Isnaini²,
Redho Yoga Nugroho², Che Abd Rahim Mohamed³

¹Environmental Management Study Program, Graduate Program, Universitas Sriwijaya
Jl. Padang Selasa No. 524, Palembang South Sumatra, 30139 Indonesia

²Department of Marine Science, Faculty of Mathematics and Natural Sciences, Universitas Sriwijaya
Jl. Raya Palembang-Prabumulih Km. 32, Ogan Ilir South Sumatra, 30862 Indonesia

³Faculty of Science and Technology, Universiti Kebangsaan Malaysia
43600 UKM Bangi, Selangor, Malaysia.

Email: rozirwan@unsri.ac.id

Abstract

Seagrass species thrive in coastal ecosystems and known for their ability to accumulate heavy metals from their surrounding environment. This study aims to evaluate the ecological risks related to the accumulation of heavy metals in seagrass roots, leaves, and sediments. The seagrass examined belong to the genera *Cymodocea* and *Thalassia*, collected from two sites: Jeralangan and Cukuh Nyai on Pahawang Island, Lampung, Indonesia. The heavy metals analyzed included Pb, Cu, Ni, and Zn, which were measured using the wet destruction method and quantified with a SHIMADZU AA-7000 Atomic Absorption Spectrophotometer (AAS). The ecological risk was evaluated through various indices, such as the Bioconcentration Factor (BCF), Translocation Factor (TF), Geoaccumulation Index (I_{geo}), Contamination Factor (Cf), and Pollution Load Index (PLI). The highest concentrations of heavy metals in sediment were detected at station 1 was Zn (15.486 mg.kg⁻¹). In the roots of *Cymodocea* was Zn (8.772 mg.kg⁻¹), while the highest concentration in leaves was Cu in *Thalassia* (10.541 mg.kg⁻¹). The ecological risk assessment revealed that $BCF < 1$ categorize an excluder, while $TF > 1$ for Pb and Zn indicate effective translocation from roots to leaves. Additionally, $I_{geo} < 0$ signify no contamination, $Cf < 1$ indicate low pollution levels, and $PLI < 0$ confirm a non-polluted status. In conclusion, the results show that the seagrass ecosystems at the study sites currently have low levels of heavy metal pollution and minimal ecological risk, suggesting they remain in a relatively safe condition.

Keywords: Accumulation, *Cymodocea*, Ecological risk assessment, Heavy metal, *Thalassia*

Introduction

Heavy metals are elements with low molecular weight that can lead to serious environmental issues when accumulated at certain concentrations (Wang et al., 2022a). Pollution from heavy metals due to tourism activities can create new problems, such as bioaccumulation and the transfer of pollutants through the food chain, a process known as biomagnification (Wang et al., 2022b). Biomagnification can occur in various organism components, including seagrass. Seagrass can absorb and accumulate pollutants like heavy metals in its tissues, which enables it to serve effectively as a bioindicator (Nel et al., 2023). Seagrass acts as a biomonitor and bioaccumulator of heavy metals because of its limited mobility, making it possible to estimate the concentrations and sources of heavy metals for analysis (Bonanno and Lo Giudice, 2010). Seagrass, which is found along the coast and in subtidal areas of Pahawang, is susceptible to direct

impacts from the presence of heavy metals (Noor et al., 2021).

Seagrass plays a crucial ecological role as an important habitat, providing abundant ecosystem services such as breeding grounds and food sources (Sun et al., 2020; Rozirwan et al., 2025a). In recent decades, seagrass has faced ecological challenges due to significant contamination, particularly from heavy metals entering coastal areas (Tu et al., 2023). Several studies have revealed the presence of heavy metals such as Pb, Cu, Zn, Cr, and Cd in various environments and organisms, including those in port areas, rivers, seagrass beds, and fish populations (Liu et al., 2019; Hosokawa et al., 2020; Fang et al., 2022; Souza-Araujo et al., 2022). Research on heavy metals in seagrass in other coastal regions has also shown that seagrass is vulnerable to accumulating all types of heavy metals, as observed in seagrass along the South China Sea (Zhang et al., 2024). Additionally, seagrass can accumulate heavy metals in its roots and leaves (Jeong et al., 2021). A study

conducted in the central Gulf of Gabes (Southeast Tunisia) indicated that seagrass not only serves as a bioindicator but can also be an effective bioremediation tool (El Zrelli *et al.*, 2023).

Seagrass and other plants can accumulate heavy metals in its roots and leaves (Jeong *et al.*, 2021; Rozirwan *et al.*, 2025b). Several previous studies have reported that heavy metal concentrations in seagrass roots and sediments are generally higher than in leaves, as observed in studies conducted in the Sydney estuaries, Australia, and Sicily, Italy (Birch *et al.*, 2018; Bonanno and Borg, 2018; Bonanno *et al.*, 2020). However, other studies have shown different results, such as research on seagrasses in the Southern Mediterranean Sea (El Zrelli *et al.*, 2023). From these findings, a hypothesis can be formulated that heavy metals are generally concentrated in the lower parts of seagrasses, such as roots and sediments, reflecting limited mobility and high sediment deposition rates. Therefore, measuring heavy metal concentrations in various seagrass organs is necessary to test this hypothesis.

The occurrence of heavy metals and their negative effects on the ecological balance of seagrass, as previously discussed, serve as a primary motivation for this research. The seagrass genus *Cymodocea* and *Thalassia* were chosen due to their dominance in the research sites. Previous studies have shown that these seagrasses have been investigated for heavy metal presence in their tissues, exhibiting relatively high levels of accumulation in various regions, including Italy, South China, India, and Florida (Bonanno *et al.*, 2020; Gopi *et al.*, 2020; Zhang *et al.*, 2021). Indonesia, research on heavy metal accumulation in *Thalassia* and *Cymodocea* has been conducted in the Seribu Islands, showing that these two seagrass species can accumulate heavy metals such as Pb, Cd, and Hg (Triyanto *et al.*, 2024). However, there has been no investigation using these seagrasses to monitor heavy metal pollution on Pahawang Island. Research on the presence of heavy metals on Pahawang Island has never been conducted, but research in the bay area adjacent to Pahawang Island (Pesawaran Chain Bay Coast) has been conducted, where the research results found heavy metal content above the quality standard in Cr of 415.86 mg.kg⁻¹, Cu of 163.312 mg.kg⁻¹, Fe of 3,339.89 mg.kg⁻¹ in marine organism (Fitrianiingsih and Widiastuti, 2021).

Pahawang Island in Lampung Province is a popular marine tourism destination with diverse ecosystems, including coral reefs, seagrass beds, and mangrove forests especially significant populations of *Cymodocea* and *Thalassia* (Mardani *et al.*, 2018; Novita *et al.*, 2022). According to the Central Bureau of Statistics of Pesawaran Regency,

tourist visits reached 448,008 in 2019 and decreased to approximately 165,342 in 2020 (BPS Statistic, 2023). The high boat activity, as the only means of transportation to the island, potentially contributes to heavy metal pollution in the (Saraswati and Rachmadiarti, 2021; Swain *et al.*, 2021; Rozirwan *et al.*, 2024).

Therefore, this study aims to provide valuable insights into the bioaccumulation patterns of heavy metals in the roots and leaves of seagrass, while also identifying the ecological risks involved. These insights can assist in developing management strategies for coastal areas and ecosystems in the region. The primary objective of this research is to analyze the ecological risks associated with the accumulation of heavy metals (Pb, Cu, Ni, and Zn) in the roots and leaves of *Cymodocea* and *Thalassia*, as well as in the sediments of Pahawang Island.

Materials and Methods

This research conducted in March 2024 off the coast of Pahawang Island in Lampung, Indonesia, at two distinct sampling locations. The sites were selected using purposive sampling within the seagrass ecosystem in Jeralangan and Cukuh Nyai (Figure 1). The first location, Jeralangan at coordinates 5°40'50.53"S and 105°13'51.15"E, is characterized by a high volume of tourism activities. The second location, Cukuh Nyai at coordinates 5°41'19.76"S and 105°13'9.28"E, has relatively low tourism activity. In addition, this location has a different residential area where the first location has a denser settlement than the second location. Tourism and residential activities can cause heavy metal pollutants to enter the waters (Purwiyanto, 2013; Purwiyanto *et al.*, 2020). Sampling was carried out during low tide, namely in morning and evening.

Two types of samples were collected at each station, consisting of seagrass and sediment. At Station One, samples of *Cymodocea* seagrass and sediment were collected, while Station Two involved the collection of *Thalassia* seagrass and sediment. The identification of seagrass species was conducted using a field guide for seagrass and dugong monitoring (Herandarudewi *et al.*, 2019). The collected of seagrass samples included both leaves and roots, gathered from shallow water habitats or intertidal zones through random sampling (Trevathan-Tackett *et al.*, 2015; Singh *et al.*, 2021). Sediment samples were taken using a grab sampler, with a total weight of 1 kg from the surface layer (0-10 cm). The collected samples were placed in plastic bags and stored in containers (Xiao *et al.*, 2022).



Figure 1. A map of seagrass sampling location

The results of this study indicate a pattern of heavy metal accumulation in seagrass at two locations with different anthropogenic pressures. While these findings provide valuable insights, it is important to note that the spatial coverage of this study is limited. Therefore, further investigation in other seagrass ecosystems is needed to determine the extent to which the observed patterns can be extrapolated to other areas with similar conditions. Additional studies with broader spatial coverage, encompassing various levels of environmental pressure and different oceanographic characteristics, will help strengthen the generalization of these findings.

Heavy metals analysis

The seagrass leaf and root samples, along with sediment, were thoroughly cleaned and subsequently dried. The dried samples were then ground to a fine powder and stored in sealed containers (Stewart *et al.*, 2021). Before the destruction, AAS calibration is carried out based on the Indonesian National Standard to ensure the accuracy and precision of heavy metal concentration measurements. The calibration process consists of the preparation of heavy metal standard solutions made by diluting certified stock solutions to various concentrations that cover the expected measurement range in the sample. These standards are used to create a calibration curve (Badan Standarisasi Nasional, 2004, 2009).

The sample digestion was carried out using the wet digestion method, where 0.5 g of powdered seagrass was mixed with 5 ml of a reagent in a ratio

of 5:2:1 (65% nitric acid: concentrated perchloric acid: sulfuric acid) on a hot plate for 10 min, along with an additional 5 ml of 2N hydrochloric acid, and heated on the hot plate for another 10 min. After reaching acid digestion, the sample was cooled and then filtered using Whatman filter paper (0.45 μm). The filtrate was then transferred into a 25 ml volumetric flask and diluted with deionized water up to the mark (FAO/SIDA, 1983; European Commission, 2011). For sediment samples, the wet digestion process involved adding 5 ml of concentrated HNO_3 to each sample, which included 50 ml of water sample and approximately 3 g of sediment sample. The sample was heated on a hot plate at a temperature of 105 $^\circ\text{C}$ - 120 $^\circ\text{C}$ until the volume was reduced to approximately 10 ml. After cooling, 5 ml of HNO_3 and 1-3 ml of perchloric acid were added, and the sample was reheated until white fumes appeared, followed by further heating for approximately 30 minutes. The sample was then filtered using quantitative filter paper with a pore size of 8.0 μm . The filtrate was transferred into a 100 ml volumetric flask and diluted with distilled water up to the mark (Badan Standarisasi Nasional, 2009; Gao *et al.*, 2021). The heavy metal analysis for each sample was performed in three replicates. The heavy metals analyzed in this study included Pb, Cu, Ni, and Zn.

The concentration of heavy metals in the samples was measured using an Atomic Absorption Spectrophotometer (AAS) SHIMADZU AA-7000. The wet digestion method is used because it is more effective in extracting heavy metals and preventing the loss of volatile elements such as Pb (Alfaro *et al.*, 2015). However, if not performed carefully, there is a possibility of contamination from reagents or

laboratory equipment that may affect the results. Therefore, quality assurance and quality control are carried out by ensuring the use of contaminant-free glassware, pure quality chemicals (p.a), and calibrated and verified atomic absorption spectrophotometers, operated by competent analysts. Quality control includes linearity of the calibration curve ($r^2 \geq 0.99$), blank analysis to control contamination with lead levels below the detection limit, and triplicate preparation to ensure accuracy with a difference in results of $\leq 20\%$ (Badan Standardisasi Nasional, 2004, 2009).

Ecological risk assessment

Ecological risk assessment can be conducted by calculating the bioconcentration factor (BCF), translocation factors (TF), geoaccumulation index (I_{geo}), contamination factor (Cf), and pollution load index (PLI) (Hakanson, 1980; Rozirwan et al., 2023).

Bioconcentration Factor (BCF)

Accumulation of heavy metals in the roots and leaves of seagrass from the surrounding sediment can be assessed using the Bioconcentration Factor (BCF) as described below (Steingraber et al., 2022).

$$BCF = \frac{\text{concentration of organs}}{\text{concentration of sediment}} \quad (1)$$

where: $BCF < 1$ is an excluder; The organism (usually a plant) does not accumulate the substance efficiently; $BCF = 1$ is an indicator; The organism takes up the substance at the same rate as it is found in the environment; $BCF > 1$ is a hyperaccumulator; The organism accumulates the substance at a much higher concentration than in the environment (Almahasheer, 2019).

Translocation Factors (TF)

The Translocation Factor (TF) is used to measure the movement of heavy metals from the roots to the leaves (Usman et al., 2019):

$$TF = \frac{BCF \text{ leaf}}{BCF \text{ of root}} \quad (2)$$

TF Value > 1 indicates that the sample functions as a phytoextractor, effectively translocating metals from the roots to the leaves, whereas TF value < 1 indicates that the sample functions as a phytostabilizer (Dinu et al., 2020).

Geoaccumulation Index (I_{geo})

The Geoaccumulation Index (I_{geo}) is used to evaluate the degree of heavy metal contamination

and classify pollution levels according to established criteria (Rozirwan et al., 2023).

$$I_{geo} = \text{Log}_2 \left(\frac{\text{concentration of sediment}}{1.5 \text{ background}} \right) \quad (3)$$

Note the I_{geo} value classification consists of not contaminated ($I_{geo} \leq 0$), not contaminated to moderately contaminated ($I_{geo} 0-1$), moderately contaminated ($I_{geo} 1-2$), moderate to highly contaminated ($I_{geo} 2-3$), highly contaminated ($I_{geo} 3-4$), highly contaminated to very high ($I_{geo} 4-5$), and highly contaminated ($I_{geo} \geq 5$).

Background concentration (Alfaro et al., 2015).

Contamination Factor (Cf)

The Contamination Factor (CF) value is used as a basis for determining the degree of heavy metal contamination (Rozirwan, Az-Zahrah et al., 2024).

$$CF = \frac{\text{concentration of sediment}}{\text{Background}} \quad (4)$$

where: contamination factor criteria according to (Shaheen et al., 2019); $Cf < 1$ = low level of contamination; $1 < Cf < 3$ = moderate level of contamination; $3 < Cf < 6$ = moderate level of contamination; $Cf > 6$ = very high level of contamination.

Pollution Load Index (PLI)

The Pollution Load Index (PLI) is used to evaluate the pollution quality (Mosa et al., 2022).

$$PLI = [Cf_1 \times Cf_2 \times Cf_3 \dots \times Cf_n]^{1/n} \quad (5)$$

where: pollution load index criteria: PLI 2 = not polluted to lightly polluted; PLI 2-4 = moderately polluted; PLI 4-6 = heavily polluted; PLI 6-8 = heavily polluted; PLI 8-10 = heavily polluted

Statistical analysis

The concentrations of heavy metals were analyzed using the Mann-Whitney test in SPSS, as the data were not normally distributed. This statistical test was applied to compare the concentrations of these metals between the two study sites based on the mean differences in the samples. The significance level (α) was set at 0.05 to determine statistical differences. If the 2-tailed significance value is greater than 0.05, it indicates no significant difference in the mean concentration between the two sites (H_0 is accepted). Conversely, if the 2-tailed significance value is less than 0.05, it suggests a significant difference in the mean concentrations (H_1 is rejected).

Result and Discussion

The seagrass samples were collected based on the dominant species found at the research site, specifically *Cymodocea* and *Thalassia* (Figure 2). *Cymodocea* was predominantly observed at Station One. This species is characterized by long, ribbon-like leaves that taper to a rounded tip, white to slightly yellowish rhizomes, and thin, brown fibrous roots. Both *Cymodocea* and *Thalassia* are commonly found in substrates ranging from muddy to sandy (Hartati et al., 2018; Thangaradjou and Bhatt, 2018), which explains their prevalence in the research area where the majority of locations feature such substrates. *Cymodocea* exhibits a strong tolerance to various environmental stresses and can thrive in diverse habitats, categorizing it as a species of least concern according to the IUCN (Terrados et al., 1998; Short et al., 2011). *Thalassia* is mainly found at Station Two and is recognized by its crescent-shaped leaves with neatly arranged margins, brown rhizomes, and thick, sheathed roots. This species allocates a greater proportion of its biomass to underground structures, such as roots and rhizomes, rather than to its leaves. This morphological variation can be seen as an adaptive response to environmental physical stressors, such as burial by sand to prevent

displacement during tidal fluctuations and storms (Westlake et al., 2022).

The root structure of *Thalassia* plays an essential role in the absorption of heavy metals in the marine environment. *Thalassia* has an extensive root and rhizome system, allowing greater contact with sediments containing heavy metals (Rosalina et al., 2022). Exposure to heavy metals can cause changes in the structure of root tissue, such as thickening or damage to the epidermis and endodermis in both seagrass roots. These changes can affect the ability of roots to absorb and translocate heavy metals (Rosalina et al., 2019). *Thalassia hemprichii* and *Cymodocea serrulata* leaves play an important role in the absorption of heavy metals from the water column. The broad and thin leaf structure increases the surface area for direct contact with water, facilitating the absorption of heavy metals (Nugraha, 2016). Additionally, the presence of aerenchyma tissue in the roots and leaves enables efficient gas exchange, which can influence the process of heavy metal absorption (Chedadi et al., 2024). According to Dilipan and Arulbalachandran (2022), both *Cymodocea* and *Thalassia* are promising candidates for bioremediation of contaminated sites and are frequently utilized as biological indicators of heavy metal accumulation.




Figure 2. Seagrass, (a) *Cymodocea*, (b) *Thalassia*

Heavy metals concentration in sediment and seagrass

The concentrations of heavy metals Pb, Cu, Zn, and Ni in the sediments, roots, and leaves of seagrasses at both stations are summarized in Figure 3. The results indicate that Station 1 (*Cymodocea*) had the highest concentrations of Pb and Zn in the sediments, roots, and leaves. Specifically, the maximum concentrations of Cu in sediments and roots were also recorded at Station 1 (*Cymodocea*), while the highest concentration in leaves was observed at Station 2 (*Thalassia*). The highest levels of Ni were found in the leaves and roots of *Cymodocea*. In contrast, the sediment at Station 2 exhibited the highest concentration of Ni. A comparative analysis of heavy metal concentrations across different seagrass species and locations is summarized in Figure 3. Based on the research results, the concentration of each sample between the two stations was analyzed using the Mann-Whitney test (as the data were not normally distributed) and showed a 2-tailed significance value of >0.05 , indicating no significant difference between the two stations.

Based on Figure 3, the concentrations of heavy metals Pb, Cu, Zn, and Ni in the sediments of this study are still below the threshold values established by ANZECC and ARMCANZ (2000) as referenced in (Ogundele *et al.*, 2015), and by the Gazzetta Ufficiale della Repubblica Italiana as noted in (Bonanno *et al.*, 2020), which are set at 50 mg.kg^{-1} , 65 mg.kg^{-1} , 50 mg.kg^{-1} , and 30 mg.kg^{-1} , respectively. Station 1 is characterized by a predominant presence of *Cymodocea*, which is directly influenced by tidal dynamics. Tidal-affected areas experience dynamic water movement, enabling heavy metals carried by the water to adhere to sediment particles (Maulana *et al.*, 2023). The concentrations of heavy metals Pb, Cu, Zn, and Ni in the sediments are higher compared to those in the roots and leaves, corroborating the findings of Khotimah *et al.* (2024) which indicate that sediment concentrations of heavy metals exceed those in the roots, implying that the roots may serve as a barrier preventing heavy metals from translocating to the upper plant structures. As noted by Flefel *et al.* (2020), sediments act as primary reservoirs for heavy metal absorption in aquatic environments. In addition, the higher concentration of heavy metals in the sediment at Station 1 compared to Station 2 can be assumed to be due to the finer sediment texture. Heavy metals tend to bind more effectively to sediment types with higher clay content or finer particle sizes (Borah *et al.*, 2018; Fitria *et al.*, 2023).

The concentrations of heavy metal Pb in the sediments, leaves, and roots of *Cymodocea* and

Thalassia differ between the two research locations (Figure 3). The higher concentration of Pb in *Cymodocea* at Station 1 may be attributed to the significant human activities in this area, including increased boat traffic and residential developments. Heavy metal Pb levels can be exacerbated by factors such as maritime traffic, fishing activities, paint on boats, and sedimentation (Irandoust *et al.*, 2021). Mitigation measures that can be implemented include strengthening regulations and waste treatment systems around coastal areas, as well as conducting long-term monitoring to identify pollution trends and their impacts on the seagrass ecosystem (Jeong *et al.*, 2021). Notably, the highest concentration of Pb is found in the leaves of *Cymodocea* when compared to its roots (figure 3). Similar findings were reported by El Zrelli *et al.* (2023), which indicated that the Pb concentration in the leaves of *Cymodocea* is greater than that in its roots. According to the WHO (1996) as cited in Ogundele *et al.* (2015), the concentration of Pb in *Cymodocea* and *Thalassia* remains below the quality standard of $2 \text{ mg.kg}^{-1}(\text{Pb})$, suggesting that these seagrasses are not currently subjected to contamination.

In this study, the highest concentration of the heavy metal Cu was observed in the leaves of *Thalassia* at Station 2. This concentration is still below the quality standard of 20 mg.kg^{-1} (Kabata-Pendias, 2011) indicating that the accumulation of Cu in seagrass does not pose any toxic effects. Copper is considered an essential metal, as it is required by organisms; however, at significantly high concentrations, Cu can be more toxic than non-essential metals such as Pb and Cd due to the active absorption mechanisms that plants employ. A similar study conducted on the leaves and roots of *Thalassia* in Xincun Bay, China, also reported the highest concentration in leaves at 11.2 mg.kg^{-1} (Zhang *et al.*, 2021). Previous research on various seagrass species has demonstrated that aboveground leaves are more efficient at accumulating Cu compared to belowground tissues (roots and rhizomes) (Lin *et al.*, 2018; Hu *et al.*, 2019). The concentration of heavy metal Cu in the leaves of *Thalassia* is higher than in *Cymodocea*. This may be due to *Thalassia* having broader leaf morphology and often exhibiting a slower growth rate compared to *Cymodocea*, allowing more heavy metals to accumulate in leaf tissues over a longer period (Nugraha, 2016). Additionally, different physiological mechanisms in each seagrass species may influence the uptake of heavy metals through the leaf surface (El Zrelli *et al.*, 2023). Previous studies on these two seagrass species, particularly in Indonesia, have shown that Cu concentrations in *Thalassia* leaves are higher than in *Cymodocea* (Bidayani *et al.*, 2017; Kholil *et al.*, 2019).

The concentration of Ni based on the research findings indicates that the highest level was found in the sediment, followed by the roots of *Cymodocea*, measuring 2.331 mg.kg⁻¹. This concentration is significantly lower than the legal limit allowed in marine waters, which is set at 20 mg.kg⁻¹. Furthermore, increases in Ni concentrations of 3 to 5 times are generally observed in environments impacted by human activities (Bonanno *et al.*, 2020). The Ni concentrations in this study are consistent with previous research that reported low Ni levels in seagrasses, such as 3.20–6.65 mg.kg⁻¹ in Italy (Bonanno *et al.*, 2017) and 1.8–3.1 mg.kg⁻¹ in South Africa (Nel *et al.*, 2023), particularly in locations with minimal human activities. This suggests that the study site is not yet contaminated with heavy metals, specifically Ni.

According to Figure 3, the highest concentration of Zn in seagrass was found in the leaves of *Cymodocea* at station 1. This concentration remains below the threshold established by the Australia and New Zealand Food Authority (ANZFA),

which is 14 mg.kg⁻¹ (Smitha *et al.*, 2010). Furthermore, the measured Zn levels are significantly lower than the concentrations that can cause phytotoxicity, which range from 500-1500 mg.kg⁻¹ (Elrashidi, 1990). Thus, the concentration of Zn detected in this study does not indicate any toxic effects. The results demonstrate that the concentrations of Zn in both the leaves and roots of the two seagrass species are higher than those of Pb and Cu (Figure 3). This suggests that Zn plays a crucial role in plant metabolism (Bonanno and Orlando-Bonaca, 2017). These findings are consistent with previous research, which reported that Zn concentrations are highest in the roots and leaves of various seagrass species, including *Cymodocea* and *Thalassia*, in locations such as Sicily, Italy, China, and India (Bonanno *et al.*, 2017; Hu *et al.*, 2019; Gopi *et al.*, 2020). After knowing the concentration of heavy metals in seagrass and sediment, the next step is to evaluate its ecological impact. Therefore, an ecological risk assessment is needed to assess the potential threat to seagrass ecosystems and organisms that depend on them.




Figure 3. Concentration of Pb, Cu, Ni and Zn in organs of *Cymodocea* and *Thalassia* (leaf and root mg.kg⁻¹), and associated sediments (mg.kg⁻¹)

Table 1. Comparison of heavy metal concentrations in *Cymodocea* and *Thalassia* with those from the open literatures

Location	Seagrass	Tissue	Heavy metals (mg.kg ⁻¹)	References
Sicily (Italy)	<i>Cymodocea</i>	Leaf	3.61–28.8 (Cu) 3.20–6.65 (Ni) 1.42–3.85 (Pb) 42.6–65.8 (Zn)	(Bonanno et al., 2017)
Sicily (Italy)	<i>Cymodocea</i>	Leaf	3.90–19.5 (Cu) 2.40–22.5 (Ni) 1.52–6.10 (Pb) 4.21–13.9 (Zn)	(Bonanno et al., 2020)
Sicily (Italy)	<i>Cymodocea</i>	Rhizome	10.6 (Cu) 1.15 (Ni) 0.73 (Pb) 21.7 (Zn)	(Bonanno and Borg, 2018)
		Root	8.83 (Cu) 4.79 (Ni) 4.81 (Pb) 38.7 (Zn)	
Florida, USA	<i>Thalassia</i>	Rhizome+root+leaf	20.23 (Pb) 9.55 (Cu) 86.68 (Zn)	(Smith et al., 2019)
Palk Bay, South Eastern India	<i>Cymodocea</i>	Rhizome+root+leaf	1.56 (Cd) 1.23 (Cd) 18.59 (Cu) 1.74 (Pb) 34.63 (Zn)	(Gopi et al., 2020)
Xincun Bay, China	<i>Thalassia</i>	Rhizome+root+leaf	11.2 (Cu) 4.05 (Ni) 2.1 (Pb) 39.1 (Zn)	(Zhang et al., 2021)
Southern Mediterranean Sea	<i>Cymodocea</i>	Rhizome	0.99 (Cu) 0.26 (Pb) 55 (Zn) 0.45 (Cd)	(El Zrelli et al., 2023)
		Root	3.51 (Cu) 3.76 (Pb) 31 (Zn) 1.33 (Cd)	
		Leaf	3.95 (Cu) 1.74 (Pb) 131.33 (Zn)	
			2.40 (Cd)	
Pahawang Island, Indonesia	<i>Cymodocea</i>	Root	0.76 (Pb) 0.54 (Cu) 8.77 (Zn) 2.33 (Ni)	This study
		Leaf	0.85 (Pb) 0.39 (Cu) 9.39 (Zn)	
	<i>Thalassia</i>	Root	1.97 (Ni) 0.47 (Pb) nd (Cu) 2.16 (Zn)	
		Leaf	0.69 (Ni) 0.65 (Pb) 10.54 (Cu) 6.77 (Zn) 1.36 (Ni)	

Note: nd (not detection)

Bioaccumulation and ecological risk assessments

Based on the research results presented in Table 2, the bioconcentration factors (BCF) for the

roots of *Cymodocea* and *Thalassia*, which accumulate heavy metals from the sediment biologically, were determined to be 0.0828 and 0.0502 for Pb, respectively. The BCF for Cu in *Cymodocea* is 0.5991,

while in *Thalassia* it is 0.0082. For Ni, the BCF values are 0.3140 for *Cymodocea* and 0.0894 for *Thalassia*, and for Zn, the BCF values are 0.5665 and 0.1670, respectively. The geoaccumulation index indicates an unpolluted status for Pb with values of -1.9696 and -1.9344, for Cu with values of -5.0658 and -4.5659, for Ni with values of -2.2167 and -2.1581, and for Zn with values of -2.6982 and -2.8687. The contamination factor (CF) suggests low contamination levels for Pb at 0.3830 and 0.3924, for Cu at 0.0448 and not detected (nd), for Ni at 0.3227 and 0.3361, and for Zn at 0.2311 and 0.2055. The pollution load index (PLI) was calculated, yielding results of 0.0013 and 0.0051, indicating that both stations are not polluted.

There are differences in the translocation factor (TF) values of metals (Table 3) between *Cymodocea* and *Thalassia*. In *Thalassia*, the TF for Cu indicates a translocation value of <1, suggesting minimal transfer of this metal to the above-ground biomass or leaves. This can also be attributed to the very low concentrations of Cu in the roots and leaves of the sampled seagrass at the study site. The metal with the highest TF value is Zn in *Thalassia*, followed by Ni and Pb. Similarly, *Cymodocea* shows the highest TF values for Zn and Pb, which are both >1. The differences in translocation values may arise from the biological characteristics of seagrasses, which have developed two distinct adaptive strategies to cope with the levels of contamination in their environment. The first strategy involves the storage of accumulated metals from the sediment in the root-rhizome system, while the second strategy entails the translocation of some contaminants from the root-rhizome to the above-ground parts of the seagrass, such as the leaves (El Zrelli *et al.*, 2023). According to the TF values presented in Table 2, *Thalassia* employs the second adaptive strategy for Zn, Ni, and Pb by translocating these contaminants to the leaves, whereas for Cu, it utilizes the first adaptive strategy. In contrast, *Cymodocea* adopts the second adaptive strategy for Pb and Zn. Previous research has also indicated that Zn undergoes significant translocation in *Cymodocea*, with a value >1 (Bonanno and Borg, 2018; Nel *et al.*, 2023).

The bioconcentration factor (BCF) of heavy metals from sediment to leaves and roots is summarized in Table 2. The BCF of zinc (Zn) in the leaves is higher than that of lead (Pb), nickel (Ni), and copper (Cu). This indicates that the accumulation level of Zn in the leaves is greater compared to the other metals, which may be influenced by several factors, including the adaptive strategies of seagrass in absorbing heavy metals and the chemical properties of the metals as well as the selectivity of the plants (Khotimah *et al.*, 2024). Zinc is an essential metal for seagrass, as it is utilized in

photosynthesis and growth (Kabata-Pendias and Pendias, 2000). This leads to higher Zn accumulation since it is more readily absorbed by both *Cymodocea* and *Thalassia*. According to Table 2, *Cymodocea* exhibits the highest BCF value for Zn. Similarly, high BCF values for Zn have also been found in the leaves of *Zostera noltei*, as well as in the roots and leaves of *Thalassia hemprichii* and in the roots and leaves of *Enhalus acoroides* (Li and Huang, 2012; Boutahar *et al.*, 2019; Jeong *et al.*, 2021). Compared to previous studies, particularly in Southeast Asia, such as research conducted in the Johor Strait, Malaysia, the accumulation of heavy metals in seagrasses and their ecological risks show a similar pattern in the accumulation of Pb and Cu, with varying levels of pollution depending on proximity to pollution sources (Sidi *et al.*, 2018). A comparison of heavy metal accumulation in seagrasses from different locations worldwide is presented in Table 1.

The values of the geoaccumulation index (I_{geo}) from this study are presented in Table 4, indicating that the sediments at both research locations are unpolluted ($I_{geo} < 0$). This suggests that the sources of heavy metal contamination at these sites are low, attributed to factors such as small residential areas, the absence of industrial and mining activities, and relatively low shipping traffic. Additionally, the physical and chemical conditions of the aquatic environment, including current patterns and tidal movements, also play a significant role (Siregar *et al.*, 2020). The contamination factor (CF) in this study indicates that each metal contributes to low or negligible contamination, as evidenced by CF values <1. These findings align with research conducted along the southern coast of Sumatra, Indonesia, which showed low CF values for each metal, except for lead (Pb), due to significant anthropogenic sources in the vicinity (Rozirwan *et al.*, 2023; Khotimah *et al.*, 2024).

The pollution status of the seagrass ecosystem at the research locations can be assessed using the pollution load index (PLI). The sediment PLI analyzed at the study sites is classified as unpolluted, with PLI values ranging from 0 to 2. The PLI values at both locations are not significantly different, indicating that both ecosystems are healthy and safe for living organisms. In contrast, other studies on seagrass ecosystems in the Southern Mediterranean Sea reported PLI values categorized as moderately contaminated and polluted, posing risks to the seagrass ecosystems, primarily due to the discharge of various industrial wastes, particularly phosphogypsum (El Zrelli *et al.*, 2023). Therefore, based on the analyses from this study, it can be concluded that all sampling locations exhibit low levels of metal pollution and ecological risk. The results of this study are limited to the surveyed location and may not fully reflect conditions in other areas.

Table 2. Values of the Bioconcentration factor (BCF)

	BCF			
	Pb	Cu	Ni	Zn
Seagrass roots				
St 1 (<i>Cymodoceae</i>)	0.0804	0.5991	0.3140	0.5665
St 2 (<i>Thalassia</i>)	0.0518	0.0082	0.0894	0.1570
Seagrass leaves				
St 1 (<i>Cymodoceae</i>)	0.0893	0.4436	0.2652	0.6064
St 2 (<i>Thalassia</i>)	0.0716	-8.3217	0.1766	0.4918

Table 3. The translocation factor (TF) from the roots to the leaves of seagrass

	TF			
	Pb	Cu	Ni	Zn
St 1 (<i>Cymodoceae</i>)	1.1112	0.7405	0.8445	1.0705
St 2 (<i>Thalassia</i>)	1.3834	-1011.6867	1.9764	3.1328

Table 4. Geoaccumulation Index (I_{geo}), Contamination Factor (Cf), and Pollution Load Index (PLI)

	I_{geo}				CF				PLI
	Pb	Cu	Ni	Zn	Pb	Cu	Ni	Zn	
St 1	-1.9696	-5.0658	-2.2167	-2.6982	0.3830	0.0448	0.3227	0.2311	0.0013
St 2	-1.9344	-4.5659	-2.1581	-2.8676	0.3924	nd	0.3361	0.2055	0.0051

The results of this study provide new insights into how tropical seagrasses, particularly *Cymodocea* and *Thalassia*, play a role in accumulating heavy metals in the marine environment. These findings contribute to understanding the distribution patterns of heavy metals in seagrass ecosystems, which can be used to monitor water quality. They also help identify seagrass species that are more effective as bioindicators of heavy metal pollution. Furthermore, this study provides a scientific basis for environmental management policies, such as the protection of seagrass beds in areas with high anthropogenic activity. However, this study is limited in its spatial coverage and does not account for broader temporal variability. Therefore, further research with a wider study area and long-term monitoring is necessary to gain a more comprehensive understanding of heavy metal accumulation patterns and to identify potential changes over time.

Conclusion

The highest concentrations of Pb, Cu, Ni, and Zn in seagrass were found in the leaves, while the sediments at Station 1 show the highest levels. Overall, the concentrations of heavy metals in the

roots and leaves of *Cymodocea* and *Thalassia* indicate relatively high levels; however, the ecological risk assessment shows that the bioaccumulation factor (BCF) is categorized as low. In contrast, the translocation factors (TF) for Pb and Zn demonstrate values greater than 1, indicating that the seagrass functions effectively in phytoremediation or translocating metals from the roots to the leaves. In summary, the ecological risk assessment of the research locations, which includes the geoaccumulation index (I_{geo}), contamination factor (CF), and pollution load index (PLI), indicates that the environment within the seagrass ecosystems at these sites has low levels of heavy metal pollution and ecological risk. The findings of this study are specific to the examined site and may not be entirely representative of conditions in other regions.

Acknowledgement

The research for this article was funded by the Ministry of Research, Technology, and Higher Education of the Republic of Indonesia (Number 090/E5/PG.02.00.PL/2024). The authors would like to express their gratitude to the Marine Bioecology Laboratory for providing the space and facilities during the course of the study.

References

- Alfaro, M.R., Montero, A., Ugarte, O.M., do Nascimento, C.W.A., de Aguiar Accioly, A.M., Biondi, C.M., & da Silva, Y.J.A.B. 2015. Background concentrations and reference values for heavy metals in soils of Cuba. *Environ. Monit. Assess.*, 187(1): <https://doi.org/10.1007/s10661-014-4198-3>
- Almahasheer, H. 2019. High levels of heavy metals in Western Arabian Gulf mangrove soils. *Mol. Biol. Rep.*, 46(2): 1585–1592. <https://doi.org/10.1007/S11033-019-04603-2/METRICS>
- Badan Standarisasi Nasional. 2004. SNI 6992.3:2004. Sediments – Chapter 3: lead (Pb) test method by acid destruction with atomic absorption spectrophotometer (AAS). Badan Standardisasi Nasional. 1-9.
- Badan Standarisasi Nasional. 2009. SNI 6989.6:2009. Chapter 6. Test method for copper (Cu) by atomic absorption spectrophotometry (AAS) in water and wastewater. In *Badan Standardisasi Nasional*. 1-9.
- Bidayani, E., Rosalina, D., & Utami, E. 2017. Heavy Metal Content of Lead (Pb) in Seagrass *Cymodocea serrulata* in Tin Mining Area of South Bangka Regency. *Maspari*, 9(2): 169–176.
- Birch, G.F., Cox, B.M., & Besley, C.H. 2018. The relationship between metal concentrations in seagrass (*Zostera capricorni*) tissue and ambient fine sediment in modified and near-pristine estuaries (Sydney estuaries, Australia). *Mar. Pollut. Bull.*, 128: 72–81. <https://doi.org/10.1016/j.marpolbul.2018.01.006>
- Bonanno, G., & Borg, J. A. 2018. Comparative analysis of trace element accumulation in seagrasses *Posidonia oceanica* and *Cymodocea nodosa*: Biomonitoring applications and legislative issues. *Mar. Pollut. Bull.*, 128: 24–31. <https://doi.org/10.1016/j.marpolbul.2018.01.013>
- Bonanno, G., Borg, J. A., & Di Martino, V. 2017. Levels of heavy metals in wetland and marine vascular plants and their biomonitoring potential: A comparative assessment. *Sci. Total Environ.*, 576: 796–806. <https://doi.org/10.1016/j.scitotenv.2016.10.171>
- Bonanno, G., & Lo Giudice, R. 2010. Heavy metal bioaccumulation by the organs of *Phragmites australis* (common reed) and their potential use as contamination indicators. *Ecol. Indic.*, 10(3): 639–645. <https://doi.org/10.1016/j.ecolind.2009.11.002>
- Bonanno, G., & Orlando-Bonaca, M. 2017. Trace elements in Mediterranean seagrasses: Accumulation, tolerance and biomonitoring. A review. *Mar. Pollut. Bull.*, 125(1–2): 8–18. <https://doi.org/10.1016/j.marpolbul.2017.10.078>
- Bonanno, G., Veneziano, V., Raccuia, S. A., & Orlando-Bonaca, M. 2020. Seagrass *Cymodocea nodosa* and seaweed *Ulva lactuca* as tools for trace element biomonitoring. A comparative study. *Mar. Pollut. Bull.*, 161: 111743. <https://doi.org/10.1016/j.marpolbul.2020.111743>
- Borah, R., Taki, K., Gogoi, A., Das, P., & Kumar, M. 2018. Contemporary distribution and impending mobility of arsenic, copper and zinc in a tropical (Brahmaputra) river bed sediments, Assam, India. *Ecotoxicol. Environ. Saf.*, 161(336): 769–776. <https://doi.org/10.1016/j.ecoenv.2018.06.038>
- Boutahar, L., Maanan, M., Bououarour, O., Richir, J., Pouzet, P., Gobert, S., Maanan, M., Zourarah, B., Benhoussa, A., & Bazairi, H. 2019. Biomonitoring environmental status in semi-enclosed coastal ecosystems using *Zostera noltei* meadows. *Ecol. Indic.*, 104: 776–793. <https://doi.org/10.1016/j.ecolind.2019.04.039>
- BPS Statistic. 2023. Badan Pusat Statistik. Number of Foreign Tourist Visits. Badan Pusat Statistik.
- Chedadi, M., Moussaoui, A. El, Bassouya, M., Jawhari, F.Z., Zoufri, I., Barnossi, A.E, Asmi, H. El, Ammari, M., Merzouki, M., & Bari, A. 2024. Capacity of an aquatic macrophyte, *Pistia stratiotes* L., for removing heavy metals from water in the Oued Fez River and their accumulation in its tissues. *Environ. Monit. Assess.*, 196(11): 1114. <https://doi.org/10.1007/S10661-024-13305-7/METRICS>
- Dinu, C., Vasile, G.G., Buleandra, M., Popa, D.E., Gheorghe, S., & Ungureanu, E.M. 2020. Translocation and accumulation of heavy metals in *Ocimum basilicum* L. plants grown in a mining-contaminated soil. *J. Soils Sediments*, 20(4): 2141–2154.
- El Zrelli, R.B., Yacoubi, L., Castet, S., Grégoire, M., Lin, Y.J., Attia, F., Ayranci, K., Baki, Z.A., Courjault-Radé, P., & Rabaoui, L.J. 2023. Compartmentation of trace metals in *Cymodocea nodosa* from a heavily polluted area (Central Gulf of Gabes; Southern Mediterranean Sea): Potential use of the seagrass as environmental monitoring and bioremediation tool. *Reg. Stud. Mar. Sci.*, 65: p. 103056. <https://doi.org/10.1016/j.rsma.2023.103056>

- Elrashidi, M. A. 1990. Inorganic Contaminants in the Vadose Zone. *J. Environ. Qual.*, 19(4): 783–784. <https://doi.org/10.2134/jeq1990.00472425001900040026x>
- European Commission. 2011. Commission Regulation (Eu). *Off. J. Eur. Union*, 2011(1131): 205–211.
- Fang, T., Wang, H., Liang, Y., Cui, K., Yang, K., Lu, W., Li, J., Zhao, X., Gao, N., Yu, Q., Li, H., & Jiang, H. 2022. Source tracing with cadmium isotope and risk assessment of heavy metals in sediment of an urban river, China. *Environ. Pollut.*, 305: 119325. <https://doi.org/10.1016/j.envpol.2022.119325>
- FAO/SIDA. 1983. Manual of Methods in Aquatic Environment Research. FAO, 62.
- Fitria, Y., Rozirwan, Fitriani, M., Nugroho, R.Y., Fauziyah, & Putri, W.A.E. 2023. Gastropods as bioindicators of heavy metal pollution in the Banyuasin estuary shrimp pond area, South Sumatra, Indonesia. *Acta Ecol. Sin.*, 43(6): 1129–1137. <https://doi.org/https://doi.org/10.1016/j.chnaes.2023.05.009>
- Fitrianiingsih, R., & Widiastuti, E. L. 2021. Heavy Metal Content of Cr, Cu, Fe and Mn in Several Biota on the Coast of Ratai Pesawaran Bay, Lampung. *Pros. Snail 2021 Semin. Nas. Ilmu Lingkungan. Tata Kelola Lingkungan. Untuk Mendukung Pembang. Berkelanjutan*, 3257(2003): 68–74.
- Flefel, H., Nokhrin, D., & Donnik, I. 2020. Determine heavy metals in water, aquatic plants, and sediment in water systems. *E3S Web Conf.*, 222: 02028. <https://doi.org/10.1051/e3sconf/202022202028>
- Gao, Y., Qiao, Y., Xu, Y., Zhu, L., & Feng, J. 2021. Assessment of the transfer of heavy metals in seawater, sediment, biota samples and determination the baseline tissue concentrations of metals in marine organisms. *Environ. Sci. Pollut. Res. Int.*, 28(22): 28764–28776. <https://doi.org/10.1007/S11356-021-12650-1>
- Gopi, S., Arulkumar, A., Ganeshkumar, A., Rajaram, R., Miranda, J.M., & Paramasivam, S. 2020. Heavy metals accumulation in seagrasses collected from Palk Bay, South-eastern India. *Mar. Pollut. Bull.*, 157: <https://doi.org/10.1016/j.marpolbul.2020.111305>
- Hakanson, L. 1980. An ecological risk index for aquatic pollution control. a sedimentological approach. *Water Res.*, 14(8): 975–1001. [https://doi.org/10.1016/0043-1354\(80\)90143-8](https://doi.org/10.1016/0043-1354(80)90143-8)
- Hartati, R., Zainuri, M., Ambariyanto, A., Widianingsih, W., Trianto, A., & Mahendrajaya, R.T. 2018. Similarity microalgal epiphyte composition on seagrass of *Enhalus acoroides* and *Thalassia hemprichii* from different waters. *IOP Conf. Ser. Earth Environ. Sci.*, 139(1): 012011. <https://doi.org/10.1088/1755-1315/139/1/012011>
- Herandarudewi, S.M., Kiswara, W., Irawan, A., Anggraeni, F., Juraij, Munandar, E., Sunudin, A., & Tania, C. 2019. Dugong and Seagrass Survey and Monitoring Guide. *ITB Press*, 43.
- Hosokawa, S., Naito, R., & Nakamura, Y. 2020. Spatial patterns of concentrations of Cu, Zn, Cd, and Pb in marine sediments from Japanese port areas. *Reg. Stud. Mar. Sci.*, 35: 101135.
- Hu, C., Yang, X., Gao, L., Zhang, P., Li, W., Dong, J., Li, C., & Zhang, X. 2019. Comparative analysis of heavy metal accumulation and bioindication in three seagrasses: Which species is more suitable as a bioindicator? *Sci. Total Environ.*, 669: 41–48. <https://doi.org/10.1016/j.scitotenv.2019.02.425>
- Irandoost, F., Agah, H., Rossi, L., Calizza, E., Careddu, G., & Costantini, M.L. 2021. Stable isotope ratios ($\delta^{13}\text{C}$ and $\delta^{15}\text{N}$) and heavy metal levels in macroalgae, sediment, and benthos from the northern parts of Persian Gulf and the Gulf of Oman. *Mar. Pollut. Bull.*, 163: 111909. <https://doi.org/10.1016/j.marpolbul.2020.111909>
- Jeong, H., Choi, J.Y., Choi, D.H., Noh, J.H., & Ra, K. 2021. Heavy metal pollution assessment in coastal sediments and bioaccumulation on seagrass (*Enhalus acoroides*) of Palau. *Mar. Pollut. Bull.*, 163: 111912. <https://doi.org/10.1016/j.marpolbul.2020.111912>
- Kabata-Pendias, A. 2011. Trace Elements in Soils and Plants, Fourth Edition. In *Taylor & Francis Group, Boca Raton, FL, USA*. <https://doi.org/10.1016/j.sajb.2012.03.008>
- Kabata-Pendias, A., & Pendias, H. 2000. Trace Elements in Soils and Plants: Third Edition. In *Trace Elements in Soils and Plants: Third Edition*. CRC Press. <https://doi.org/10.1201/9781420039900>
- Kholil, R., Eryati, R., & Sari, L. I. 2019. Heavy Metal Content of Pb (Lead) and Cu (Copper) in Seagrass (*Thalassia hemprichii*) in Melahing and Sapa Segajah Hamlets, Bontang, East Kalimantan. *J. Aquarime*, 6(2): 12–20.

- Khotimah, N.N., Rozirwan, Putri, W.A.E., Fauziyah, Aryawati, R., Isnaini, & Nugroho, R. Y. 2024. Bioaccumulation and Ecological Risk Assessment of Heavy Metal Contamination (Lead and Copper) Build Up in the Roots of *Avicennia alba* and *Excoecaria agallocha*. *J. Ecol. Eng.*, 25(5): 101–113.
- Li, L., & Huang, X. 2012. Three tropical seagrasses as potential bio-indicators to trace metals in Xincun Bay, Hainan Island, South China. *Chinese J. Oceanol. Limnol.*, 30(2): 212–224. <https://doi.org/10.1007/S00343-012-1092-0>/METRICS
- Lin, H., Sun, T., Adams, M. P., Zhou, Y., Zhang, X., Xu, S., & Gu, R. 2018. Seasonal dynamics of trace elements in sediment and seagrass tissues in the largest *Zostera japonica* habitat, the Yellow River Estuary, northern China. *Mar. Pollut. Bull.*, 134: 5–13. <https://doi.org/10.1016/j.marpolbul.2018.02.043>
- Liu, M., Chen, J., Sun, X., Hu, Z., & Fan, D. 2019. Accumulation and transformation of heavy metals in surface sediments from the Yangtze River estuary to the East China Sea shelf. *Environ. Pollut.*, 245: 111–121. <https://doi.org/10.1016/j.envpol.2018.10.128>
- Mardani, A., Purwanti, F., & Rudiyanti, S. 2018. Community-based ecotourism development strategy on Pahawang Island, Lampung Province. *Manag. Aquat. Resour. J.*, 6(1): 1–9. <https://doi.org/10.14710/marj.v6i1.19804>
- Maulana, F., Muhiddin, A.H., Lanuru, M., Samad, W., & Ukkas, M. 2023. Distribution of Bottom Sediment Before and After Reclamation At Center Point of Indonesia (CPI) Makassar City. *J. Ilmu Kelaut. Spermonde*, 9(1): 10–19. <https://doi.org/10.20956/jiks.v9i1.19929>
- Mosa, A., Selim, E.M.M., El-Kadi, S.M., Khedr, A.A., Elnaggar, A.A., Hefny, W.A., Abdelhamid, A.S., El Kenawy, A.M., El-Naggar, A., Wang, H., & Shaheen, S.M. 2022. Ecotoxicological assessment of toxic elements contamination in mangrove ecosystem along the Red Sea coast, Egypt. *Mar. Pollut. Bull.*, 176: 113446. <https://doi.org/10.1016/J.MARPOLBUL.2022.113446>
- Nel, M.A., Adams, J.B., Rubidge, G., & Human, L.R.D. 2023. Heavy metal compartmentalisation in salt marsh and seagrass of the urbanised Swartkops estuary, South Africa. *Mar. Pollut. Bull.*, 192: <https://doi.org/10.1016/j.marpolbul.2023.115007>
- Noor, N. M., Febriani, D., & Ali, M. 2021. Seagrass of *Enhalus Acoroides* as a Traditional Body Scrubs in Preventing Malarial Bites by Pahawang Island Community in Indonesia. *IOP Conf. Ser. Earth Environ. Sci.*, 1012(1): 012037. <https://doi.org/10.1088/17551315/1012/1/012037>
- Novita, S., Abidin, Z., & Kasymir, E. 2022. Economic valuation using the travel cost method for biodiversity park tourism in Mesuji Regency. *J. Ilmu-Ilmu Agribisnis*, 10(2): 217. <https://doi.org/10.23960/jiia.v10i2.6012>
- Nugraha, A.H. 2016. Physiological Response of Seagrass *Thalassia hemprichii* and *Cymodocea rotundata* to Anthropogenic Pressure in the Pari Island Cluster of the Seribu Islands. IPB Scientific Repository.
- Ogundele, D. T., Adio, A., & Oludele, O. E. 2015. Heavy Metal Concentrations in Plants and Soil along Heavy Traffic Roads in North Central Nigeria. *J. Environ. Anal. Toxicol.*, 5(6): 1-6.
- Purwiyanto, A. I. S., Suteja, Y., Ningrum, P. S., Putri, W. A. E., Agustriani, F., Cordova, M. R., & Koropitan, A. F. 2020. Concentration and adsorption of Pb and Cu in microplastics: case study in aquatic environment. *Mar. Pollut. Bull.*, 158: 111380.
- Rosalina, D., Herawati, E. Y., Musa, M., Sofarini, D., Amin, M., & Risjani, Y. 2019. Lead accumulation and its histological impact on *Cymodocea serrulata* seagrass in the laboratory. 48(4): 813–822. <https://doi.org/10.17576/JSM-2019-4804-13>
- Rosalina, D., Rombe, K. H., Jamil, K., & Surachmat, A. 2022. Analysis of heavy metals (Pb and Cd) in seagrasses *Thalassia hemprichii* and *Enhalus acoroides* from Pulau Sembilan, South Sulawesi Province, Indonesia. *Biodiversitas J. Biol. Divers.*, 23(4): 2130–2136. <https://doi.org/10.13057/BIODIV/D230448>
- Rozirwan, Adefta, A., Apri, R., Fauziyah, Putri, W. A. E., Melki, Iskandar, I., Novianti, E., Mustopa, A. Z., Fatimah, & Nugroho, R. Y. 2024a. The Ecological Health of Coral Reefs In The Waters of Enggano Island, The Outermost Island In The Eastern Indian Ocean. *Int. J. Conserv. Sci.*, 15(4): 1947–1958. <https://doi.org/10.36868/IJCS.2024.04>
- Rozirwan, Az-Zahrah, S. A. F., Khotimah, N. N., Nugroho, R. Y., Putri, W. A. E., Fauziyah, Melki, Agustriani, F., & Siregar, Y. I. 2024b. Ecological Risk Assessment of Heavy Metal Contamination in Water, Sediment, and Polychaeta (*Neoleanira Tetragona*) from Coastal Areas Affected by Aquaculture, Urban Rivers, and Ports in South Sumatra. *J. Ecol. Eng.*, 25(1): 303–319.

- Rozirwan, Khotimah, N. N., Putri, W. A. E., Fauziyah, Aryawati, R., Damiri, N., Isnaini, & Nigr. 2023. Environmental risk assessment of Pb, Cu, Zn, and Cd concentrations accumulated in selected mangrove roots and surrounding their sediment. *24(12)*: 6733–6742. <https://doi.org/10.13057/biodiv/d241236>
- Rozirwan, Khotimah, N. N., Putri, W. A. E., Fauziyah, Aryawati, R., Diansyah, G., & Nugroho, R. Y. 2025a. Biomarkers of heavy metals pollution in mangrove ecosystems: Comparative assessment in industrial impact and conservation zones. *Toxicol. Reports*, 14: 102011. <https://doi.org/10.1016/j.toxrep.2025.102011>
- Rozirwan, Nopriani, Fauziyah, Putri, W. A. E., Melki, Iskandar, I., Novianti, E., Mustopa, A. Z., Fatimah, & Nugroho, R. Y. 2025b. The Natural Calcium and Phosphorus Content in Two Fishbone of Commercial Species from Enggano Waters. *IOP Conf. Ser. Earth Environ. Sci.*, 1496(1): 12013. <https://doi.org/10.1088/1755-1315/1496/1/012013>
- Saraswati, A. R., & Rachmadiarti, F. 2021. Analysis of the heavy metal content of lead (Pb) in seaweed at Sendang Biru Beach, Malang. *LenteraBio Berk. Ilm. Biol.*, 10(1): 67–76. <https://doi.org/10.26740/lenterabio.v10n1.p67-76>
- Shaheen, S. M., Abdelrazek, M. A. S., Elthoth, M., Moghanm, F. S., Mohamed, R., Hamza, A., El-Habashi, N., Wang, J., & Rinklebe, J. 2019. Potentially toxic elements in saltmarsh sediments and common reed (*Phragmites australis*) of Burullus coastal lagoon at North Nile Delta, Egypt: A survey and risk assessment. *Sci. Total Environ.*, 649: 1237–1249. <https://doi.org/10.1016/J.scitotenv.2018.08.359>
- Short, F. T., Polidoro, B., Livingstone, S. R., Carpenter, K. E., Bandeira, S., Bujang, J. S., Calumpang, H. P., Carruthers, T. J. B., Coles, R. G., Dennison, W. C., Erftemeijer, P. L. A., Fortes, M. D., Freeman, A. S., Jagtap, T. G., Kamal, A. H. M., Kendrick, G. A., Judson Kenworthy, W., La Nafie, Y. A., Nasution, I. M., Zieman, J. C. 2011. Extinction risk assessment of the world's seagrass species. *Biol. Conserv.*, 144(7): 1961–1971. <https://doi.org/10.1016/J.BIOCON.2011.04.010>
- Sidi, N., Aris, A. Z., Mohamat Yusuff, F., Looi, L. J., & Mokhtar, N. F. 2018. Tape seagrass (*Enhalus acoroides*) as a bioindicator of trace metal contamination in Merambong shoal, Johor Strait, Malaysia. *Mar. Pollut. Bull.*, 126: 113–118. <https://doi.org/10.1016/j.marpolbul.2017.10.041>
- Singh, S., Lal, M. M., Southgate, P. C., Wairiu, M., & Singh, A. 2021. Trace metal content in sediment cores and seagrass biomass from a tropical southwest Pacific Island. *Mar. Pollut. Bull.*, 171: 112745. <https://doi.org/10.1016/j.marpolbul.2021.112745>
- Siregar, A. S., Sulisty, I., & Prayogo, N. A. 2020. Heavy metal contamination in water, sediments and *Planiliza subviridis* tissue in the Donan River, Indonesia. *J. Water L. Dev.*, 45: 157–164. <https://doi.org/10.24425/jwld.2020.133057>
- Smith, E.M., Giarikos, D., Daniels, A., & Hirons, A.C. 2019. Heavy Metal Accumulation in Seagrasses in Southeastern Florida. *J. Mar. Biol. Oceanogr.*, 8(2):1-8.
- Smitha, J. L., Summers, G., & Wong, R. 2010. Nutrient and heavy metal content of edible seaweeds in New Zealand. *New Zeal. J. Crop Hortic. Sci.*, 38(1): 19–28. <https://doi.org/10.1080/01140671003619290>
- Souza-Araujo, J. de, Hussey, N. E., Hauser-Davis, R. A., Rosa, A. H., Lima, M. de O., & Giarrizzo, T. 2022. Human risk assessment of toxic elements (As, Cd, Hg, Pb) in marine fish from the Amazon. *Chemosphere*, 301: 134575.
- Steingraber, L. F., Ludolphy, C., Metz, J., Kierdorf, H., & Kierdorf, U. 2022. Uptake of lead and zinc from soil by blackberry plants (*Rubus fruticosus* L. agg.) and translocation from roots to leaves. *Environ. Adv.*, 9: 100313. <https://dx.doi.org/10.1016/j.envadv.2022.100313>
- Stewart, B. D., Jenkins, S. R., Boig, C., Sinfield, C., Kennington, K., Brand, A. R., Lart, W., & Kröger, R. 2021. Metal pollution as a potential threat to shell strength and survival in marine bivalves. *Sci. Total Environ.*, 755: 143019. <https://doi.org/10.1016/j.scitotenv.2020.143019>
- Sun, Y., Song, Z., Zhang, H., Liu, P., & Hu, X. 2020. Seagrass vegetation affect the vertical organization of microbial communities in sediment. *Mar. Environ. Res.*, 162: 105174.
- Swain, S., Pattanayak, A.A., Sahu, B.K., Satapathy, D. R., & Panda, C.R. 2021. Time-series monitoring and ecological risk assessment of heavy metal pollution in Mahanadi estuary, east coast of India. *Reg. Stud. Mar. Sci.*, 47: 101923. <https://doi.org/10.1016/J.RSMA.2021.101923>
- Terrados, J., Duarte, C. M., Fortes, M. D., Borum, J., Agawin, N. S. R., Bach, S., Thampanya, U., Kamp-Nielsen, L., Kenworthy, W. J., Geertz-Hansen, O., & Vermaat, J. 1998. Changes in Community

- Structure and Biomass of Seagrass Communities along Gradients of Siltation in SE Asia. *Estuar. Coast. Shelf Sci.*, 46(5): 757–768. <https://doi.org/10.1006/ECSS.1997.0304>
- Thangaradjou, T., & Bhatt, J. R. 2018. Status of seagrass ecosystems in India. *Ocean Coast. Manag.*, 159: 7–15. <https://doi.org/10.1016/j.ocecoaman.2017.11.025>
- Trevathan-Tackett, S. M., Lane, A. L., Bishop, N., & Ross, C. 2015. Metabolites derived from the tropical seagrass *Thalassia testudinum* are bioactive against pathogenic *Labyrinthula* sp. *Aquat. Bot.*, 122: 1–8. <https://doi.org/10.1016/j.aquabot.2014.12.005>
- Triyanto, H. C., Azizah, R., Nuraini, T., & Haryanti, D. 2024. Accumulation of Heavy Metal Lead (Pb) in Water, Sediment, and Seagrass (*Thalassia hemprichii*) on Kelapa Dua Island, Seribu Islands. *J. Mar. Res.*, 13(1): 37–44. <https://doi.org/10.14710/JMR.V13I1.35263>
- Tu, Y. J., Luo, P. C., Li, Y. L., Liu, J., Sun, T. T., Li, G. J., & Duan, Y. P. 2023. Seasonal heavy metal speciation in sediment and source tracking via Cu isotopic composition in Huangpu River, Shanghai, China. *Ecotoxicol. Environ. Saf.*, 260(100): 115068.
- Usman, K., Al-Ghouthi, M. A., & Abu-Dieyeh, M. H. 2019. The assessment of cadmium, chromium, copper, and nickel tolerance and bioaccumulation by shrub plant *Tetraena qataranse*. *Sci. Rep.*, 9(1): 1–11.
- Wang, C., Ju, J., Zhang, H., Liu, P., Zheng, Q., & Hu, X. 2022. Disclosing the ecological implications of heavy metal disturbance on the microbial N-transformation process in the ocean tidal flushing urban estuary. *Ecol. Indic.*, 144: 109504. <https://doi.org/10.1016/J.ECOLIND.2022.109504>
- Wang, Z., Lin, K., & Liu, X. 2022. Distribution and pollution risk assessment of heavy metals in the surface sediment of the intertidal zones of the Yellow River Estuary, China. *Mar. Pollut. Bull.*, 174: 113286. <https://doi.org/10.1016/J.marpolbul.2021.113286>
- Westlake, E. L., Keesing, J. K., Hardiman, L. K., Tonks, M., & Olsen, Y. 2022. Growth, biomass and productivity of the seagrass *Thalassia hemprichii* at Ashmore Reef, Australia. *Aquat. Bot.*, 183: 1–5. <https://doi.org/https://doi.org/10.1016/j.aquabot.2022.103557>
- Xiao, J., Wang, Y., Fu, S., Yang, S., Liu, G., Luo, X., & Zhang, C. 2022. Distribution of heavy metals and the exploration of potential indicators and hyperaccumulators in Jiang'an River, Chengdu, PR China. *Ecol. Indic.*, 145: 109665. <https://doi.org/10.1016/j.ecolind.2022.109665>
- Zhang, L., Ni, Z., Cui, L., Li, J., He, J., Jiang, Z., & Huang, X. 2021. Heavy metal accumulation and ecological risk on four seagrass species in South China. *Mar. Pollut. Bull.*, 173: 113153. <https://doi.org/10.1016/j.marpolbul.2021.113153>
- Zhang, L., Wu, Y., Jiang, Z., Ren, Y., Li, J., Lin, J., Ni, Z., & Huang, X. 2024. Identification of anthropogenic source of Pb and Cd within two tropical seagrass species in South China: Insight from Pb and Cd isotopes. *Ecotoxicol. Environ. Saf.*, 270: 115917.