

# dr Ramzi URINE ALBUMIN CREATININE RATIO AMONG DIABETIC (1) - Copy

*by* Dr. Ramzi, Spm

---

**Submission date:** 04-Dec-2018 10:38AM (UTC+0700)

**Submission ID:** 1050103909

**File name:** Ramzi\_URINE\_ALBUMIN\_CREATININE\_RATIO\_AMONG\_DIABETIC\_1\_-\_Copy.pdf (525.81K)

**Word count:** 2694

**Character count:** 14751

## URINE ALBUMIN CREATININE RATIO AMONG DIABETIC RETINOPATHY PATIENT WITH AND WITHOUT DIABETIC MACULAR EDEMA IN MOH. HOESIN HOSPITAL PALEMBANG

Rina Astuti, AK Ansyori, Ramzi Amin

Department of Ophthalmology, Universitas Sriwijaya, Palembang, Indonesia

### ABSTRACT

**Introduction:** Diabetic Macular Edema is a diffuse thickening in macula which can be found in various severity of Diabetic Retinopathy. There are issue about relationship between Diabetic Macular Edema and albuminuria caused by chronic renal failure. The aim of this study is to know and compare urine albumin creatinine ratio among Diabetic Retinopathy patients with and without Diabetic Macular Edema in Moh. Hoesin Hospital.

**Methods:** Cross sectional study with 25 sample was conducted. Diabetic Retinopathy and Diabetic Macular Edema was classified base on Early Treatment Of Diabetic Retinopathy Study (ETDRS) criteria. T-test, odd ratio and multiple logistic regression analysis was used to analysed sociodemography characteristic (age and gender), clinical characteristic (duration of DM, hipertension, treatment, body mass indeks and antioksidan consumption), ophthalmology characteristic (visus, anterior segment anomaly and posterior segment/severity of Diabetic Retinopathy), laboratory characteristic (HbA1c, ureum, creatinine, urine albumin creatinine ratio, and lipid profile).

**Result:** Urine albumin creatinine ratio mean ( $2146.77 \pm 3796.19$ ) in Diabetic Macular Edema and ( $49.0 \pm 45.35$ ) in non-Diabetic Macular Edema; cutoff point 62.00 mg/dL. Odd ratio adjusted for urine albumin creatinine ratio = 18,8. In this research, risk factors which has significantly were urine albumin creatinine ratio ( $p=0.047$ ) and High-Density Lipoprotein/HDL ( $p=0.028$ ) with odd ratio 8.571 and 6.67 respectively. Urine albumin creatinine ratio showed significantly high Mann whitney analysis 0.02 ( $p<0.005$ ).

**Conclusion:** Urine albumin creatinine ratio in Diabetic Retinopathy with Diabetic Macular Edema was higher than without Diabetic Macular Edema. Urine albumin creatinine ratio and High Density Lipoprotein (HDL) are the two important risk factors associated with Diabetic Macular Edema.

**Keywords:** Risk factor, Urine albumin creatinine ratio, Diabetic Macular Edema, Diabetic Retinopathy

**Cite This Article:** ASTUTI, Rina; ANSYORI, AK; AMIN, Ramzi. Urine Albumin Creatinine Ratio Among Diabetic Retinopathy Patient With And Without Diabetic Macular Edema In Moh. Hoesin Hospital Palembang. International Journal of Retina, [S.l.], v. 1, n. 1, july 2018. ISSN 2614-8536. Available at: <<https://www.ijretina.com/index.php/ijretina/article/view/32>>.

### INTRODUCTION

Diabetic retinopathy is a chronic progressive the leading cause of new blindness in adults 20- 70 years of age caused by long hiperglicemic status. Among Diabetic retinopathy patients, Diabetic Macular edema can be found. Diabetic Macular Edema is a diffuse or focal thickening in macula. In addition to visual loss, the negative impact of Diabetic Macular Edema especially on patients health-related quality of life (QoL). Base on

Banu Turgut Ozturk, patient with Diabetic Macular Edema consume significantly more healthcare resources, incur higher costs, and have low work efficiency compared to diabetic patients without retinal complications.<sup>1-6</sup>

Diabetic Macular Edema can accompany every stage of Diabetic Retinopathy. The incidence is 3% for mild non proliferative Diabetic Retinopathy, 38% for moderate and severe non proliferative Diabetic Retinopathy,

\*Correspondence to:

Rina Astuti,  
Department of Ophthalmology,  
Universitas Sriwijaya  
[rinabertawan@gmail.com](mailto:rinabertawan@gmail.com)

and 71% for proliferative Diabetic Retinopathy. About one diabetic patient in four can be expected to develop Diabetic Macular Edema in a lifetime. Once a patient's medical management has been optimized, ocular therapies should be considered to prevent progression vision loss and optimize the patient's vision.<sup>1,2,7</sup>

The major microvascular complications, retinopathy and nephropathy are the more important causes of blindness and end-stage renal disease. Chronic Kidney Disease and Diabetic Retinopathy are reflected in the clinical manifestations of similar microvascular lesions in the glomerular and retinal vessels.<sup>3,7</sup>

Many study found relationship between urine albumin creatinine ratio with diabetic retinopathy. High urine albumin creatinine ratio shows endothelial dysfunction and may influence microvasculature of the kidney and retina. Previous studies have shown that urine albumin creatinine ratio not only is important clinical marker for chronic kidney disease but also is closely associated to progression of diabetic retinopathy.<sup>8-12</sup>

The aim of this research is to know and compare urine albumin creatinine ratio among Diabetic Retinopathy patients with and without Diabetic Macular Edema in Moh. Hoesin Hospital.

## METHODS

Cross sectional study with 25 subject diabetic retinopathy was conducted in polyclinic/ward ophthalmology and internist of RSUP Moh. Hoesin

Palembang on October 2016 until October 2017. Inclusion criteria : subject was diagnosed diabetic retinopathy with and without diabetic macular edema and willing to follow this research. Exclusion criteria : high intraocular pressure, infection in eye and infection urinary tract or nephrolithiasis/ureterolithiasis. Diabetic Retinopathy and Diabetic Macular Edema was classified base on Early Treatment Of Diabetic Retinopathy Study (ETDRS) criteria. Subject was interviewed and examined (body mass, height, blood pressure, ophthalmology and blood/urine withdrawal).

## RESULT

In interim analysis with 25 subject, urine albumin creatinine ratio was significantly higher in patient diabetic retinopathy with diabetic macular edema just than without diabetic macular edema. So the research can be terminated. Risk factor of Diabetic macular edema was divided in many characteristic : sociodemography characteristic (age and gender), clinical characteristic (duration of DM, hipertension, treatment, body mass indeks and antioksidan consumption), ophthalmology characteristic (visus, anterior segment anomaly and posterior segment/ severity of Diabetic Retinopathy), laboratory characteristic (HbA1c, ureum, creatinine, urine albumin creatinine ratio, and lipid profile). There were no significantly between diabetic macular edema and sociodemography characteristic.

Table 1. Sociodemography Characteristic

Sociodemography	N	%	Average
<b>Age</b>			
<40 year	1	4	
40-49 year	8	32	
50-59 year	10	40	53.80 ± 8.25
≥60 year	6	24	
Total	25	100	
<b>Gender</b>			
Male	10	40	
Female	15	60	
Total	25	100	
<b>Occupation</b>			
Entrepreneur	10	40.0	
Civil servant	5	20.0	
Jobless	9	36.0	
Pension	1	4.0	
Total	25	100	
<b>Education</b>			
Elementary school	6	24.0	
Secondary school	5	20.0	

Senior high school	3	12.0
Collage	11	44.0
Total	25	100

In this research, there were 13 subject with diabetic macular edema and 12 without diabetic macular edema. There were no significantly between diabetic macular edema and clinical characteristic.

Table 2. **Clinical Characteristic**

Clinical	DME Positive		DME Negative		Total	P
	N	%	N	%		
	<b>Hipertension Complication</b>					
Yes	6	60	4	40	10	*0,513
No	7	46,67	8	53.3	15	
<b>Treatment</b>						
OHO	4	36.4	7	63.6	11	*0,165
Injection	9	64.3	5	35.7	14	
<b>BMI</b>						
Underweight	0	0	0	0	0	*0,552
Normoweight	5	62.5	3	37.5	8	
Overweight	3	50.0	3	50.0	6	
Obesity	5	45.5	6	54.5	11	
<b>Antioxdant Consumption</b>						
Yes	9	50	9	50	18	*0,748
No	4	57,1	3	42,9	7	

In this research, risk factors which has significant odd ratio were urine albumin creatinine ratio and High Density Lipoprotein (HDL) 8.571 and 6.67 respectively. Urine albumin creatinine ratio showed significantly high Mann whitney analysis 0.02 ( $p < 0.005$ ). But, there were no

relationship between ophthalmology characteristic and in laboratory characteristic, only urine albumin creatinine ratio ( $p = 0,047$ ) and High Density lipoprotein/HDL ( $p = 0,028$ ) were significantly with odd ratio 8.571 and 6.67.

Table 3. **Laboratory Characteristic**

Laboratory Characteristic	DME Positive		DME Negative		p	OR
	N	%	N	%		
	<b>HbA1C</b>					
Increased	11	61.1	7	38.9	18	
Normal	2	28.6	5	71.4	7	0.144
<b>Kreatinin</b>						
Increased /Critical	9	60.0	6	40.0	15	
Normal	4	40.0	6	60.0	10	0.284
<b>Ureum</b>						
Increased	7	77.8	2	22.2	9	
Normal	6	37.5	10	62.5	16	0.063+

Urine Albumin Creatinine Ratio						
Increased	12	63,2	7	36,8	19	8,57
Normal	1	16,7	5	83,3	6	0.047
Total Lipid						
Increased	9	64,3	5	35,7	14	
Normal	4	36,4	7	63,6	11	0.165
LDL						
Increased	10	50	10	50	20	
Normal	3	60	2	40	5	*1.00
Triglyceride						
Increased	9	69,2	4	30,8	13	
Normal	4	33,3	8	66,7	12	*0.073
HDL						
Decreased	10	71,4	4	28,6	14	6,67
Normal	3	27,3	8	72,7	11	0.028
Dislipidemia						
Yes	13	54,2	11	45,8	24	0.480
No	0	0,0	1	100	1	

#### Differentiation between Urine Albumin Creatinine Ratio among Diabetic Retinopathy Patients with and without Diabetic Macular Edema

Base on normality test, the data of diabetic macular edema on urine albumin creatinine ratio distribution were abnormal so writer use independent sampels Mann whitney test.

Table 4. Differentiation Urine Albumin Creatinine Rasio among Diabetic Retinopathy with and Without Diabetic Macular Edema

	Mean ± SD	Median (Min – Max)	p*
Positive DME	2146.77 ± 3796.19	356 (38 – 13628)	0.002
Negative DME	49.0 ± 45.35	38.5 (5 – 169)	

\*Mann-Whitney test

Urine albumin creatinine ratio mean (2146.77 ± 3796.19) in Diabetic Macular Edema and (49.0 ± 45.35) in non Diabetic Macular Edema; independent sampels Mann whitney test highly significant ( $p=0,02$ ); cutoff point 62.00 mg/dL. Sensitivity 76.9% and specificity 75.0%, predictive positif value 76.9%, predictive negaitif value 75.0%. Odd ratio adjusted for urine albumin creatinine ratio = 18,8.

Multivariat analytic show signifantly among diabetic macular edema with urine albumin creatinine and High Density lipoprotein. The formula :

$Y$  (Diabetic Macular Edema) = -0.696 + 2.934 urine albumin creatinine - 0,64 HDL.

Explanation :

- Urine albumin creatinine ratio :
  - 0 if the value under 62 mg/dL
  - 1 if the value above 62 mg/dL
- High Density Lipoprotein in mg/dL
- Odd ratio adjusted urine albumin creatinine ratio = 18,8, means patient diabetic retinopathy with urine albumin creatinine ratio > 62 mg/dL has

probability dibetic macular edema 18,8 times than they are with urine albumin creatinine ratio < 62 mg/dL.

#### DISCUSSION

Many research had been done to connect diabetic macular edema and chronic kidney disease. According to Kang et al., in patients with serous-type diabetic macular edema, albumin and other body fluids leak into the subretinal space through an external limiting membrane because of destruction of the inner blood retinal barrier. Nam Kyun Koo results show a significantly higher frequency of serous-type macular edema in patients with albuminuria. Despite a lack of statistical significance, serous-type macular edema usually occurred in patients with a lower level of serum albumin. These results indicate that serous-type macular edema might have a relationship with the excretion of albumin into the urine due to decreased renal function. It is also presumed that the excretion of albumin and the decreased concentration

of serum albumin cause an overload of body fluids and decreased intravascular osmotic pressure, which leads to the occurrence of serous- type macular edema.<sup>13</sup>

Akihiro Ishibawa et al (2013) about relationship among diabetic macular edema and chronic kidney disease in patients with type 2 diabetes shows serous retinal detachment developed 28,2% of the chronic kidney disease (-) group and 63,6% of the chronic kidney disease (+) group, a difference that reached significance ( $p=0.015$ ).<sup>14</sup>

Gagan Sathya Prakash (2016) did research which related to Clinically Significant Macular Edema (CSME) in Diabetes Melitus type 2. In this study, the incidence of microalbuminuria is significantly associated with CSME. 81,2% patients with CSME had macroalbuminuria and 12,9% patients with CSME had microalbuminuria.<sup>15</sup>

Ong ming Jew et al (2012) the urea and creatinine level were significantly higher in the CSME group. The urea level in the CSME and non-CSME ( $6.530\pm 3.171$ )mmol/L and ( $5.078\pm 1.747$ ) mmol/L. Whereas the creatinine level were ( $125.00\pm 86.02$ )mmol/L and ( $90.94\pm 36.76$ ) mmol/L respectively. The albumin level in the CSME and non-CSME were ( $40.24\pm 4.47$ ) mmol/L and ( $44.12\pm 2.32$ ) mmol/L respectively.<sup>16</sup>

In this research, urine albumin creatinine level higher in diabetic retinopathy with diabetic macular edema than without diabetic macular edema. And it is showed chronic kidney disease significantly higher with diabetic macular edema. Hypoalbumin, caused by renal leakage. Hypoalbumin effect osmotic pressure (decrease) in plasma and this factor cause macular edema. Decrease osmotic pressure cause the liquid extravasation leave the vascular to fill around tissue.

Many studies had been done to identify the risk factors or association of diabetic macular edema. However, the association varied from one study to another. The variation in the result may be due among other things to epidemiological differences. In this research, urine albumin creatinine and High Density Lipoprotein/HDL significantly high with diabetic macular edema.

In this research, there were limitation : we didn't do Optical Coherence Tomography (OCT) to know macular thickness quantitatively and albumin serum to know hypoalbuminemia which caused edema generalisation. Besides, only 25 subject participated so we suggested to add the research subject.

## CONCLUSION

Urine albumin creatinine ratio in Diabetic Retinopathy with Diabetic Macular Edema was higher than without Diabetic Macular Edema. Urine albumin creatinine ratio and High Density Lipoprotein (HDL) are the two important risk factors associated with Diabetic Macular Edema.

## REFERENCES

1. Retinal Vascular Disease : Diabetic Retinopathy in Retina and Vitreous American Academy of Ophthalmology. 2014-2015. Page 89-111.
2. Yanuzzi, Lawrence Y. Retinal vascular disease in The retinal atlas. Elsevier. 2010. Page 441-451.
3. Wessler, Michael J. Microvascular and macrovascular complications of diabetes. Clinical diabetes volume 26 number 2. 2008. Page 75-82.
4. Jousen, Antonia M et al. Pathology, clinical course and treatment of retinal vascular disease in Retinal vascular disease. Springer, Jerman. 2007. Page 291-390.
5. Sitompul, Ratna. Retinopati Diabetik. Jurnal Indonesia Medical Association, Volume: 61, Nomor: 8, Agustus 2011. Halaman 337-341.
6. Ozturk, Banu Turgut. Who gets Diabetic Macular Edema; When & why? Pathogenesis and risk factor ? Pathogenesis and risk factors. European Medical Journal October 2014. Page 105-111.
7. Mathew, Christopher et al. Article Review Updates in The Management of Diabetic Macular Edema. Journal of Diabetes Research Volume 2015, Article ID 794036, 8 pages. Hindawi Publishing Corporation. <http://dx.doi.org/10.1155/2015/794036>
8. Wessler, Michael J. Microvascular and macrovascular complications of diabetes. Clinical diabetes volume 26 number 2. 2008. Page 77-82.
9. Poncelas, Antonio R, et al. Chronic kidney disease and diabetic retinopathy in patients with type 2 diabetes. Journal of Diabetes Research Volume 2016, Article ID 149448 February, 17 2016. Page 1-10.
10. Natalie, et al. Retinopathy and clinical outcomes in patients with type 2 diabetes mellitus, chronic kidney disease, and anemia. British medical journal open diabetes research and care 2014; 2:e000011.DOI:10.1136/bmjdr-2013-000011.
11. Nagaoka, Taiji and Yoshida, Okitoshi. Relationship between retinal blood flow and renal function in DM type 2 patients and chronic kidney disease. Journal of diabetes care volume 36 2013. Page 957-961
12. Haneda, Masakazu et al. A new classification of diabetic nephropathy 2014 : a report from joint committee on diabetic nephropathy. Journal of diabetes investigation volume 6 2015. Page 242-246
13. Koo, Nam Kyun et al. Relationship between the morphology of diabetic macular edema and renal dysfunction in diabetes. Korean journal of ophthalmology 2013 volume 27 (4) page 98-102
14. Ishibazawa, Akihiro et al. Association between diabetic macular edema and chronic kidney disease in patients with type 2 diabetes. Investigative ophthalmology and visual science, Volume 54, issue 15 June 2013.

15. Prakash, Gagan Satya and Kothari, Minal. Risk Factors Associated with Clinically Significant Macular Edema in Patients with Type 2 Diabetes Mellitus. International Journal of Scientific Study Vol 3 Issue 12 March 2016.
16. Jew, Ong Ming, etc. Risk Factors for Clinically Significant Macular Edema in A Multi-Ethnic Population with Type 2 Diabetes. Internationaal Journal Ophthalmology, Volume 5 No 4 August 2012.



This work licensed under Creative Commons Attribution

# dr Ramzi URINE ALBUMIN CREATININE RATIO AMONG DIABETIC (1) - Copy

## ORIGINALITY REPORT

15%

SIMILARITY INDEX

%

INTERNET SOURCES

13%

PUBLICATIONS

6%

STUDENT PAPERS

## PRIMARY SOURCES

- 1** Nam Kyun Koo, Hyun Cheol Jin, Kwang Soo Kim, Yu Cheol Kim. "Relationship between the Morphology of Diabetic Macular Edema and Renal Dysfunction in Diabetes", Korean Journal of Ophthalmology, 2013 **5%**

Publication
- 2** Rodríguez-Poncelas, Antonio, Xavier Mundet-Tudurí, Sonia Miravet-Jiménez, Aina Casellas, Joan F. Barrot-De la Puente, Josep Franch-Nadal, and Gabriel Coll-de Tuero. "Chronic Kidney Disease and Diabetic Retinopathy in Patients with Type 2 Diabetes", PLoS ONE, 2016. **2%**

Publication
- 3** Bello, N. A., M. A. Pfeffer, H. Skali, J. B. McGill, J. Rossert, K. A. Olson, L. Weinrauch, M. E. Cooper, D. de Zeeuw, P. Rossing, J. J. V. McMurray, and S. D. Solomon. "Retinopathy and clinical outcomes in patients with type 2 diabetes mellitus, chronic kidney disease, and

**1%**



anemia", BMJ Open Diabetes Research & Care, 2014.

Publication

4

Submitted to Flinders University

Student Paper

1%

5

Michael Larsen. "Grading of Diabetic Retinopathy", Retinal Vascular Disease, 2007

Publication

1%

6

Submitted to Cranfield University

Student Paper

1%

7

Submitted to University of South Florida

Student Paper

1%

8

Submitted to October University for Modern Sciences and Arts (MSA)

Student Paper

1%

9

"Study results from University Hospital, Department of Ophthalmology provide new insights into diabet", Blood Weekly, April 29 2010 Issue

Publication

1%

10

Einar Stefánsson. "Diabetic macular edema", Saudi Journal of Ophthalmology, 2009

Publication

1%

11

Lally, David R., Chirag P. Shah, and Jeffrey S. Heier. "Vascular endothelial growth factor and diabetic macular edema", Survey of

1%

# Ophthalmology, 2016.

Publication

---

12

"Evaluation of systemic risk factors in different optical coherence tomographic patterns of diabetic macular edema", International Journal of Ophthalmology, 2018

Publication

---

1%

---

Exclude quotes Off

Exclude matches < 15 words

Exclude bibliography Off