

Bukti Koresponding

RESEARCH J. PHARM. AND TECH. 15(8), 2022.

S1 Farmasi FMIPA UNSRI

Regarding submission of final version of article.

Eksternal

Kotak Masuk x



Anvpublications@anvpublication.org

kepada dianto_72@yahoo.com; saya ▾

Jum, 29 Okt 2021, 16.23



[Terjemahkan ke Indonesia](#)



Research Journal of Pharmacy and Technology

Paper ID: 21713171640045385 Date of Submission: 13-Jul-2021

Paper Title: FORMULATION AND IN-VITRO ANTIBACTERIAL ACTIVITY OF GEL CONTAINING ETHANOLIC EXTRACT OF PURPLE SWEET POTATO LEAVES (Ipomoea batatas (L.)) LOADED POLY LACTIC CO-GLYCOLIC ACID SUBMICROPARTICLES AGAINST Staphylococcus aureus

Authors: Mardiyanto Mardiyanto;Elsa Fitria Apriani;Muhammad Hafizhaldi Alfarizi

Dear Author(s),

Editorial board has considered your article titled "FORMULATION AND IN-VITRO ANTIBACTERIAL ACTIVITY OF GEL CONTAINING ETHANOLIC EXTRACT OF PURPLE SWEET POTATO LEAVES (Ipomoea batatas (L.)) LOADED POLY LACTIC CO-GLYCOLIC ACID SUBMICROPARTICLES AGAINST Staphylococcus aureus". for publication in Year : 2022, Vol: 15, Issue: 8. You are requested to send the final version of your article by clearly mentioning the *Paper Title* and *Author(s) name, affiliation and email address* in proper sequence.

Note:- Please submit the final version of article with in 7 Days.

Please Login to account from which you have submitted the article for more details. (To login now click : http://www.anvpublication.org/Subscribed_User_Login.aspx)

Thank you.

Editor

This is auto generated mail. Please do not reply.

Regarding submission of final version of article. Eksternal Kotak Masuk x



Anvpublications@anvpublication.org

kepada dianto_72@yahoo.com, saya ▾

Jum, 29 Okt 2021, 16.23



Terjemahkan ke Indonesia



Research Journal of Pharmacy and Technology

Paper ID: 21713171640045385 Date of Submission: 13-Jul-2021

Paper Title: FORMULATION AND IN-VITRO ANTIBACTERIAL ACTIVITY OF GEL CONTAINING ETHANOLIC EXTRACT OF PURPLE SWEET POTATO LEAVES (Ipomoea batatas (L.)) LOADED POLY LACTIC CO-GLYCOLIC ACID SUBMICROPARTICLES AGAINST Staphylococcus aureus

Authors: Mardiyanto Mardiyanto;Elsa Fitria Apriani;Muhammad Hafizhaldi Alfarizi

Dear Author(s),

Editorial board has considered your article titled "FORMULATION AND IN-VITRO ANTIBACTERIAL ACTIVITY OF GEL CONTAINING ETHANOLIC EXTRACT OF PURPLE SWEET POTATO LEAVES (Ipomoea batatas (L.)) LOADED POLY LACTIC CO-GLYCOLIC ACID SUBMICROPARTICLES AGAINST Staphylococcus aureus". for publication in Year : 2022, Vol: 15, Issue: 8. You are requested to send the final version of your article by clearly mentioning the *Paper Title* and *Author(s) name, affiliation and email address* in proper sequence.

Note:- Please submit the final version of article with in 7 Days.

Please Login to account from which you have submitted the article for more details. (To login now click : http://www.anvpublication.org/Subscribed_User_Login.aspx)

Thank you.

Editor

This is auto generated mail. Please do not reply.

Reviewer 1

1. Please include lines in each sentence states.
2. In the abstract, use italics for the name of the bacteria.
3. In the abstract, use italics for the name of the plant.
4. In the abstract, is there any quantitative data to state antibacterial activity?
5. On page 4, how do you store the product because it is affected by light?
6. On page 5, part B should be made into sentences with neat spacing.
7. In the method section, is there no need for an analysis data section?
8. On page 7 please arrange the position of the image.
9. At the end of the results and discussion, please provide a summary work.
10. Please put a space between the conflict of interest and the reference.
11. In the conclusion section, please include what are the benefits for the future of this research.

Reviewer 2

1. Italics is used for the name of the bacteria and plants.
2. In the abstract, use quantitative data of antibacterial activity?
3. Use the correct method to avoid light during storage.
4. Include an analysis data in method.
5. The position of the image should be properly.
6. Add a summary work in end of results.

Formulation and *In-vitro* Antibacterial Activity of Gel containing Ethanolic extract of Purple Sweet Potato Leaves (*Ipomoea batatas* (L.) Loaded Poly Lactic Co-Glycolic Acid Submicroparticles against *Staphylococcus aureus*

Mardiyanto Mardiyanto, Elsa Fitria Apriani*, Muhammad Hafizhaldi Alfarizi

Department of Pharmacy, Faculty of Mathematics and Natural Sciences, Sriwijaya University, South Sumatra, Indonesia.

*Corresponding Author E-mail: elsafitria@mipa.unsri.ac.id

ABSTRACT:

Staphylococcus aureus has been resistant to various antibiotics including erythromycin, clindamycin, penicillin, trimethoprim-sulfamethoxazole, tetracyclines, chloramphenicol, and piperacillin-tazobactam so that an alternative treatment is needed. The purple sweet potato leaves (*Ipomoea batatas* (L.) Poir) contain flavonoid compounds that have antibacterial activity by inhibiting nucleic acid, protein synthesis, cell membrane, and energy metabolism in bacteria. In this study, ethanolic extract of purple sweet potato leaves is loaded to poly lactic-co-glycolic acid submicroparticles to increase the stability of flavonoids and the antibacterial effect. Submicroparticle gel was prepared with various concentrations of hydroxypropyl methylcellulose ie F1, F2, and F3 respectively 3%, 5%, and 7%. The antibacterial activity of submicroparticles gel compared with a gel containing extracts without submicroparticle and erythromycin gel as a positive control. Phytochemical test results that the ethanolic extract of purple sweet potato leaves contains flavonoids. Based on the research results, the best formula was F1(3%) with pH, homogeneity, viscosity, dispersibility, adhesion, and washability, respectively 7.4 ± 0.0361 ; homogeneous; 8358.9 ± 228.1391 cps; 4.2667 ± 0.3005 cm; 45.333 ± 2.5166 seconds; 11.6667 ± 1.5275 mL. F1 was also shown to have strong antibacterial activity.

KEYWORDS: Purple sweet potato leaves (*Ipomoea batatas* (L.) Poir), Flavonoids, Hydroxypropyl Methylcellulose, *Staphylococcus aureus*, Antibacterial, Submicroparticles.

INTRODUCTION:

Bacterial resistance is a growing problem throughout the world, including in Indonesia. One of the bacteria that has been resistant to various antibiotics is *Staphylococcus aureus*. *Staphylococcus aureus* has been resistant to penicillin, methicillin, oxacillin, vancomycin, daptomycin, erythromycin, clindamycin, trimethoprim- sulfamethoxazole, tetracycline, chloramphenicol, ampicillin-sulbactam, and piperacillin-tazobactam.^{1,2} So that, alternative treatment is needed to avoid this problem using a natural compound.³

The plant that is known to have antibacterial activity is purple sweet potato (*Ipomoea batatas* (L.) Poir). Purple sweet potato leaves contain secondary metabolite compounds such as alkaloids, tannins, saponins, and flavonoids.⁴

Research conducted by Osuntokun et al.⁵ stated that the ethanolic extract of purple sweet potato leaves has antibacterial activity against several bacteria, one of them is the *Staphylococcus aureus*. A concentration of 50mg/mL of ethanolic extract of purple sweet potato leaves act as a moderate antibacterial by producing an inhibition zone

diameter of $\pm 10\mu\text{m}$ and a minimum inhibitory concentration of 12.5mg/ml.

Secondary metabolites that act as antibacterial agents in purple sweet potato leaves are flavonoids.^{6,7,8} Flavonoids have antibacterial activity by inhibiting nucleic acid synthesis, cell membrane function, energy metabolism, and protein synthesis.⁹ Flavonoids are compounds that are unstable against the effects of oxidation, light, and chemical changes, so that when oxidized their structure will change and their function as an active ingredient will decrease.^{10,11,12} The stability of flavonoids can be improved by making them into the form of submicroparticles.

Submicroparticle preparations have the advantage such as can protect the active substance, being enhancers, and can be efficient in drug dosage.^{13,14} The polymer used in the manufacture of submicroparticle preparations is PLGA (Poly - Lactic - co - Glycolic Acid) with PVA (PolyVynil Alcohol)¹⁵ as a stabilizer using the Emulsion solvent evaporation method. PLGA was chosen because it is biodegradable, biocompatible, non-toxic, and has been approved by the FDA.¹⁶ PLGA polymers are capable of delivering drugs and producing small particle sizes ranging from 180-200nm.¹⁷ The submicroparticles formulated into a gel preparation so that it is easy for the patient to use.

A gel is topical preparation which is usually applied through the skin or mucous membranes. The gel contains a lot of water components and has good drug delivery capabilities.¹⁸ The gelling agent used in this study was Hydroxypropyl methylcellulose (HPMC). HPMC can form a clear, neutral gel and has a stable viscosity.¹⁹ The concentration of HPMC is playing important role in topical preparation. A high concentration of HPMC will produce a viscous gel. When the viscosity of gel is high, the gel will be more resistant to dilution or erosion.²⁰ This will have an impact on drug release. Thus, gel with high viscosity will exhibited slower drug release.²¹

Based on the description above, in this research, ethanolic extract of purple sweet potato leaves will be formulated into submicroparticle and submicroparticle will be formulated into gel. Submicroparticles of ethanolic extract of purple sweet potato leaves will be seen antibacterial activity against *S. aureus* by determining the inhibition zone. Antibacterial activity testing was divided into 5 groups, namely negative control (gel containing submicroparticles without active substances), positive control (erythromycin gel), and treatment groups F1, F2, F3 with HPMC variations of 3%, 5%, and 7%.

MATERIALS AND METHODS:

Materials:

The materials used in this study were purple sweet potato leaves (South Sumatra), 96% ethanol (DiraSonita®), 10% Pb Acetate (Bratachem®), 20% NaOH (Bratachem®), PLGA (SigmaAldrich®), PVA (Bratachem®), Aquadest. (DiraSonita®), ethyl acetate (Bratachem®), hydroxypropyl methylcellulose (Bratachem®), propylene glycol (Bratachem®), methylparaben (Bratachem®), propylparaben (Bratachem®), *Staphylococcus aureus* (Biomeriux®), nutrient broth (Merck®), nutrient agar (Merck®), erythromycin (Erymed®), and McFarland's solution (Merck®).

Plant Collection and Determination:

The plant was collected from Pagar Alam, South Sumatra, Indonesia. The determination of this plant was conducted in Herbarium of Andalas University Andalas, Padang, Indonesia.

Preparation of Ethanolic Extract:

The leaves taken are greenish. The leaves that have been picked then washed under running water. The drying process is protected from direct sunlight for 7 days. The dried leaves are powdered using a grinding machine. The extract was

made using the maceration method. The simplicia powder of purple sweet potato leaves was dissolved in 96% ethanol solvent with a ratio of simplicia and solvent of 1:20. The filtrate obtained is evaporated at a temperature of 60°C and 80 rpm so that a thick extract is obtained.

Phytochemical Test for Flavonoid Compound:

A 1 ml of test solution was inserted into three test tubes. Tube 1 as a control. Tube 2 was added with 1 mL of 10% Pb acetate (lead acetate) solution, the positive result for flavonoids if there is a yellow precipitate. Tube 3 added with a few drops of 20% NaOH forms a yellow color if it contains flavonoids.²²

Preparation of Submicroparticles:

Submicroparticles were made using the Emulsion Solvent Evaporation Method. The ethanolic extract of purple sweet potato leaves was dispersed with 96% ethanol, Poly Lactic-Co-Glycolic Acid (PLGA) was dissolved in ethyl acetate and PolyVinyl Alcohol (PVA) was dissolved in distilled water. Extract solution containing 150mg of purple sweet potato leaf extract was dissolved with PLGA solution containing 50 mg of PLGA in a beaker glass using a Magnetic Stirrer at 750 rpm (oil phase). PVA solution containing 50mg of PVA act as a water phase. Emulsify the oil phase into the water phase by dropping it using a 50µL Micropipette (O/W emulsion) on a Magnetic Stirrer. The emulsification process is left for 1 hour at a speed of 750 rpm. In the next step, the solution was homogenized using ultra turrax at a speed of 12.0 x 1000rpm for 5 minutes. The particle solution was evaporated to remove the remaining organic solvent on the Magnetic Stirrer for 24 hours.²³

Preparation of Submicroparticles Gel:

The gel was made of four formulas. The three formulas containing submicroparticle of the ethanolic extract with variations in the concentration of HPMC, namely 3%, 5%, and 7% and one formula containing ethanolic extract without submicroparticles. The formulas for each are shown in Table 1.

Table 1. Formula Gel

Materials	Quantity			
	F1	F2	F3	F4
Submicroparticle of the ethanolic extract of purple sweet potato leaves	150 mg	150 mg	150 mg	-
Ethanolic extract of purple sweet potato leaves	-	-	-	150 mg
HPMC	3 g	5 g	10 g	*
Propylene Glycol	15 mL	15 mL	15 mL	15 mL
Methylparaben	0.18 g	0.18 g	0.18 g	0.18 g
Propylparaben	0.02 g	0.02 g	0.02 g	0.02 g
Aquadest	Ad. 100 mL	Ad. 100 mL	Ad. 100 mL	Ad 100 mL

Note : * Refers to optimum formula between F1, F2, F3

HPMC is dissolved in hot water in a ratio of 1:20 until it forms a gel mass. Methylparaben and propylparaben were dissolved in propylene glycol and added little by little to the gel mass and then stirred until homogeneous. Submicroparticles of ethanolic extract of purple sweet potato leaves are added gradually to the gel mass, then add 100mL of distilled water, stirring until homogeneous.

Evaluation of Submicroparticles Gel:

a. Organoleptic Test:

The organoleptic test in this study was carried out directly observing the texture, color, and smell.

b. pH Test:

The pH of the gel was determined using a digital pH meter. An amount of submicroparticles gel was stirred in distilled water and the pH of the solution was measured.

c. Homogeneity Test:

The homogeneity test was carried out by applying the gel to a transparent glass preparation. It was observed visually. If there were no particles so the gel was homogenous.

d. Viscosity Test:

Measurement of the viscosity of the submicroparticle gel was carried out using a viscometer. A total of 30mL of gel preparation in a beaker is placed on the viscometer by inserting the rotor into the beaker and setting the speed to 60rpm. Observe the needle manually on the tool pointing to the number on the viscosity scale. The viscosity value is characterized by a stable needle movement and is expressed in cP.

e. Spreadability Test:

An excess of gel was placed in between two glass slides and then 125 gram weight was placed on slides for 1 min and then the diameter formed is measured Weight.

f. Adhesion Test:

An 0.25-gram gel was placed in between two glass s and then 1 kg weight was placed on slides for 5 min. Then added weight 80gram to pan and determine the time required to separate the two slides.

g. Washable Test:

The gel is applied on the surface of human skin, then flattene, and then run with water until the stains of the gel

preparation disappear. Record the volume of water used.

h. Cycling Tests:

The submicroparticles gel was kept in the refrigerator at $4\pm 2^{\circ}\text{C}$ for 24 hours and subsequently moved to $40\pm 2^{\circ}\text{C}$ for 24 hours. The organoleptic, pH, homogeneity, and syneresis phenomenon were determined.²⁴

In vitro Antibacterial Activity Test:

Antibacterial activity was evaluated using the disk diffusion method to determine the value of the inhibition zone.

The bacterial suspension was made by diluting some colonies in stock bacterial to the nutrient broth. The bacterial suspension of *Staphylococcus aureus* was compared visually with 1.5 McFarland Standard. The treatments were divided into 6 groups as shown in Table 2. Erythromycin gel was selected as a positive control and submicroparticle placebo gel as a negative control.

Table 2. Treatment Group

No.	Group	Treatment
1	Negative Control	Submicroparticles placebo Gel
2	Positive Control	Erythromycin Gel
3	F1	Submicroparticles Gel with 3 % HPMC
4	F2	Submicroparticles Gel with 5 % HPMC
5	F3	Submicroparticles Gel with 7 % HPMC
6	F4	Ethanollic Extract Gel with optimum concentration of HPMC

First, the 100 μL of bacterial inoculums containing were spread over plates containing Nutrient agar. The disc with a diameter of 6 mm impregnated to agar and the gel was placed on the disc. The plates were incubated for 24 h at 37°C , and the experiments were performed in triplicate. The diameters of inhibition zones were measured.

RESULT:

Preparation of Ethanolic Extract:

The purple sweet potato leaves were macerated using 96% ethanol. This solvent is used because it is universal so that it can dissolve polar and non-polar compounds and has a good ability to extract low molecular weight compounds such as flavonoids.²⁵ The filtrate produced during the maceration process was concentrated using a rotary evaporator at 60°C and obtained 56.36 g of thick and blackish-brown extract with a yield of 9.39%.

Phytochemical Test for Flavonoid Compound:

The phytochemical test carried out on the extract was the flavonoid test. This is because flavonoid compounds play a role in antibacterial activity. The ethanolic extract of purple sweet potato leaves positively contained flavonoids as evidenced by the formation of a yellowish precipitate when Pb acetate was added and the formation of a brick red color when 20% NaOH was added.

Preparation of Submicroparticles:

Submicroparticles were prepared using the emulsion solvent evaporation method. Submicroparticles are made by mixing the extract solution into the PLGA solution, this aims to form an oil phase which will then be mixed into the PVA solution (water phase). The oil phase is mixed into the water phase with a micropipette to form an O/W emulsion. Then the emulsion

is evaporated. During the evaporation process, the emulsion will become a suspension and the remaining content after the evaporation process is only the active substance of the ethanolic extract of purple sweet potato leaves encapsulated by PLGA and PVA as a stabilizer in the water phase. The submicroparticle formulation is expected to increase the stability of flavonoids and increase their antibacterial activity so that the resulting effect is maximized. The submicroparticles solution obtained in this study can be seen in Figure 1.



Figure 1. Submicroparticles Solution

Table 3. The Result of Evaluation of Submicroparticles Gel

Parameter	Formula		
	F1	F2	F3
Organoleptic	Light yellow, characteristic smell of extract, semi-solid	Light yellow, characteristic smell of extract, semi-solid	Light yellow, characteristic smell of extract, semi-solid
pH	7.40±0.04	7.44±0.06	7.48±0.05
Homogeneity	Homogenous	Homogenous	Homogenous
Viscosity	9093.9±248.2 cps	12020.4±379.1 cps	14546.3±156.5 cps
Spreadability	4.3±0.3 cm	3.6 ±0.1 cm	2.5±0.2 cm
Adhesion	45.3±2.5 s	724±2.6 s	1794.3±5.5 s
Washability	11.7±1.5 mL	15.3±1.5 mL	21.3±2.1 mL

Preparation of Submicroparticles Gel:

Preparation of submicroparticle gel of ethanolic extract of purple sweet potato leaves requires ingredients such as HPMC as a gelling agent, propylene glycol as a humectant, methylparaben and propylparaben as a preservative and aquadest as a solvent. HPMC is able to form a gel by absorbing solvents and forming a compact liquid mass. HPMC is used as a gelling agent because HPMC can form a clear and neutral gel and has a stable viscosity and has good drug delivery capability.¹⁸ In this study, three gel formulas (F1, F2, F3) were made by variation concentration of HPMC ie 3%, 5%, and 7%.




Figure 2. Submicroparticles Gel

Evaluation of Submicroparticles Gel:

Gel evaluation carried out in this study was an organoleptic test, pH test, homogeneity test, viscosity test, spreadability test, adhesion test, washability test, and cycling test. The results of the gel evaluation can be seen in Table 3.

Based on the result in Table 3 that the F1 was chosen as optimum formula because of the viscosity value is not too high so the resulting gel has a soft texture than the others.

***In-vitro* Antibacterial Activity Test:**

Antibacterial activity test was carried out by measuring the diameter of the inhibition zone. The results of the antibacterial activity test can be seen in Table 4.

Table 4. Inhibition Zone Measurement Results for Gel Preparations

Group	Average Inhibition Zone Diameter±SD (mm)
Negative Control	0±0*
Positive Control	13±1.73
F1	10.23 ± 0.69**
F2	5.78 ± 1.39*
F3	4.0 ± 0.0*
F4	1.89±1.54*

Notes: Statistical significance for each condition compared to positive control (*P<0.05; **P>0.05)

Based on the results of the antibacterial activity test above, F1 has antibacterial activity that is not significantly different from the positive control while the other two formulas had significant differences. This proves that the use of HPMC concentration has an effect on antibacterial activity.

DISCUSSION:

Ethanol extract of purple sweet potato leaves obtained in this study was thick and blackish-brown extract with a yield of 9.39%. Ethanol extract in this study is proven to contain flavonoid compounds. This is in accordance with research conducted by Kim et al²⁶, who succeeded in isolating the ethanol extract of purple sweet potato leaves and obtained 3 types of flavonoids, namely quercetin-3-O-β-D-glucopyranoside, isoquercitrin, and kaempferol 3-O-β-glucopyranoside. Quercetin-3-O-β-D-glucopyranoside has been researched and proven to have antimicrobial activity.^{27,28} But, quercetin are compounds that has low stability, poor water solubility and also low bioavailability.²⁹ Quercetin is thermally unstable, has low stability when in aqueous solution so that it is easily hydrolyzed, and unstable in the presence of light and oxygen which causes oxidation.³⁰ The stability of flavonoids can be improved by making them into the form of submicroparticles. In this study, PLGA used as polymer and PVA used as stabilizer in submicroparticles. PLGA will encapsulate the ethanol extract of purple sweet potato leaves due to ionic interactions. Ionic interactions between the drug and polymer will result in higher incorporation in the non- end-capped polymers while if the interaction shows hydrophobic features, the end-capped polymers will show greater incorporation.³¹ The presence of PVA will increase the stability of submicroparticles because PVA acts as an emulsifying agent. During the manufacture of submicroparticles, hydrophobic bonds will occur between the acetyl group in PLGA and the hydrolyzed PVA.³² The presence of PVA will strengthen the encapsulation of PLGA-ethanol extract of purple sweet potato leaves.

The submicroparticles formulated into a gel preparation. In this study, submicroparticles formulated into three formula gel with various concentrations of hydroxypropyl methylcellulose ie F1, F2, and F3 respectively 3%, 5%, and 7%. Based on the results of the gel evaluation in Table 3, the organoleptic test, pH test, homogeneity test and cycling test between the three formulas did not show a significant difference (p > 0.05). The gel produced from the three formulas is light yellow, has the distinctive smell of the extract, has a semi-solid, and homogenous texture. The pH results of the three formulas show that the pH is in the range of 7. Ideally, topical preparations have the same or similar pH value to the skin pH, namely 4.5-6.5 to avoid irritation of the skin. However, the gel at a pH of 7 can still be tolerated by the skin and has not irritated. The resulting gel pH tends to be high due to the influence of PVA on submicroparticles. PVA has a pH range of 5-8, the higher concentration of PVA used, the more hydroxyl ions (OH-) from PVA are likely to be released and will shift the pH value to higher.³³ In the viscosity test, the spreadability test, the adhesion test, and the washability test, there was a significant difference (p<0.05) which indicated that there was an effect of the HPMC concentration used on the test. Increasing the amount of gelling agent can strengthen the gel constituent matrix so the viscosity will increase.

In addition, the presence of PVA also affects the viscosity value. HPMC and PVA are polymers that can enter the cavity formed by water molecules so that hydrogen bonds occur between the hydroxyl (-OH) groups of the polymer and water molecules.^{34,35} This hydrogen bond plays a role in hydration in the swelling process.³⁶ The results of this viscosity test are inversely proportional to the results of the spreadability and washability tests but are directly proportional to the results of the adhesion test. The higher the viscosity result, the dispersibility and washability's result will be smaller, but the adhesion will be stronger. The physical stability test (cycling test) also showed no significant difference between the three formulas ($p > 0.05$). The parameters observed in this test included organoleptic (color, smell, and texture) and changes in pH. The results of the observations obtained were that there was no organoleptic change in the three formulas, each formula had a light yellow color, the distinctive odor of the extract, semi-solid texture, and homogeneous. The three formulas do not indicate a syneresis phenomenon. Based on the gel evaluation and stability results, the F1 formula with a concentration of 3% HPMC was chosen as the optimum formula in making submicroparticle gel and used as the basis for making purple sweet potato leaves ethanolic extract gel which not formulated into submicroparticles for antibacterial activity testing.

Based on the results of the antibacterial activity test above, the three formulas gel containing the ethanolic extract of purple sweet potato leaves in the form of submicroparticles had antibacterial activity. F1 with the smallest HPMC concentration of 3% has the largest inhibition zone diameter compared to F2 and F3. The antibacterial activity result also depends on the viscosity of the gel. Viscosity of the gel can affect the drug release from the vesicle. Increase in viscosity will result in slower drug release and the effect of antibacterial will be reduced.^{37,38} Viscosity of gel was directly proportional to the concentration of gelling agent used so in this study F1 with the smallest concentration had the largest inhibition zone diameter. The negative control did not produce inhibition because the component of submicroparticle gel without an active substance did not have antibacterial activity. The positive control of erythromycin gel had the largest inhibition zone diameter was 13 ± 1.73 mm. Erythromycin has antibacterial activity by binding to the ribosome subunit 50s so that protein synthesis is inhibited in bacteria.^{39,40}

Based on the data in Table 4, it can be seen that there is an increase in antibacterial activity after the ethanolic extract of purple sweet potato leaves is formed into submicroparticles. F4 or gel contain extract which not formulated into submicroparticles with an optimum concentration of HPMC (3%) can only cause inhibition of 1.67 ± 2.08 mm. In the research of Osuntokun et al.⁵ ethanolic extract of purple sweet potato leaves with a concentration of 50 mg/ml gives an inhibition zone diameter of 10 mm. In this study, the amount of extract used was 150 mg/100 ml and gave an inhibition zone of 1.67 ± 2.08 mm. However, when it was formed into submicroparticles, it was able to provide an inhibition zone diameter of 10.23 ± 0.69 mm. This proves that the submicroparticle can reduce dose usage.

Submicroparticle preparations can also increase the contact surface area because the resulting particle size is microparticle size so that the penetration of the active substance will increase and the antibacterial effect increases.⁴¹ Flavonoids can act as an antibacterial by inhibiting protein synthesis. Protein synthesis from bacteria occurs in the cytoplasm so that by increasing the penetration power of flavonoids into bacterial cells, more flavonoid molecules can inhibit protein synthesis in bacteria. In addition, the flavonoids formulated into submicroparticle preparations have better stability. Submicroparticle polymers protect flavonoids by encapsulating flavonoid molecules so they are protected from chemical disturbances. The mechanism for the release of active substances from PLGA submicroparticles can go through 4 ways, namely diffusion through water-filled pores, diffusion through the polymer, osmotic pumping, and erosion. In

general, the mechanism for releasing the active substance from PLGA submicroparticles is by diffusion through water-filled pores. The water molecules around the submicroparticle will enter the system and cause swelling so that the formation of pores which allows the encapsulated active substance to leave the submicroparticle system constantly until the erosion stage where the PLGA polymer as a submicroparticle maker begins to degrade.^{42,43} Submicroparticles reach the target site of bacterial also assisted by propylene glycol used in the gel. Propylene glycol as a penetration enhancer works by extracting lipids and proteins from the stratum corneum to form pores in the stratum corneum. Propylene glycol also changes the solubility parameters of the stratum corneum so that it affects the thermodynamic activity of the submicroparticles contained in the gel formula. This pushes the submicroparticles into the stratum corneum thereby increasing the penetration of the submicroparticles and increase antibacterial activity.^{44,45,46}

The results of the shapiro-wilk statistical analysis showed that the data for all groups were normally distributed. One Way ANOVA statistical analysis showed the significance value $p < 0.05$ so it can be concluded that there is a significant difference in the results of the antibacterial activity test of all treatments. In the post hoc test, it can be seen that there is a significant difference in the antibacterial activity of the positive control group with all treatment groups, except F1.

CONCLUSION:

Based on the results of the research, there was an effect of the HPMC concentration on the results of the physical evaluation of the gel preparation. The higher concentration of HPMC used will cause the resulting gel to be viscous so that it will affect the releasing drug from the gel matrix. F1 with a concentration of 3% HPMC was chosen as the optimum formula because physically good, stable, and produces the best antibacterial effect among the three formulas. In addition, the use of submicroparticle preparations as a carrier for ethanolic extract of purple sweet potato leaves has also been shown to increase antibacterial activity that be able to efficiently use dosages.

CONFLICT OF INTEREST:

The authors have no conflicts of interest regarding this investigation.

REFERENCES:

1. Che Hamzah AM, Yeo CC, Puah SM, Chua KH, Chew CH. Staphylococcus aureus Infections in Malaysia: A Review of Antimicrobial Resistance and Characteristics of the Clinical Isolates, 1990-2017. Antibiotics (Basel). 2019; 8(3):128. doi:10.3390/antibiotics8030128
2. Foster TJ. Antibiotic resistance in Staphylococcus aureus. Current status and future prospects. FEMS Microbiology Reviews. 2017; 41 (3): 430 – 449. doi: 10.1093/femsre/fix007
3. Bezlon G, Shanmugha SD, Rinu ERE. Design and Stabilization of Natural Antibacterial Compound Allicin against Methicillin-Resistant Staphylococcus Aureus for Treatment as a Novel Antibiotic. Research J. Engineering and Tech. 2013; 4(4): 179-181.
4. Pochapski MT, Fosquiera EC, Esmerino LA, dos Santos EB, Farago PV, Santos FA, Groppo FC. Phytochemical screening, antioxidant, and antimicrobial activities of the crude leaves' extract from Ipomoea batatas (L.) Lam. Pharmacognosy Magazine. 2011; 7 (26): 165-170. doi: 10.4103/0973-1296.80682
5. Osuntokun OT, Babantude Y, Oledale MA, Fasila O. Components and bioactivity of Ipomoea batatas (L.) (Sweet Potato) ethanolic leaf extract. Asian Journal of Advanced Research and Reports. 2020; 10 (1): 10-26. doi:10.9734/AJARR/2020/v10i1/130232
6. Kusuma SAF, Wahyuni UT, Zuhrotun A. Evaluation of Antibacterial Activity of Indonesian Varieties Sweet Potato Leaves Extract from Cilembu Against Shigella Dysenteriae Atcc 13313. Asian J Pharm Clin Res. 2017; 10 (2): 377-380. doi:10.22159/ajpcr.2017v10i2.15773
Manvar MN. Antibacterial Activity of Leaves and Flowers of Ipomoea aquatica Forsk. (Convolvulaceae). Asian J. Pharm. Res. 2018; 8(2): 94-98. doi: 10.5958/2231-5691.2018.00016.3
7. Patil SD, Hafizur MA, Priti A, Shelke PB, Yardi S. Synthesis and evaluation of novel Flavonoid derivatives for Antibacterial activity. Asian J. Pharm.

- Res. 2016; 6 (1): 27-30. doi: 10.5958/2231-5691.2016.00005.8
8. Dzoyem JP, Hamamoto H, Ngameni B, Ngadjui BT, Sekimizu K. Antimicrobial action mechanism of flavonoids from *Dorstenia* species. *Drug Discoveries & Therapeutics*. 2013; 7 (2): 66-72. doi: 10.5582/ddt.2013.v7.2.66
 9. Takashi K, Yoshinobu T, Takahisa N, Takashi N, Hiroshi T, Shigetaka O. Acceptor specificity of cyclodextrin glucanotransferase from an alkalophilic *Bacillus* species and synthesis of glycosyl rhamnose. *Bioscience, Biotechnology, and Biochemistry*. 1996; 60 (7): 1176-1178. doi: 10.1271/bbb.60.1176
 10. Pourcel L, Routaboul JM, Cheynier V, Lepiniec L, Debeaujon I. Flavonoid oxidation in plants: from biochemical properties to physiological functions. *Trends in Plant Science*. 2007; 12 (1): 29-36. doi:10.1016/j.tplants.2006.11.006
 11. Zulham G, Wilar Y, Susilawati A, Subarnas AY, Chaerunisaa. Microparticles of Herbal Extracts with Antioxidant Activity. *Pharmacog J*. 2021; 13 (1): 285-295. doi: 10.5530/pj.2021.13.38
 12. Sehar S, Khan AIH. Role of ZnO Nanoparticles for improvement of Antibacterial Activity in Food Packaging. *Asian Journal of Pharmaceutical Research*. 2021; 11(2):128-131. doi: 10.52711/2231-5691.2021.00024
 13. Buzea C, Blandino IIP, Robbie K. Nanomaterial and nanoparticles: sources and toxicity. *Biointerphase*. 2007; 2: 170-172. doi: 10.1116/1.2815690
 14. Somayadineshraj D, Shanmugapriyan R, Lakshmanaswamy R, Nishiha CS, Dooslin MBV, Muralidharan P. Synthesis and Characterization of MgO nanoparticles Coated by polyvinyl alcohol for Biomedical applications. *Research J. Pharm. and Tech*. 2018; 11(11): 4997-5000. doi: 10.5958/0974-360X.2018.00911.3
 15. Chetan MP, Manan AP, Nikhil PP, Prajapati PH, Patel CN. Poly Lactic Glycolic Acid (PLGA) As Biodegradable Polymer. *Research J. Pharm. and Tech*. 2020; 3(2): 353-360.
 16. Irfan M, Saiyyad DS, Bhambere, Sanjay K. Formulation and Optimization of Silymarin Loaded PLGA Nanoparticle for liver targeting. *Asian J. Pharm. Tech*. 2017; 7 (4): 209-220. doi: 10.5958/2231-5713.2017.00032.0
 17. Verma A, Singh VS, Kaur R, Jain UK. Formulation and Evaluation of Clobetasol Propionate Gel. *Asian Journal of Pharmaceutical and Clinical Research*. 2013; 6 (5): 15-18. ISSN - 0974-2441
 18. Rowe RC, Sheskey PJ, Quinn ME. Handbook of pharmaceutical excipients, Sixth Edition. UK, London: Pharmaceutical Press and American Pharmacists Association; 2009.
 19. Ghosal K. and Nanda A. Development of diclofenac potassium gel from hydrophobically modified HPMC. *Iranian Polymer Journal*. 2013; 22 (6): 1-24. doi: 10.1007/s13726-013-0145-3
 20. Binder L, Mazal J, Petz R, Klang V, Valenta C. The role of viscosity on skin penetration from cellulose ether-based hydrogels. *Skin Research and Technology*. 2019; 25 (5): 725-734. doi: 10.1111/srt.12709
 21. Indarti K, Apriani EF, Wibowo AE, Simanjuntak P. Antioxidant Activity of Ethanolic Extract and Various Fractions from Green Tea (*Camellia sinensis* L.) Leaves. *Pharmacognosy Journal*. 2019; 11 (4): 771-776. doi: 10.5530/pj.2019.11.122
 22. Mardiyanto. Investigation of nanoparticulate formulation intended for caffeine delivery into hair follicle. Dissertation. Department of Pharmacy, Faculty of Science, Saarland University, Saarbruecken. Germany; 2013.
 23. Apriani EF, Rosana Y, Iskandarsyah I. Formulation, characterization, and in vitro testing of azelaic acid ethosome-based cream against *Propionibacterium acnes* for the treatment of acne. *J Adv Pharm Technol Res*. 2019; 10 (2): 75-80. doi: 10.4103/japtr.JAPTR_289_18
 24. Ahmad I, Owais M, Shahid M, Aqil F. *Combating Fungal Infections* New York, USA : Springer; 2010.
 25. Kim HJ, Jin C, Lee YS. Isolation and Antioxidative Activities of Caffeoylquinic Acid Derivates and Flavonoid Glycosides from Leaves of Sweet Potato (*Ipomoea batatas* L.). *The Journal of Applied Pharmacology*. 2007; 15: 46-51. doi: 10.4062/biomolther.2007.15.1.046
 26. Islam MM, Akter S, Al-Amin M, Haque MM. Isolation of Quercetin-3- O-beta-d glucopyranoside from the Leaves of *Azadirachta Indica* and Antimicrobial and Cytotoxic screening of the Crude Extracts. *Dhaka University Journal of Science*. 2012; 60 (1): 11-14. doi:10.3329/dujs.v60i1.10328
 27. Sholkamy EN, Ahmed MS, Yasser MM, Mostafa AA. Antimicrobial quercetin 3-O-glucoside derivative isolated from *Streptomyces antibioticus* strain ess_amA8. *Journal of King Saud University*. 2020; 32 (3): 1838-1844. doi: 10.1016/j.jksus.2020.01.026
 28. Wang W, Sun C, Mao L, Ma P, Liu F, Yang J, Gao Y. The biological activities, chemical stability, metabolism and delivery systems of quercetin: A review. *Trends in Food Science & Technology*. 2016; 56: 21-38. doi: 10.1016/j.tifs.2016.07.004
 29. Cunico LP, Cobo AM, Al-Hamimi S, Turner C. Solubility and Thermal Degradation of Quercetin in CO₂-Expanded Liquids. *Molecules*. 2020; 25: 5582-5591. doi: 10.3390/molecules25235582
 30. Lagreca E, Onesto V, Natale CD, Manna SL, Netti PA, Vecchione R. Recent advances in the formulation of PLGA microparticles for controlled drug delivery. *Progress in Biomaterials*. 2020; 9: 153–174. doi: 10.1007/s40204-020-00139-y
 31. Stromberg ZR, Phipps ML, Magurudeniya HD, Pedersen CA, Rajale T, Sheehan CJ, Courtney SJ, Bradfute SB, Hraber P, Rush MN, Kubicek-

- Sutherland JZ, Martinez JS. Formulation of stabilizer-free, nontoxic PLGA and elastin-PLGA nanoparticle delivery systems. *International Journal of Pharmaceutics*. 2021; 597: 1-11. doi: 10.1016/j.ijpharm.2021.120340
32. Wiśniewska M, Chibowski S, Urban T. Effect of the type of polymer functional groups on the structure of its film formed on the alumina surface—suspension stability. *React Funct Polym*. 2012; 72: 791–798. doi: 10.1016/j.reactfunctpolym.2012.08.005
 33. Sharma N, Madan P, Lin S. Effect of process and formulation variables on the preparation of parenteral paclitaxel-loaded biodegradable polymeric nanoparticles: A co-surfactant study. *Asian journal of pharmaceutical sciences*. 2016; 11 (3): 404-416. doi: 10.1016/j.ajps.2015.09.004
 34. Zolnik BS, Burgess DJ. Effect of Acidic pH on PLGA Microsphere Degradation and Release. *Journal of Controlled Release*. 2007; 122: 338-344. doi: 10.1016/j.jconrel.2007.05.034
 35. Das N, Bose C, Tripathi N, Maitra S. Progress in the development of gelling agents for improved culturability of microorganisms. *Frontiers in Microbiology*. 2015; 6 (698): 1-7. doi: 10.3389/fmicb.2015.00698
 36. Pacharne PS, Patil JR, Chaudhari SP, Ratnaparkhi MP. Design, Development and formulation of Topical anti arthritic gel using different gelling agents. *Elixir Pharmacy*. 2013; 62: 17599-17603.
 37. Yener G, Dal O, Üner M. Effect of Vehicles on Release of Meloxicam from Various Topical Formulations. *The Open Drug Delivery Journal*. 2009; 3: 19-23. doi: 10.2174/1874126600903010019
 38. Jednacak T, Mikulandra I, Novak P. Advanced Methods for Studying Structure and Interactions of Macrolide Antibiotics. *International Journal of Molecular Sciences*. 2020; 21 (7799): 1 – 26. doi: 10.3390/ijms21207799
 39. Singh RK, Garg R. Formulation and Evaluation of Erythromycin Estolate Loaded Drug Balls. *Research J. Pharm. and Tech*. 2020; 13(1): 282-286 doi: 10.5958/0974-360X.2020.00057.8
 40. Mohanraj VJ, Chen Y. Nanoparticles-A review. *Trop J Pharm Res*. 2006; 5(1): 561-573. doi: 10.4314/tjpr.v5i1.14634
 41. Fredenberg S, Wahlgren M, Reslow M, Axelsson A. The mechanisms of drug release in poly(lactic-co-glycolic acid)-based drug delivery systems-A review. *International Journal of Pharmaceutics*. 2011; 415 (2011): 34-52. doi: 10.1016/j.ijpharm.2011.05.049.
 42. Vimal P, Bhavesh A, Anand D, Manish G, Adil P. A Review on Long Acting PLGA Based in Situ Forming Microparticles Formulation for a Novel Drug Delivery System. *Res. J. Pharm. Dosage Form. and Tech*. 2016; 8(2):127-134. doi: 10.5958/0975-4377.2016.00017.3
 43. Hakue T, Talukder MMU. Chemical Enhancer: A simplisitic way to modulate barrier function of the stratum comeum. *Adv. Pharm. Bull*. 2018; 8 (2):169-179. doi: 10.15171/apb.2018.021
 44. Vinod K, Aditi S. Study of Antimicrobial activity of Star Anise loaded poly (DL-lactide-co-glycolide) nanoparticles. *Research J. Pharm. and Tech*. 2019; 12 (2): 499-507. doi: 10.5958/0974-360X.2019.00088.X
 45. Vinod K, Aditi S. Synthesis, Characterization, Antimicrobial activity and Release Study of Cinnamon loaded poly (DL-lactide-co-glycolide) Nanoparticles. *Research J. Pharm. and Tech*. 2019; 12(4):1529-1535. doi: 10.5958/0974-360X.2019.00253.1

Formulation and *In-vitro* Antibacterial Activity of Gel containing Ethanolic extract of Purple Sweet Potato Leaves (*Ipomoea batatas* (L.) Loaded Poly Lactic Co-Glycolic Acid Submicroparticles against *Staphylococcus aureus*

Mardiyanto Mardiyanto, Elsa Fitria Apriani*, Muhammad Hafizhaldi Alfarizi

Department of Pharmacy, Faculty of Mathematics and Natural Sciences, Sriwijaya University, South Sumatra, Indonesia.

*Corresponding Author E-mail: elsafitria@mipa.unsri.ac.id

ABSTRACT:

Staphylococcus aureus has been resistant to various antibiotics including erythromycin, clindamycin, penicillin, trimethoprim-sulfamethoxazole, tetracyclines, chloramphenicol, and piperacillin-tazobactam so that an alternative treatment is needed. The purple sweet potato leaves (*Ipomoea batatas* (L.) Poir) contain flavonoid compounds that have antibacterial activity by inhibiting nucleic acid, protein synthesis, cell membrane, and energy metabolism in bacteria. In this study, ethanolic extract of purple sweet potato leaves is loaded to poly lactic-co-glycolic acid submicroparticles to increase the stability of flavonoids and the antibacterial effect. Submicroparticle gel was prepared with various concentrations of hydroxypropyl methylcellulose ie F1, F2, and F3 respectively 3%, 5%, and 7%. The antibacterial activity of submicroparticles gel compared with a gel containing extracts without submicroparticle and erythromycin gel as a positive control. Phytochemical test results that the ethanolic extract of purple sweet potato leaves contains flavonoids. Based on the research results, the best formula was F1(3%) with pH, homogeneity, viscosity, dispersibility, adhesion, and washability, respectively 7.4 ± 0.0361 ; homogeneous; 8358.9 ± 228.1391 cps; 4.2667 ± 0.3005 cm; 45.333 ± 2.5166 seconds; 11.6667 ± 1.5275 mL. F1 was also shown to have strong antibacterial activity with an inhibition zone value of 13.67 ± 4.04 mm.

KEYWORDS: Purple sweet potato leaves (*Ipomoea batatas* (L.) Poir), Flavonoids, Hydroxypropyl Methylcellulose, *Staphylococcus aureus*, Antibacterial, Submicroparticles.

INTRODUCTION:

Bacterial resistance is a growing problem throughout the world, including in Indonesia. One of the bacteria that has been resistant to various antibiotics is *Staphylococcus aureus*. *Staphylococcus aureus* has been resistant to penicillin, methicillin, oxacillin, vancomycin, daptomycin, erythromycin, clindamycin, trimethoprim- sulfamethoxazole, tetracycline, chloramphenicol, ampicillin-sulbactam, and piperacillin-tazobactam.^{1,2} So that, alternative treatment is needed to avoid this problem using a natural compound.³

The plant that is known to have antibacterial activity is purple sweet potato (*Ipomoea batatas* (L.) Poir). Purple sweet potato leaves contain secondary metabolite compounds such as alkaloids, tannins, saponins, and flavonoids.⁴ Research conducted by Osuntokun et al.⁵ stated that the ethanolic extract of purple sweet potato leaves has antibacterial activity against several bacteria, one of them is the *Staphylococcus aureus*. A concentration of 50mg/mL of ethanolic extract of purple sweet potato leaves act as a moderate antibacterial by producing an inhibition zone diameter of ± 10 mm and a

minimum inhibitory concentration of 12.5mg/ml. Secondary metabolites that act as antibacterial agents in purple sweet potato leaves are flavonoids.^{6,7,8} Flavonoids have antibacterial activity by inhibiting nucleic acid synthesis, cell membrane function, energy metabolism, and protein synthesis.⁹ Flavonoids are compounds that are unstable against the effects of oxidation, light, and chemical changes, so that when oxidized their structure will change and their function as an active ingredient will decrease.^{10,11,12} The stability of flavonoids can be improved by making them into the form of submicroparticles.

Submicroparticle preparations have the advantage such as can protect the active substance, being enhancers, and can be efficient in drug dosage.^{13,14} The polymer used in the manufacture of submicroparticle preparations is PLGA (Poly - Lactic - co - Glycolic Acid) with PVA (PolyVynil Alcohol)¹⁵ as a stabilizer using the Emulsion solvent evaporation method. PLGA was chosen because it is biodegradable, biocompatible, non-toxic, and has been approved by the FDA.¹⁶ PLGA polymers are capable of delivering drugs and producing small particle sizes ranging from 180-200nm.¹⁷ The submicroparticles formulated into a gel preparation so that it is easy for the patient to use.

A gel is topical preparation which is usually applied through the skin or mucous membranes. The gel contains a lot of water components and has good drug delivery capabilities.¹⁸ The gelling agent used in this study was Hydroxypropyl methylcellulose (HPMC). HPMC can form a clear, neutral gel and has a stable viscosity.¹⁹ The concentration of HPMC is playing important role in topical preparation. A high concentration of HPMC will produce a viscous gel. When the viscosity of gel is high, the gel will be more resistant to dilution or erosion.²⁰ This will have an impact on drug release. Thus, gel with high viscosity will exhibited slower drug release.²¹

Based on the description above, in this research, ethanolic extract of purple sweet potato leaves will be formulated into submicroparticle and submicroparticle will be formulated into gel. Submicroparticles of ethanolic extract of purple sweet potato leaves will be seen antibacterial activity against *S. aureus* by determining the inhibition zone. Antibacterial activity testing was divided into 5 groups, namely negative control (gel containing submicroparticles without active substances), positive control (erythromycin gel), and treatment groups F1, F2, F3 with HPMC variations of 3%, 5%, and 7%.

MATERIALS AND METHODS:

Materials:

The materials used in this study were purple sweet potato leaves (South Sumatra), 96% ethanol (DiraSonita®), 10% Pb Acetate (Bratachem®), 20% NaOH (Bratachem®), PLGA (SigmaAldrich®), PVA (Bratachem®), Aquadest. (DiraSonita®), ethyl acetate (Bratachem®), hydroxypropyl methylcellulose (Bratachem®), propylene glycol (Bratachem®), methylparaben (Bratachem®), propylparaben (Bratachem®), *Staphylococcus aureus* (Biomeriux®), nutrient broth (Merck®), nutrient agar (Merck®), erythromycin (Erymed®), and McFarland's solution (Merck®).

Plant Collection and Determination:

The plant was collected from Pagar Alam, South Sumatra, Indonesia. The determination of this plant was conducted in Herbarium of Andalas University Andalas, Padang, Indonesia.

Preparation of Ethanolic Extract:

The leaves taken are greenish. The leaves that have been picked then washed under running water. The drying process is protected from direct sunlight for 7 days. The dried leaves are powdered using a grinding machine. The extract was made

using the maceration method. The simplicia powder of purple sweet potato leaves was dissolved in 96% ethanol solvent with a ratio of simplicia and solvent of 1:20. The filtrate obtained is evaporated at a temperature of 60°C and 80 rpm so that a thick extract is obtained.

Phytochemical Test for Flavonoid Compound:

A 1 ml of test solution was inserted into three test tubes. Tube 1 as a control. Tube 2 was added with 1 mL of 10% Pb acetate (lead acetate) solution, the positive result for flavonoids if there is a yellow precipitate. Tube 3 added with a few drops of 20% NaOH forms a yellow color if it contains flavonoids.²²

Preparation of Submicroparticles:

Submicroparticles were made using the Emulsion Solvent Evaporation Method. The ethanolic extract of purple sweet potato leaves was dispersed with 96% ethanol, Poly Lactic-Co-Glycolic Acid (PLGA) was dissolved in ethyl acetate and PolyVinyl Alcohol (PVA) was dissolved in distilled water. Extract solution containing 150mg of purple sweet potato leaf extract was dissolved with PLGA solution containing 50 mg of PLGA in a beaker glass using a Magnetic Stirrer at 750 rpm (oil phase). PVA solution containing 50mg of PVA act as a water phase. Emulsify the oil phase into the water phase by dropping it using a 50µL Micropipette (O/W emulsion) on a Magnetic Stirrer. The emulsification process is left for 1 hour at a speed of 750 rpm. In the next step, the solution was homogenized using ultra turrax at a speed of 12.0 x 1000rpm for 5 minutes. The particle solution was evaporated to remove the remaining organic solvent on the Magnetic Stirrer for 24 hours.²³

Preparation of Submicroparticles Gel:

The gel was made of four formulas. The three formulas containing submicroparticle of the ethanolic extract with variations in the concentration of HPMC, namely 3%, 5%, and 7% and one formula containing ethanolic extract without submicroparticles. The formulas for each are shown in Table 1.

Table 1. Formula Gel

Materials	Quantity			
	F1	F2	F3	F4
Submicroparticle of the ethanolic extract of purple sweet potato leaves	150 mg	150 mg	150 mg	-
Ethanolic extract of purple sweet potato leaves	-	-	-	150 mg
HPMC	3 g	5 g	10 g	*
Propylene Glycol	15 mL	15 mL	15 mL	15 mL
Methylparaben	0.18 g	0.18 g	0.18 g	0.18 g
Propylparaben	0.02 g	0.02 g	0.02 g	0.02 g
Aquadest	Ad. 100 mL	Ad. 100 mL	Ad. 100 mL	Ad 100 mL

Note : * Refers to optimum formula between F1, F2, F3

HPMC is dissolved in hot water in a ratio of 1:20 until it forms a gel mass. Methylparaben and propylparaben were dissolved in propylene glycol and added little by little to the gel mass and then stirred until homogeneous. Submicroparticles of ethanolic extract of purple sweet potato leaves are added gradually to the gel mass, then add 100mL of distilled water, stirring until homogeneous, then stored in a tightly closed container.

Evaluation of Submicroparticles Gel:

a. Organoleptic Test:

The organoleptic test in this study was carried out directly observing the texture, color, and smell.

b. pH Test:

The pH of the gel was determined using a digital pH meter. An amount of submicroparticles gel was stirred in distilled water and the pH of the solution was measured.

c. Homogeneity Test:

The homogeneity test was carried out by applying the gel to a transparent glass preparation. It was observed visually. If there were no particles so the gel was homogenous.

d. Viscosity Test:

Measurement of the viscosity of the submicroparticle gel was carried out using a viscometer. A total of 30mL of gel preparation in a beaker is placed on the viscometer by inserting the rotor into the beaker and setting the speed to 60rpm. Observe the needle manually on the tool pointing to the number on the viscosity scale. The viscosity value is characterized by a stable needle movement and is expressed in cP.

e. Spreadability Test:

An excess of gel was placed in between two glass slides and then 125 gram weight was placed on slides for 1 min and then the diameter formed is measured Weight.

f. Adhesion Test:

An 0.25-gram gel was placed in between two glass s and then 1 kg weight was placed on slides for 5 min. Then added weight 80gram to pan and determine the time required to separate the two slides.

g. Washable Test:

The gel is applied on the surface of human skin, then flattene, and then run with water until the stains of the gel

preparation disappear. Record the volume of water used.

h. Cycling Tets:

The submicroparticles gel was kept in the refrigerator at $4\pm 2^{\circ}\text{C}$ for 24 hours and subsequently moved to $40\pm 2^{\circ}\text{C}$ for 24 hours. The organoleptic, pH, homogeneity, and syneresis phenomenal were determined.²⁴

***In vitro* Antibacterial Activity Test:**

Antibacterial activity was evaluated using the disk diffusion method to determine the value of the inhibition zone. The bacterial suspension was made by diluting some colonies in stock bacterial to the nutrient broth. The bacterial suspension of *Staphylococcus aureus* was compared visually with 1.5 Mc Farland Standard. The treatments were divided into 6 groups as shown in Table 2. Erythromycin gel was selected as a positive control and submicroparticle placebo gel as a negative control.

Table 2. Treatment Group

No.	Group	Treatment
1	Negative Control	Submicroparticles placebo Gel
2	Positive Control	Erythromycin Gel
3	F1	Submicroparticles Gel with 3 % HPMC
4	F2	Submicroparticles Gel with 5 % HPMC
5	F3	Submicroparticles Gel with 7 % HPMC
6	F4	Ethanollic Extract Gel with optimum concentration of HPMC

First, the 100 μL of bacterial inoculums containing were spread over plates containing Nutrient agar. The disc with a diameter of 6 mm impregnated to agar and the gel was placed on the disc. The plates were incubated for 24 h at 37°C , and the experiments were performed in triplicate. The diameters of inhibition zones were measured.

Data Analysis:

The data were analyzed by Shapiro-Wilk to see the normality of the distribution. If the data is normally distributed, then the test is carried out using ANOVA One Way. From this test, then proceed with the Post Hoc Test if the p-value is < 0.05 .

RESULT:

Preparation of Ethanolic Extract:

The purple sweet potato leaves were macerated using 96% ethanol. This solvent is used because it is universal so that it can dissolve polar and non-polar compounds and has a good ability to extract low molecular weight compounds such as flavonoids.²⁵ The filtrate produced during the maceration process was concentrated using a rotary evaporator at 60°C and obtained 56.36 g of thick and blackish-brown extract with a yield of 9.39%.

Phytochemical Test for Flavonoid Compound:

The phytochemical test carried out on the extract was the flavonoid test. This is because flavonoid compounds play a role in antibacterial activity. The ethanolic extract of purple sweet potato leaves positively contained flavonoids as evidenced by the formation of a yellowish precipitate when Pb acetate was added and the formation of a brick red color when 20% NaOH was added.

Preparation of Submicroparticles:

Submicroparticles were prepared using the emulsion solvent evaporation method. Submicroparticles are made by mixing

the extract solution into the PLGA solution, this aims to form an oil phase which will then be mixed into the PVA solution (water phase). The oil phase is mixed into the water phase with a micropipette to form an O/W emulsion. Then the emulsion is evaporated. During the evaporation process, the emulsion will become a suspension and the remaining content after the evaporation process is only the active substance of the ethanolic extract of purple sweet potato leaves encapsulated by PLGA and PVA as a stabilizer in the water phase. The submicroparticle formulation is expected to increase the stability of flavonoids and increase their antibacterial activity so that the resulting effect is maximized. The submicroparticles solution obtained in this study can be seen in Figure 1.



Figure 1. Submicroparticles Solution

Table 3. The Result of Evaluation of Submicroparticles Gel

Parameter	Formula		
	F1	F2	F3
Organoleptic	Light yellow, characteristic smell of extract, semi-solid	Light yellow, characteristic smell of extract, semi-solid	Light yellow, characteristic smell of extract, semi-solid
pH	7.40±0.04	7.44±0.06	7.48±0.05
Homogeneity	Homogenous	Homogenous	Homogenous
Viscosity	9093.9±248.2 cps	12020.4±379.1 cps	14546.3±156.5 cps
Spreadability	4.3±0.3 cm	3.6 ±0.1 cm	2.5±0.2 cm
Adhesion	45.3±2.5 s	724±2.6 s	1794.3±5.5 s
Washability	11.7±1.5 mL	15.3±1.5 mL	21.3±2.1 mL

Preparation of Submicroparticles Gel:

Preparation of submicroparticle gel of ethanolic extract of purple sweet potato leaves requires ingredients such as HPMC as a gelling agent, propylene glycol as a humectant, methylparaben and propylparaben as a preservative and aquadest as a solvent. HPMC is able to form a gel by absorbing solvents and forming a compact liquid mass. HPMC is used as a gelling agent because HPMC can form a clear and neutral gel and has a stable viscosity and has good drug delivery capability.¹⁸ In this study, three gel formulas (F1, F2, F3) were made by variation concentration of HPMC ie 3%, 5%, and 7%.




Figure 2. Submicroparticles Gel

Evaluation of Submicroparticles Gel:

Gel evaluation carried out in this study was an organoleptic test, pH test, homogeneity test, viscosity test, spreadability test, adhesion test, washability test, and cycling test. The results of the gel evaluation can be seen in Table 3.

Based on the result in Table 3, F1 was chosen as optimum formula because of the viscosity value is not too high so the resulting gel has a soft texture than the others.

***In-vitro* Antibacterial Activity Test:**

Antibacterial activity test was carried out by measuring the diameter of the inhibition zone. The results of the antibacterial activity test can be seen in Table 4.

203 **Table 4. Inhibition Zone Measurement Results for Gel Preparations**

Group	Average Inhibition Zone Diameter±SD (mm)
Negative Control	0±0*
Positive Control	13±1.73
F1	10.23 ± 0.69**
F2	5.78 ± 1.39*
F3	4.0 ± 0.0*
F4	1.89±1.54*

Notes: Statistical significance for each condition compared to positive control (*P<0.05; **P>0.05)

204
205
206
207
208 Based on the results of the antibacterial activity test above, F1 has antibacterial activity that is not significantly different from
209 the positive control while the other two formulas had significant differences. This proves that the use of HPMC concentration
210 has an effect on antibacterial activity.

211 **DISCUSSION:**

212 Ethanolic extract of purple sweet potato leaves obtained in this study was thick and blackish-brown extract with a yield of
213 9.39%. Ethanol extract in this study is proven to contain flavonoid compounds. This is in accordance with research conducted
214 by Kim et al²⁶, who succeeded in isolating the ethanolic extract of purple sweet potato leaves and obtained 3 types of
215 flavonoids, namely quercetin-3-O-β-D-glucopyranoside, isoquercitrin, and kaempferol 3-O-β-glucopyranoside. Quercetin-3-
216 O-β-D- glucopyranoside has been researched and proven to have antimicrobial activity.^{27,28} But, quercetin are compounds
217 that has low stability, poor water solubility and also low bioavailability.²⁹ Quercetin is thermally unstable, has low stability
218 when in aqueous solution so that it is easily hydrolyzed, and unstable in the presence of light and oxygen which causes
219 oxidation.³⁰ The stability of flavonoids can be improved by making them into the form of submicroparticles. In this study,
220 PLGA used as polymer and PVA used as stabilizer in submicroparticles. PLGA will encapsulate the ethanolic extract of
221 purple sweet potato leaves due to ionic interactions. Ionic interactions between the drug and polymer will result in higher
222 incorporation in the non- end-capped polymers while if the interaction shows hydrophobic features, the end-capped polymers
223 will show greater incorporation.³¹ The presence of PVA will increase the stability of submicroparticles because PVA acts as
224 an emulsifying agent. During the manufacture of submicroparticles, hydrophobic bonds will occur between the acetyl group
225 in PLGA and the hydrolyzed PVA.³² The presence of PVA will strengthen the encapsulation of PLGA-ethanolic extract of
226 purple sweet potato leaves.

227 The submicroparticles formulated into a gel preparation. In this study, submicroparticles formulated into three formula gel
228 with various concentrations of hydroxypropyl methylcellulose ie F1, F2, and F3 respectively 3%, 5%, and 7%. Based on the
229 results of the gel evaluation in Table 3, the organoleptic test, pH test, homogeneity test and cycling test between the three
230 formulas did not show a significant difference (p > 0.05). The gel produced from the three formulas is light yellow, has the
231 distinctive smell of the extract, has a semi-solid, and homogenous texture. The pH results of the three formulas show that the
232 pH is in the range of 7. Ideally, topical preparations have the same or similar pH value to the skin pH, namely 4.5-6.5 to avoid
233 irritation of the skin. However, the gel at a pH of 7 can still be tolerated by the skin and has not irritated. The resulting gel
234 pH tends to be high due to the influence of PVA on submicroparticles. PVA has a pH range of 5-8, the higher concentration
235 of PVA used, the more hydroxyl ions (OH-) from PVA are likely to be released and will shift the pH value to higher.³³ In the
236 viscosity test, the spreadability test, the adhesion test, and the washability test, there was a significant difference (p<0.05)
237 which indicated that there was an effect of the HPMC concentration used on the test. Increasing the amount of gelling agent
238 can strengthen the gel constituent matrix so the viscosity will increase. In addition, the presence of PVA also affects the

viscosity value. HPMC and PVA are polymers that can enter the cavity formed by water molecules so that hydrogen bonds occur between the hydroxyl (-OH) groups of the polymer and water molecules.^{34,35} This hydrogen bond plays a role in hydration in the swelling process.³⁶ The results of this viscosity test are inversely proportional to the results of the spreadability and washability tests but are directly proportional to the results of the adhesion test. The higher the viscosity result, the dispersibility and washability's result will smaller, but the adhesion will be stronger. The physical stability test (cycling test) also showed no significant difference between the three formulas ($p > 0.05$). The parameters observed in this test included organoleptic (color, smell, and texture) and changes in pH. The results of the observations obtained were that there was no organoleptic change in the three formulas, each formula had a light yellow color, the distinctive odor of the extract, semi-solid texture, and homogeneous. The three formulas do not indicate a syneresis phenomenon. Based on the gel evaluation and stability results, the F1 formula with a concentration of 3% HPMC was chosen as the optimum formula in making submicroparticle gel and used as the basis for making purple sweet potato leaves ethanolic extract gel which not formulated into submicroparticles for antibacterial activity testing.

Based on the results of the antibacterial activity test above, the three formulas gel containing the ethanolic extract of purple sweet potato leaves in the form of submicroparticles had antibacterial activity. F1 with the smallest HPMC concentration of 3% has the largest inhibition zone diameter compared to F2 and F3. The antibacterial activity result also depends on the viscosity of the gel. Viscosity of the gel can affect to the drug release from the vesicle. Increase in viscosity will resulted in slower drug release and the effect of antibacterial will be reduced.^{37,38} Viscosity of gel was directly proportional to the concentration of gelling agent used so in this study F1 with the smallest concentration had the largest inhibition zone diameter. The negative control did not produce inhibition because the component of submicroparticle gel without an active substance did not have antibacterial activity. The positive control of erythromycin gel had the largest inhibition zone diameter was 13 ± 1.73 mm. Erythromycin has antibacterial activity by binding to the ribosome subunit 50s so that protein synthesis is inhibited in bacteria.^{39,40}

Based on the data in Table 4, it can be seen that there is an increase in antibacterial activity after the ethanolic extract of purple sweet potato leaves is formed into submicroparticles. F4 or gel contain extract which not formulated into submicroparticles with an optimum concentration of HPMC (3%) can only cause inhibition of 1.67 ± 2.08 mm. In the research of Osuntokun et al.⁵ ethanolic extract of purple sweet potato leaves with a concentration of 50mg/ml gives an inhibition zone diameter of 10mm. In this study, the amount of extract used was 150mg/100ml and gave an inhibition zone of 1.67 ± 2.08 mm. However, when it was formed into submicroparticles, it was able to provide an inhibition zone diameter of 10.23 ± 0.69 mm. This proves that the submicroparticle can reduce dose usage.

Submicroparticle preparations can also increase the contact surface area because the resulting particle size is microparticle size so that the penetration of the active substance will increase and the antibacterial effect increases.⁴¹ Flavonoids can act as an antibacterial by inhibiting protein synthesis. Protein synthesis from bacteria occurs in the cytoplasm so that by increasing the penetration power of flavonoids into bacterial cells, more flavonoid molecules can inhibit protein synthesis in bacteria. In addition, the flavonoids formulated into submicroparticle preparations have better stability. Submicroparticle polymers protect flavonoids by encapsulating flavonoid molecules so they are protected from chemical disturbances. The mechanism for the release of active substances from PLGA submicroparticles can go through 4 ways, namely diffusion through water-filled pores, diffusion through the polymer, osmotic pumping, and erosion. In general, the mechanism for releasing the active substance from PLGA submicroparticles is by diffusion through water-filled pores. The water molecules around the

submicroparticle will enter the system and cause swelling so that the formation of pores which allows the encapsulated active substance to leave the submicroparticle system constantly until the erosion stage where the PLGA polymer as a submicroparticle maker begins to degrade.^{42,43} Submicroparticles reach the target site of bacterial also assisted by propylene glycol used in the gel. Propylene glycol as a penetration enhancer works by extracting lipids and proteins from the stratum corneum to form pores in the stratum corneum. Propylene glycol also changes the solubility parameters of the stratum corneum so that it affects the thermodynamic activity of the submicroparticles contained in the gel formula. This pushes the submicroparticles into the stratum corneum thereby increasing the penetration of the submicroparticles and increase antibacterial activity.^{44,45,46}

The results of the shapiro-wilk statistical analysis showed that the data for all groups were normally distributed. One Way ANOVA statistical analysis showed the significance value $p < 0.05$ so it can be concluded that there is a significant difference in the results of the antibacterial activity test of all treatments. In the post hoc test, it can be seen that there is a significant difference in the antibacterial activity of the positive control group with all treatment groups, except F1. From this result, we can conclude that F1 has similar antibacterial activity to positive control.

CONCLUSION:

Based on the results of the research, there was an effect of the HPMC concentration on the results of the physical evaluation of the gel preparation. The higher concentration of HPMC used will cause the resulting gel to be viscous so that it will affect the releasing drug from the gel matrix. F1 with a concentration of 3% HPMC was chosen as the optimum formula because physically good, stable, and produces the best antibacterial effect among the three formulas. In addition, the use of submicroparticle preparations as a carrier for ethanolic extract of purple sweet potato leaves has also been shown to increase antibacterial activity, be able to efficiently use dosages, and can be used as an alternative treatment for infection to prevent resistance.

CONFLICT OF INTEREST:

The authors have no conflicts of interest regarding this investigation.

REFERENCES:

1. Che Hamzah AM, Yeo CC, Puah SM, Chua KH, Chew CH. Staphylococcus aureus Infections in Malaysia: A Review of Antimicrobial Resistance and Characteristics of the Clinical Isolates, 1990-2017. Antibiotics (Basel). 2019; 8(3):128. doi:10.3390/antibiotics8030128
2. Foster TJ. Antibiotic resistance in Staphylococcus aureus. Current status and future prospects. FEMS Microbiology Reviews. 2017; 41 (3): 430 – 449. doi: 10.1093/femsre/fux007
3. Bezlon G, Shanmugha SD, Rinu ERE. Design and Stabilization of Natural Antibacterial Compound Allicin against Methicillin-Resistant Staphylococcus Aureus for Treatment as a Novel Antibiotic. Research J. Engineering and Tech. 2013; 4(4): 179-181.
4. Pochapski MT, Fosquiera EC, Esmerino LA, dos Santos EB, Farago PV, Santos FA, Groppo FC. Phytochemical screening, antioxidant, and antimicrobial activities of the crude leaves' extract from Ipomoea batatas (L.) Lam. Pharmacognosy Magazine. 2011; 7 (26): 165-170. doi: 10.4103/0973-1296.80682
5. Osuntokun OT, Babantude Y, Oledale MA, Fasila O. Components and bioactivity of Ipomoea batatas (L.) (Sweet Potato) ethanolic leaf extract. Asian Journal of Advanced Research and Reports. 2020; 10 (1): 10-26. doi:10.9734/AJARR/2020/v10i1/130232
6. Kusuma SAF, Wahyuni UT, Zuhrotun A. Evaluation of Antibacterial Activity of Indonesian Varieties Sweet Potato Leaves Extract from Cilembu Against Shigella Dysenteriae Atcc 13313. Asian J Pharm Clin Res. 2017; 10 (2): 377-380. doi:10.22159/ajpcr.2017v10i2.15773

- Manvar MN. Antibacterial Activity of Leaves and Flowers of *Ipomoea aquatica* Forsk. (Convolvulaceae). *Asian J. Pharm. Res.* 2018; 8(2): 94-98. doi: 10.5958/2231-5691.2018.00016.3
7. Patil SD, Hafizur MA, Priti A, Shelke PB, Yardi S. Synthesis and evaluation of novel Flavonoid derivatives for Antibacterial activity. *Asian J. Pharm. Res.* 2016; 6 (1): 27-30. doi: 10.5958/2231-5691.2016.00005.8
8. Dzoyem JP, Hamamoto H, Ngameni B, Ngadjui BT, Sekimizu K. Antimicrobial action mechanism of flavonoids from *Dorstenia* species. *Drug Discoveries & Therapeutics.* 2013; 7 (2): 66-72. doi: 10.5582/ddt.2013.v7.2.66
9. Takashi K, Yoshinobu T, Takahisa N, Takashi N, Hiroshi T, Shigetaka O. Acceptor specificity of cyclodextrin glucanotransferase from an alkalophilic *Bacillus* species and synthesis of glycosyl rhamnose. *Bioscience, Biotechnology, and Biochemistry.* 1996; 60 (7): 1176-1178. doi: 10.1271/bbb.60.1176
10. Pourcel L, Routaboul JM, Cheynier V, Lepiniec L, Debeaujon I. Flavonoid oxidation in plants: from biochemical properties to physiological functions. *Trends in Plant Science.* 2007; 12 (1): 29-36. doi:10.1016/j.tplants.2006.11.006
11. Zulham G, Wilar Y, Susilawati A, Subarnas AY, Chaerunisaa. Microparticles of Herbal Extracts with Antioxidant Activity. *Pharmacog J.* 2021; 13 (1): 285-295. doi: 10.5530/pj.2021.13.38
12. Sehar S, Khan AIH. Role of ZnO Nanoparticles for improvement of Antibacterial Activity in Food Packaging. *Asian Journal of Pharmaceutical Research.* 2021; 11(2):128-131. doi: 10.52711/2231-5691.2021.00024
13. Buzea C, Blandino IIP, Robbie K. Nanomaterial and nanoparticles: sources and toxicity. *Biointerphase.* 2007; 2: 170-172. doi: 10.1116/1.2815690
14. Somayadineshraj D, Shanmugapriyan R, Lakshmanaswamy R, Nishiha CS, Dooslin MBV, Muralidharan P. Synthesis and Characterization of MgO nanoparticles Coated by polyvinyl alcohol for Biomedical applications. *Research J. Pharm. and Tech.* 2018; 11(11): 4997-5000. doi: 10.5958/0974-360X.2018.00911.3
15. Chetan MP, Manan AP, Nikhil PP, Prajapati PH, Patel CN. Poly Lactic Glycolic Acid (PLGA) As Biodegradable Polymer. *Research J. Pharm. and Tech.* 2020; 3(2): 353-360.
16. Irfan M, Saiyyad DS, Bhambere, Sanjay K. Formulation and Optimization of Silymarin Loaded PLGA Nanoparticle for liver targeting. *Asian J. Pharm. Tech.* 2017; 7 (4): 209-220. doi: 10.5958/2231-5713.2017.00032.0
17. Verma A, Singh VS, Kaur R, Jain UK. Formulation and Evaluation of Clobetasol Propionate Gel. *Asian Journal of Pharmaceutical and Clinical Research.* 2013; 6 (5): 15-18. ISSN - 0974-2441
18. Rowe RC, Sheskey PJ, Quinn ME. Handbook of pharmaceutical excipients, Sixth Edition. UK, London: Pharmaceutical Press and American Pharmacists Association; 2009.
19. Ghosal K, and Nanda A. Development of diclofenac potassium gel from hydrophobically modified HPMC. *Iranian Polymer Journal.* 2013; 22 (6): 1-24. doi: 10.1007/s13726-013-0145-3
20. Binder L, Mazal J, Petz R, Klang V, Valenta C. The role of viscosity on skin penetration from cellulose ether-based hydrogels. *Skin Research and Technology.* 2019; 25 (5): 725-734. doi: 10.1111/srt.12709
21. Indarti K, Apriani EF, Wibowo AE, Simanjuntak P. Antioxidant Activity of Ethanolic Extract and Various Fractions from Green Tea (*Camellia sinensis* L.) Leaves. *Pharmacognosy Journal.* 2019; 11 (4): 771-776. doi: 10.5530/pj.2019.11.122
22. Mardiyanto. Investigation of nanoparticulate formulation intended for caffeine delivery into hair follicle. Dissertation. Department of Pharmacy, Faculty of Science, Saarland University, Saarbruecken. Germany; 2013.
23. Apriani EF, Rosana Y, Iskandarsyah I. Formulation, characterization, and in vitro testing of azelaic acid ethosome-based cream against *Propionibacterium acnes* for the treatment of acne. *J Adv Pharm Technol Res.* 2019; 10 (2): 75-80. doi: 10.4103/japtr.JAPTR_289_18
24. Ahmad I, Owais M, Shahid M, Aqil F. Combating Fungal Infections New York, USA : Springer; 2010.
25. Kim HJ, Jin C, Lee YS. Isolation and Antioxidative Activities of Caffeoylquinic Acid Derivates and Flavonoid Glycosides from Leaves of Sweet Potato (*Ipomoea batatas* L.). *The Journal of Applied Pharmacology.* 2007; 15: 46-51. doi: 10.4062/biomolther.2007.15.1.046
26. Islam MM, Akter S, Al-Amin M, Haque MM. Isolation of Quercetin-3-O-beta-D-glucopyranoside from the Leaves of *Azadirachta indica* and Antimicrobial and Cytotoxic screening of the Crude Extracts. *Dhaka University Journal of Science.* 2012; 60 (1): 11-14. doi:10.3329/dujs.v60i1.10328
27. Sholkamy EN, Ahmed MS, Yasser MM, Mostafa AA. Antimicrobial quercetin 3-O-glucoside derivative isolated from *Streptomyces antibioticus* strain *ess_amA8*. *Journal of King Saud University.* 2020; 32 (3): 1838-1844. doi: 10.1016/j.jksus.2020.01.026
28. Wang W, Sun C, Mao L, Ma P, Liu F, Yang J, Gao Y. The biological activities, chemical stability, metabolism and delivery systems of quercetin: A review. *Trends in Food Science & Technology.* 2016; 56: 21-38. doi: 10.1016/j.tifs.2016.07.004
29. Cunico LP, Cobo AM, Al-Hamimi S, Turner C. Solubility and Thermal Degradation of Quercetin in CO₂-Expanded Liquids. *Molecules.* 2020; 25: 5582-5591. doi: 10.3390/molecules25235582

30. Lagreca E. Onesto V. Natale CD. Manna SL. Netti PA. Vecchione R. Recent advances in the formulation of PLGA microparticles for controlled drug delivery. *Progress in Biomaterials*. 2020; 9: 153–174. doi: 10.1007/s40204-020-00139-y
31. Stromberg ZR. Phipps ML. Magurudeniya HD. Pedersen CA. Rajale T. Sheehan CJ. Courtney SJ. Bradfute SB. Hrabec P. Rush MN. Kubicek-Sutherland JZ. Martinez JS. Formulation of stabilizer-free, nontoxic PLGA and elastin-PLGA nanoparticle delivery systems. *International Journal of Pharmaceutics*. 2021; 597: 1-11. doi: 10.1016/j.ijpharm.2021.120340
32. Wiśniewska M. Chibowski S. Urban T. Effect of the type of polymer functional groups on the structure of its film formed on the alumina surface—suspension stability. *React Funct Polym*. 2012; 72: 791–798. doi: 10.1016/j.reactfunctpolym.2012.08.005
33. Sharma N. Madan P. Lin S. Effect of process and formulation variables on the preparation of parenteral paclitaxel-loaded biodegradable polymeric nanoparticles: A co-surfactant study. *Asian journal of pharmaceutical sciences*. 2016; 11 (3): 404-416. doi: 10.1016/j.ajps.2015.09.004
34. Zolnik BS. Burgess DJ. Effect of Acidic pH on PLGA Microsphere Degradation and Release. *Journal of Controlled Release*. 2007; 122: 338-344. doi: 10.1016/j.jconrel.2007.05.034
35. Das N. Bose C. Tripathi N. Maitra S. Progress in the development of gelling agents for improved culturability of microorganisms. *Frontiers in Microbiology*. 2015; 6 (698): 1-7. doi: 10.3389/fmicb.2015.00698
36. Pacharne PS. Patil JR Chaudhari SP. Ratnaparkhi MP. Design, Development and formulation of Topical anti arthritic gel using different gelling agents. *Elixir Pharmacy*. 2013; 62: 17599-17603.
37. Yener G. Dal O. Üner M. Effect of Vehicles on Release of Meloxicam from Various Topical Formulations. *The Open Drug Delivery Journal*. 2009; 3: 19-23. doi: 10.2174/1874126600903010019
38. Jednacak T. Mikulandra I. Novak P. Advanced Methods for Studying Structure and Interactions of Macrolide Antibiotics. *International Journal of Molecular Sciences*. 2020; 21 (7799): 1 – 26. doi: 10.3390/ijms21207799
39. Singh RK. Garg R. Formulation and Evaluation of Erythromycin Estolate Loaded Drug Balls. *Research J. Pharm. and Tech*. 2020; 13(1): 282-286 doi: 10.5958/0974-360X.2020.00057.8
40. Mohanraj VJ. Chen Y. Nanoparticles-A review. *Trop J Pharm Res*. 2006; 5(1): 561-573. doi: 10.4314/tjpr.v5i1.14634
41. Fredenberg S. Wahlgren M. Reslow M. Axelsson A. The mechanisms of drug release in poly(lactic-co-glycolic acid)-based drug delivery systems-A review. *International Journal of Pharmaceutics*. 2011; 415 (2011): 34-52. doi: 10.1016/j.ijpharm.2011.05.049.
42. Vimal P. Bhavesh A. Anand D. Manish G. Adil P. A Review on Long Acting PLGA Based in Situ Forming Microparticles Formulation for a Novel Drug Delivery System. *Res. J. Pharm. Dosage Form. and Tech*. 2016; 8(2):127-134. doi: 10.5958/0975-4377.2016.00017.3
43. Hakue T. Talukder MMU. Chemical Enhancer: A simplistic way to modulate barrier function of the stratum corneum. *Adv. Pharm. Bull*. 2018; 8 (2):169-179. doi: 10.15171/apb.2018.021
44. Vinod K. Aditi S. Study of Antimicrobial activity of Star Anise loaded poly (DL-lactide-co-glycolide) nanoparticles. *Research J. Pharm. and Tech*. 2019; 12 (2): 499-507. doi: 10.5958/0974-360X.2019.00088.X
45. Vinod K. Aditi S. Synthesis, Characterization, Antimicrobial activity and Release Study of Cinnamon loaded poly (DL-lactide-co-glycolide) Nanoparticles. *Research J. Pharm. and Tech*. 2019; 12(4):1529-1535. doi: 10.5958/0974-360X.2019.00253.1

403

404

405

406

407

408



avpublication avpublication <avpublications@gmail.com>

kepada saya ▾

Rab, 15 Sep 2021, 19.33



Terjemahkan ke Indonesia



Dear Author,
Your article has been Sent to Reviewers. Comments still awaited.
thanks
editor.

A and V Publications,

RJPT House,

Lokmanya Grih Nirman Society, Rohanipuram, In-front of Sector- 1,

Pt. Deendayal Upadhyay Nagar,

Raipur 492 010. (CG) India.

Phone No. 9406051618.

E. mail: avpublications@gmail.com;

www.anvpublications.org

www.ajrconline.org

www.rjptonline.org

www.asianpharmaonline.org

www.enggresearch.net

Date: November 03, 2021

Research Journal of Pharmacy and Technology

Paper ID:21713171640045385

Author's Name:Mardiyanto Mardiyanto, Elsa Fitria Apriani, Muhammad Hafizhaldi Alfarizi

Paper Title: FORMULATION AND IN-VITRO ANTIBACTERIAL ACTIVITY OF GEL CONTAINING ETHANOLIC EXTRACT OF PURPLE SWEET POTATO LEAVES (*Ipomoea batatas* (L.)) LOADED POLY LACTIC CO-GLYCOLIC ACID SUBMICROPARTICLES AGAINST *Staphylococcus aureus*



Acceptance of Manuscript

With reference to your article titled '**FORMULATION AND IN-VITRO ANTIBACTERIAL ACTIVITY OF GEL CONTAINING ETHANOLIC EXTRACT OF PURPLE SWEET POTATO LEAVES (*Ipomoea batatas* (L.)) LOADED POLY LACTIC CO-GLYCOLIC ACID SUBMICROPARTICLES AGAINST *Staphylococcus aureus***' Author by **Mardiyanto Mardiyanto, Elsa Fitria Apriani, Muhammad Hafizhaldi Alfarizi**. We wish to bring to your kind notice the following:

- We acknowledge the receipt of the above mentioned article.
- The above mentioned article has been sent to the reviewer of expert comments.
- The above mentioned article have been accepted for publication in Year : 2022, Vol: 15, Issue: 8 of the journal.

Thanking you for submission of manuscript.

Yours sincerely,



Sir Baharwal
Editor in Chief

Note: -

1. To verify the originality of this certificate please visit the following link:

http://www.anvpublication.org/Download_Acceptance.aspx?PID=21713171640045385

Aplikasi Setoran/Transfer/Kliring/Inkaso
Deposit/Transfer/Clearing/Collection Application



Bismillaahirrahmaanirrahiim

Kepada to PT Bank Syariah Indonesia

Harap dilakukan transaksi berikut Please do this transaction:

Cabang Branch Palembang

No.: C 000997

Tanggal Date 15-11-2021

Jenis transaksi
Transaction

☐ Setoran
Deposit

☐ Pemindahbukuan
Overbooking

☐ RTGS
RTGS

☐ Kliring-Inkaso
Clearing-Collection

☒ SWIFT
SWIFT

☐ Wesel
Demand Draft

☐ Lainnya
Others

harap ditulis dengan huruf cetak fill in with block letters

VALIDASI Validation

ACB4 D 0001/11/20
ACB4 K USD1341300010019
204 K PL52710
KC PALEMBANG BSM199716526
15 NOV 2021 15 NOV 2021 12:19

ELSA FITRIA APRIANI
Outgoing TT
Fee Atas Transaksi SWIFT
FT21319D81LS 0000997

IDR 2,877,000.00
USD 200.00
IDR 355,875.00

PENERIMA
Beneficiary

☐ Penduduk
Resident

☒ Bukan Penduduk
Non Resident

Nama
Name

A and V Publications

Nomor Rekening
Account Number

208636 20032

Bank
Bank

Hala Bank of India

Alamat & Nomor Telepon
Address & Telephone Number

Ravi Shankar University Raipur
Swift code: SPININ BB 646

MATA UANG
Currency

☐ Rupiah
Local Currency

☒ Valuta Asing
Foreign Currency

USD

BERITA UNTUK PENERIMA Message for Beneficiary

Article Charge No 2173171640045385

diisi oleh Bank filled out by bank

Jumlah Transfer Amount of Tranfer	
Komisi Commission	
Biaya Pengiriman (SWIFT/RTGS/SKN)	
Biaya Koresponden Correspondent Change	
Sub Total	PT BANK SYARIAH INDONESIA
Kurs Rate	KC Palembang Damang
Total	

Pemohon telah membaca berita memahami dan dengan ini menyetujui sepenuhnya syarat-syarat dan ketentuan yang tercantum dibalik formulir transaksi ini Applicant have read and understand, and with this applicant unconditionally accept all term and condition on the reserve of this transaction form

Pengesahan Bank Bank's Authorization

Tanda Tangan Pemohon Application Signature

Teller

PENGIRIM
Applicant

☒ Penduduk
Resident

☐ Bukan Penduduk
Non Resident

☐ Nasabah
Customer

☐ Bukan Nasabah
Walk in Customer

Nama
Name

Elsa Fitria Apriani

Nomor Identitas
Identity Number

3201015404920004

Alamat & Nomor Telepon
Address & Telephone Number

Paramarua Nagan Puth No 83
082261067221

SUMBER DANA
TRANSAKSI
Source of Fund

☐ Tunai
Cash

☐ Cek/Bilyet Giro
Cheque

☒ Debet Rekening :
Debet Account :

9481714700

Bank Tertarik Drawee Bank	Nomor Cek/BG Cheque Number	Valuta Currency	Nominal Amount

Jumlah
Amount

200 USD

Terbilang
In Words

Two Hundred US Dollars

BIAYA TRANSAKSI
Handing Charge

☐ Tunai
Cash

☒ Debet Rekening :
Debet Account :

9481714700

Biaya Bank Koresponden
Correspondent Change

☐ Pengirim
Applicant

☐ Penerima
Beneficiary

☐ Lainnya
Others

TUJUAN TRANSAKSI Underlying Transaction

Perhatian : Apabila penerima adalah bukan Nasabah/Walk In Customer maka wajib mengisi alamat dan nomor telepon penerima

AG 001
Lembar 1 : untuk Teller

