

Hasil IThenticate_JAS_Fitra Yosi

By Fitra Yosi

WORD COUNT

9757

TIME SUBMITTED

14-JUL-2025 03:33PM

PAPER ID

117212760

Lactation-related dynamics of bacterial and fungal microbiomes in feces of sows and gut colonization in suckling and newly weaned piglets

Fitra Yosi,^{1,†,‡} Frederike Lerch,^{1,†} Julia C. Vötterl,^{1,†} Simone Koger,^{1,§} Doris Verhovsek,¹ and Barbara U. Metzler-Zebeli^{1,†,‡,¶}

¹Centre for Veterinary Systems Transformation and Sustainability, Clinical Department for Farm Animals and Food System Science, University of Veterinary Medicine Vienna, Vienna 1210, Austria

²Christian-Doppler Laboratory for Innovative Gut Health Concepts of Livestock, Institute of Animal Nutrition and Functional Plant Compounds, Centre for Animal Nutrition and Welfare, University of Veterinary Medicine Vienna, Vienna 1210, Austria

³Department of Animal Science, Faculty of Agriculture, University of Sriwijaya, Palembang 30662, Indonesia

⁴Centre for Animal Nutrition and Welfare, Clinical Department for Farm Animals and Food System Science, University of Veterinary Medicine Vienna, Vienna 1210, Austria

⁵Clinical Centre for Population Medicine in Fish, Pig and Poultry, Clinical Department for Farm Animals and Food System Science, University of Veterinary Medicine Vienna, Vienna 1210, Austria

[†]Corresponding author: barbara.metzler@vetmeduni.ac.at

Abstract

Changes in the gut microbial composition of sows during lactation may influence the gut microbial colonization in their offspring, for which less information was available in the literature. This study aimed to assess: 1) the changes that occur in the bacterial and fungal communities in sow feces during the 28-d lactation period as well as in gastric and cecal digesta of piglets until one week after weaning, and 2) bacterial and fungal taxa in cecal digesta of the piglets postweaning that associate with fecal consistency. Aside from sow milk, piglets had access to creep feed from day of life (DoL) 3. Fecal samples from sows for microbial analysis were collected ($n = 20$) on days postpartum (DPP) 1, 6, 13, 20, and 27, as well as from weaned piglets for fecal scoring on DoL 30 and 34. Gastric and cecal digesta of piglets was collected on DoL 3, 7, 14, 21, 28, 31, and 35 ($n = 5/\text{sex}/\text{DoL}$). Progressing lactation affected bacterial and fungal communities in sow feces, including 10.3- and 3.0-fold increases in the relative abundances of *Lactobacillus* from DPP1 to 6 and *Kazachstania* from DPP1 to 13, respectively ($P < 0.001$). Although time- and gut-site-related differences existed, bacterial and fungal taxa found in sow feces were also present in gastric and cecal digesta of piglets, which supports their role in gut colonization in neonatal piglets. In piglets, bacterial and fungal alpha-diversities showed certain fluctuations during the suckling period, whereby weaning affected the fungal than bacterial diversity at both gut sites ($P < 0.05$). At both gut sites, *Lactobacillus* largely increased from DoL3 to 7 and remained a dominating taxon until DoL35 ($P < 0.05$). Postweaning, plant-glycan fermenters (e.g., *Prevotella*) seemed to replace milk-glycan fermenting *Fusobacterium* and *Bacteroides* ($P < 0.05$). In gastric and cecal digesta, *Kazachstania*, *Tausonia*, *Candida*, and *Blumeria* were dominating fungi from DoL3 to 35, with *Kazachstania* becoming even more dominant postweaning ($P < 0.001$). Fecal consistency was softer on DoL34 than 30 ($P < 0.05$). Correlation analysis identified that softer feces were linked to the relative abundances of plant-glycan and proteolytic bacterial taxa including pathobionts (e.g., *Clostridium sensu stricto*) in the cecum on DoL34. However, the potential association between cecal mold and plant-pathogenic fungi *Talaromyces*, *Mrkia*, and *Blumeria* and softer feces are worth investigating in the future in relation to (gut) health of piglets.

Lay Summary

During the suckling phase, piglets are commonly housed in the same pen as their mother sows until weaning. Consequently, the mother's feces are a rich source of microbes for the colonization of the neonatal gut. In the past, mainly maturational changes in the bacterial community of piglet's gut have been investigated, whereas other microbial groups, such as fungi, received less attention. In the present study, we could show that the bacterial and fungal communities in sow feces changed with progressing lactation and that bacterial and fungal taxa present in sow feces were also present in gastric and cecal digesta of the suckling piglets. These findings support the importance of maternal microbes for the gut colonization of neonatal piglets. In both, bacterial and fungal communities, weaning left a characteristic mark on the overall community structures and relative taxa abundances, which can be explained by the withdrawal of sow milk and low feed intake. Correlation analysis also revealed potential bacterial and fungal marker candidates for softer and firmer feces.

Keywords: bacteriome, lactation, mycobiome, piglet, sow, weaning

Abbreviations: ASV, amplicon sequence variant; DADA2, fastq Amplicon Denoising Algorithm 2; DoL, day of life; DPP, day postpartum; NMDS, non-metric multidimensional scaling; pdiff, probability of difference; PERMANOVA, permutational multivariate analysis of variance; w/v, weight/volume; w/w, weight/weight

Received May 9, 2024 Accepted October 25, 2024.

© The Author(s) 2024. Published by Oxford University Press on behalf of the American Society of Animal Science.

This is an Open Access article distributed under the terms of the Creative Commons Attribution License (<https://creativecommons.org/licenses/by/4.0/>), which permits unrestricted reuse, distribution, and reproduction in any medium, provided the original work is properly cited.

Introduction

Postnatal microbial colonization is a substantial contributor to gut maturation, including immune and barrier functions (Everaert et al., 2017). Microbes that colonize the piglet's gastrointestinal tract after birth originate from the sow (i.e., vagina, skin/nipple surface, colostrum, milk, and feces), environment, and handling farm staff (Jost et al., 2014; Chen et al., 2018). During the suckling period, piglets are commonly housed in farrowing pens with their sows until weaning (Nowland et al., 2019). Hence, changes in the gut microbial composition of the sow that occur during lactation may influence the gut microbial colonization in their offspring. Despite increasing knowledge on the gut microbial development in piglets, alterations in the gut microbial composition of the dam over the course of lactation have rarely been investigated so far.

Moreover, knowledge of the postnatal fungal colonization of the piglet's gut and whether this development is driven by the fecal mycobiome of the sow is still scarce. The stomach of suckling and weaned piglets contains a diverse and complex microbiome (Mann et al., 2014; Lerch et al., 2023), which contributes to the degradation of nutrients and the production of fermentation metabolites. Although some data for the postnatal development of bacteria in the stomach exists (Lerch et al., 2023), the development of gastric fungal communities has been poorly described. Evidence from our group supports the importance of investigating fungal communities in the neonatal period (Lerch et al., 2023), as some of the fungi found in cecal digesta represented plant pathogens and mold fungi (Yeh et al., 2021), which may compromise the development of the immune system (Wang et al., 2023). This indicates the need for monitoring the development of the gastrointestinal microbial communities in healthy piglets from pre- and postweaning more closely.

Scoring the fecal consistency in piglets is also helpful for identifying eubiotic and dysbiotic gut conditions. Recently, we have shown that feces of different colors and consistencies differ in their bacterial composition in suckling and newly weaned piglets (Metzler-Zebeli et al., 2023). However, feces are mostly representative of the distal large intestine of piglets; therefore, targeting the microbial composition in the more proximal parts of the gut (e.g., cecum) and linking it to the fecal score may provide further marker taxa for eubiotic and dysbiotic gut conditions. In particular, the relationship between intestinal fungi and fecal score has been little investigated so far.

The objectives of this study were to assess 1) the changes that occur in the bacterial and fungal communities in sow feces during the lactation period as well as in gastric and cecal digesta of piglets from the day of life (DoL) 3 until one week after weaning; and 2) bacterial and fungal taxa in cecal digesta of piglets postweaning that associate with fecal consistency. Our research was based on the following hypothesis: 1) microbial communities in the piglet's stomach and cecum would comprise similar bacterial and fungal taxa to that of the sow feces during the suckling phase but would diverge postweaning and 2) the association of fecal scores with microbial taxa in cecal digesta would allow the identification of taxa linked to a lower (dysbiosis) and higher (eubiotics) gut homeostatic state postweaning.

92

Materials and Methods

Ethical statement

All procedures requiring animal handling and treatment have been approved by the institutional ethics committee of

the University of Veterinary Medicine Vienna and National authority in accordance with the Law for Animal Experiments in Austria (GZ 2020-0.437.208).

Animals, housing, and experimental procedure

The pig experiment was conducted at the pig facility of the University of Veterinary Medicine Vienna (VetFarm) under production conditions, consisting of 2 consecutive replicate batches. In each replicate batch, 10 sows (Large White) and their litters (Large White × Piétrain) were used. The average litter size was 13.9 ± 1.7 (SD) live piglets at birth across both replicate batches. The total number of piglets born alive was 143 and 134 in replicate batches 1 and 2, respectively.

The experiment lasted from 26 d prior to farrowing, throughout the 28-d lactation period to 7 d after weaning. At the pig facility, sows and their litters were handled according to standard procedures at the pig facility. Sows were group-housed in pens during gestation but had access to individual feeders. Five days before farrowing, sows were transferred to separate farrowing pens (BeFree, Schauer Agrotonic, Prambachkirchen, Austria; 2.3×2.6 m in size), which were equipped with a feeder, bowl drinker, and hayrack for the sow (for environmental enrichment and nesting behavior), as well as a round feeder, small bowl drinker, and nest with heated flooring for the piglets. All sows gave birth within 48 h and were not restrained during the farrowing process and the whole lactation period. Piglets were supplemented with iron by injection on DoL4 (2 mL of Ferriphor 100 mg/mL, OGRIS Pharma Vertriebs-GmbH, Wels, Austria), followed by castration of male piglets on DoL11 (general sedation with Stresnil 40 mg/mL, 0.025 mL/kg body weight, Elanco Tiergesundheit AG, Basel, Switzerland and Narketan 100 mg/mL, 0.1 mL/kg body weight, Vetoquinol Österreich GmbH, Vienna, Austria). On DoL17, piglets were vaccinated (1 mL Ingelvac CircoFLEX and 1 mL Ingelvac MycoFLEX, both from Boehringer Ingelheim GmbH, Ingelheim/Rhein, Germany). On DoL28, sows were removed from the farrowing pens and all remaining piglets were transferred to rearing pens in an outdoor climate house with a heated lying area. Each rearing pen had a size of 3.3×4.6 m each and was equipped with one round feeder, nipple and bowl drinkers, and a heated nest. The animal's health was monitored daily throughout the experiment. Piglets with health problems were removed from the experiment. Water was freely available to sows and piglets throughout the experiment.

Feed and feeding

The feeding protocol of gestating and lactating sows as well as of suckling and weaned piglets corresponded to the standard protocol at the pig facility. From 26 to 5 d prior to farrowing, sows were provided with a gestation diet in the morning (08:00 hours) and afternoon (14:30 hours; 3 to 4 kg/meal; Königshofer GmbH, Ebergassing, Austria; Supplementary Table S1 and S2). After moving the sows to the farrowing pens 5 d before the farrowing date, the sows were offered a lactation diet (approximately 3 kg/meal; Königshofer GmbH, Ebergassing, Austria; Supplementary Table S1 and S2) in the morning and afternoon (08:00 and 14:30 hours). Additionally, sows received 500 g of linseeds soaked in water once a day for 5 d prior to farrowing to avoid constipation. After farrowing, the feed amount of the sows was gradually increased (approximately 4 to 9 kg/meal) according to the regular feeding protocol.

Litters received creep feed that was manually prepared at least twice daily (08:00 and 15:00 hours) from DoL3 to 28. In the first 3 wk of life, the creep feed was a commercial milk replacer (NuriStart Sweet, BIOMIN Holding GmbH, Part of DSM-Firmenich, Getzersdorf 71 Austria; Supplementary Table S2), which was prepared according to the manufacturer's instructions. The milk replacer was offered in liquid form, by mixing the powder with warm water (45 °C) at a ratio of 500 g/L (w/v) by hand. The milk replacer was always prepared freshly before feeding and the stainless steel feeders were cleaned before re-filling them. Each litter received a minimum amount of 1,000 mL per day (500 mL at 08:00 and 500 mL at 15:00 hours), and more when piglets finished their portion. From DoL3 to 23, the piglets were given 100% of milk replacer. Then, the milk replacer was gradually mixed with the prestarter diet from DoL24 to 26 (Königshofer GmbH, Ebergassing, Austria; Supplementary Table S1 and S2), starting with a ratio of 70:30 (w/w) on DoL24, 50:50 (w/w) on DoL25, and 30:70 (w/w) on DoL26, respectively, and provided in mash form. After that, the litters were fed 100% of the prestarter diet as mashed on DoL27 and in dry form 69 weaning from DoL28 to 35. All diets used in the study met or exceeded the current recommendations for nutrient requirements (NRC, 2012). Leftover creep feed and spills were collected to estimate the creep feed intake during the suckling phase. The feed intake after weaning could not be estimated due to the set-up of the round feeders in the rearing barn.

Body weight measurement, fecal score, and collection

Piglets were weighed at 8 time points, immediately after birth, DoL4, 6, 13, 20, 27, 30, and 34. Freshly defecated feces of sows or feces after rectal stimulation were collected on days postpartum (DPP) 1, 6, 13, 20, and 27 for microbiome analysis. To avoid contamination from the floor, only the inside part of the defecated feces was used and homogenized with a sterile spatula before placing them 10 to cryo tubes. The tubes were kept on ice before storage at -80 °C until further analysis. Fecal samples from weaned piglets were collected by means of rectal stimulation on DoL30 and DoL34 to assess fecal consistency. To obtain the samples, the inner anal sphincter was stimulated by inserting a sterile cotton tip and gently rotating it. The consistency was scored according to 0 (hard and dry balls), 1 (clearly defined shape with cracks), 2 (pasty; moist feces with no cracks), 3 (very wet but not liquid feces), 4 (liquid feces with minimal consistency), and 5 (entirely liquid feces). Fecal scores of 4 and 5 were considered as clinical signs of diarrhea.

Gut sampling

In each of the 2 replicate batches, gut sampling took place on DoL3, 7, 14, 21, 28, 31, and 35. On each sampling day, 5 female and 5 male piglets were used for invasive sampling in each replicate batch. From each litter, 1 piglet (with alternating sexes on the consecutive sampling days) was selected per sampling day based on having average body weight within the litter. Prior to slaughter, piglets were weighed and anesthetized 44 the ear vein with azaperone (Stresnil 40 mg/mL, 0.025 mL/kg body weight, Elanco Tiergesundheit AG, Bad 27 mburg, Germany) and ketamine (Narketan 100 mg/mL, 0.1 mL/kg body weight, Vetoquinol Österreich GmbH, Vienna, Austria 27). Afterward, piglets were euthanized with embutramide (161,

0.1 mL/kg body weight, Intervet GesmbH, Vienna, Austria) via intracardiac 51 injection. Piglets were bled by cutting the neck. Then, the abdomen was opened and the entire gut was removed aseptically. The stomach and cecum were identified, clamped, and separated. Digesta samples were then collected from both gut segments. 30 homogenized digesta samples for microbiome analysis were snap-frozen in liquid nitrogen and stored at -80 °C.

DNA extraction, 16S rRNA and ITS sequencing, and bioinformatics

For DNA extraction 3, the protocol as described in Lerch et al. (2023) was used (DNeasy PowerSoil Pro Kit; Qiagen, Hilden, Germany) with the same modifications including a heating step and mechanical lysis. The concentration of DNA 19 each extract was measured with a Qubit fluorometer (Qubit 4 Fluorometer, Thermo Fisher Scientific, USA) using the Qubit dsDNA HS Assay Kit 8 Thermo Fisher Scientific Inc., Waltham, MA, USA). Targeted 16S rRNA gene (V3-V4 hypervariable region) and ITS2 amplicon sequencing were performed in an external laboratory (Novogene, Cambridge, UK). In order to do so, aliquots of the DNA extracts were sent for library preparation (NEBNext Ultra 4 DNA Library Prep Kit, Illumina, San Diego, CA, USA). The 16S rRNA gene amplicon was amplified using primers 341F-ill (5'-CCTACGGGNGG-CWGCAG-3') and 805R-ill (5'-GACTA 29 VGGGTATCTA-ATCC-3'; Klindworth et al., 2013), and the ITS2 region was amplified using primers ITS3 (5'-GCATCGATGAAGAACG-74 GC-3') and ITS4 (5'-TCCTCCGCTTATTGATATGC-3'; Op De Bee 55 et al., 2014). Equimolar pools of samples were sequenced to generate 250 bp paired-end raw reads in the Novaseq 6000 platform (Illumina). Demultiplexing and trimming of the raw sequences were performed by Novogene.

Trimmed reads for the 16S rRNA and fungal ITS amplicons 5 were processed, aligned, and classified independently using the Divisive Amplicon Denoising Algorithm 1 2 (DADA2; version 1.26.0) in R studio (version 1.4.1106; Callahan et al., 2016). The forward and reverse read quality profiles were separately examined. To account for the decrease in the quality score of the subsequent nucleotides, the total length of forward and reverse reads was truncated to 220 nucleotides with a maximum error rate of 5 for both forward and reverse reads (truncQ=5) using the 'filterAndTrim' function. This function was used to pre-filter sequences in order to remove reads with ambiguous bases for both bacterial and fungal sequences. For the fungal ITS amplicons, the first 10 nucleotides of each read were trimmed to account for the decrease 65 quality score of the following nucleotides, and a minimum length of 50 nucleotides was enforced to eliminate very short sequences. Reads with ambiguities were removed from both amplicon sets, as were reads that exceeded the probabilistic estimated error of 2 nucleotides for the ITS reads. For both the 16S rRNA and ITS amplicons, amplicon sequence variants were inferred after 61 replication of the filtered data and estimation of error rates (Callahan et al., 2016). The inferred forward and reverse sequences were then merged, with paired sequences that did not perfectly match removed to control for residual errors, and a sequence table was constructed. The removeBimeraDe-novo () function was used to remove chimeras, and taxonomy was assigned using the SILVA 138.1 ribosomal RNA database for bacteria (Quast et al., 2012) and the UNITE ITS database (version 9.0) for fungi (Nilsson et al., 2019) with a 3% dissimilarity threshold. The raw sequence counts from the taxa

tables were collapsed and compositionally normalized such that each sample summed to 1. The relative abundances at genus rank were statistically analyzed as described below. Alpha-diversity (Shannon, Simpson, observed ASV) analysis was performed using phyloseq (version 1.42.0). For beta-diversity analysis, statistical assessment of dissimilarity matrices (Bray–Curtis) was performed using the ‘adonis2’ function in the R package ‘vegan’ (version 2.6.4; Oksanen et al., 2022), separately for the bacterial and fungal composition. The permutational multivariate analysis of variance (PERMANOVA) was used on the Bray–Curtis distance matrices to assess the dissimilarities between the bacterial and fungal community structures in sows’ feces postpartum and in the gastric and cecal digesta of piglets on the various sampling days during the suckling period. The statistical significance was determined after 999 random permutations. The 2-dimensional non-metric multidimensional scaling (NMDS) ordination plots were generated using the ‘metaMDS’ function and the ggplot2 package was used to visualize the clustering of bacteriomes and mycobiomes in gastric and cecal digesta according to sample type and age as well as between sow feces and piglet digesta samples of piglets.

The datasets generated for this study were deposited into the NCBI Bioproject databank under accession number PRJNA1103974.

Statistical analyses

The Shapiro–Wilk test with the UNIVARIATE procedure in SAS (version 9.4; SAS Institute, Inc., Cary, NC) was used to test the normal distribution of the residuals of all variables. The residuals were transformed using the Boxcox method and the Transreg procedure in SAS if they were not normally distributed. All data, both from sows, i.e., the gut microbiome (relative abundance of genera), and from piglets, i.e., body weight, the gut microbiome (relative abundance of genera), and fecal score, were subjected to ANOVA using the MIXED procedure in SAS. Across datasets for sow feces and digesta samples from piglets, bacterial >0.2% of all reads and fungal taxa >0.05% of all reads were taken into consideration. The ranked relative abundances were analyzed separately for each genus in SAS. Repeated measures were used to investigate effects on the diversity estimators and relative genera abundances with progressing lactation of the sows (DPP) and increasing age of the piglets (DoL). For data for the diversity estimators and relative genera abundances of the sow, the model included the fixed effects of DPP, replicate batch, litter, and the respective 2- and 3-way interactions. The model for data for the diversity estimators and relative genera abundances of the piglets included the fixed effects of sex, replicate batch, DoL, litter, and the respective 2- and 3-way interactions. For most parameters in piglets, differences between sexes were not detected and were excluded from the final model. For the growth performance parameters of the piglets, the fixed effects were replicate batch, sex, and their interaction, as well as litter size on the weighing day and date of birth were considered covariates. The sow and piglet represented the experimental unit. Degrees of freedom were approximated by the Kenward–Rogers method (ddfm = kr). Data were reported as the least-square means \pm standard errors of the mean (SEM). Multiple pairwise comparisons among least-square means were performed using the pdiff statement. A significant difference was defined at $P \leq 0.05$ and trends at $0.05 < P \leq 0.10$. Descriptive statistics using the PROC MEANS procedure in SAS were applied to calculate the feed intake of sows during

late gestation and lactation, as well as the creep feed intake of piglets during the suckling period. PROC CORR in SAS was used to calculate Pearson correlation coefficients between cecal bacterial and fungal taxa and postweaning fecal scores on DoL30 and 34. To visualize the obtained correlations, heat maps were generated using the ‘levelplot()’ function in the lattice package in R Studio (version 2023.06.0).

Results

Feed intake of sows and piglets and growth performance of piglets

The average feed intake during the late gestation and lactation periods was 3.5 and 6.6 kg/d/sow, respectively (Supplementary Table S3). The creep feed intake was on average 19.6 g/d/piglet during the suckling period (Supplementary Table S4). Except for the higher birth weight of male piglets compared to female piglets ($P = 0.014$; Supplementary Table S5), body weight and average daily weight gain were similar during the suckling and early postweaning period. In total, 23 piglets were removed from the experiment due to illness or died (either crushing or sudden death, mostly in the first week of life).

Changes in the relative bacterial and fungal abundance and composition in the feces of lactating sows

The PERMANOVA based on Bray–Curtis dissimilarities showed significant separation between animal (i.e., sows and piglets), age (i.e., DoL and DPP), and gut segment (i.e., feces, stomach, and cecum) for the bacterial and fungal communities ($P < 0.001$; Supplementary Table S6). The NMDS (Bray–Curtis dissimilarity) demonstrated separate clustering for the bacterial and fungal communities in sow feces, gastric, and cecal digesta. Time-related clustering of the microbial communities was visible, especially for the communities in gastric and cecal digesta of the piglets but also for the fungal communities in sow feces with increasing DPP (Figure 1A and B).

Day postpartum did not influence the bacterial species richness (observed ASV) in the feces of sows (Table 1), whereas the bacterial alpha-diversity (Shannon and Simpson) increased from DPP1 to 6 and decreased thereafter until DPP27 ($P < 0.05$). The DPP affected the fungal species richness (observed ASV) and alpha-diversity (Shannon and Simpson; $P < 0.05$; Table 1) but in an opposite manner compared to the bacterial community. The fungal species richness and diversity decreased from DPP1 to 13 but increased thereafter until DPP27 ($P < 0.05$). These observations were supported by the results for the relatively most abundant genera (Figure 2A and B). As an example, the relatively most abundant genus *Lactobacillus* increased by 10.3-fold from DPP1 to 6 ($P < 0.001$), whereas the relative abundance of *Streptococcus* decreased by 3.5-fold from DPP1 to 13 and increased thereafter until DPP27 ($P < 0.001$). The relative abundance of *Rikenellaceae* RC9 gut group increased by 1.6-fold from DPP1 to 6 but decreased afterward on DPP13 by 1.5-fold ($P < 0.001$), whereas the relative abundance of *Terrisporobacter* increased from DPP1 to 20 by up to 1.4-fold ($P = 0.034$). As examples for fungal taxa, the relative abundance of *Kazachstania* as the dominant fungal genus in sow feces increased by 3.0-fold from DPP1 to 13 but decreased thereafter until DPP27 by up to 1.9-fold ($P < 0.001$). The relative abundance of *Tausonia*, which was the dominant genus on DPP1, declined by 28.9-fold




Figure 1. Non-metric multidimensional scaling (NMDS) plot of pairwise Bray-Curtis dissimilarities between communities of bacteriome (A) and mycobiome (B) in sow feces across days postpartum (DPP) and in the gastric and cecal digesta of piglets across days of life (DoL). Ellipses represent the standard deviation. Feces on DPP3 (dark blue), 7 (yellow), 14 (green), 21 (purple), and 28 (dark gray). Gastric digesta on DoL3 (green), 7 (purple), 14 (red), 21 (blue), and 28 (gray). Cecal digesta on DoL3 (red), 7 (blue), 14 (dark gray), 21 (dark blue), and 28 (yellow). Weaning took place on DoL28. Number 1 indicates sow feces, 2 is piglet's gastric digesta, and 3 is piglet's cecal digesta.

Table 1. Development in species richness (Observed ASV) and alpha diversity (Shannon and Simpson) indices for the bacterial and fungal community in sow feces after farrowing ($n = 20/d$ postpartum)

Day postpartum	1	6	13	20	27	SEM	P-value
Bacterial community							
Observed ASV	1195	1234	1251	1233	1219	37.2	0.909
Shannon	4.96 ^{bc}	5.24 ^a	5.14 ^{ab}	5.05 ^{abc}	4.92 ^c	0.079	0.020
Simpson	0.957 ^b	0.976 ^a	0.973 ^a	0.969 ^{ab}	0.961 ^b	0.004	0.007
Fungal community							
Observed ASV	178 ^a	99 ^b	95 ^b	211 ^a	174 ^a	23.3	<0.001
Shannon	1.65 ^a	1.06 ^{bc}	0.83 ^c	1.38 ^{ab}	1.67 ^a	0.195	0.007
Simpson	0.546 ^a	0.393 ^{ab}	0.265 ^b	0.401 ^{ab}	0.549 ^a	0.059	0.005

Values are presented as least squares means \pm standard error of the mean (SEM). Weaning took place on day 28 postpartum. ASV, amplicon sequence variants.

Means without a common superscript in the same row differ ($P < 0.05$).

DPP1 to 13 but remained stable thereafter until DPP27 ($P < 0.001$). The relative abundance of *Geotrichum* was low on DPP1 and 6, whereas it increased to DPP27 by up to 20.4-fold, becoming the second dominant fungal genus.

Age-related changes in bacterial and fungal communities in gastric and cecal digesta of piglets

Day of life did not affect bacterial and fungal species richness (observed ASV) but influenced bacterial (Simpson) and fungal (Shannon, Simpson) diversity ($P < 0.05$) in gastric digesta (Table 2). Accordingly, the bacterial diversity decreased from DoL14 to 31, whereas fungal diversity decreased from DoL28 to 35 in gastric digesta ($P < 0.05$). In cecal digesta, DoL affected bacterial and fungal species richness (observed ASV) and diversity (Shannon, Simpson; $P < 0.05$; Table 2), whereby the effect of the DoL were different for bacteria and fungi. In terms of

relative abundances in gastric digesta (Figure 3A), the relative abundances of the predominant *Lactobacillaceae* genera, including *Lactobacillus*, *Limosilactobacillus*, HT002, and *Ligilactobacillus*, together with *Streptococcus* altered from DoL3 to 7 and/or from DoL28 to 31 and 35 ($P < 0.05$). Alterations in relative abundances of fungal genera in gastric digesta were similarly visible (Figure 3B). For instance, gastric relative abundances of *Candida* and *Blumeria* largely increased from DoL7 to 14, whereas those of an unclassified genus and *Kazachstania* greatly increased from DoL28 to 31 ($P < 0.05$).

In cecal digesta, the relative abundance of *Fusobacterium* and *Bacteroides* were the predominant bacterial genera on DoL3, whereas the relative abundance of *Lactobacillus* became the dominant genus from DoL7 to 31 (Figure 3C). Postweaning, the relative abundance of *Prevotella-9* became dominant, whereas the relative abundances of *Fusobacterium*




Figure 2. Differences in the relative abundance (%) of the most abundant bacterial (A) and fungal (B) genera in sow feces across days postpartum ($n = 20/d$ postpartum). Weaning took place on day 28 postpartum. Effect ($P < 0.05$) of day postpartum is indicated by '*'.

Table 2. Age-related development of species richness (Observed ASV) and alpha diversity (Shannon and Simpson) for the bacterial and fungal communities in the gastric and cecal digesta of suckling and newly weaned piglets ($n = 20/d$ of life)

Day of life	3	7	14	21	28	31	35	SEM	P-value
Bacterial community in gastric digesta									
Observed ASV	468	434	491	483	897	562	581	132.4	0.180
Shannon	3.54	3.49	3.61	3.38	3.57	3.12	3.08	0.150	0.070
Simpson	0.913 ^a	0.907 ^a	0.923 ^a	0.893 ^{ab}	0.887 ^{ab}	0.860 ^b	0.855 ^b	0.013	0.002
Fungal community in gastric digesta									
Observed ASV	68	71	58	71	71	68	54	6.2	0.309
Shannon	2.23 ^{ab}	2.47 ^a	2.02 ^b	2.25 ^{ab}	2.18 ^{ab}	1.88 ^{bc}	1.53 ^c	0.135	<0.001
Simpson	0.772 ^{ab}	0.825 ^a	0.740 ^{ab}	0.763 ^{ab}	0.758 ^{ab}	0.718 ^{bc}	0.622 ^c	0.038	0.020
Bacterial community in cecal digesta									
Observed ASV	749 ^c	836 ^{ab}	821 ^{abc}	800 ^{abc}	875 ^a	867 ^{ab}	786 ^{bc}	27.7	0.038
Shannon	4.21 ^b	4.88 ^a	4.87 ^a	4.81 ^a	4.93 ^a	4.72 ^a	4.71 ^a	0.095	<0.001
Simpson	0.939 ^b	0.971 ^a	0.974 ^a	0.970 ^a	0.974 ^a	0.962 ^a	0.964 ^a	0.005	<0.001
Fungal community in cecal digesta									
Observed ASV	240 ^a	136 ^{cd}	213 ^{ab}	173 ^{bc}	183 ^{abc}	161 ^{bc}	107 ^d	19.8	<0.001
Shannon	3.08 ^a	2.74 ^a	3.02 ^a	2.51 ^a	2.54 ^a	1.77 ^b	0.66 ^c	0.201	<0.001
Simpson	0.816 ^a	0.866 ^a	0.865 ^a	0.788 ^a	0.790 ^a	0.530 ^b	0.207 ^c	0.048	<0.001

Values are presented as least squares means \pm standard error of the mean (SEM). Weaning took place on day 28 of life. ASV, amplicon sequence variants. Means without a common superscript in the same row differ ($P < 0.05$).

and *Bacteroides* largely dropped compared to the time points during the suckling phase ($P < 0.05$). The relative abundance of *Tausonia* decreased postweaning compared to the suckling phase, whereas the relative abundance of *Kazachstania* largely increased from DoL28 to 35 ($P < 0.001$; Figure 3D).

Postweaning fecal scores

Feces for scoring were obtained from 110 piglets each on DoL30 and 34. The fecal score was 0.67 on DoL30 and increased by 3.1-fold on DoL34 ($P < 0.001$; Figure 4A). Fecal

scores of the piglets ($n = 20$) from which gut samples were obtained were correlated with the bacterial abundances. Fecal scores correlated positively ($P < 0.05$) with the relative abundance of *Holdemanella* ($r = 0.55$) and negatively with the relative abundance of *Turicibacter* ($r = -0.54$) and *Odoribacter* ($r = -0.52$) but not with fungal genera on DoL30 (Figure 4B). On DoL34, the fecal score positively correlated with the relative abundance of 3 fungal genera (*Talaromyces*, *Mrakia*, and *Blumeria* ($r > 0.5$; $P < 0.05$; Figure 4C), as well as with that of 9 bacterial genera (*Clostridium sensu stricto*




Figure 3. Differences in the relative abundance (%) of the most abundant bacterial genera in gastric (A) and cecal (C) digesta as well as the most abundant fungal genera in gastric (B) and cecal (D) digesta of suckling and newly weaned piglets across days of life ($n = 20$ /d of life). Weaning took place on day 28 of life. Effect ($P < 0.05$) of the day of life is indicated by *.

1, *Prevotellaceae* NK3B31 group, *Parabacteroides*, *Terrisporobacter*, *Romboutsia*, *Turicibacter*, dgA-11 gut group, *Prevotellaceae* UCG-001, and *Elusimicrobium* ($r > 0.5$; $P < 0.05$; Figure 4D).

Discussion

The present study provides novel information about the alterations of the bacterial and fungal communities in sow feces during the lactation period as well as about the fungal diversity that establishes in the piglet's gastrointestinal tract after birth. There was a large overlap in bacterial and fungal

taxa between sow feces and gastric and cecal digesta of piglets during the suckling period, especially in the early days postpartum, which supports that the microbial community in sow feces is important for the neonatal gut colonization of piglets. Notably, the large variety of fungal taxa present and their relative abundance in the gastric and cecal digesta of the piglets on DoL3 resembled to a certain degree the fungal abundances in sow feces on DPP1. The uptake of feces by the piglets is similar to the concept of 'fecal transplants' (McCormack et al., 2018). Therefore, our results may be useful for future formulation of transition and lactation diets for sows to modulate their fecal microbiome in order to target




Figure 4. Differences in fecal scores of weaned piglets between days 30 and 34 of life (A; $n = 110$ piglets). Pearson's correlation heat map shows significant associations ($P < 0.05$) between fecal scores of the weaned piglets from which gut digesta was collected and cecal bacterial genera (B) on day 30 of life ($n = 20$ piglets), as well as between fecal scores and cecal fungal (C) and bacterial (D) genera on day 34 of life ($n = 20$ piglets). The fecal scoring system consisted of 0 (hard and dry balls), 1 (clearly defined shape with cracks), 2 (pasty; moist feces with no cracks), 3 (very soft but not liquid feces), 4 (liquid feces with minimal consistency), and 5 (entirely liquid feces). Piglets were weaned on day 28 of life. Fecal score values are least squares mean and standard error of the mean.

specific developmental stages in the gut microbiome of suckling piglets. Nevertheless, the differences in relative abundances indicated that the establishment of the microbes in the gut segments depended on the 'local' conditions (e.g., lower stomach pH in the stomach and nutrient availability) and other sources of microbes (e.g., sow colostrum/transient milk, skin, and environment). This assumption was supported by the beta-diversity, demonstrating different clustering bacterial and fungal communities for sow feces, and gastric and cecal digesta of piglets. With regard to environmental microbes, the milk replacer was based on bovine whey, which probably contained microbes from cheese-making that could have contributed to gut colonization.

Different types of fungi were present in sow feces and piglets' digesta including common gut inhabitants (e.g., *Kazachstania*; Harlow et al., 2024), plant-related fungi (Pietrusińska and Trat-

wal, 2020), yeasts (e.g., *Candida*; Pérez, 2021) and mold fungi (Liew and Mohd-Redzwan, 2018). Metabolically, these fungi probably filled different niches, breaking down residual dietary carbohydrates (Luo et al., 2021) as well as being involved in epithelial glucose turnover (Hu et al., 2023). Some plant-related fungi—some of them plant pathogens (e.g., *Blumeria*; Mapuranga et al., 2022)—may have been transient, colonizing the plant feed particles. Their role in the gut microbe-microbe/host interactions needs further research. Furthermore, the abundance of *Cladosporium*, *Alternaria*, *Fusarium*, and *Aspergillus* in sow feces and piglet's digesta showed that the animals may have been exposed to mold and ergot fungi from the environment and feed components (i.e., milk replacer and prestarter diet, lactation diet and hay) and hence potentially to mycotoxins and ergot alkaloids which may compromise their health (Coulal-Majewski et al., 2016; Deligeorgakis et al., 2023).

Results for the beta-diversity and ANOVA demonstrated that the bacterial and fungal communities in sow feces were not stable throughout lactation but continuously changed in relative abundance and diversity. This was probably a result of stress and dietary changes (i.e., discontinuation of feeding linseeds) around farrowing as well as the gradual increase in feed intake level during lactation (Jašarević et al., 2017; Lu et al., 2022). More specifically, the bacterial community changed more between DPP1 and 6, whereas the 4 dominant fungal taxa showed large changes in their abundances throughout the lactation phase. The fungal community was more susceptible to changes in fermentable substrate quality and quantity that reached the distal parts of the hindgut and/or substrate-related microbial interactions with progressing lactation than the bacterial community. Sows had access to hay as environmental enrichment; as a slowly fermentable fiber, it likely acted as a substrate for fibrolytic bacterial (e.g., *Treponema* and *Fibrobacter*; Xie et al., 2018) and fungal taxa (e.g., *Mucor*; Karimi and Zamani, 2013) in feces. Unfortunately, we do not have exact data for the hay intake of sows. Of note, the species richness and diversity of the fungi were greater in the cecal digesta of piglets than in cow feces in the first 2 wk of life, which may be linked to the immature immune system and microbe-to-microbe interactions.

The piglets showed a similar increase in *Lactobacillaceae* in gastric and cecal digesta to sow feces from DoL3 to 7, which may be advantageous due to their multiple beneficial effects on the composition of gut homeostasis and immune development (Valeriano et al., 2017; Zhang et al., 2022). From the relative abundance patterns, it can be assumed that sow feces were not the only source of *Lactobacillaceae* for the colonization of the piglet's stomach and distal gut. However, their predominance throughout the suckling phase was likely supported by the milk glycans. Of note, *Lactobacillus* was similarly relatively abundant in gastric and cecal digesta, whereas other milk glycan fermenters, such as *Limosilactobacillus*, HT002, and *Ligilactobacillus*, *Actinobacillus*, *Fusobacterium* and *Bacteroides* (García-Alija et al., 2022), were differently abundant in gastric and cecal digesta. Different substrate availabilities and subsequent cross-feeding relationships were probably behind these observations, also explaining the higher bacterial diversity in cecal compared to gastric digesta. Moreover, the bacterial and fungal composition probably reflected the slowly increasing amounts of creep feed as well as the change in the type of creep feed from the milk replacer to the pre-starter from DoL21 to 28. Even in very small amounts plant polysaccharides, such as starch, probably supported the growth of starch-degrading taxa such as *Turicibacter*, *Isiporobacter*, *Porphyromonas*, and *Prevotella* in digesta (Sun et al., 2015; Umu et al., 2015; Trachsel et al., 2019). As an example, the starch component may have proportionally lowered the relative abundance of milk glycan-fermenting bacteria in gastric digesta on DoL21. Other bacteria, such as *Clostridium sensu stricto*, may have thrived on milk peptides, either from sow milk or milk replacer.

Kazachstania is a commensal yeast in the porcine gut (Summers et al., 2021), which per se may explain its high presence in sow feces and gastric and cecal digesta of piglets. They feed on dietary or host-derived sugars, such as glucose and galactose (Kondybayev et al., 2023). Previously, *Kazachstania slooffiae* has been linked to the provision of amino acids and energy to other bacteria, such as *Lactobacillus* and *Prevotella*, as well as the host piglet (Summers et al., 2021;

Hu et al., 2023). The relative abundance of the yeast *Candida* increased more in gastric than in cecal digesta of piglets during the suckling phase, which may be linked to the glycan intake from milk and creep feed. *Geotrichum*, which was another relatively abundant fungus in sow feces but low abundant in gastric and cecal digesta of the piglets, may exert some positive effects on the host. In this respect, *Geotrichum* has been used as a probiotic in ruminant nutrition to promote productivity and bacterial diversity in dairy cattle (Zaman et al., 2022), whereas very little is known for pigs. According to its abundance pattern in feces throughout lactation, the abundance of *Geotrichum* seemed to be related to the feed intake level of the sows.

The diversity and relative abundance patterns of the bacterial and fungal communities allowed distinguishing the time point of weaning. It can be assumed that the weaning-related removal of sow milk and low feed intake as well as changes in the bacterial crosstalk may be behind the decreasing fungal diversity in cecal digesta from DoL28 (postweaning) to DoL35 (postweaning). It is difficult to relate the changes in the fungal community postweaning to the action of certain bacteria as our understanding is only advancing. Weaning changed the relative bacterial and fungal abundances in the sense of milk glycan fermenters being replaced by plant glycan utilizers. For instance, the removal of the milk glycans reduced the abundance of *Limosilactobacillus* and HT002 in gastric digesta. By contrast, other *Lactobacillaceae*, such as *Lactobacillus* and *Ligilactobacillus*, in gastric and cecal digesta remained stable or increased in their abundance postweaning, indicating that they were capable of utilizing the glycans in the pre-starter diet (i.e., starch and lactose) and/or those derived from the host. Of note, *Bifidobacterium*, another genus known for its milk glycan-fermenting capabilities (Jang and Kim, 2022), only raised in its abundance in gastric digesta on DoL31. This taxon possibly benefited from the opening niches of *Lactobacillaceae* but seemed to be outcompeted again on DoL35, e.g., by HT002 and *Ligilactobacillus*. In cecal digesta, typical plant glycan-utilizing taxa, like *Prevotellaceae* genera and *Succinivibrio* (Tan and Nie, 2020), became dominant on DoL31 and 35. Regarding the fungi, *Kazachstania* potentially benefited from increased glucose release from starch fermentation or mucosal glycolysis in both gastric and cecal digesta. From the abundance of pathobionts, the increase in *Campylobacter* in cecal digesta postweaning may be worth mentioning. Although it mostly does not cause enteritis in pigs, *Campylobacter* is an important zoonotic agent that is shed in feces, with the potential to be transmitted to humans (Ishengoma et al., 2023). By contrast, the gut conditions postweaning seemed to lower the relative abundance of mold and ergot fungi in gastric and cecal digesta. The piglets from which the gut samples were collected were not diarrhetic, as indicated by the fecal scores. This may be the reason for the few relationships between cecal microbial genera and fecal consistency on DoL30. Piglets had firm feces on DoL30. Accordingly, the fecal score only moderately correlated with 3 starch-fermenting and short-chain fatty acid-producing bacteria, namely *Holdemanella*, *Turicibacter*, and *Odoribacter*, but not with *Campylobacter*. On DoL34, only the positive association between fecal score and *Clostridium sensu stricto* 1 may relate moisture feces to bacterial toxins. The other positive correlations with bacterial taxa (e.g., *Prevotellaceae* NK3B31 group, *Prevotellaceae* UCG-001, and *Terrisporobacter*) may have been indicative of increased complex carbohydrate

35

fermentation and short-chain fatty acid production in the large intestine, which are osmotically active and increase the fecal water content (Amat et al., 2020; Niu et al., 2023). The 3 identified positive correlations of the fecal score with fungi were only with low abundant genera. Nevertheless, as these taxa were mold and plant pathogenic fungi, an activation of secretory functions in the large intestine may be thinkable.

In conclusion, this study demonstrated that progressing lactation affected bacterial and fungal community structure and bacterial richness in sows. Our results also support that sow feces were a contributing source of microbes for the gut colonization of piglets, as indicated by the shared bacterial and fungal taxa in sow feces and in the gastric and cecal digesta of piglets. We could identify microbial taxa in cecal digesta that are associated with the fecal scores on DoL30 and DoL34. However, fecal scores indicated that the feces were firm to moist but not entirely liquid feces. Therefore, the relationships miss information for a low gut homeostatic state post-weaning.

Supplementary Data

Supplementary data are available at *Journal of Animal Science* online.

Acknowledgments

This research was supported by the Austrian Federal Ministry for Digital and Economic Affairs and the National Foundation for Research, Technology and Development. BIOMIN Holding GmbH, which is part of dsm-firmenich, financially supported the Christian Doppler Laboratory for Innovative Gut Health Concepts of Livestock. We thank J. Ehmig, T. Enzinger (Clinical Department for Farm Animals and Food Systems Science), S. Posseth and T. Strini (VetFarm) for their excellent help with laboratory analysis, animal care and assistance during sampling. The first author thanks Austria's Agency for Education and Internationalization, Mobility Programs and Cooperation (OeAD-GmbH) in cooperation with the ASEAN-European Academic University Network for awarding Ernst-Mach-ASEA UNINET scholarship, which is fully funded by the Austrian Federal Ministry of Education, Science and Research (BMBWF).

Conflict of interest statement

All authors declare no conflict of interest other than the fact that BIOMIN Holding, which is part of dsm-firmenich provided partial funding for this research but had no influence on data analysis, data interpretation and preparation of the manuscript.

Author Contributions

Fitra Yosi (Data curation, Investigation, Writing—original draft), Frederike Lerch (Data curation, Investigation, Writing—review & editing), Julia Voetterl (Data curation, Investigation, Writing—review & editing), Simone Koger (Data curation, Investigation, Methodology, Writing—review & editing), Doris Verhovsek (Investigation, Writing—review & editing), and Barbara Metzler-Zebeli (Conceptualization, Data curation, Funding acquisition, Investigation, Methodology, Writing—review & editing).

Literature Cited

- Amat, S., H. Lantz, P. M. Munyaka, and B. P. Willing. 2020. *Prevotella* in pigs: the positive and negative associations with production and health. *Microorganisms*. 8:1584. doi: 10.3390/microorganisms8101584
- Callahan, B. J., P. J. McMurdie, M. J. Rosen, A. W. Han, A. J. A. Johnson, and S. P. Holmes. 2016. DADA2: high-resolution sample inference from Illumina amplicon data. *Nat. Methods* 13:581–583. doi: 10.1038/nmeth.3869
- Chen, X., J. Xu, E. Ren, Y. Su, and W. Zhu. 2018. Co-occurrence of early gut colonization in neonatal piglets with microbiota in the maternal and surrounding delivery environments. *Anaerobe* 49:30–40. doi: 10.1016/j.anaerobe.2017.12.002
- Coufal-Majewski, S., K. Stanford, T. McAllister, B. Blakley, J. McKinnon, A. V. Chaves, and Y. Wang. 2016. Impacts of cereal ergot in food animal production. *Front. Vet. Sci.* 3:15. doi: 10.3389/fvets.2016.00015
- Deligeorgakis, C., C. Magro, A. Skendi, H. H. Gebrehiwot, V. Valdramidis, and M. Papageorgiou. 2023. Fungal and toxin contaminants in cereal grains and flours: systematic review and meta-analysis. *Foods*. 12:4328. doi: 10.3390/foods12234328
- Everaert, N., S. Van Cruchten, B. Weström, M. Bailey, C. Van Ginneken, T. Thymann, and R. Pieper. 2017. A review on early gut maturation and colonization in pigs, including biological and dietary factors affecting gut homeostasis. *Anim. Feed Sci. Technol.* 233:89–103. doi: 10.1016/j.anifeedsci.2017.06.011
- García-Alíja, M., J. J. Du, I. Ordóñez, A. Diz-Vallenilla, A. Moraleda-Montoya, N. Sultana, C. G. Huynh, C. Li, T. C. Donahue, L.-X. Wang, et al. 2022. Mechanism of cooperative N-glycan processing by the multi-modular endoglycosidase EndoE. *Nat. Commun.* 13:1137. doi: 10.1038/s41467-022-28722-w
- Harlow, K., K. L. Summers, W. T. Oliver, J. E. Wells, M. Crouse, B. W. Neville, L. A. Rempel, I. Rivera, T. G. Ramsay, and C. P. Davies. 2024. Weaning transition, but not the administration of probiotic candidate *Kazachstania slooffiae*, shaped the gastrointestinal bacterial and fungal communities in nursery piglets. *Front. Vet. Sci.* 10:1–22. doi: 10.3389/fvets.2023.1303984
- Hu, J., J. Chen, Q. Hou, X. Xu, J. Ren, L. Ma, and X. Yan. 2023. Core-predominant gut fungus *Kazachstania slooffiae* promotes intestinal epithelial glycolysis via lysine desuccinylation in pigs. *Microbiome*. 11:31. doi: 10.1186/s40168-023-01468-3
- Ishengoma, V. L., R. G. Amachawadi, M. D. Tokach, X. Shi, Q. Kang, R. D. Goodband, J. DeRouchey, J. Woodworth, and T. G. Nagaraja. 2023. Impact of in-feed versus in-water chlortetracycline and tiamulin administrations on fecal prevalence and antimicrobial susceptibilities of *Campylobacter* in a population of nursery pigs. *Microorganisms*. 11:2876. doi: 10.3390/microorganisms11122876
- Jang, K. B., and S. W. Kim. 2022. Role of milk carbohydrates in intestinal health of nursery pigs: a review. *J. Anim. Sci. Biotechnol.* 13:6. doi: 10.1186/s40104-021-00650-7
- Jašarević, E., C. D. Howard, A. M. Misić, D. P. Beiting, and T. L. Bale. 2017. Stress during pregnancy alters temporal and spatial dynamics of the maternal and offspring microbiome in a sex-specific manner. *Sci. Rep.* 7:1–13. doi: 10.1038/srep44182
- Jost, T., C. Lacroix, C. P. Braegger, F. Rochat, and C. Chassard. 2014. Vertical mother–neonate transfer of maternal gut bacteria via breastfeeding. *Environ. Microbiol.* 16:2891–2904. doi: 10.1111/1462-2920.12238
- Karimi, K., and A. Zamani. 2013. *Mucor indicus*: biology and industrial application perspectives: a review. *Biotechnol. Adv.* 31:466–481. doi: 10.1016/j.biotechadv.2013.01.009
- Klindworth, A., E. Pruesse, T. Schweer, J. Peplies, C. Quast, M. Horn, and F. O. Glöckner. 2013. Evaluation of general 16S ribosomal RNA gene PCR primers for classical and next-generation sequencing-based diversity studies. *Nucleic Acids Res.* 41:e1. doi: 10.1093/nar/gks808
- Kondybayev, A., N. Achir, C. Mestres, I. Collombel, C. Strub, J. Grabulos, N. Akhmetadykov, A. Aubakirova, U. Kamidinkyzy, W. Ghanmi, et al. 2023. Growth kinetics of *Kazachstania unispora* and

- its interaction with lactic acid bacteria during qymyz production. *Fermentation*. 9:101. doi: [10.3390/fermentation9020101](https://doi.org/10.3390/fermentation9020101)
- Lerch, F., F. Yosi, J. C. Vötterl, S. Koger, J. Ehlig, S. Sharma, D. Verhovsek, and B. U. Metzler-Zebeli. 2023. An insight into the temporal dynamics in the gut microbiome, metabolite signaling, immune response, and barrier function in suckling and weaned piglets under production conditions. *Front. Vet. Sci.* 10:1184277. doi: [10.3389/fvets.2023.1184277](https://doi.org/10.3389/fvets.2023.1184277)
- Liew, W.-P.-P., and S. Mohd-Redzwan. 2018. Mycotoxin: its impact on gut health and microbiota. *Front. Cell. Infect. Microbiol.* 8:60. doi: [10.3389/fcimb.2018.00060](https://doi.org/10.3389/fcimb.2018.00060)
- Lu, D., Y. Pi, H. Ye, Y. Wu, Y. Bai, S. Lian, D. Han, D. Ni, X. Zou, J. Zhao, et al. 2022. consumption of dietary fiber with different physicochemical properties during late pregnancy alters the gut microbiota and relieves constipation in sow model. *Nutrients*. 14:2511. doi: [10.3390/nu14122511](https://doi.org/10.3390/nu14122511)
- Luo, Y., J. Li, H. Zhou, B. Yu, J. He, A. Wu, Z. Huang, P. Zheng, X. Mao, J. Yu, et al. 2021. The nutritional significance of intestinal fungi: alteration of dietary carbohydrate composition triggers colonic fungal community shifts in a pig model. *Appl. Environ. Microbiol.* 87:1–19. doi: [10.1128/AEM.00038-21](https://doi.org/10.1128/AEM.00038-21)
- Mann, E., S. Schmitz-Esser, Q. Zebeli, M. Wagner, M. Ritzmann, and B. U. Metzler-Zebeli. 2014. Mucosa-associated bacterial microbiome of the gastrointestinal tract of weaned pigs and dynamics linked to dietary calcium-phosphorus. *PLoS One* 9:e86950. doi: [10.1371/journal.pone.0086950](https://doi.org/10.1371/journal.pone.0086950)
- Mapuranga, J., J. Chang, and W. Yang. 2022. Combating powdery mildew: advances in molecular interactions between *Blumeria graminis* f. sp. *tritici* and wheat. *Front. Plant Sci.* 13:1–19. doi: [10.3389/fpls.2022.1102908](https://doi.org/10.3389/fpls.2022.1102908)
- McCormack, U. M., T. Curião, T. Wilkinson, B. U. Metzler-Zebeli, H. Reyer, T. Ryan, J. A. Calderon-Diaz, F. Crispie, P. D. Cotter, C. J. Creevey, et al. 2018. Fecal microbiota transplantation in gestating sows and neonatal offspring alters lifetime intestinal microbiota and growth in offspring. *mSystems* 3:1–20. doi: [10.1128/mSystems.00134-17](https://doi.org/10.1128/mSystems.00134-17)
- Metzler-Zebeli, B. U., F. Lerch, F. Yosi, J. Vötterl, J. Ehlig, S. Koger, and D. Verhovsek. 2023. Temporal microbial dynamics in feces discriminate by nutrition, fecal color, consistency and sample type in suckling and newly weaned piglets. *Animals*. 13:2251. doi: [10.3390/ani13142251](https://doi.org/10.3390/ani13142251)
- National Research Council (NRC). 2012. Nutrient requirements of swine. 10th ed. Washington, DC, USA: National Academy Press.
- Nilsson, R. H., K. -H. Larsson, A. F. S. Taylor, J. Bengtsson-Palme, T. S. Jeppesen, D. Schigel, P. Kennedy, K. Picard, F. O. Glöckner, L. Tedersoo, et al. 2019. The UNITE database for molecular identification of fungi: handling dark taxa and parallel taxonomic classifications. *Nucleic Acids Res.* 47:D259–D264. doi: [10.1093/nar/gky1022](https://doi.org/10.1093/nar/gky1022)
- Niu, J., X. Liu, J. Xu, F. Li, J. Wang, X. Zhang, X. Yang, L. Wang, S. Ma, D. Li, et al. 2023. Effects of silage diet on meat quality through shaping gut microbiota in finishing pigs. *Microbiol. Spectr.* 11:e0241622. doi: [10.1128/spectrum.02416-22](https://doi.org/10.1128/spectrum.02416-22)
- Nowland, T., K. Plush, M. Barton, and R. Kirkwood. 2019. Development and function of the intestinal microbiome and potential implications for pig production. *Animals*. 9:76. doi: [10.3390/ani9030076](https://doi.org/10.3390/ani9030076)
- Oksanen, J., F. G. Blanchet, R. Kindt, P. Legendre, P. R. Minchin, R. B. O'Hara, G. L. Simpson, P. Sölymos, M. H. H. Stevens, and H. Wagne. 2022. *vegan: Community Ecology Package* R Package Version 2.6–4.
- Op De Beeck, M., B. Lievens, P. Busschaert, S. Declercq, J. Vangronsveld, and J. V. Colpaert. 2014. Comparison and validation of some ITS primer pairs useful for fungal metabarcoding studies. *PLoS One* 9:e97629. doi: [10.1371/journal.pone.0097629](https://doi.org/10.1371/journal.pone.0097629)
- Pérez, J. C. 2021. The interplay between gut bacteria and the yeast *Candida albicans*. *Gut. Microbes* 13:1–11. doi: [10.1080/19490976.2021.1979877](https://doi.org/10.1080/19490976.2021.1979877)
- Pietrusińska, A., and A. Tratwal. 2020. Characteristics of powdery mildew and its importance for wheat grown in Poland. *Plant Prot. Sci.* 56:141–153. doi: [10.17221/99/2019-pps](https://doi.org/10.17221/99/2019-pps)
- Quast, C., E. Pruesse, P. Yilmaz, J. Gerken, T. Schweer, P. Yarza, J. Peplies, and F. O. Glöckner. 2012. The SILVA ribosomal RNA gene database project: improved data processing and web-based tools. *Nucleic Acids Res.* 41:D590–D596. doi: [10.1093/nar/gks1219](https://doi.org/10.1093/nar/gks1219)
- Summers, K. L., J. Foster Frey, and A. M. Arfken. 2021. Characterization of *Kazachstania slooffiae*, a proposed commensal in the porcine gut. *J. Fungi*. 7:146. doi: [10.3390/jof7020146](https://doi.org/10.3390/jof7020146)
- Sun, Y., L. Zhou, L. Fang, Y. Su, and W. Zhu. 2015. Responses in colonic microbial community and gene expression of pigs to a long-term high resistant starch diet. *Front. Microbiol.* 6:1–10. doi: [10.3389/fmicb.2015.00877](https://doi.org/10.3389/fmicb.2015.00877)
- Tan, H., and S. Nie. 2020. Deciphering diet-gut microbiota-host interplay: investigations of pectin. *Trends Food Sci. Technol.* 106:171–181. doi: [10.1016/j.tifs.2020.10.010](https://doi.org/10.1016/j.tifs.2020.10.010)
- Trachsel, J., C. Briggs, N. K. Gabler, H. K. Allen, and C. L. Loving. 2019. Dietary resistant potato starch alters intestinal microbial communities and their metabolites, and markers of immune regulation and barrier function in swine. *Front. Immunol.* 10:1–13. doi: [10.3389/fimmu.2019.01381](https://doi.org/10.3389/fimmu.2019.01381)
- Umu, C. O., J. A. Frank, J. U. Fangel, M. Oostindjer, C. S. da Silva, E. J. Bolhuis, G. Bosch, W. G. T. Willats, P. B. Pope, and D. B. Diep. 2015. Resistant starch diet induces change in the swine microbiome and a predominance of beneficial bacterial populations. *Microbiome*. 3:16. doi: [10.1186/s40168-015-0078-5](https://doi.org/10.1186/s40168-015-0078-5)
- Valeriano, V. D. V., M. P. Balolong, and D. -K. Kang. 2017. Probiotic roles of *Lactobacillus* sp. in swine: insights from gut microbiota. *J. Appl. Microbiol.* 122:554–567. doi: [10.1111/jam.13364](https://doi.org/10.1111/jam.13364)
- Wang, K., V. Espinosa, and A. Rivera. 2023. Commander-in-chief: monocytes rally the troops for defense against aspergillosis. *Curr. Opin Immunol.* 84:102371. doi: [10.1016/j.coi.2023.102371](https://doi.org/10.1016/j.coi.2023.102371)
- Xie, X., C. Yang, L. L. Guan, J. Wang, M. Xue, and J. X. Liu. 2018. Persistence of cellulolytic bacteria *Fibrobacter* and *Treponema* after short-term corn stover-based dietary intervention reveals the potential to improve rumen fibrolytic function. *Front. Microbiol.* 9:1–15. doi: [10.3389/fmicb.2018.01363](https://doi.org/10.3389/fmicb.2018.01363)
- Yeh, Y. -W., P. -Y. Chou, H. -Y. Hou, and R. Kirschner. 2021. First records of powdery mildew fungi (Erysiphales) on medicinal plants in Taiwan. *Bot. Stud.* 62:1. doi: [10.1186/s40529-020-00307-0](https://doi.org/10.1186/s40529-020-00307-0)
- Zaman, S., M. Gohar, H. Kanwal, A. Chaudhary, and M. Imran. 2022. Impact of probiotic *Geotrichum candidum* QAUGC01 on health, productivity, and gut microbial diversity of dairy cattle. *Curr. Microbiol.* 79:376. doi: [10.1007/s00284-022-03074-2](https://doi.org/10.1007/s00284-022-03074-2)
- Zhang, Q., R. Vasquez, J. M. Yoo, S. H. Kim, D. -K. Kang, and I. H. Kim. 2022. Dietary supplementation of *Limosilactobacillus mucosae* LM1 enhances immune functions and modulates gut microbiota without affecting the growth performance of growing pigs. *Front. Vet. Sci.* 9:1–12. doi: [10.3389/fvets.2022.918114](https://doi.org/10.3389/fvets.2022.918114)

17%

SIMILARITY INDEX

PRIMARY SOURCES

1	vtechworks.lib.vt.edu Internet	97 words — 1%
2	Natalie E Diether, Seo Lin Nam, Janelle Foughse, Bich V Le Thanh et al. "Dietary benzoic acid and supplemental enzymes alter fiber-fermenting taxa and metabolites in the cecum of weaned pigs", Journal of Animal Science, 2022 Crossref	60 words — 1%
3	www.biorxiv.org Internet	46 words — 1%
4	Guevara Guillen, Fiama Estefany. "Characterization of the Diversity and Functional Potential of Bacterial and Fungal Communities in Hydroponic Systems.", The Ohio State University ProQuest	39 words — < 1%
5	journals.plos.org Internet	39 words — < 1%
6	www.animbiosci.org Internet	39 words — < 1%
7	Leonie Bettin, Maria Stadler, Christine Unterweger, Maximiliane Dippel et al. "A Cyclic-di-AMP Adjuvanted CPAF Protein Vaccine Is Immunogenic in Swine, but It Fails to Reduce Genital Chlamydia trachomatis Burden", Vaccines, 2025 Crossref	37 words — < 1%
8	krex.k-state.edu Internet	35 words — < 1%

9	assets.researchsquare.com Internet	29 words — < 1%
10	Qianqian Zhang, Robie Vasquez, Jeong Min Yoo, Sang Hoon Kim, Dae-Kyung Kang, In Ho Kim. "Dietary Supplementation of Limosilactobacillus mucosae LM1 Enhances Immune Functions and Modulates Gut Microbiota Without Affecting the Growth Performance of Growing Pigs", Frontiers in Veterinary Science, 2022 Crossref	27 words — < 1%
11	escholarship.org Internet	27 words — < 1%
12	Bo Bo, Seul-Ah Kim, Nam Soo Han. "Bacterial and fungal diversity in Laphet, traditional fermented tea leaves in Myanmar, analyzed by culturing, DNA amplicon-based sequencing, and PCR-DGGE methods", International Journal of Food Microbiology, 2020 Crossref	23 words — < 1%
13	Zebeli, Q., K. Ghareeb, E. Humer, B.U. Metzler-Zebeli, and U. Besenfelder. "Nutrition, rumen health and inflammation in the transition period and their role on overall health and fertility in dairy cows", Research in Veterinary Science, 2015. Crossref	21 words — < 1%
14	mdpi-res.com Internet	20 words — < 1%
15	tel.archives-ouvertes.fr Internet	20 words — < 1%
16	www.msard-journal.com Internet	20 words — < 1%
17	inspectapedia.com Internet	19 words — < 1%
18	www.jkfs.or.kr Internet	19 words — < 1%

19	genomemedicine.biomedcentral.com Internet	18 words — < 1 %
20	minds.wisconsin.edu Internet	17 words — < 1 %
21	repository.kaust.edu.sa Internet	17 words — < 1 %
22	You-Lin Tain, Chi-I Chang, Chih-Yao Hou, Guo-Ping Chang-Chien, Sufan Lin, Chien-Ning Hsu. "Dietary Resveratrol Butyrate Monoester Supplement Improves Hypertension and Kidney Dysfunction in a Young Rat Chronic Kidney Disease Model", <i>Nutrients</i> , 2023 Crossref	16 words — < 1 %
23	fvtm.stafpu.bu.edu.eg Internet	16 words — < 1 %
24	unbscholar.dspace.lib.unb.ca Internet	16 words — < 1 %
25	Jinlong Li, Yong Tuo, Linjiao He, Yan Ma, Zhijun Zhang, Zhiqiang Cheng, Changjiang Zang, Tongjun Guo. "Effects of chili straw on rumen fermentation, meat quality, amino acid and fatty acid contents, and rumen bacteria diversity in sheep", <i>Frontiers in Microbiology</i> , 2025 Crossref	15 words — < 1 %
26	www.pubfacts.com Internet	15 words — < 1 %
27	Hiroshi Iseki, Eri Furukawa, Tomoya Shimasaki, Shogo Higaki. "Automatic monitoring of activity intensity in a chicken flock using a computer vision-based background image subtraction technique: an experimental infection study with fowl adenovirus", <i>Smart Agricultural Technology</i> , 2025 Crossref	14 words — < 1 %
28	Mathilde Le Sciellour, Olivier Zemb, Isabelle Hochu, Juliette Riquet et al. "Effect of chronic and	14 words — < 1 %

acute heat challenges on fecal microbiota composition,
production, and thermoregulation traits in growing pigs^{1,2}",
Journal of Animal Science, 2019

Crossref

29 ktisis.cut.ac.cy 14 words — < 1 %
Internet

30 www.science.org 14 words — < 1 %
Internet

31 www.tdx.cat 14 words — < 1 %
Internet

32 ijal.se 13 words — < 1 %
Internet

33 animalmicrobiome.biomedcentral.com 11 words — < 1 %
Internet

34 dissertations.umi.com 11 words — < 1 %
Internet

35 mtaisociety.weebly.com 11 words — < 1 %
Internet

36 Anne H Lee, Ching-Yen Lin, Sungho Do, Patricia M Oba, Sara E Belchik, Andrew J Steelman, Amy Schauwecker, Kelly S Swanson. "Dietary supplementation with fiber, "biotics," and spray-dried plasma affects apparent total tract macronutrient digestibility and the fecal characteristics, fecal microbiota, and immune function of adult dogs", Journal of Animal Science, 2022 10 words — < 1 %
Crossref

37 Feng Gao, Yongqing Du, Haiyang Liu, Hongwei Ding, Wentao Zhang, Zhongyu Li, Baoming Shi. "Maternal supplementation with konjac glucomannan and κ-carrageenan promotes sow performance and benefits the gut barrier in offspring", Animal Nutrition, 2024 10 words — < 1 %
Crossref

38 Marcos Elias Duarte, Sung Woo Kim. "Intestinal microbiota and its interaction to intestinal health in nursery pigs", Animal Nutrition, 2021 10 words — < 1 %
Crossref

39 Thomas H Naragon, Joani W Viliunas, Mina Yousefelahiyeh, Adrian Bruckner et al. "Symbiotic entrenchment through ecological Catch-22", Cold Spring Harbor Laboratory, 2025 10 words — < 1 %
Crossref Posted Content

40 link.springer.com 10 words — < 1 %
Internet

41 pure.mpg.de 10 words — < 1 %
Internet

42 rcastoragev2.blob.core.windows.net 10 words — < 1 %
Internet

43 "Sustainable Swine Nutrition", Wiley, 2022 9 words — < 1 %
Crossref

44 Annika Mohr, Jens Peter Hölzen, Sandra Stöppeler, Hans-Ullrich Spiegel et al. "Development and evaluation of a surgical skills lab for trainee surgeons:a 10-year experience at the Münster University Hospital", BMC Medical Education, 2025 9 words — < 1 %
Crossref

45 Bianca Sampaio Dotto Fiuza, Candace Machado de Andrade, Pedro Milet Meirelles, Jorley Santos da Silva et al. "Gut microbiome signature and nasal lavage inflammatory markers in young asthmatics", Journal of Allergy and Clinical Immunology: Global, 2024 9 words — < 1 %
Crossref

46 Ida Romano, Valeria Ventorino, Giuseppina Magaraci, Fatima Ezzahra Chouyia, Mauro Mori, Olimpia Pepe. "Application of an optimized pectinolytic bacterial consortium to boost hemp water retting", Industrial Crops and Products, 2025 9 words — < 1 %
Crossref

47 Poudel, Prakash . "Effects of Commercial Feed Additives on the Gut Microbiota of Food Animals.", South Dakota State University, 2020
ProQuest 9 words — < 1 %

48 Xiangyu Bian, Tala Shi, Yawen Wang, Yuying Ma, Yijing Yu, Weina Gao, Changjiang Guo. "Gut dysbiosis induced by antibiotics is improved by tangerine pith extract in mice", Nutrition Research, 2022
Crossref 9 words — < 1 %

49 dr.lib.iastate.edu
Internet 9 words — < 1 %

50 harvest.usask.ca
Internet 9 words — < 1 %

51 mspace.lib.umanitoba.ca
Internet 9 words — < 1 %

52 researcherslinks.com
Internet 9 words — < 1 %

53 researchoutput.csu.edu.au
Internet 9 words — < 1 %

54 theses.gla.ac.uk
Internet 9 words — < 1 %

55 uu.diva-portal.org
Internet 9 words — < 1 %

56 www.wur.nl
Internet 9 words — < 1 %

57 Anu Anupma, Jyoti Prakash Tamang. "Diversity of Filamentous Fungi Isolated From Some Amylase and Alcohol-Producing Starters of India", Frontiers in Microbiology, 2020
Crossref 8 words — < 1 %

58 Hao-Yu Liu, Shicheng Li, Kennedy Jerry Ogamune, Abdelkareem A. Ahmed, In Ho Kim, Yunzeng Zhang, Demin Cai. "Fungi in the Gut Microbiota: Interactions, Homeostasis, and Host Physiology", Microorganisms, 2025 8 words — < 1%
Crossref

59 Israel Rivera, KaLynn Harlow, Robert N. Cole, Robert O'Meally et al. "A metaproteomic analysis of the piglet fecal microbiome across the weaning transition", Frontiers in Microbiology, 2025 8 words — < 1%
Crossref

60 Kane, Jennifer Lynne. "Cross-Kingdom Interactions Shape Soil Biogeochemistry in Natural and Agricultural Ecosystems", West Virginia University, 2023 8 words — < 1%
ProQuest

61 M. Saladrigas-García, M. D'Angelo, H. L. Ko, S. Traserra et al. "Early socialization and environmental enrichment of lactating piglets affects the caecal microbiota and metabolomic response after weaning", Scientific Reports, 2021 8 words — < 1%
Crossref

62 Madeeha Gohar, Nida Shaheen, Sagar M. Goyal, Sunil Kumar Mor, Luis M. Rodriguez-R, Muhammad Imran. "Probiotic Potential of Yeast, Mold, and Intermediate Morphotypes of Geotrichum candidum in Modulating Gut Microbiota and Body Physiology in Mice", Probiotics and Antimicrobial Proteins, 2025 8 words — < 1%
Crossref

63 Meinrad Drexel, Günther Sperk. "Seizure-induced overexpression of NPY induces epileptic tolerance in a mouse model of spontaneous recurrent seizures", Frontiers in Molecular Neuroscience, 2022 8 words — < 1%
Crossref

64 Panagiotis A. Karas, Sotirios Vasileiadis, Christina Papazlatani, Anna Batsaki et al. "Depuration of fungicide-contaminated effluents from fruit-packaging plants in 8 words — < 1%

a full-scale modular bioreactor inoculated with tailor-made microbial inocula", Journal of Hazardous Materials, 2025

Crossref

- 65 Qing Xia, Thomas Rufty, Wei Shi. "Soil microbial diversity and composition: Links to soil texture and associated properties", Soil Biology and Biochemistry, 2020 8 words — < 1%

Crossref

- 66 Thapa, Soni. "Ecophysiology of Thermophilic Bacterial Endospores in Deep Marine Sediments.", Idaho State University 8 words — < 1%

ProQuest

- 67 Wei Chen, Jingyun Ma, Yiming Jiang, Li Deng et al. "Selective Maternal Seeding and Rearing Environment From Birth to Weaning Shape the Developing Piglet Gut Microbiome", Frontiers in Microbiology, 2022 8 words — < 1%

Crossref

- 68 Xiao, Jin, Barbara Metzler-Zebeli, and Qendrim Zebeli. "Gut Function-Enhancing Properties and Metabolic Effects of Dietary Indigestible Sugars in Rodents and Rabbits", Nutrients, 2015. 8 words — < 1%

Crossref

- 69 Xiaofan Wang, Tsungcheng Tsai, Bin Zuo, Xiaoyuan Wei et al. "Donor age and body weight determine the effects of fecal microbiota transplantation on growth performance, and fecal microbiota development in recipient pigs", Journal of Animal Science and Biotechnology, 2022 8 words — < 1%

Crossref

- 70 Xue Chen, Jumei Xu, Yong Su, Weiyun Zhu. "Effects of Intravenous Infusion With Sodium Butyrate on Colonic Microbiota, Intestinal Development- and Mucosal Immune-Related Gene Expression in Normal Growing Pigs", Frontiers in Microbiology, 2018 8 words — < 1%

Crossref

- 71 Y. Sun, K. Yu, L. Zhou, L. Fang, Y. Su, W. Zhu. "Metabolomic and transcriptomic responses 8 words — < 1%

induced in the livers of pigs by the long-term intake of resistant starch1", Journal of Animal Science, 2016

Crossref

72	atrium.lib.uoguelph.ca Internet	8 words — < 1 %
73	core.ac.uk Internet	8 words — < 1 %
74	cris.maastrichtuniversity.nl Internet	8 words — < 1 %
75	dspace.umh.es Internet	8 words — < 1 %
76	eprints.qut.edu.au Internet	8 words — < 1 %
77	era.library.ualberta.ca Internet	8 words — < 1 %
78	iris.uniss.it Internet	8 words — < 1 %
79	pmc.ncbi.nlm.nih.gov Internet	8 words — < 1 %
80	refubium.fu-berlin.de Internet	8 words — < 1 %
81	services.phaidra.univie.ac.at Internet	8 words — < 1 %
82	www.frontiersin.org Internet	8 words — < 1 %
83	www.midwestwinenutritionconference.com Internet	8 words — < 1 %
84	www.nwfirescience.org Internet	8 words — < 1 %

-
- 85 www.pharmazie.com 8 words — < 1 %
Internet
-
- 86 www.researchsquare.com 8 words — < 1 %
Internet
-
- 87 "Gut Microbiota, Immunity, and Health in Production Animals", Springer Science and Business Media LLC, 2022 7 words — < 1 %
Crossref
-
- 88 Hammers, Kelsey Lynn. "Feeding High Dietary Zinc to Gestating Sows and Implications on Sow and Piglet Performance, Their Biological Responses, and the Environment", University of Minnesota, 2023 7 words — < 1 %
ProQuest
-
- 89 Wilbert Gutiérrez-Sarmiento, Betsy Anaid Peña-Ocaña, Anayancy Lam-Gutiérrez, Jorge Martín Guzmán-Albores et al. "Microbial community structure, physicochemical characteristics and predictive functionalities of the Mexican tepache fermented beverage", Microbiological Research, 2022 7 words — < 1 %
Crossref
-
- 90 "Production Diseases in Farm Animals", Springer Science and Business Media LLC, 2024 6 words — < 1 %
Crossref
-
- 91 B.U. Metzler-Zebeli. "Interactive effects of the microbiome and nutrition on mucosal physiology and functions in the porcine intestinal tract", Animal - science proceedings, 2022 6 words — < 1 %
Crossref
-
- 92 Beatriz Albuixech-Crespo, Laura López-Blanch, Demian Burguera, Ignacio Maeso, Luisa Sánchez-Arrones, Juan Antonio Moreno-Bravo, Ildiko Somorjai, Juan Pascual-Anaya, Eduardo Puelles, Paola Bovolenta, Jordi Garcia-Fernàndez, Luis Puelles, Manuel Irimia, José Luis Ferran. "Molecular regionalization of the developing amphioxus neural

- 93 Choudhury, Raka. "Early-Life Feeding in Piglets: the Impact on Intestinal Microbiota and Mucosal Development", Wageningen University and Research, 2022 6 words — < 1%

ProQuest

- 94 Dongmei Wu, Xingqian Ye, Robert J. Linhardt, Xuwei Liu, Kai Zhu, Chengxiao Yu, Tian Ding, Donghong Liu, Qiaojun He, Shiguo Chen. "Dietary pectic substances enhance gut health by its polycomponent: A review", Comprehensive Reviews in Food Science and Food Safety, 2021 6 words — < 1%

Crossref

- 95 Eudald Llauradó-Calero, Eric Climent, Empar Chenoll, Maria Ballester et al. "Influence of dietary n-3 long-chain fatty acids on microbial diversity and composition of sows' feces, colostrum, milk, and suckling piglets' feces", Frontiers in Microbiology, 2022 6 words — < 1%

Crossref

- 96 Fawen Dai, Tao Lin, Muqu Jin, Xia Huang, Lu Wang, Jing Ma, Hang Yu, Xianlin Fan, Xiang Nong, Jianjun Zuo. "Bamboo fiber improves piglet growth performance by regulating the microbial composition of lactating sows and their offspring piglets", Frontiers in Microbiology, 2024 6 words — < 1%

Crossref

- 97 Mireia Saladrigas-García, Mario Durán, Matilde D'Angelo, Jaume Coma, José Francisco Pérez, Susana María Martín-Orúe. "An insight into the commercial piglet's microbial gut colonization: from birth towards weaning", Animal Microbiome, 2022 6 words — < 1%

Crossref

- 98 Moamen M. Elmassry, Ahmed Zayed, Mohamed A. Farag. "Gut homeostasis and microbiota under attack: impact of the different types of food contaminants on gut health", Critical Reviews in Food Science and Nutrition, 2020 6 words — < 1%

99 Nuria Canibe, Ole Højberg, Hanne Kongsted, Darya Vodolazska, Charlotte Lauridsen, Tina Skau Nielsen, Anna A. Schönherz. "Review on Preventive Measures to Reduce Post-Weaning Diarrhoea in Piglets", Animals, 2022

Crossref

6 words — < 1%

100 Steven McLoughlin, Charles Spillane, Noel Claffey, Paul E. Smith, Tommy O'Rourke, Michael G. Diskin, Sinéad M. Waters. "Rumen Microbiome Composition Is Altered in Sheep Divergent in Feed Efficiency", Frontiers in Microbiology, 2020

Crossref

6 words — < 1%

101 Érica B da Silva, Rebecca M Savage, Amy S Biddle, Stephanie A Polukis, Megan L Smith, Limin Kung. "Effects of a chemical additive on the fermentation, microbial communities, and aerobic stability of corn silage with or without air stress during storage", Journal of Animal Science, 2020

Crossref

6 words — < 1%

EXCLUDE QUOTES OFF
EXCLUDE BIBLIOGRAPHY ON

EXCLUDE SOURCES OFF
EXCLUDE MATCHES OFF