


#### 4. Bukti konfirmasi review ketiga dan hasil revisi ketiga

13 Oktober 2023



## COVERING LETTER

Dear **Editor-in-Chief**,

I herewith enclosed a research article,

- ☒ The submission has not been previously published, nor is it before another journal for consideration (or an explanation has been provided in Comments to the Editor).
- ☒ The submission file is in OpenOffice, Microsoft Word (DOC, not DOCX), or RTF document file format.
- ☒ The text is single-spaced; uses a 10-point font; employs italics, rather than underlining (except with URL addresses); and all illustrations, figures, and tables are placed within the text at the appropriate points, rather than at the end.
- ☒ The text adheres to the stylistic and bibliographic requirements outlined in the Author Guidelines.
- ☒ Most of the references come from current scientific journals (c. 80% published in the last 10 years), except for taxonomic papers.
- ☒ Where available, DOIs for the references have been provided.
- ☒ When available, a certificate for proofreading is included.

## SUBMISSION CHECKLIST

Ensure that the following items are present:

The first corresponding author must be accompanied with contact details:

- ☒ E-mail address
- ☒ Full postal address (incl street name and number (location), city, postal code, state/province, country)
- ☒ Phone and facsimile numbers (incl country phone code)

All necessary files have been uploaded, and contain:

- ☒ Keywords
- ☒ Running titles
- ☒ All figure captions
- ☒ All tables (incl title and note/description)

Further considerations

- ☒ Manuscript has been "spell & grammar-checked" Better, if it is revised by a professional science editor or a native English speaker
- ☒ References are in the correct format for this journal
- ☒ All references mentioned in the Reference list are cited in the text, and vice versa
- ☒ Colored figures are only used if the information in the text may be losing without those images
- ☒ Charts (graphs and diagrams) are drawn in black and white images; use shading to differentiate

**Title:**

Species of Aphids Found in Ornamental and Wild Plants in Highland, Pagar Alam, South Sumatra

**Author(s) name:**

Chandra Irsan<sup>a\*</sup>, Erise Anggraini<sup>a,b</sup>, Siti Herlinda<sup>a</sup>, Wenny Ramadhani<sup>c</sup>, M. Umar Harun<sup>d</sup>,

**Address**

(Fill in your institution's name and address, your personal cellular phone and email)

<sup>a</sup> Department of Plant Pests and Diseases, Faculty of Agriculture, Universitas Sriwijaya, Indralaya, Ogan Ilir, South Sumatra, Indonesia

<sup>b</sup> Agroecotechnology Study Program, Faculty of Agriculture, Universitas Sriwijaya, Indralaya, Ogan Ilir, South Sumatra, Indonesia

<sup>c</sup> Plant Quarantine, Palembang, Sumatera, Indonesia

<sup>d</sup> Department of Agronomy, Faculty of Agriculture, Universitas Sriwijaya, Indralaya, Ogan Ilir, South Sumatra, Indonesia

**For possibility publication on the journal:**

(fill in *Biodiversitas* or *Nusantara Bioscience* or mention the others)

- |  |   |
|--|---|
| <input checked="" type="checkbox"/> Biodiversitas Journal of Biological Diversity      | <input type="checkbox"/> Nusantara Bioscience                       |
| <input type="checkbox"/> Prosiding Seminar Nasional Masyarakat Biodiversitas Indonesia | <input type="checkbox"/> Asian Journal of Agriculture               |
| <input type="checkbox"/> Asian Journal of Ethnobiology                                 | <input type="checkbox"/> Asian Journal of Forestry                  |
| <input type="checkbox"/> Asian Journal of Natural Product Biochemistry                 | <input type="checkbox"/> Asian Journal of Tropical Biotechnology    |
| <input type="checkbox"/> International Journal of Bonorowo Wetlands                    | <input type="checkbox"/> Cell Biology and Development               |
| <input type="checkbox"/> Indo Pacific Journal of Ocean Life                            | <input type="checkbox"/> International Journal of Tropical Drylands |

**Novelty:**

(state your claimed novelty of the findings versus current knowledge)

This paper described the alternative host of aphids in high land, South Sumatera. The knowledge regarding the alternative of insect pest could be beneficial resource for basic control of aphids.

**Statements:**

This manuscript has not been published and is not under consideration for publication to any other journal or any other type of publication (including web hosting) either by me or any of my co-authors.

Author(s) has been read and agree to the Ethical Guidelines.

**List of five potential reviewers**

(Fill in names of five potential reviewers **that agree to review your manuscript** and their **email** addresses. He/she should have Scopus ID and come from different institution with the authors; and from at least three different countries)

1. Dr. Koko Dwi Sutanti (email: [ksutanto@ksu.edu.sa](mailto:ksutanto@ksu.edu.sa) )
2. Dr. Lau Wei Hong (email: [lauweih@upm.edu.my](mailto:lauweih@upm.edu.my))
3. Prof. Dr. Dra. Asni Johari, M.Si. ([johari\\_asni@yahoo.com](mailto:johari_asni@yahoo.com) )
4. Dr. Mahesh Gunasena ([mahesh.gunasena@gmail.com](mailto:mahesh.gunasena@gmail.com))
5. Dr. Hasber Salim ([hasbersalim@usm.my](mailto:hasbersalim@usm.my))

**Place and date:**

Palembang, 5 October 2023

**Sincerely yours,**

(fill in your name, no need scanned autograph)

Dr. Chandra Irsan

# Species of Aphids Found in Ornamental and Wild Plants in Highland, Pagar Alam, South Sumatra

## Abstract

Aphids are one of the crucial pests in the tropical and sub-tropical regions. The presence of aphids in a plant can be very detrimental due to their role as vectors. Aphids exhibit species diversity, but not much information has been reported about the species diversity of aphids associated with essential crops such as ornamental plants. Furthermore, many aphid species were found on plants that were not actually hosts such as wild plants. Therefore, this study reported the species of aphids found in ornamental plants and the wild plants. The field research employed a purposive and direct observation approach to inventory cultivated or wild plants hosting aphids and collecting aphids. The plant selection process included cultivated plants encompassing ornamental plants, as well as wild plants or weeds. The collection and identification of host plants, and aphids, involved systematic searches for the selected plants and subsequent examination for the presence of aphids. Observations were made to all existing plant species to find those colonized by aphids. This study revealed that 15 species of aphids were found in Pagaralam, namely *Aphis gossypii*, *Uroleucon* sp., *Toxoptera odinae*, *Macrosiphum rosae*, *Aphis citricola*, *Aphis craccivora*, *Toxoptera aurantii*, *Pentalonia nigronervosa*, *Hystenura* sp., *Aphis glycine*, *Greenidae* sp., *Rhopalosiphum padi*, *Rhopalosiphum maidis*, *Hyperomyzus* sp. *Lipaphis erysimi*.

**Keywords:** aphids, ornamental plants, wild plants

**Running title:** Aphids Found in Ornamental and Wild Plants

## INTRODUCTION

Aphids are one of the crucial pests in the tropics and sub-tropics, exhibiting various polyphagous, oligophagous, and monophagous characteristics (Kennedy & Stroyan, 1959). One species of aphids can host more than 400 species from 40 families (Bass et al., 2014). In addition to pests, aphids can also be vectors of plant viral diseases (Gadhawe et al., 2020). Aphids can transmit 275 viruses (Ertunc, 2020). In tropical areas, aphids can always be found throughout the year due to their parthenogenetic nature of reproduction (Blackman & Eastop, 2017). Aphids consume young leaves sap, which can deplete essential nutrients for healthy growth (Cao et al., 2018). Moreover, when aphids transmit viral diseases from one plant to another, this can further weaken and stunt the growth of infected plants (Jones, 2022). According to Kinley et al. (2021), aphids cause yield losses directly (35 - 40%) by sucking the plant sap or indirectly (20 - 80%) through viral transmission. Therefore, aphid infestations can have adverse effects on crop yields and overall plant health (Sarwan Kumar, 2019).

Due to their function as vectors, the presence of aphids on a plant can be highly damaging (Jaouannet et al., 2014). They feed by piercing the plant's tissues and consuming its sap, which can reduce the plant's growth and productivity, ultimately leading to weakness and possible death (Chandel et al., 2022). Additionally, as vectors, aphids can transmit a variety of plant diseases. They are as carriers for various plant viruses, and when they move from infected to healthy plants, these viruses can rapidly spread and cause extensive damage (Guo et al., 2019). In addition, the honeydew that aphids secrete can lead to the growth of sooty mold, a black fungus that can prevent sunlight from reaching the plant's

**Commented [Ma1]:** This paper has too long Introduction. It should consist about 600-700 words

leaves, thereby impairing photosynthesis, the process by which plants produce food (Singh & Singh, 2021). Therefore, it is crucial to control aphid populations in gardens and crops.

Understanding the species diversity of aphids is fundamental to effective aphid control, as it facilitates the development of measures to keep their populations in check. In addition, understanding the diversity of aphid species can provide valuable insights into potential plant diseases, as different aphid species carry distinct viruses. Methods used to control aphids often encompass various techniques, including the use of natural enemies such as predators (like ladybugs, lacewings, and parasitic wasps) (Singh & Singh, 2021; Völkl et al., 2023), parasitoids (Boivin et al., 2012), entomopathogens (Hullé et al., 2020), the use of essential oils as botanical pesticides to control aphids (Ikbal & Pavela, 2019), and crop rotation techniques (Degani et al., 2019). Regular monitoring of aphid populations and diversity can help in detecting when population sizes may be reaching harmful levels, allowing for prompt implementation of the necessary countermeasures.

Many aphid species were found on plants that were not their actual hosts (Maharani et al., 2018). Aphids have one or more secondary, or "alternative," host plants in addition to their primary host plants, which are the types of plants they feed on most frequently (Clarke et al., 2020). An alternative host can also be a collateral host belonging to the same plant family as the primary host, helping crop pests to survive when the primary hosts are unavailable (Kumar et al., 2021). These secondary hosts may offer less adequate nutrition for insects (Mo & Smilanich, 2023). However, they may provide a means of survival when primary hosts are unavailable, during certain seasons, or under certain environmental conditions (Kumar et al., 2021). According to Liu et al. (2017), since hibiscus serves as an overwintering host for cotton-specialized aphids but not for cucurbit-specialized aphids, it is evident that host-specialized aphids have refuges during times of food shortage. The life cycles of numerous aphid species exhibit such complexity (Jousselin et al., 2010). They maintain a cycle of host alternation, shifting between their primary hosts (typically a woody plant) and secondary hosts (often herbaceous plants) (Yamamoto et al., 2020). Weeds pose a continuous threat in both cropped and non-crop areas, providing food, shelter and reproductive sites for various pest organisms (Kumar et al., 2021). This indicates that weeds can serve as alternative hosts for aphids.

A study of aphid species on horticultural plants has been conducted (Maharani et al., 2018). However, information about aphid species on ornamental and wild plants has not received as much attention and remains largely unexplored. In South Sumatra, particularly in the highland areas like Pagar Alam, there are numerous ornamental and native plants. The research on the diversity of aphid species in ornamental and wild plants has received little attention. Therefore, this study was conducted in Pagar Alam, a highland region of South Sumatra, with the aim of obtaining information on the diversity of aphid species found in ornamental and wild plants. The findings from this study can serve as a valuable resource for aphid management.

## MATERIALS AND METHODS

The field research employed a purposive and direct observation approach to inventory cultivated or wild plants hosting aphids and collecting aphids. The plant selection process included cultivated plants encompassing ornamental plants, as well as wild plants or weeds. The collection and identification of host plants, and aphids, involved systematic searches for the selected plants and subsequent examination for the presence of aphids. Observations were made to all existing plant species to find those colonized by aphids. Any plants colonized by aphids were documented as aphid hosts. Aphids, along with their natural enemies within the aphid colonies, were systematically collected. All components of the collected observations were then identified.

Aphid identification was conducted using identification keys (Blackman & Eastop, 2008). Identification of aphid species took place in the Laboratory of Entomology, Faculty of Agriculture, Universitas Sriwijaya. Identification relied on morphological characteristics. The host plants were identified using weed identification hand book (Kallas, 2010; Meuninck, 2023; Naidu, 2012). The location and size of aphid colonies, morphology of aphids including their shape and color, as well as any symptoms observed in the host plants were recorded, and photographs of the aphid colonies and their host plants were taken.

RESULT AND DISCUSSION

Result

The results showed that 15 aphid species were found in Pagaram, namely *Aphis gossypii*, *Uroleucon* sp., *Toxoptera odinae*, *Macrosiphum rosae*, *Aphis citricola*, *Aphis craccivora*, *Toxoptera aurantii*, *Pentalonia nigronervosa*, *Hystenura* sp., *Aphis glycine*, *Greenidae* sp., *Rhopalosiphum padi*, *Rhopalosiphum maidis*, *Hyperomyzus* sp. *Lipapis erysimi*. Based on the observation, these aphids were found on various ornamental plants (Table 1). The primary colony locations were generally in flowers, and this study documented these colony locations in ornamental plants (Figure 1).

Table 1: Aphid species found in ornamental plants and their colony locations.

No	Host Plant	Aphid Species	Colony location
1	<i>Aster alpinus</i>	<i>Sitobion luteum</i>	flower
2	<i>Brugmansia suaveolens</i>	<i>Aulacorthum solani</i>	flower
		<i>Neomyzus circumflexus</i>	
		<i>Myzus persicae</i>	
3	<i>Caladium</i> sp.	<i>Pentalonia</i> sp	flower
4	<i>Cananga odoratum</i>	<i>Aphis gossypii</i>	flower
5	<i>Canna indica</i>	<i>Pentalonia nigronervosa</i>	flower
6	<i>Catharanthus roseus</i>	<i>Aphis citricola</i>	flower
7	<i>Cestrum</i> sp.	<i>Aphis gossypii</i>	flower
		<i>Neomyzus circumflexus</i>	
8	<i>Clitoria ternatea</i>	<i>Aphis craccivora</i>	flower
9	<i>Cosmos caudatus</i>	<i>Uroleucon</i> sp.	flower
10	<i>Dahlia Kelvin</i>	<i>Aphis gossypii</i>	flower
11	<i>Dendrobium</i> sp.	<i>Sinemogoura citricola</i>	flower
12	<i>Duranta</i> sp.	<i>Aphis gossypii</i>	flower
13	<i>Helianthus</i> sp.	<i>Aphis glycines</i>	flower
		<i>Hyperomyzus</i> sp.	
14	<i>Hibiscus rosinensis</i>	<i>Aphis gossypii</i>	flower
15	<i>Ixora paludosa</i>	<i>Aphis gossypii</i> ,	flower
		<i>Toxoptera aurantii</i>	
16	<i>Ixora</i> sp.	<i>Aphis citricola</i>	flower
		<i>Aphis gossypii</i>	
		<i>Toxoptera aurantii</i>	
17	<i>Murraya paniculata</i>	<i>Aphis craccivora</i>	flower
		<i>Toxoptera citricidus</i>	
18	<i>Mussaenda frondosa</i>	<i>Aphis citricola</i>	flower
		<i>Toxoptera odinae</i>	
19	<i>Rosa indica</i>	<i>Macrosiphum rosae</i>	flower
20	<i>Spondiras dulcissoland</i>	<i>Aphis citricola</i>	flower
		<i>Hysteroneura setariae</i>	



Fig 1. The location of aphid colonization on various plant parts. a) *A. gossypii* in *D. Kelvin* flower b) *A. gossypii* in *H. rosasinensis* flower c) *A. gossypii* in tuberose flower, d) *A. craccivora* in *Clitoria ternatea* flower, e) *A. citricola* in *Helianthus* sp., f) *A. aurantii* on the *M. paniculata* flower, g) *T. odinae* in the *S. dulcissoland*, h) *Uroleucon* sp. in chrysanthemums, i) *Macrosiphum rosae* in *R. indica* flower, j) *Pentalonia nigronervosa* in *C. indica* leaves

In addition, this study documented the presence of weeds, which might serve as alternative hosts for aphids (Table 2). The location of aphid colonies also varied, namely on flowers, stalks, plant tops, young leaves and old leaves of wild plants (Figure 2). The presence of specific plants or host plants within a habitat influenced the types of aphids found. Many aphid species are found on a broad range of plants or host plants, while others are highly specialized and are only found on specific plants or host plants. This is closely related to the polyphagous, oligophagous or monophagous nature of aphids (Blackman & Eastop 2000).

124 Table 2: Species of aphids found in wild plants and their colony locations.

No	Host Plant	Aphid species	Colony location
1	<i>Ageratum conyzoides</i>	<i>Aphis gossypii</i>	Shoots, young leaves, old leaves, flowers
2	<i>Alternanthera philoxeroides</i>	<i>Aphis gossypii</i>	Shoots, buds
3	<i>Alternanthera sessilis</i>	<i>Aphis gossypii</i>	Shoots, buds
4	<i>Amaranthus gracilis</i>	<i>Aphis craccivora</i>	Flowers, shoots, young leaves, old leaves
5	<i>Blumea lacera</i>	<i>Lipaphis erysimi</i>	Flowers, shoots, and buds
6	<i>Croton hirtus</i>	<i>Aphis gossypii</i>	Flowers, shoots, young leaves, old leaves young twigs
7	<i>Cynodon dactylon</i>	<i>Schizaphis rotundiventris</i>	Flower, flower stalks
8	<i>Cyperus rotundus</i>	<i>Schizaphis rotundiventris</i>	Flower, flower stalks, leaf axils
9	<i>Cyperus compressus</i>	<i>Schizaphis rotundiventris</i>	Flower, flower stalks, leaf axils
10	<i>Digitaria ciliaris</i>	<i>Hysteroneura setariae</i>	Flower, flower stalks
11	<i>Echinocloa crusgali</i>	<i>Hiperomyzus</i> sp.	Young leaves, old leaves
12	<i>Ecliptica prostrata</i>	<i>Aphis gossypii</i>	Shoots, young leaves
13	<i>Eleusin indica</i>	<i>Hysteroneura setariae</i> <i>Rhopalosiphum maidis</i>	Flower, flower stalks, leaf axils
14	<i>Emilia sonchifolia</i>	<i>Aphis gossypii</i>	Flower, flower stalks, shoots
15	<i>Eragrostis tenella</i>	<i>Hysteroneura setariae</i>	Flower, flower stalks, seeds
16	<i>Euphorbia hirta</i>	<i>Aphis gossypii</i>	Young leaves, old leaves
17	<i>Eupotarium odoratum</i>	<i>Aphis gossypii</i> , <i>Aphis glycine</i>	Young leaves, old leaves, young twigs
18	<i>Hymenochera acutigluma</i>	<i>Hysteroneura setariae</i>	Flowers, flower stalks, leaf axils
19	<i>Lagerstromia</i> Sp.	<i>Greenidea</i> sp.	Young leaves
20	<i>Lophatherum gracile</i>	<i>Hysteroneura setariae</i> <i>Rhopalosiphum maidis</i>	Young leaves, old leaves, leaf axils
21	<i>Melastoma affine</i>	<i>Aphis gossypii</i>	Shoots, young leaves
22	<i>Mikania mikranta</i>	<i>Aphis gossypii</i> <i>Aphis glycine</i>	Shoots, young leaves, old leaves
23	<i>Mimosa invisa</i>	<i>Aphis craccivora</i>	Shoots, pods
24	<i>Mimosa pudica</i>	<i>Aphis craccivora</i>	Shoots, pods, flowers
25	<i>Mimosa vigra</i>	<i>Aphis craccivora</i>	Shoots, pods
26	<i>Oryza rufipogon</i>	<i>Rhopalosiphum padi</i> , <i>Rhopalosiphum maidis</i>	Old leaves, young leaves (pupus), leaf axils
27	<i>Oxonopus compressus</i>	<i>Hysteroneura setariae</i>	Flower, flower stalk, leaf axils
28	<i>Paspalum conjugatum</i>	<i>Hysteroneura setariae</i>	Flower, flower stalk, seeds
29	<i>Phyllanthus neruri</i>	<i>Aphis citricola</i>	Shoot, young leaves, old leaves, young twigs petioles
30	<i>Portulaca oleraceae</i>	<i>Aphis craccivora</i>	Shoots, young leaves, flower
31	<i>Physalis angulata</i>	<i>Aphis craccivora</i> , <i>A. gossypii</i>	Shoots, young leaves, old leaves
32	<i>Rorippa indica</i>	<i>Lipapis erysimi</i>	Flower, fruit, shoots, young leaves
33	<i>Sida rhombifolia</i>	<i>Aphis gossypii</i>	Shoots, young leaves, old leaves, fruit/seeds
34	<i>Sonchus arventris</i>	<i>Lipapis erysimi</i>	Young leaves, fruit stalks, flower, fruit

125

126

127





Figure 2. Aphids found on wild plants a) *A. gossypii* on the weed *Ageratum conyzoides*, b) *A. gossypii* on Croton weed *hirtus* c) *A. gossypii* on the weed *Eupatorium odoratum*, d) *A. gossypii* on plants *Pachystochys* sp., e) *A. gossypii* on plants *Caladium* sp., f) *A. gossypii* on the weed *Alternanthera sessilis*, g) *A. gossypii* in *Portulaca oleraceae* weeds, h) *A. gossypii* on the weed *Euphorbia hirta*, i) *A. citricola* on the weed *Phyllanthus nerruri*, j) *A. citricola* on *Sida rhombifolia* weed, k) *A. citricola* on plants *Annona muricata*, l) *A. citricola* on the weed *Ludwigia peruviana*, m) *A. craccivora* on *Mimosa pudica* weed, n) *A. craccivora* on weeds *Amaranthus gracilis*, o) *A. glycine* in *Mikania micrantha* weed, p) *Hysteneura* sp. in Eleusine weeds, q) *Greenidae* sp. in kenidai trees (shrubs) *indica*, r) *Hyperomyzus* sp. in *Echinocloa crusgali* Weed, s) *L. erysimi* on weed *sonchus arvensis*, t) *Rhopalosiphum* rice on the weed *Oryza rufipogon*, u) *Rhopalosiphum* *Maidis* on the weed *Oryza rufipogon*.

## Discussion

The plant species or host plant influences the distribution of aphids. There are aphid species that can be found on a wide range of host plants, which is closely related to the polyphagous nature of aphids, allowing them to colonize many different species of host plants. Host plants can also affect the distribution of aphids, as evidenced by the presence of aphid species exclusively found on certain host plants (Santiago et al., 2017). But there are some species of aphids found only on one particular host and are not found on other host plants (Döring, 2014). *A. gossypii*, and *Aphis aurantii* have been found on many host plants because both aphids are classified as polyphagous aphids (Alotaibi et al., 2023).

Aphids can commonly be found infesting a variety of ornamental plants. They are attracted to these plants due to the rich nutrient content in the plant sap (Braham et al., 2023). In this present study, some aphid species were found on some ornamental plants in Pagaralam. The location of aphid colonization on the plants varied. On *Adiantum predatum* plants, aphids formed colonies on young leaf stalks and on newly emerging leaves. The aphids displayed brown and black coloration. The aphid colonies found were small, and the colonized plant parts showed no signs of disease. The identification results showed that the aphids were *Neotoxoptera* sp., and notably, they were not associated with ants. On *Aster alpinus*, aphids were found to form colonies on the stems or young leaf shoots, and the colonies were relatively large. The color of the aphids was dark brown to black. The colonized plant parts showed symptoms of stunting. The identification results showed that the aphids were *Uroleucon* sp., and they were associated with ants.

On the *Brugmansia suaveolens*, *M. persicae* were found on the undersides of old leaves or leaves that have started to turn yellow. The colonies were relatively small. The aphids found were green and large bodies. The colonized plant parts did not show any signs of disease. On *Caladium* sp. (taro) was found one species of aphids: *A. gossypii*. The aphids formed colonies under the surface of young and older leaves. The occupied leaf areas did not display severe symptoms. The aphids were yellow green to dark green. The wingless adult aphids often had a white, flour-like appearance on their bodies. On the *Cananga odoratum* (ylang-ylang), colonies of *T. aurantii* were found on the undersides of the leaves, the shoots, buds, and unopened flower petals. The *T. aurantii* colonies found were relatively large. Colonized parts, especially shoots, showed signs of stunting. The aphids found were brown to black in color. The colonies of *T. aurantii* were found to be associated with black ants.

Aphids on *C. indica* (Indian shot, African arrowroot) were found to form colonies in the leaf axils and under the leaf surface near the leaf base. The colonies were quite large. The aphids were dark brown to dark red coloring with a medium-sized body. The identification results showed that the aphids were *P. nigronervosa*. The colonies of *P. nigronervosa* were found to be associated with ants. In the *Catharanthus roseus* (periwinkle), *A. citricola* aphids were found. The aphids were yellow-green, sifunculi, and black cauda. The aphids formed colonies on flowers and shoots, and the colonized plant parts did not show any symptoms of disease. On *Cestrum* sp. (Bastard jasmine), aphids formed colonies on the undersides of young leaves, shoots, and within flower parts, especially between petals or flower stalks that had not fully bloomed. The colonies were quite large. The body color of aphids was green to dark green with small to medium-sized bodies. The colonized plant parts, especially leaves, showed stunting symptoms. The identification results showed that the aphids were *A. gossypii*. The aphid colonies found were consistently associated with ants.

Aphids on *Clitoria ternatea* were found to form colonies on flower parts, flower crowns, stems and young leaves. The aphids were brown to black in color. Colonized plant parts, especially shoots and young leaves, showed stunting symptoms. The identification results showed that the aphids were *A. craccivora*. These colonies were consistently associated with ants. On the plant *Cosmos caudatus*, aphids were found on the flower petals. The colonies were not very large. The body color was green and light green. The identification results showed that the aphids were *A. gossypii*, and

they were also associated with ants. The aphids on the *Dahlia kelin plant* formed colonies on unopened flower buds, with a significant population among the blooming petals. The body color was green to dark green. The identification results showed that the aphids were *A. gossypii*. Aphids on *Datura metel* (amethyst) were found to form colonies on the undersides of old leaves. The aphids were medium sized with a green body color. The colonized plant parts did not show any symptoms of disease. The identification results showed that the aphids were *Myzus ornatus*. The aphid colonies were not associated with ants. Within *Dendrobium* sp., aphid colonies were found on the young leaves. The aphids were yellow, green to dark green. The colonized plants did not show any disease symptoms. The identification results showed that the aphids were *A. gossypii*, and they were associated with ants. On *Duranta* sp. (bonsai), colonies of aphids were located on the undersides of young leaves. The colonized plant parts showed stunting symptoms. The colonies were very large. The aphids were green in color. The identification results showed that the aphids were *A. gossypii*. The aphid colonies were consistently associated with ants.

On the *Helianthus annuus* (sunflower) plants, aphid colonies were found between the flower petals. The colonized flowers, especially the crowns, exhibited a tendency to fall off easily. The aphids were green and yellow in color. The colonies were small. The identification results showed that the aphids were *A. gossypii*. These aphid colonies were associated with ants. Aphid colonies on *Helianthus* sp. were found on the undersides of old leaves. These colonies were small in size. The aphids were green with a medium body size. The colonized plant parts did not show any disease symptoms. The identification results showed that the aphids were *M. ornatus*. The aphid colonies were not associated with ants. Within the colonies, mummified aphids that were parasitized by Aphidiidae were found.

On the *Hibiscus rosa-sinensis*, aphids ranging in color from yellow to dark green were found. The aphids formed colonies on flower buds, unopened flower crowns, and the undersides of aging leaves. The colonies grew to be very large. The identification results showed that the aphids were *A. gossypii*. The aphid colonies were consistently associated with ants. Two types of aphids were found on the flowering plant *Ixora paludosa*. First, the aphids formed colonies on the undersides of young leaves that were still red or light green and sometimes on flower stalks that had not yet bloomed. The occupied plant parts showed symptoms such as stunted leaf growth, leaf shrinkage, necrotic spots on the leaf surface, and slightly downward-curved leaf edges. The upper leaf surface looked wet and sticky, similar to sugar. The aphids had yellow, green, or slightly dark green bodies, with some wingless adults having a powdery white upper surface. The identification results showed that the aphids were *A. gossypii*, and they were almost always associated with ants. The second type of aphids on *Ixora paludosa* formed colonies under the surface of young and older leaves. The colonies could also be found on newly emerging flowers and leaves. The plant parts occupied by these aphids did not show obvious signs of illness. These aphids were dark red to black, with once-branched stigma and venation in their black wings. The identification results showed that the aphids were *T. aurantii*. These aphids were also associated with ants.

In *Ixora* sp. flower plants, two forms of aphids were discovered. These aphids occupied the shoots, young leaves and unopened flowers. The affected plant parts did not show obvious symptoms. The aphids exhibited colors ranging from yellow and green to a slightly darker green. Sometimes the upper surface of the wingless imago's body appeared white, resembling flour. The identification results showed that these aphids were *A. gossypii*. These aphid colonies were almost always associated with ants. Another species of aphids was founded and formed colonies on flower stalks that had not yet bloomed and on newly emerging shoots or leaves. The presence of these aphids on the plant did not induce any symptoms of plant disease. The aphids were yellow or yellow green, with black cauda and siphunculi. Their bodies were very small to small. The identification results showed that the aphids were *A. citricola*. The colonies of *A. citricola* were also frequently found in association with ants.

Two types of aphids were found on *Mussaenda frondosa*, each forming colonies in different locations. The first type formed colonies on young leaves, shoots, and flowers. The plant parts they occupied showed no obvious disease symptoms. The aphids were yellow, green, and some with dark green. The identification results showed that the aphids were *A. gossypii*. The second type of aphids formed colonies on the stems or young twigs, appearing densely clustered as if piled up. The aphid colonies could also be found on young leaves, shoots and within flower parts. The plant parts they infested showed no signs of diseases. The aphids were yellow or yellow green, with black cauda and siphunculi. They had tiny to small bodies. The identification results showed that the aphids were *A. citricola*.

The results showed that 34 species of wild plants, including weeds, were growing in the yard colonized by aphids. This indicated that multiple species of aphids colonized various host plants. The aphid species colonizing these plants were generally consistent within the same taxon. *Ageratum conyzoides* was infested by *Aphis gossypii*. These aphids formed colonies on the flower sections, shoots, lower surfaces of young leaves, or leaves turning yellow. The aphids were green, yellow-green to dark green, often forming large colonies. *Alternanthera philoxeroides* or alligator grass was also colonized by *Aphis gossypii*. Small colonies were found on shoots or stems. These aphids had small bodies and were green, ranging from yellow-green to dark green. *Alternanthera sessilis* was colonized by *Aphis gossypii*, forming colonies on shoots, flowers, and fruit. The colonies were typically large, and they were often associated with tiny brown ants. *Amaranthus gracilis* was infested by *Aphis craccivora*. These aphids established colonies on shoots, flowers and young and old leaves. They were dark brown to black in color, with shiny black wingless imagoes. Colonies of these aphids were associated with both black and red ants. *Blumea lacera* was colonized by *Lipaphis erysimi* aphids. These aphids were bright green, and of medium size. The colonies formed on flowers, flower stalks and the undersides of the leaves at the top. The aphid colonies were not associated with ants. *Croton hirtus* or fire grass was infested by *Aphis gossypii*. The aphids were yellow-green to dark green. The colonies were found on the stems, leaves, buds and flowers, often forming large colonies. *Cynodon dactylon* or Bermuda grass was colonized by *Schizaphis rotundiventris*. The aphids colonized the flowers, flower stalks and sometimes in the leaf axils of the plant. Small colonies were formed. The aphids were brown to red-brown. They were associated with ants. *Cyperus rotundus* or nut grass was infested by *Schizaphis rotundiventris* aphids. The colonies were found on flower stalks, flowers, and leaf axils. The colonies were quite large and associated with both black and red ants. The aphids were dark brown in color. *Cyperus compressus* or grass puzzle was colonized by *Schizaphis rotundiventris* aphids, forming colonies in the flowers, flower stalks and sometimes in the axils and leaves of the shoots or buds. Small colonies were observed. *Digitaria ciliaris* was infested by *Hysteroneura setariae* aphids, with small colonies scattered on the flowers and flower stalks. These aphids were light brown to brown in color. *Echinocloa crusgali* or water hyacinth plants were colonized by *Hipermizus* sp. aphids. These aphids were dark brown to black and formed large colonies on the undersides of both young and old leaves. The aphid colonies were never found in association with ants. *Ecliptica prostrata* or urang aring was colonized by *Aphis gossypii*, forming small colonies on the shoots and flowers. The aphids were bright green to blackish green. The aphid colonies were also consistently associated with ants.

*Eleusine indica* was colonized by two species of aphids: *Hysteroneura setariae* and *Rhopalosiphum maidis*. *H. setariae* formed colonies in flower parts, flower stalks and leaf axils resulting in quite large colonies. *H. setariae* body color ranged from red brown to dark brown. The colonies were consistently associated with ants. The aphids of *R. maidis* formed colonies in the leaf axils and undersides of leaves and on leaf shoots that had not yet opened. The colonies were not densely packed. The leaf aphids of *R. maidis* were green in color, with distinct black siphunculi and cauda. These aphids had relatively large bodies with a slightly elongated shape. *R. maidis* colonies were always associated with ants.

291 The plant *Emilia sonchifolia*, characterized by its purple flowers, was colonized by *Aphis gossypii*. The aphids were yellow  
292 to green in colour. The colonies formed near flowers, flower stalks, and shoot leaves.

293 *Eragrostis tenella* was infested by *Hysteroneura setariae* aphids. The aphids were brown to red brown. Small  
294 colonies formed on flowers near the seeds, with groups of aphids surrounding the plant's seeds. The aphids of *H. setariae*  
295 were consistently associated with ants. *Euphorbia hirta* or wart grass was colonized by *Aphis gossypii*. The aphids formed  
296 colonies on the undersides of leaves, resulting in stunted growth of the leaves. The aphids were yellow to dark green in  
297 color. *A. gossypii* colonies on *E. hirta* plants were consistently associated with ants. *Eupotarium odoratum* was colonized  
298 by both *Aphis gossypii* and *Aphis citricola*. *A. gossypii* formed colonies in the buds, young leaves, old leaves, and young  
299 twigs. Young leaves that were colonized by *A. gossypii* became stunted with an irregular shape. *A. gossypii* found in this  
300 plant showed yellow-green to dark green in body colour. The colonies of *A. citricola* formed on the young twigs near the  
301 shoots, with these aphids displaying yellow-green coloration and having black siphunculi and cauda. Aphid colonies of  
302 both *A. gossypii* and *A. citricola* on *E. odoratum* plants were associated with either black or red ants.

303 *Hymenochera acutigluma* or hair axis was colonized by *Hysteroneura setariae*, which formed colonies on the  
304 flower stalks and flowers. The colonized parts of the plants did not display any noticeable symptoms. *Lagerstromia sp.* or  
305 *kenidai*, was infested by *Greenidae* sp. These aphids had bright green bodies and distinctive elongated siphunculi with  
306 thorns. The aphids formed colonies on the undersides of leaves, especially on young leaves. The colonized leaves did not  
307 show any disease symptoms. *Lophatherum gracile* or bamboo grass plants were colonized by two species of aphids:  
308 *hysteroneura setariae* and *Rhopalosiphum maidis*. The aphids of *H. setariae* formed colonies on the undersides of leaves,  
309 leaf shoots, and leaf axils. The colonized leaves did not show any disease symptoms. *H. setariae* aphids were brown to  
310 red-brown. *R. maidis* aphids also formed colonies on the undersides of leaves, but the colonies were small. *R. maidis*  
311 aphids were green to bright green in color, with black siphunculi and cauda. It was possible for colonies of the two species  
312 of aphids on *L. gracile* to mix.

313 *Melastoma affine* was colonized by *Aphis gossypii*. The colonies formed on shoots, particularly near newly  
314 emerging shoots and on newly emerging fruits and flowers. The body colour of aphids ranged from yellow to green. The  
315 colonized plant parts did not show any disease symptoms. *Mikania miranta* was colonized by *Aphis gossypii* and *Aphis*  
316 *glycine*. *A. gossypii* formed colonies on the shoots, especially on the undersides of the leaves, resulting in stunted and  
317 curled leaves. *A. glycine* formed colonies on the branches. The colonies were densely populated. *A. Glycine* aphids were  
318 light green to green in color. The colonized plant parts became distorted. The two species of aphids could mix to form a  
319 single colony.

320 *Mimosa invisa* (cater-grass) was colonized by *Aphis craccivora*. The aphids of *A. craccivora* on *M. invisa* plants  
321 formed colonies only on the shoots with small colonies. The aphids appeared dark black with wingless imagoes. *Mimosa*  
322 *pudica* was observed to be colonized by *Aphis craccivora*. The aphids formed colonies on shoots, especially young shoots,  
323 and occasionally on flowers and pods. The aphids were black and of medium size, resulting in stunted growth of the  
324 colonized plant parts. The colonies were quite large. *Mimosa vigra* was colonized by *Aphis craccivora*. The colonies of  
325 aphids occupied the pods and shoots with small colonies. The nymphs of aphids were black, and wingless adults were  
326 shiny black. The colonized plant parts did not show any disease symptoms.

327 *Oryza rufipogon* was colonized by two species of aphids: *Rhopalosiphum rice* and *Rhopalosiphum maidis*. Both  
328 aphids colonized the same plant parts, namely the unopened leaves and the leaf axils with large colonies. The two species  
329 could be distinguished by their body color. *R. maidis* appeared green with black sifunculi and cauda, while *R. rice*  
330 appeared white. The colonies of *R. maidis* and *R. rice* in *O. rufipogon* plants were associated with the presence of red ants.

331 *Oxonopus compressus* or *pait* grass was colonized by *Hysteroneura setariae* aphids. The colonies occupied flowers, flower  
332 stalks, seeds, and sometimes in the leaf axils. The aphids were brown to dark brown in color. Small colonies were formed,  
333 and they were also consistently associated with ants.

334 *Paspalum conjugatum* was colonized by *H. setariae* aphids. The colonies occupied flower parts, especially the  
335 seeds and flower stalks. Aphids had brown to dark brown bodies. *Phyllanthus niruri* was colonized by *Aphis citricola*. The  
336 colonies formed on the shoots and the undersides of leaves and petioles. The colonized parts became distorted, stunted,  
337 and wrinkled. The aphids had yellow bodies with black sifunculi and cauda, and the colonies formed were quite large.  
338 *Portulaca oleraceae* plants were colonized by *Aphis craccivora*. The aphids of *A. craccivora* in *P. oleraceae* plants  
339 formed colonies on the undersides of leaves, especially young leaves, shoots and in flowers. The colonized plant parts  
340 became stunted, and leaf edges curled downward. The aphids had dark brown to black bodies, with wingless imagoes that  
341 appeared glossy black.

342 *Physalis angulata* plants were colonized by *Aphis craccivora*. The aphids had dark green to black bodies, with  
343 glossy black wingless imagoes. *A. craccivora* formed colonies on the shoots or near the leaf buds. The colonized plant  
344 parts did not show any symptoms of disease. *Rorippa indica* or mustard land was colonized by *Lipaphis erysimi*. The  
345 colonies formed on the flowers, fruits, flower stalks and the lower surface of leaves. The colonized plant parts showed  
346 symptoms such as curling and stunting. *Sida rhombifolia* or cacabean was colonized by *Aphis gossypii*. The aphids had  
347 green-yellow to green body colors. The colonies formed on the surface of lower leaves, stalks and flower petals. The  
348 colonized plant parts, especially the shoots, showed curling, and the leaf edges curled downward. *Sonchus arvensis* plants  
349 were colonized by *L. erysimi*. The aphids had green to whitish green body colours, and the colonies formed on flower  
350 stalks, under petals, and on young shoots or leaves. The colonized plant parts became stunted over time.

351 In general, aphids observed on ornamental and wild plants formed colonies. The colonized plant parts typically  
352 displayed typical symptoms of damage, but some did not show any symptoms. Generally, the symptoms of the plants  
353 caused by aphid colonies were relatively the same, such as stunted growth, abnormal shape, and stunted or curly leaves.  
354 These characteristic symptoms serve as indicators of aphid infestations. However, some plants or plant parts did not show  
355 symptoms when colonized by aphids. This condition happened because the colonized parts had reached their maximum  
356 growth or development. It indicated that the colonized part was not currently undergoing a growth phase. The colonies that  
357 did not induce symptoms typically occurred when the colonized leaves had reached their maximum growth or when the  
358 leaves and plant parts were old. The old leaves or twigs might not show the typical symptoms associated with aphid  
359 infestations.

360 The part of the plant exhibiting characteristic symptoms when colonized by aphids also often experienced a  
361 cessation in growth due to the piercing by the aphids. In contrast, the areas surrounding the puncture site continued to  
362 grow, resulting in some parts developing normally while others become stunted (Pettersson et al., 2017). This condition  
363 could lead to the bending of shoots or young stems, curling of leaves, downward curling of leaf edges, or stunted leaf  
364 growth. In this observation, monocot plants or groups of grasses with narrow leaves generally did not display any  
365 distinctive symptoms when colonized by aphids. This might be because the growth or development of their leaves differed  
366 from that of dicot plants. Therefore, the presence of aphids in monocot plants or plants was often easier to recognize  
367 through the presence of ants. If a plant was found to have a significant number of ants, there was a possibility that aphids  
368 had colonized the plant (Tegelaar et al., 2012). Therefore, the presence of ants could serve as an indicator of the presence  
369 of aphid colonies.

Throughout their life cycle, aphids exhibited host alternation by switching between two distinct host plants (Yamamoto et al., 2020). They overwintered on woody plants, reproduced in the spring, and migrated to herbaceous plants during the summer before returning to their primary host in the autumn (Yamamoto et al., 2020). This allowed aphids to maximize resource utilization, avoid congestion and competition, evade predators and parasites, circumvent plant defenses, and colonize new areas. Aphids could distribute their population efficiently, thereby avoiding overcrowding, predators and parasites, and plant defenses developing over time through host switching (Yamamoto et al., 2020). This behavior was essential for the survival and environmental adaptation of aphids.

Aphids colonized flowers because they may offer an accessible and rich food source, sugary plant sap found in new growth or reproductive parts of plants. Flowers contain a nutrient-rich nature and easy access to sap, therefore aphids were attractive to sap the flowers. Some aphid species were drawn to certain colors (Jakubczyk et al., 2022). Herbs served as an alternative host for aphids in this present study. Aphids consume sugar-rich liquid in plants, known as "sap". Aphids considered herbs and other green vegetation as abundant food sources. Aphids utilize needle-like mouthparts to penetrate plant tissues and access this fluid (Brožek et al., 2015). Several aphids colonized herbs such as Indian mustards, *Lipaphis erysimi*, and *Myzus persicae* are the most devastating insects, infesting leaves, stems, and floral parts (Jayaswal et al., 2022).

Due to a symbiotic relationship, the prevalence of aphids and ants was frequently correlated. Aphids produced a delicious substance known as honeydew as a waste product, which ants found highly attractive as a food source (Nelson & Mooney, 2022). The honeydew contained an abundance of sugars, extracted by aphids from the plant juice (Zheng et al., 2022). Ants were drawn to this nutrient-rich food source and would often 'farm' aphids for it. In exchange for honeydew, ants provided aphids with protection from other insects and predators, such as ladybugs, lacewing larvae, and parasitic wasps (Karami-jamouri et al., 2018). Certain species of ants would transport aphids to new host plants for improved foraging opportunities, ensuring that aphids had a continuous food source (Giannetti et al., 2021). Honeydew not only nourished the ant colony, but its high sugar content also supported the development of their fungus farms (in certain species) and provided energy for the growth of their own progeny (Biedermann & Vega, 2020).

#### CONCLUSION

15 species of aphids were found in ornamental and wild plants in Pagaram, namely *Aphis gossypii*, *Uroleucon* sp., *Toxoptera odinae*, *Macrosiphum rosae*, *Aphis citricola*, *Aphis craccivora*, *Toxoptera aurantii*, *Pentalonia nigronervosa*, *Hystenura* sp., *Aphis glycine*, *Greenidae* sp., *Rhopalosiphum padi*, *Rhopalosiphum maidis*, *Hyperomyzus* sp. *Lipaphis erysimi*.

#### ACKNOWLEDGMENTS

The authors thank Universitas Sriwijaya, that supported this research.

#### REFERENCES

- Alotaibi, N. J., Alsufyani, T., M'sakni, N. H., Almalki, M. A., Alghamdi, E. M., & Spittler, D. (2023). Rapid Identification of Aphid Species by Headspace GC-MS and Discriminant Analysis. *Insects*, 14(589). <https://doi.org/https://doi.org/10.3390/insects14070589>
- Bass, C., Puinean, A. M., Zimmer, C. T., Denholm, I., Field, L. M., Foster, S. P., Gutbrod, O., Nauen, R., Slater, R., & Williamson, M. S. (2014). The evolution of insecticide resistance in the peach potato aphid, *Myzus persicae*. *Insect Biochemistry and Molecular Biology*, 51, 41–51. <https://doi.org/10.1016/j.ibmb.2014.05.003>
- Biedermann, P. H. W., & Vega, F. E. (2020). Ecology and evolution of insect–fungus mutualisms. *Annual Review of Entomology*, 65, 431–455. <https://doi.org/https://doi.org/10.1146/annurev-ento-011019-024910>
- Blackman, R. L., & Eastop, V. F. (2008). *Aphids on the world's herbaceous plants and shrubs, 2 volume set*. John Wiley

**Commented [Ma2]:** Please follow the guidance for reference writing (<https://smujo.id/biodiv/guidance-for-author>)



413 & Sons.

414 Blackman, R. L., & Eastop, V. F. (2017). Taxonomic issues. In *Aphids as crop pests* (pp. 1–36). CABI Wallingford UK.

415 Boivin, G., Hance, T., & Brodeur, J. (2012). Aphid parasitoids in biological control. *Canadian Journal of Plant Science*,  
416 92(1), 1–12. <https://doi.org/DOI: 10.4141/cjps2011-045>

417 Braham, M., Boulahia-kheder, S., Kahia, M., & Nouria, S. (2023). Aphids and citrus responses to nitrogen fertilization.  
418 *Journal of the Saudi Society of Agricultural Sciences*, 22(6), 374–383. <https://doi.org/10.1016/j.jssas.2023.03.003>

419 Brożek, J., Mróz, E., Wylęzek, D., Depa, Ł., & Węgierek, P. (2015). The structure of extremely long mouthparts in the  
420 aphid genus *Stomaphis* Walker (Hemiptera: Sternorrhyncha: Aphididae). *Zoomorphology*, 134, 431–445.  
421 <https://doi.org/https://doi.org/10.1007/s00435-015-0266-7>

422 Cao, H., Zhang, Z., Wang, X., & Liu, T. (2018). Nutrition versus defense : Why *Myzus persicae* ( green peach aphid )  
423 prefers and performs better on young leaves of cabbage. *PloS One*, 13(4), 1–16.  
424 <https://doi.org/https://doi.org/10.1371/journal.pone.0196219>

425 Chandel, R. S., Chandla, V. K., Verma, K. S., & Pathania, M. (2022). *Chapter 21 - Insect pests of potato in India: biology*  
426 *and management* (A. Alyokhin, S. I. Rondon, & Y. B. T.-I. P. of P. (Second E. Gao (eds.); pp. 371–400). Academic  
427 Press. <https://doi.org/https://doi.org/10.1016/B978-0-12-821237-0.11001-7>

428 Clarke, R., Kehoe, M. A., Broughton, S., & Jones, R. A. C. (2020). Host plant affiliations of aphid vector species found  
429 in a remote tropical environment. *Virus Research*, 281(December 2019), 197934.  
430 <https://doi.org/10.1016/j.virusres.2020.197934>

431 Degani, E., Leigh, S. G., Barber, H. M., Jones, H. E., Lukac, M., Sutton, P., & Potts, S. G. (2019). Crop rotations in a  
432 climate change scenario: short-term effects of crop diversity on resilience and ecosystem service provision under  
433 drought. *Agriculture, Ecosystems & Environment*, 285, 106625.  
434 <https://doi.org/https://doi.org/10.1016/j.agee.2019.106625>

435 Döring, T. F. (2014). How aphids find their host plants, and how they don't. *Annals of Applied Biology*, 165(1), 3–26.  
436 <https://doi.org/https://doi.org/10.1111/aab.12142>

437 Ertunc, F. (2020). *Chapter 46 - Emerging Plant Viruses* (M. M. B. T.-E. and R. V. P. Ennaji (ed.); pp. 1041–1062).  
438 Academic Press. <https://doi.org/https://doi.org/10.1016/B978-0-12-819400-3.00046-6>

439 Gadhave, K. R., Gautam, S., Rasmussen, D. A., & Srinivasan, R. (2020). Aphid transmission of Potyvirus: the largest  
440 plant-infecting RNA virus genus. *Viruses*, 12(7), 773. <https://doi.org/doi: 10.3390/v12070773>

441 Giannetti, D., Mandrioli, M., Schifani, E., Castracani, C., Spotti, F. A., Mori, A., & Grasso, D. A. (2021). First report on  
442 the acrobat ant *Crematogaster scutellaris* storing live aphids in its oak-gall nests. *Insects*, 12(2), 108.  
443 <https://doi.org/https://doi.org/10.3390/insects12020108>

444 Guo, H., Gu, L., Liu, F., Chen, F., Ge, F., & Sun, Y. (2019). Aphid-borne Viral Spread Is Enhanced by Virus-induced  
445 Accumulation of Plant Reactive Oxygen Species 1. *Plant Physiol*, 179(January), 143–155.  
446 <https://doi.org/10.1104/pp.18.00437>

447 Hullé, M., Chaubet, B., Turpeau, E., & Simon, J.-C. (2020). Encyclop'Aphid: A website on aphids and their natural  
448 enemies. *Entomologia Generalis*, 40(1). <https://doi.org/DOI: 10.1127/entomologia/2019/0867>

449 Ikbal, C., & Pavela, R. (2019). Essential oils as active ingredients of botanical insecticides against aphids. *Journal of Pest*  
450 *Science*, 92, 971–986. <https://doi.org/DOI: 10.1007/s10340-019-01089-6>

451 Jakubczyk, K., Koprowska, K., Gottschling, A., & Janda-Milczarek, K. (2022). Edible Flowers as a Source of Dietary  
452 Fibre (Total, Insoluble and Soluble) as a Potential Athlete's Dietary Supplement. *Nutrients*, 14(12).  
453 <https://doi.org/10.3390/nu14122470>

454 Jaouannet, M., Rodriguez, P. A., Thorpe, P., Lenoir, C. J. G., & Macleod, R. (2014). Plant immunity in plant – aphid  
455 interactions. *Front Plant Sci.*, 5(December), 1–10. <https://doi.org/10.3389/fpls.2014.00663>

456 Jayaswal, D., Mainkar, P., Kumar, K., Agarwal, Y., & Prabha, R. (2022). Pyramiding and evaluation of segregating lines  
457 containing lectin and protease inhibitor genes for aphid resistance in Brassica juncea. *Indian Journal of Biochemistry*  
458 *& Biophysics*, 59(August), 800–807. <https://doi.org/10.56042/ijbb.v59i8.62319>

459 Jones, R. A. C. (2022). Alteration of plant species mixtures by virus infection: Managed pastures the forgotten dimension.  
460 *Plant Pathology*, 71(6), 1255–1281. <https://doi.org/DOI: 10.1111/ppa.13571>

461 Jousselin, E., Gwenaelle, G., & Armelle, C. D. A. (2010). Evolutionary lability of a complex life cycle in the aphid genus  
462 *Brachycaudus*. *BMC Evolutionary Biology*, 10(1). <https://doi.org/10.1186/1471-2148-10-295>

463 Kallas, J. (2010). *Edible wild plants*. Gibbs Smith.

464 Karami-jamouri, T., Mirmoayed, A., Zamani, A., & Khajehzadeh, Y. (2018). The impact of ant attendance on protecting  
465 *Aphis gossypii* against two aphidophagous predators and it's role on the intraguild predation between them. *Journal*  
466 *of Insect Behavior*, 31, 222–239. <https://doi.org/DOI: 10.1007/s10905-018-9670-4>

467 Kennedy, J. S., & Stroyan, H. L. G. (1959). Biology of aphids. *Annual Review of Entomology*, 4(1), 139–160.

468 Kinley, C., Banu, A. N., Raut, A. M., Wahengbam, J., & Jamtsho, T. (2021). A review on past, present and future  
469 approaches for Aphids management. *Journal of Entomological Research*, 45(2), 336–346.  
470 <https://doi.org/10.5958/0974-4576.2021.00053.0>

471 Kumar, Sarwan. (2019). Aphid-Plant Interactions: Implications for Pest Management. In M. T. Oliveira, F. Candan, & A.  
472 Fernandes-Silva (Eds.), *Plant Communities and Their Environment* (p. Ch. 7). IntechOpen.



<https://doi.org/10.5772/intechopen.84302>  
 Kumar, Sushil, Bhowmick, M. K., & Ray, P. (2021). Weeds as alternate and alternative hosts of crop pests. *Indian Journal of Weed Science*, 53(1), 14–29. <https://doi.org/10.5958/0974-8164.2021.00002.2>  
 Liu, X. D., Xu, T. T., & Lei, H. X. (2017). Refuges and host shift pathways of host-specialized aphids *Aphis gossypii*. *Scientific Reports*, 7(1), 1–9. <https://doi.org/10.1038/s41598-017-02248-4>  
 Maharani, Y., Hidayat, P., Rauf, A., & Maryana, N. (2018). Short communication: New records of aphid species subfamily aphidinae (Hemiptera: Aphididae) in West Java, Indonesia. *Biodiversitas*, 19(2), 460–465. <https://doi.org/10.13057/biodiv/d190219>  
 Margaritopoulos, J. T., Tzortzi, M., Zarpas, K. D., Tsitsipis, J. A., & Blackman, R. L. (2006). Morphological discrimination of *Aphis gossypii* (Hemiptera: Aphididae) populations feeding on Compositae. *Bulletin of Entomological Research*, 96(2), 153–165. <https://doi.org/10.1079/ber2005410>  
 Meuninck, J. (2023). *Basic Illustrated Edible Wild Plants and Useful Herbs*. Rowman & Littlefield.  
 Mo, C., & Smilanich, A. M. (2023). Feeding on an exotic host plant enhances plasma levels of phenoloxidase by modulating feeding efficiency in a specialist insect herbivore. *Frontiers in Physiology*, 14(February), 1–10. <https://doi.org/10.3389/fphys.2023.1127670>  
 Naidu, V. (2012). *Hand book on weed identification*. Dr. VSGR Naidu.  
 Nelson, A. S., & Mooney, K. A. (2022). The evolution and ecology of interactions between ants and honeydew-producing hemipteran insects. *Annual Review of Ecology, Evolution, and Systematics*, 53, 379–402. <https://doi.org/https://doi.org/10.1146/annurev-ecolsys-102220-014840>  
 Pettersson, J., Tjallingii, W. F., & Hardie, J. (2017). Host-plant selection and feeding. In *Aphids as crop pests* (pp. 173–195). CABI Wallingford UK. <https://doi.org/DOI: 10.1079/9780851998190.0087>  
 Piron, P., de Haas, M., & Sonnemans, M. (2019). The presence of *Aphis* (Toxoptera) *aurantii* (Homoptera: Aphididae) in the Netherlands. *Entomologische Berichten*, 79(5), 162–164.  
 Santiago, G. S., Fonseca, L., Canedo-ju, E. O., Ribas, C. R., Pereira, R., & Alves, G. P. (2017). Isolated and Community Contexts Produce Distinct Responses by Host Plants to the Presence of Ant-Aphid Interaction : Plant Productivity and Seed Viability. *Plos One*, 12(1), 1–12. <https://doi.org/10.1371/journal.pone.0170915>  
 Singh, R., & Singh, G. (2021). Aphids. *Polyphagous Pests of Crops*, 105–182. [https://doi.org/DOI: 10.1007/978-981-15-8075-8\\_3](https://doi.org/DOI: 10.1007/978-981-15-8075-8_3)  
 Tegelaar, K., Hagman, M., Glinwood, R., Pettersson, J., & Leimar, O. (2012). Ant–aphid mutualism: the influence of ants on the aphid summer cycle. *Oikos*, 121(1), 61–66. <https://doi.org/https://doi.org/10.1111/j.1600-0706.2011.19387.x>  
 Völkl, W., Mackauer, M., Pell, J. K., & Brodeur, J. (2023). Predators, parasitoids and pathogens. In *CABI Books*. CABI Books. <https://doi.org/10.1079/9780851998190.0187>  
 Yamamoto, T., Hattori, M., & Itino, T. (2020). *Seasonal Migration in the Aphid Genus Stomaphis ( Hemiptera : Aphididae ) : Discovery of Host Alternation Between Woody Plants in Subfamily Lachninae*. 20. <https://doi.org/10.1093/jisesa/ieaa103>  
 Zheng, Z., Zhao, M., Zhang, Z., Hu, X., Xu, Y., & Wei, C. (2022). Lactic Acid Bacteria Are Prevalent in the Infrabuccal Pockets and Crops of Ants That Prefer Aphid Honeydew. *Front. Microbiol.*, 12(January), 1–17. <https://doi.org/10.3389/fmicb.2021.785016>

**Commented [Ma3]:** Not italic, use the journal abbreviation

**Commented [Ma4]:** Mind the period and coma

16 October 2023

Dear Editor,  
Biodiversitas

As requested, this is our response to reviewers' comments and suggestions.

Thank you so much for the very kind attention and great helps provided by editorial team of Journal of Biodiversitas

**"Letter on responses to reviewers' comments and suggestions from Reviewer 1"**

No.	Reviewers' suggestion	Our response	Location in revised manuscript
1	The Introduction has more than 700 words	The Introduction has been revised	Line 32-83
2	The reference must be revised following Biodiversitas guidelines.	The references have been updated	Line 394

Sincerely  
Corresponding author,



Chandra Irsan

## COVERING LETTER

Dear **Editor-in-Chief**,

I herewith enclosed a research article,

- ☐ The submission has not been previously published, nor is it before another journal for consideration (or an explanation has been provided in Comments to the Editor).
- ☐ The submission file is in OpenOffice, Microsoft Word (DOC, not DOCX), or RTF document file format.
- ☐ The text is single-spaced; uses a 10-point font; employs italics, rather than underlining (except with URL addresses); and all illustrations, figures, and tables are placed within the text at the appropriate points, rather than at the end.
- ☐ The text adheres to the stylistic and bibliographic requirements outlined in the Author Guidelines.
- ☐ Most of the references come from current scientific journals (c. 80% published in the last 10 years), except for taxonomic papers.
- ☐ Where available, DOIs for the references have been provided.
- ☐ When available, a certificate for proofreading is included.

## SUBMISSION CHECKLIST

Ensure that the following items are present:

The first corresponding author must be accompanied with contact details:

- ☐ E-mail address
- ☐ Full postal address (incl street name and number (location), city, postal code, state/province, country)
- ☐ Phone and facsimile numbers (incl country phone code)

All necessary files have been uploaded, and contain:

- ☐ Keywords
- ☐ Running titles
- ☐ All figure captions
- ☐ All tables (incl title and note/description)

Further considerations

- ☐ Manuscript has been "spell & grammar-checked" Better, if it is revised by a professional science editor or a native English speaker
- ☐ References are in the correct format for this journal
- ☐ All references mentioned in the Reference list are cited in the text, and vice versa
- ☐ Colored figures are only used if the information in the text may be losing without those images
- ☐ Charts (graphs and diagrams) are drawn in black and white images; use shading to differentiate

**Title:**

**Species of Aphids Found in Ornamental and Wild Plants in Highland, Pagar Alam, South Sumatra**

**Author(s) name:**

Chandra Irsan<sup>a\*</sup>, Erise Anggraini<sup>a,b</sup>, Siti Herlinda<sup>a</sup>, Wenny Ramadhani<sup>c</sup>, M. Umar Harun<sup>d</sup>,

**Address**

(Fill in your institution's name and address, your personal cellular phone and email)

<sup>a</sup> Department of Plant Pests and Diseases, Faculty of Agriculture, Universitas Sriwijaya, Indralaya, Ogan Ilir, South Sumatra, Indonesia

<sup>b</sup> Agroecotechnology Study Program, Faculty of Agriculture, Universitas Sriwijaya, Indralaya, Ogan Ilir, South Sumatra, Indonesia

<sup>c</sup> Plant Quarantine, Palembang, Sumatera, Indonesia

<sup>d</sup> Department of Agronomy, Faculty of Agriculture, Universitas Sriwijaya, Indralaya, Ogan Ilir, South Sumatra, Indonesia

**For possibility publication on the journal:**

(fill in *Biodiversitas* or *Nusantara Bioscience* or mention the others)

☐ Biodiversitas Journal of Biological Diversity

☐ Nusantara Bioscience

☐ Prosiding Seminar Nasional Masyarakat Biodiversitas Indonesia

☐ Asian Journal of Agriculture

☐ Asian Journal of Ethnobiology

☐ Asian Journal of Forestry

☐ Asian Journal of Natural Product Biochemistry

☐ Asian Journal of Tropical Biotechnology

☐ International Journal of Bonorowo Wetlands

☐ Cell Biology and Development

☐ Indo Pacific Journal of Ocean Life

☐ International Journal of Tropical Drylands

**Novelty:**

(state your claimed novelty of the findings versus current knowledge)

This paper described the alternative host of aphids in high land, South Sumatera. The knowledge regarding the alternative of insect pest could be beneficial resource for basic control of aphids.

**Statements:**

This manuscript has not been published and is not under consideration for publication to any other journal or any other type of publication (including web hosting) either by me or any of my co-authors.

Author(s) has been read and agree to the Ethical Guidelines.

**List of five potential reviewers**

(Fill in names of five potential reviewers **that agree to review your manuscript** and their **email** addresses. He/she should have Scopus ID and come from different institution with the authors; and from at least three different countries)

1. Dr. Koko Dwi Sutanti (email: [ksutanto@ksu.edu.sa](mailto:ksutanto@ksu.edu.sa) )
2. Dr. Lau Wei Hong (email: [lauweih@upm.edu.my](mailto:lauweih@upm.edu.my))
3. Prof. Dr. Dra. Asni Johari, M.Si. ([johari\\_asni@yahoo.com](mailto:johari_asni@yahoo.com) )
4. Dr. Mahesh Gunasena ([mahesh.gunasena@gmail.com](mailto:mahesh.gunasena@gmail.com))
5. Dr. Hasber Salim ([hasbersalim@usm.my](mailto:hasbersalim@usm.my))

**Place and date:**

Palembang, 5 October 2023

**Sincerely yours,**

(fill in your name, no need scanned autograph)

Dr. Chandra Irsan

# Species of Aphids Found in Ornamental and Wild Plants in Highland, Pagar Alam, South Sumatra

## Abstract

Aphids are one of the crucial pests in the tropical and sub-tropical regions. The presence of aphids in a plant can be very detrimental due to their role as vectors. Aphids exhibit species diversity, but not much information has been reported about the species diversity of aphids associated with essential crops such as ornamental plants. Furthermore, many aphid species were found on plants that were not actually hosts such as wild plants. Therefore, this study reported the species of aphids found in ornamental plants and the wild plants. The field research employed a purposive and direct observation approach to inventory cultivated or wild plants hosting aphids and collecting aphids. The plant selection process included cultivated plants encompassing ornamental plants, as well as wild plants or weeds. The collection and identification of host plants, and aphids, involved systematic searches for the selected plants and subsequent examination for the presence of aphids. Observations were made to all existing plant species to find those colonized by aphids. This study revealed that 15 species of aphids were found in Pagaralam, namely *Aphis gossypii*, *Uroleucon* sp., *Toxoptera odinae*, *Macrosiphum rosae*, *Aphis citricola*, *Aphis craccivora*, *Toxoptera aurantii*, *Pentalonia nigronervosa*, *Hystenura* sp., *Aphis glycine*, *Greenidae* sp., *Rhopalosiphum padi*, *Rhopalosiphum maidis*, *Hyperomyzus* sp. *Lipaphis erysimi*.

**Keywords:** aphids, ornamental plants, wild plants

**Running title:** Aphids Found in Ornamental and Wild Plants

## INTRODUCTION

Aphids are one of the crucial pests in the tropics and sub-tropics, exhibiting various polyphagous, oligophagous, and monophagous characteristics (Kennedy and Stroyan 1959). One species of aphids can host more than 400 species from 40 families (Bass et al. 2014). In addition to pests, aphids can also be vectors of plant viral diseases (Gadhawe et al. 2020). Aphids can transmit 275 viruses (Ertunc 2020). In tropical areas, aphids can always be found throughout the year due to their parthenogenetic nature of reproduction (Blackman and Eastop 2017). Aphids consume young leaves sap, which can deplete essential nutrients for healthy growth (Cao et al. 2018). Moreover, when aphids transmit viral diseases from one plant to another, this can further weaken and stunt the growth of infected plants (Jones 2022). According to Kinley et al. (2021), aphids cause yield losses directly (35 - 40%) by sucking the plant sap or indirectly (20 - 80%) through viral transmission. Therefore, aphid infestations can have adverse effects on crop yields and overall plant health (Kumar 2019).

Due to their function as vectors, the presence of aphids on a plant can be highly damaging (Jaouannet et al. 2014). They feed by piercing the plant's tissues and consuming its sap, which can reduce the plant's growth and productivity, ultimately leading to weakness and possible death (Chandel et al. 2022). Additionally, as vectors, aphids can transmit a variety of plant diseases. They are as carriers for various plant viruses, and when they move from infected to healthy plants, these viruses can rapidly spread and cause extensive damage (Guo et al. 2019). In addition, the honeydew that aphids secrete can lead to the growth of sooty mold, a black fungus that can prevent sunlight from reaching the plant's

leaves, thereby impairing photosynthesis, the process by which plants produce food (Singh and Singh 2021). Therefore, it is crucial to control aphid populations in gardens and crops.

Many aphid species were found on plants that were not their actual hosts (Maharani et al. 2018). Aphids have one or more secondary, or "alternative," host plants in addition to their primary host plants, which are the types of plants they feed on most frequently (Clarke et al. 2020). An alternative host can also be a collateral host belonging to the same plant family as the primary host, helping crop pests to survive when the primary hosts are unavailable (Kumar et al., 2021). These secondary hosts may offer less adequate nutrition for insects (Mo and Smilanich 2023), However, they may provide a means of survival when primary hosts are unavailable, during certain seasons, or under certain environmental conditions (Kumar et al., 2021). According to Liu et al. (2017), since hibiscus serves as an overwintering host for cotton-specialized aphids but not for cucurbit-specialized aphids, it is evident that host-specialized aphids have refuges during times of food shortage. The life cycles of numerous aphid species exhibit such complexity (Jousselin, Gwenaëlle, and Armelle 2010). They maintain a cycle of host alternation, shifting between their primary hosts (typically a woody plant) and secondary hosts (often herbaceous plants) (Yamamoto, Hattori, and Itino 2020). Weeds pose a continuous threat in both cropped and non-crop areas, providing food, shelter and reproductive sites for various pest organisms (Kumar et al., 2021). This indicates that weeds can serve as alternative hosts for aphids.

A study of aphid species on horticultural plants has been conducted (Maharani et al. 2018), However, information about aphid species on ornamental and wild plants has not received as much attention and remains largely unexplored. In South Sumatra, particularly in the highland areas like Pagar Alam, there are numerous ornamental and native plants. The research on the diversity of aphid species in ornamental and wild plants has received little attention. Therefore, this study was conducted in Pagar Alam, a highland region of South Sumatra, with the aim of obtaining information on the diversity of aphid species found in ornamental and wild plants. The findings from this study can serve as a valuable resource for aphid management.

## MATERIALS AND METHODS

The field research employed a purposive and direct observation approach to inventory cultivated or wild plants hosting aphids and collecting aphids. The plant selection process included cultivated plants encompassing ornamental plants, as well as wild plants or weeds. The collection and identification of host plants, and aphids, involved systematic searches for the selected plants and subsequent examination for the presence of aphids. Observations were made to all existing plant species to find those colonized by aphids. Any plants colonized by aphids were documented as aphid hosts. Aphids, along with their natural enemies within the aphid colonies, were systematically collected. All components of the collected observations were then identified.

Aphid identification was conducted using identification keys (Blackman and Eastop 2008). Identification of aphid species took place in the Laboratory of Entomology, Faculty of Agriculture, Universitas Sriwijaya. Identification relied on morphological characteristics. The host plants were identified using weed identification hand book (Kallas, 2010; Meuninck, 2023; Naidu, 2012). The location and size of aphid colonies, morphology of aphids including their shape and color, as well as any symptoms observed in the host plants were recorded, and photographs of the aphid colonies and their host plants were taken.

## RESULT AND DISCUSSION

### Result

The results showed that 15 aphid species were found in Pagaralam, namely *Aphis gossypii*, *Uroleucon* sp., *Toxoptera odinae*, *Macrosiphum rosae*, *Aphis citricola*, *Aphis craccivora*, *Toxoptera aurantii*, *Pentalonia nigronervosa*,

88 *Hystenura* sp., *Aphis glycine*, *Greenidae* sp., *Rhopalosiphum padi*, *Rhopalosiphum maidis*, *Hyperomyzus* sp. *Lipapis*  
89 *erysimi*. Based on the observation, these aphids were found on various ornamental plants (Table 1). The primary colony  
90 locations were generally in flowers, and this study documented these colony locations in ornamental plants (Figure 1).  
91

92 Table 1: Aphid species found in ornamental plants and their colony locations.

No	Host Plant	Aphid Species	Colony location
1	<i>Aster alpinus</i>	<i>Sitobion luteum</i>	flower
2	<i>Brugmansia suaveolens</i>	<i>Aulacorthum solani</i> <i>Neomyzus circumflexus</i> <i>Myzus persicae</i>	flower
3	<i>Caladium</i> sp.	<i>Pentalonia</i> sp.	flower
4	<i>Cananga odoratum</i>	<i>Aphis gossypii</i>	flower
5	<i>Canna indica</i>	<i>Pentalonia nigronervosa</i>	flower
6	<i>Catharanthus roseus</i>	<i>Aphis citricola</i>	flower
7	<i>Cestrum</i> sp.	<i>Aphis gossypii</i> <i>Neomyzus circumflexus</i>	flower
8	<i>Clitoria ternatea</i>	<i>Aphis craccivora</i>	flower
9	<i>Cosmos caudatus</i>	<i>Uroleucon</i> sp.	flower
10	<i>Dahlia Kelvin</i>	<i>Aphis gossypii</i>	flower
11	<i>Dendrobium</i> sp.	<i>Sinemogoura citricola</i>	flower
12	<i>Duranta</i> sp.	<i>Aphis gossypii</i>	flower
13	<i>Helianthus</i> sp.	<i>Aphis glycines</i> <i>Hyperomyzus</i> sp.	flower
14	<i>Hibiscus rosasinensis</i>	<i>Aphis gossypii</i>	flower
15	<i>Ixora paludosa</i>	<i>Aphis gossypii</i> , <i>Toxoptera aurantii</i>	flower
16	<i>Ixora</i> sp.	<i>Aphis citricola</i> <i>Aphis gossypii</i> <i>Toxoptera aurantii</i>	flower
17	<i>Murraya paniculata</i>	<i>Aphis craccivora</i> <i>Toxoptera citricidus</i>	flower
18	<i>Mussaenda frondosa</i>	<i>Aphis citricola</i> <i>Toxoptera odinae</i>	flower
19	<i>Rosa indica</i>	<i>Macrosiphum rosae</i>	flower
20	<i>Spondiras dulcissoland</i>	<i>Aphis citricola</i> <i>Hysteroneura setariae</i>	flower





99

Fig 1. The location of aphid colonization on various plant parts. a) *A. gossypii* in *D. Kelvin* flower b) *A. gossypii* in *H. rosasinensis* flower c) *A. gossypii* in tuberose flower, d) *A. craccivora* in *Clitoria ternatea* flower, e) *A. citricola* in *Helianthus* sp., f) *A. aurantii* on the *M. paniculata* flower, g) *T. odinae* in the *S. dulcssoland*, h) *Uroleucon* sp. in chrysanthemums, i) *Macrosiphum rosae* in *R. indica* flower, j) *Pentalonia nigronervosa* in *C. indica* leaves

In addition, this study documented the presence of weeds, which might serve as alternative hosts for aphids (Table 2). The location of aphid colonies also varied, namely on flowers, stalks, plant tops, young leaves and old leaves of wild plants (Figure 2). The presence of specific plants or host plants within a habitat influenced the types of aphids found. Many aphid species are found on a broad range of plants or host plants, while others are highly specialized and are only found on specific plants or host plants. This is closely related to the polyphagous, oligophagous or monophagous nature of aphids (Blackman & Eastop 2000).



Table 2: Species of aphids found in wild plants and their colony locations.

No	Host Plant	Aphid species	Colony location
1	<i>Ageratum conyzoides</i>	<i>Aphis gossypii</i>	Shoots, young leaves, old leaves, flowers
2	<i>Alternanthera philoxeroides</i>	<i>Aphis gossypii</i>	Shoots, buds
3	<i>Alternanthera sessilis</i>	<i>Aphis gossypii</i>	Shoots, buds
4	<i>Amaranthus gracilis</i>	<i>Aphis craccivora</i>	Flowers, shoots, young leaves, old leaves
5	<i>Blumea lacera</i>	<i>Lipaphis erysimi</i>	Flowers, shoots, and buds
6	<i>Croton hirtus</i>	<i>Aphis gossypii</i>	Flowers, shoots, young leaves, old leaves, young twigs
7	<i>Cynodon dactylon</i>	<i>Schizaphis rotundiventris</i>	Flower, flower stalks
8	<i>Cyperus rotundus</i>	<i>Schizaphis rotundiventris</i>	Flower, flower stalks, leaf axils
9	<i>Cyperus compressus</i>	<i>Schizaphis rotundiventris</i>	Flower, flower stalks, leaf axils
10	<i>Digitaria ciliaris</i>	<i>Hysteronura setariae</i>	Flower, flower stalks
11	<i>Echinocloa crusgali</i>	<i>Hiperomyzus</i> sp.	Young leaves, old leaves
12	<i>Ecliptica prostrata</i>	<i>Aphis gossypii</i>	Shoots, young leaves
13	<i>Eleusin indica</i>	<i>Hysteronura setariae</i> <i>Rhopalosiphum maidis</i>	Flower, flower stalks, leaf axils
14	<i>Emilia sonchifolia</i>	<i>Aphis gossypii</i>	Flower, flower stalks, shoots
15	<i>Eragrostis tenella</i>	<i>Hysteronura setariae</i>	Flower, flower stalks, seeds
16	<i>Euphorbia hirta</i>	<i>Aphis gossypii</i>	Young leaves, old leaves
17	<i>Eupotarium odoratum</i>	<i>Aphis gossypii</i> , <i>Aphis glycine</i>	Young leaves, old leaves, young twigs
18	<i>Hymenochera acutigluma</i>	<i>Hysteronura setariae</i>	Flowers, flower stalks, leaf axils
19	<i>Lagerstromia</i> Sp.	<i>Greenidea</i> sp.	Young leaves
20	<i>Lophatherum gracile</i>	<i>Hysteronura setariae</i> <i>Rhopalosiphum maidis</i>	Young leaves, old leaves, leaf axils
21	<i>Melastoma affine</i>	<i>Aphis gossypii</i>	Shoots, young leaves
22	<i>Mikania mikranta</i>	<i>Aphis gossypii</i> <i>Aphis glycine</i>	Shoots, young leaves, old leaves
23	<i>Mimosa invisa</i>	<i>Aphis craccivora</i>	Shoots, pods
24	<i>Mimosa pudica</i>	<i>Aphis craccivora</i>	Shoots, pods, flowers
25	<i>Mimosa vigra</i>	<i>Aphis craccivora</i>	Shoots, pods
26	<i>Oryza rufipogon</i>	<i>Rhopalosiphum padi</i> , <i>Rhopalosiphum maidis</i>	Old leaves, young leaves (pupus), leaf axils
27	<i>Oxonopus compressus</i>	<i>Hysteronura setariae</i>	Flower, flower stalk, leaf axils
28	<i>Paspalum conjugatum</i>	<i>Hysteronura setariae</i>	Flower, flower stalk, seeds
29	<i>Phylanthus neruri</i>	<i>Aphis citricola</i>	Shoot, young leaves, old leaves, young twigs, petioles
30	<i>Portulaca oleraceae</i>	<i>Aphis craccivora</i>	Shoots, young leaves, flower
31	<i>Physalis angulata</i>	<i>Aphis craccivora</i> , <i>A. gossypii</i>	Shoots, young leaves, old leaves
32	<i>Rorippa indica</i>	<i>Lipapis erysimi</i>	Flower, fruit, shoots, young leaves
33	<i>Sida rhombifolia</i>	<i>Aphis gossypii</i>	Shoots, young leaves, old leaves, fruit/seeds
34	<i>Sonchus arventris</i>	<i>Lipapis erysimi</i>	Young leaves, fruit stalks, flower, fruit



Figure 2. Aphids found on wild plants a) *A. gossypii* on the weed *Ageratum conyzoides*, b) *A. gossypii* on *Croton* weed *hirtus* c) *A. gossypii* on the weed *Eupatorium odoratum*, d) *A.gossypii* on plants *Pachystochys* sp., e) *A.gossypii* on plants *Caladium* sp., f) *A. gossypii* on the weed *Alternanthera sessilis*, g) *A.gossypii* in *Portulaca oleraceae* weeds, h) *A.gossypii* on the weed *Euphorbia hirta*, i) *A. citricola* on the weed *Phyllanthus nerruri*, j) *A. citricola* on *Sida rhombifolia* weed, k) *A. citricola* on plants *Annona muricata*, l) *A.citricola* on the weed *Ludwigia peruviana*, m) *A. craccivora* on *Mimosa pudica* weed, n) *A.craccivora* on weeds *Amaranthus gracilis*, o) *A. glycine* in *Mikania micrantha* weed, p) *Hysteneura* sp. in *Eleusine indica* weeds, q) *Greenidae* sp. in kenidai trees (shrubs) *indica*, r) *Hyperomyzus* sp. in *Echinocloa crusgali* Weed, s) *L. erysimi* on weed *sonchus arvensis*, t) *Rhopalosiphum* rice on the weed *Oryza rufipogon*, u) *Rhopalosiphum* *Maidis* on the weed *Oryza rufipogon*.

## Discussion

The plant species or host plant influences the distribution of aphids. There are aphid species that can be found on a wide range of host plants, which is closely related to the polyphagous nature of aphids, allowing them to colonize many different species of host plants. Host plants can also affect the distribution of aphids, as evidenced by the presence of aphid species exclusively found on certain host plants (Santiago et al. 2017). But there are some species of aphids found only on one particular host and are not found on other host plants (Döring 2014). *A. gossypii*, and *Aphis aurantii* have been found on many host plants because both aphids are classified as polyphagous aphids (Alotaibi et al., 2023).

Aphids can commonly be found infesting a variety of ornamental plants. They are attracted to these plants due to the rich nutrient content in the plant sap (Braham et al. 2023). In this present study, some aphid species were found on some ornamental plants in Pagaralam. The location of aphid colonization on the plants varied. On *Adiantum predatum* plants, aphids formed colonies on young leaf stalks and on newly emerging leaves. The aphids displayed brown and black coloration. The aphid colonies found were small, and the colonized plant parts showed no signs of disease. The identification results showed that the aphids were *Neotoxoptera* sp., and notably, they were not associated with ants. On *Aster alpinus*, aphids were found to form colonies on the stems or young leaf shoots, and the colonies were relatively large. The color of the aphids was dark brown to black. The colonized plant parts showed symptoms of stunting. The identification results showed that the aphids were *Uroleucon* sp., and they were associated with ants.

On the *Brugmansia suaveolens*, *M. persicae* were found on the undersides of old leaves or leaves that have started to turn yellow. The colonies were relatively small. The aphids found were green and large bodies. The colonized plant parts did not show any signs of disease. On *Caladium* sp. (taro) was found one species of aphids: *A. gossypii*. The aphids formed colonies under the surface of young and older leaves. The occupied leaf areas did not display severe symptoms. The aphids were yellow green to dark green. The wingless adult aphids often had a white, flour-like appearance on their bodies. On the *Cananga odoratum* (ylang-ylang), colonies of *T. aurantii* were found on the undersides of the leaves, the shoots, buds, and unopened flower petals. The *T. aurantii* colonies found were relatively large. Colonized parts, especially shoots, showed signs of stunting. The aphids found were brown to black in color. The colonies of *T. aurantii* were found to be associated with black ants.

Aphids on *C. indica* (Indian shot, African arrowroot) were found to form colonies in the leaf axils and under the leaf surface near the leaf base. The colonies were quite large. The aphids were dark brown to dark red coloring with a medium-sized body. The identification results showed that the aphids were *P. nigronervosa*. The colonies of *P. nigronervosa* were found to be associated with ants. In the *Catharanthus roseus* (periwinkle), *A. citricola* aphids were found. The aphids were yellow-green, sifunculi, and black cauda. The aphids formed colonies on flowers and shoots, and the colonized plant parts did not show any symptoms of disease. On *Cestrum* sp. (Bastard jasmine), aphids formed colonies on the undersides of young leaves, shoots, and within flower parts, especially between petals or flower stalks that had not fully bloomed. The colonies were quite large. The body color of aphids was green to dark green with small to medium-sized bodies. The colonized plant parts, especially leaves, showed stunting symptoms. The identification results showed that the aphids were *A. gossypii*. The aphid colonies found were consistently associated with ants.

Aphids on *Clitoria ternatea* were found to form colonies on flower parts, flower crowns, stems and young leaves. The aphids were brown to black in color. Colonized plant parts, especially shoots and young leaves, showed stunting symptoms. The identification results showed that the aphids were *A. craccivora*. These colonies were consistently associated with ants. On the plant *Cosmos caudatus*, aphids were found on the flower petals. The colonies were not very large. The body color was green and light green. The identification results showed that the aphids were *A. gossypii*, and

203 they were also associated with ants. The aphids on the *Dahlia kellyi* plant formed colonies on unopened flower buds, with  
204 a significant population among the blooming petals. The body color was green to dark green. The identification results  
205 showed that the aphids were *A. gossypii*. Aphids on *Datura metel* (amethyst) were found to form colonies on the  
206 undersides of old leaves. The aphids were medium sized with a green body color. The colonized plant parts did not show  
207 any symptoms of disease. The identification results showed that the aphids were *Myzus ornatus*. The aphid colonies were  
208 not associated with ants. Within *Dendrobium* sp., aphid colonies were found on the young leaves. The aphids were yellow,  
209 green to dark green. The colonized plants did not show any disease symptoms. The identification results showed that the  
210 aphids were *A. gossypii*, and they were associated with ants. On *Duranta* sp. (bonsai), colonies of aphids were located on  
211 the undersides of young leaves. The colonized plant parts showed stunting symptoms. The colonies were very large. The  
212 aphids were green in color. The identification results showed that the aphids were *A. gossypii*. The aphid colonies were  
213 consistently associated with ants.

214 On the *Helianthus annuus* (sunflower) plants, aphid colonies were found between the flower petals. The  
215 colonized flowers, especially the crowns, exhibited a tendency to fall off easily. The aphids were green and yellow in  
216 color. The colonies were small. The identification results showed that the aphids were *A. gossypii*. These aphid colonies  
217 were associated with ants. Aphid colonies on *Helianthus* sp. were found on the undersides of old leaves. These colonies  
218 were small in size. The aphids were green with a medium body size. The colonized plant parts did not show any disease  
219 symptoms. The identification results showed that the aphids were *M. ornatus*. The aphid colonies were not associated with  
220 ants. Within the colonies, mummified aphids that were parasitized by Aphidiidae were found.

221 On the *Hibiscus rosa-sinensis*, aphids ranging in color from yellow to dark green were found. The aphids formed  
222 colonies on flower buds, unopened flower crowns, and the undersides of aging leaves. The colonies grew to be very large.  
223 The identification results showed that the aphids were *A. gossypii*. The aphid colonies were consistently associated with  
224 ants. Two types of aphids were found on the flowering plant *Ixora paludosa*. First, the aphids formed colonies on the  
225 undersides of young leaves that were still red or light green and sometimes on flower stalks that had not yet bloomed. The  
226 occupied plant parts showed symptoms such as stunted leaf growth, leaf shrinkage, necrotic spots on the leaf surface, and  
227 slightly downward-curved leaf edges. The upper leaf surface looked wet and sticky, similar to sugar. The aphids had  
228 yellow, green, or slightly dark green bodies, with some wingless adults having a powdery white upper surface. The  
229 identification results showed that the aphids were *A. gossypii*, and they were almost always associated with ants. The  
230 second type of aphids on *Ixora paludosa* formed colonies under the surface of young and older leaves. The colonies could  
231 also be found on newly emerging flowers and leaves. The plant parts occupied by these aphids did not show obvious signs  
232 of illness. These aphids were dark red to black, with once-branched stigma and venation in their black wings. The  
233 identification results showed that the aphids were *T. aurantii*. These aphids were also associated with ants.

234 In *Ixora* sp. flower plants, two forms of aphids were discovered. These aphids occupied the shoots, young leaves  
235 and unopened flowers. The affected plant parts did not show obvious symptoms. The aphids exhibited colors ranging from  
236 yellow and green to a slightly darker green. Sometimes the upper surface of the wingless imago's body appeared white,  
237 resembling flour. The identification results showed that these aphids were *A. gossypii*. These aphid colonies were almost  
238 always associated with ants. Another species of aphids was founded and formed colonies on flower stalks that had not yet  
239 bloomed and on newly emerging shoots or leaves. The presence of these aphids on the plant did not induce any symptoms  
240 of plant disease. The aphids were yellow or yellow green, with black cauda and siphunculi. Their bodies were very small  
241 to small. The identification results showed that the aphids were *A. citricola*. The colonies of *A. citricola* were also  
242 frequently found in association with ants.



243 Two types of aphids were found on *Mussaenda frondosa*, each forming colonies in different locations. The first  
244 type formed colonies on young leaves, shoots, and flowers. The plant parts they occupied showed no obvious disease  
245 symptoms. The aphids were yellow, green, and some with dark green. The identification results showed that the aphids  
246 were *A. gossypii*. The second type of aphids formed colonies on the stems or young twigs, appearing densely clustered as  
247 if piled up. The aphid colonies could also be found on young leaves, shoots and within flower parts. The plant parts they  
248 infested showed no signs of diseases. The aphids were yellow or yellow green, with black cauda and siphunculi. They had  
249 tiny to small bodies. The identification results showed that the aphids were *A. citricola*.

250 The results showed that 34 species of wild plants, including weeds, were growing in the yard colonized by aphids.  
251 This indicated that multiple species of aphids colonized various host plants. The aphid species colonizing these plants were  
252 generally consistent within the same taxon. *Ageratum conyzoides* was infested by *Aphis gossypii*. These aphids formed  
253 colonies on the flower sections, shoots, lower surfaces of young leaves, or leaves turning yellow. The aphids were green,  
254 yellow-green to dark green, often forming large colonies. *Alternanthera philoxeroides* or alligator grass was also colonized  
255 by *Aphis gossypii*. Small colonies were found on shoots or stems. These aphids had small bodies and were green, ranging  
256 from yellow-green to dark green. *Alternanthera sessilis* was colonized by *Aphis gossypii*, forming colonies on shoots,  
257 flowers, and fruit. The colonies were typically large, and they were often associated with tiny brown ants. *Amaranthus*  
258 *gracilis* was infested by *Aphis craccivora*. These aphids established colonies on shoots, flowers and young and old leaves.  
259 They were dark brown to black in color, with shiny black wingless imagoes. Colonies of these aphids were associated with  
260 both black and red ants. *Blumea lacera* was colonized by *Lipaphis erysimi* aphids. These aphids were bright green, and of  
261 medium size. The colonies formed on flowers, flower stalks and the undersides of the leaves at the top. The aphid colonies  
262 were not associated with ants. *Croton hirtus* or fire grass was infested by *Aphis gossypii*. The aphids were yellow-green to  
263 dark green. The colonies were found on the stems, leaves, buds and flowers, often forming large colonies. *Cynodon*  
264 *dactylon* or Bermuda grass was colonized by *Schizaphis rotundiventris*. The aphids colonized the flowers, flower stalks  
265 and sometimes in the leaf axils of the plant. Small colonies were formed. The aphids were brown to red-brown. They were  
266 associated with ants. *Cyperus rotundus* or nut grass was infested by *Schizaphis rotundiventris* aphids. The colonies were  
267 found on flower stalks, flowers, and leaf axils. The colonies were quite large and associated with both black and red ants.  
268 The aphids were dark brown in color. *Cyperus compressus* or grass puzzle was colonized by *Schizaphis rotundiventris*  
269 aphids, forming colonies in the flowers, flower stalks and sometimes in the axils and leaves of the shoots or buds. Small  
270 colonies were observed. *Digitaria ciliaris* was infested by *Hysteroneura setariae* aphids, with small colonies scattered on  
271 the flowers and flower stalks. These aphids were light brown to brown in color. *Echinocloa crusgali* or water hyacinth  
272 plants were colonized by *Hiperomyzus* sp. aphids. These aphids were dark brown to black and formed large colonies on  
273 the undersides of both young and old leaves. The aphid colonies were never found in association with ants. *Ecliptica*  
274 *prostrata* or urang aring was colonized by *Aphis gossypii*, forming small colonies on the shoots and flowers. The aphids  
275 were bright green to blackish green. The aphid colonies were also consistently associated with ants.

276 *Eleusin indica* was colonized by two species of aphids: *Hysteroneura setariae* and *Rhopalosiphum maidis*. *H.*  
277 *setariae* formed colonies in flower parts, flower stalks and leaf axils resulting in quite large colonies. *H. setariae* body  
278 color ranged from red brown to dark brown. The colonies were consistently associated with ants. The aphids of *R. maidis*  
279 formed colonies in the leaf axils and undersides of leaves and on leaf shoots that had not yet opened. The colonies were  
280 not densely packed. The leaf aphids of *R. maidis* were green in color, with distinct black siphunculi and cauda. These  
281 aphids had relatively large bodies with a slightly elongated shape. *R. maidis* colonies were always associated with ants.

282 The plant *Emilia sonchifolia*, characterized by its purple flowers, was colonized by *Aphis gossypii*. The aphids were yellow  
283 to green in colour. The colonies formed near flowers, flower stalks, and shoot leaves.

284 *Eragrostis tenella* was infested by *Hysteroneura setariae* aphids. The aphids were brown to red brown. Small  
285 colonies formed on flowers near the seeds, with groups of aphids surrounding the plant's seeds. The aphids of *H. setariae*  
286 were consistently associated with ants. *Euphorbia hirta* or wart grass was colonized by *Aphis gossypii*. The aphids formed  
287 colonies on the undersides of leaves, resulting in stunted growth of the leaves. The aphids were yellow to dark green in  
288 color. *A. gossypii* colonies on *E. hirta* plants were consistently associated with ants. *Eupatorium odoratum* was colonied  
289 by both *Aphis gossypii* and *Aphis citricola*. *A. gossypii* formed colonies in the buds, young leaves, old leaves, and young  
290 twigs. Young leaves that were colonized by *A. gossypii* became stunted with an irregular shape. *A. gossypii* found in this  
291 plant showed yellow-green to dark green in body colour. The colonies of *A. citricola* formed on the young twigs near the  
292 shoots, with these aphids displaying yellow-green coloration and having black siphunculi and cauda. Aphid colonies of  
293 both *A. gossypii* and *A. citricola* on *E. odoratum* plants were associated with either black or red ants.

294 *Hymenochera acutigluma* or hair axis was colonized by *Hysteroneura setariae*, which formed colonies on the  
295 flower stalks and flowers. The colonized parts of the plants did not display any noticeable symptoms. *Lagerstromia* sp. or  
296 *kenidai*, was infested by *Greenidae* sp. These aphids had bright green bodies and distinctive elongated siphunculi with  
297 thorns. The aphids formed colonies on the undersides of leaves, especially on young leaves. The colonized leaves did not  
298 show any disease symptoms. *Lophatherum gracile* or bamboo grass plants were colonized by two species of aphids:  
299 *hysteroneura setariae* and *Rhopalosiphum maidis*. The aphids of *H. setariae* formed colonies on the undersides of leaves,  
300 leaf shoots, and leaf axils. The colonized leaves did not show any disease symptoms. *H. setariae* aphids were brown to  
301 red-brown. *R. maidis* aphids also formed colonies on the undersides of leaves, but the colonies were small. *R. maidis*  
302 aphids were green to bright green in color, with black siphunculi and cauda. It was possible for colonies of the two species  
303 of aphids on *L. gracile* to mix.

304 *Melastoma affine* was colonized by *Aphis gossypii*. The colonies formed on shoots, particularly near newly  
305 emerging shoots and on newly emerging fruits and flowers. The body colour of aphids ranged from yellow to green. The  
306 colonized plant parts did not show any disease symptoms. *Mikania miranta* was colonized by *Aphis gossypii* and *Aphis*  
307 *glycine*. *A. gossypii* formed colonies on the shoots, especially on the undersides of the leaves, resulting in stunted and  
308 curled leaves. *A. glycine* formed colonies on the branches. The colonies were densely populated. *A. Glycine* aphids were  
309 light green to green in color. The colonized plant parts became distorted. The two species of aphids could mix to form a  
310 single colony.

311 *Mimosa invisa* (cater-grass) was colonized by *Aphis craccivora*. The aphids of *A. craccivora* on *M. invisa* plants  
312 formed colonies only on the shoots with small colonies. The aphids appeared dark black with wingless imagoes. *Mimosa*  
313 *pudica* was observed to be colonized by *Aphis craccivora*. The aphids formed colonies on shoots, especially young shoots,  
314 and occasionally on flowers and pods. The aphids were black and of medium size, resulting in stunted growth of the  
315 colonized plant parts. The colonies were quite large. *Mimosa vigra* was colonized by *Aphis craccivora*. The colonies of  
316 aphids occupied the pods and shoots with small colonies. The nymphs of aphids were black, and wingless adults were  
317 shiny black. The colonized plant parts did not show any disease symptoms.

318 *Oryza rufipogon* was colonized by two species of aphids: *Rhopalosiphum rice* and *Rhopalosiphum maidis*. Both  
319 aphids colonized the same plant parts, namely the unopened leaves and the leaf axils with large colonies. The two species  
320 could be distinguished by their body color. *R. maidis* appeared green with black sifunculi and cauda, while *R. rice*  
321 appeared white. The colonies of *R. maidis* and *R. rice* in *O. rufipogon* plants were associated with the presence of red ants.

322 *Oxonopus compressus* or pait grass was colonized by *Hysteroneura setariae* aphids. The colonies occupied flowers, flower  
323 stalks, seeds, and sometimes in the leaf axils. The aphids were brown to dark brown in color. Small colonies were formed,  
324 and they were also consistently associated with ants.

325 *Paspalum conjugatum* was colonized by *H. setariae* aphids. The colonies occupied flower parts, especially the  
326 seeds and flower stalks. Aphids had brown to dark brown bodies. *Phylanthus niruri* was colonized by *Aphis citricola*. The  
327 colonies formed on the shoots and the undersides of leaves and petioles. The colonized parts became distorted, stunted,  
328 and wrinkled. The aphids had yellow bodies with black sifunculi and cauda, and the colonies formed were quite large.  
329 *Portulaca oleraceae* plants were colonized by *Aphis craccivora*. The aphids of *A. craccivora* in *P. oleraceae* plants  
330 formed colonies on the undersides of leaves, especially young leaves, shoots and in flowers. The colonized plant parts  
331 became stunted, and leaf edges curled downward. The aphids had dark brown to black bodies, with wingless imagoes that  
332 appeared glossy black.

333 *Physalis angulata* plants were colonized by *Aphis craccivora*. The aphids had dark green to black bodies, with  
334 glossy black wingless imagoes. *A. craccivora* formed colonies on the shoots or near the leaf buds. The colonized plant  
335 parts did not show any symptoms of disease. *Rorippa indica* or mustard land was colonized by *Lipaphis erysimi*. The  
336 colonies formed on the flowers, fruits, flower stalks and the lower surface of leaves. The colonized plant parts showed  
337 symptoms such as curling and stunting. *Sida rhombifolia* or cacabean was colonized by *Aphis gossypii*. The aphids had  
338 green-yellow to green body colors. The colonies formed on the surface of lower leaves, stalks and flower petals. The  
339 colonized plant parts, especially the shoots, showed curling, and the leaf edges curled downward. *Sonchus arvensis* plants  
340 were colonized by *L. erysimi*. The aphids had green to whitish green body colours, and the colonies formed on flower  
341 stalks, under petals, and on young shoots or leaves. The colonized plant parts became stunted over time.

342 In general, aphids observed on ornamental and wild plants formed colonies. The colonized plant parts typically  
343 displayed typical symptoms of damage, but some did not show any symptoms. Generally, the symptoms of the plants  
344 caused by aphid colonies were relatively the same, such as stunted growth, abnormal shape, and stunted or curly leaves.  
345 These characteristic symptoms serve as indicators of aphid infestations. However, some plants or plant parts did not show  
346 symptoms when colonized by aphids. This condition happened because the colonized parts had reached their maximum  
347 growth or development. It indicated that the colonized part was not currently undergoing a growth phase. The colonies that  
348 did not induce symptoms typically occurred when the colonized leaves had reached their maximum growth or when the  
349 leaves and plant parts were old. The old leaves or twigs might not show the typical symptoms associated with aphid  
350 infestations.

351 The part of the plant exhibiting characteristic symptoms when colonized by aphids also often experienced a  
352 cessation in growth due to the piercing by the aphids. In contrast, the areas surrounding the puncture site continued to  
353 grow, resulting in some parts developing normally while others become stunted (Pettersson, Tjallingii, and Hardie 2017).  
354 This condition could lead to the bending of shoots or young stems, curling of leaves, downward curling of leaf edges, or  
355 stunted leaf growth. In this observation, monocot plants or groups of grasses with narrow leaves generally did not display  
356 any distinctive symptoms when colonized by aphids. This might be because the growth or development of their leaves  
357 differed from that of dicot plants. Therefore, the presence of aphids in monocot plants or plants was often easier to  
358 recognize through the presence of ants. If a plant was found to have a significant number of ants, there was a possibility  
359 that aphids had colonized the plant (Tegelaar et al. 2012). Therefore, the presence of ants could serve as an indicator of the  
360 presence of aphid colonies.

Throughout their life cycle, aphids exhibited host alternation by switching between two distinct host plants (Yamamoto et al. 2020). They overwintered on woody plants, reproduced in the spring, and migrated to herbaceous plants during the summer before returning to their primary host in the autumn (Yamamoto et al. 2020). This allowed aphids to maximize resource utilization, avoid congestion and competition, evade predators and parasites, circumvent plant defenses, and colonize new areas. Aphids could distribute their population efficiently, thereby avoiding overcrowding, predators and parasites, and plant defenses developing over time through host switching (Yamamoto et al. 2020). This behavior was essential for the survival and environmental adaptation of aphids.

Aphids colonized flowers because they may offer an accessible and rich food source, sugary plant sap found in new growth or reproductive parts of plants. Flowers contain a nutrient-rich nature and easy access to sap, therefore aphids were attractive to sap the flowers. Some aphid species were drawn to certain colors (Jakubczyk et al. 2022). Herbs served as an alternative host for aphids in this present study. Aphids consume sugar-rich liquid in plants, known as "sap". Aphids considered herbs and other green vegetation as abundant food sources. Aphids utilize needle-like mouthparts to penetrate plant tissues and access this fluid (Brožek et al. 2015). Several aphids colonized herbs such as Indian mustards, *Lipaphis erysimi*, and *Myzus persicae* are the most devastating insects, infesting leaves, stems, and floral parts (Jayaswal et al. 2022).

Due to a symbiotic relationship, the prevalence of aphids and ants was frequently correlated. Aphids produced a delicious substance known as honeydew as a waste product, which ants found highly attractive as a food source (Nelson and Mooney 2022). The honeydew contained an abundance of sugars, extracted by aphids from the plant juice (Zheng et al. 2022). Ants were drawn to this nutrient-rich food source and would often 'farm' aphids for it. In exchange for honeydew, ants provided aphids with protection from other insects and predators, such as ladybugs, lacewing larvae, and parasitic wasps (Karami-jamour et al. 2018). Certain species of ants would transport aphids to new host plants for improved foraging opportunities, ensuring that aphids had a continuous food source (Giannetti et al. 2021). Honeydew not only nourished the ant colony, but its high sugar content also supported the development of their fungus farms (in certain species) and provided energy for the growth of their own progeny (Biedermann and Vega 2020).

## CONCLUSION

15 species of aphids were found in ornamental and wild plants in Pagaralam, namely *Aphis gossypii*, *Uroleucon* sp., *Toxoptera odinae*, *Macrosiphum rosae*, *Aphis citricola*, *Aphis craccivora*, *Toxoptera aurantii*, *Pentalonia nigronervosa*, *Hystenura* sp., *Aphis glycine*, *Greenidae* sp., *Rhopalosiphum padi*, *Rhopalosiphum maidis*, *Hyperomyzus* sp. *Lipaphis erysimi*.

## ACKNOWLEDGMENTS

The authors thank Universitas Sriwijaya, that supported this research.

## REFERENCES

- Alotaibi, N. J., T. Alsufyani, N. H. M'sakni, M. A. Almalki, E. M. Alghamdi, and D. Spiteller. 2023. "Rapid Identification of Aphid Species by Headspace GC-MS and Discriminant Analysis." *Insects* 14(589). doi: <https://doi.org/10.3390/insects14070589>.
- Bass, Chris, Alin M. Puinean, Christoph T. Zimmer, Ian Denholm, Linda M. Field, Stephen P. Foster, Oliver Gutbrod, Ralf Nauen, Russell Slater, and Martin S. Williamson. 2014. "The Evolution of Insecticide Resistance in the Peach Potato Aphid , *Myzus Persicae*." *Insect Biochemistry and Molecular Biology* 51:41–51. doi: 10.1016/j.ibmb.2014.05.003.
- Biedermann, Peter H. W., and Fernando E. Vega. 2020. "Ecology and Evolution of Insect–Fungus Mutualisms." *Annual Review of Entomology* 65:431–55. doi: <https://doi.org/10.1146/annurev-ento-011019-024910>.



404 Blackman, Roger L., and Victor F. Eastop. 2008. *Aphids on the World's Herbaceous Plants and Shrubs, 2 Volume Set*.  
 405 John Wiley & Sons.

406 Blackman, Roger L., and Victor F. Eastop. 2017. "Taxonomic Issues." Pp. 1–36 in *Aphids as crop pests*. CABI  
 407 Wallingford UK.

408 Braham, Mohamed, Synda Boulahia-kheder, Mouna Kahia, and Siwar Nouria. 2023. "Aphids and Citrus Responses to  
 409 Nitrogen Fertilization." *Journal of the Saudi Society of Agricultural Sciences* 22(6):374–83. doi:  
 410 10.1016/j.jssas.2023.03.003.

411 Brożek, Jolanta, Ewa Mróz, Dominika Wyleżek, Łukasz Depa, and Piotr Węgierek. 2015. "The Structure of Extremely  
 412 Long Mouthparts in the Aphid Genus *Stomaphis* Walker (Hemiptera: Sternorrhyncha: Aphididae)." *Zoomorphology*  
 413 134:431–45. doi: <https://doi.org/10.1007/s00435-015-0266-7>.

414 Cao, He-he, Zhan-feng Zhang, Xiao-feng Wang, and Tong-xian Liu. 2018. "Nutrition versus Defense : Why *Myzus*  
 415 *Persicae* ( Green Peach Aphid ) Prefers and Performs Better on Young Leaves of Cabbage." *PloS One* 13(4):1–16.  
 416 doi: <https://doi.org/10.1371/journal.pone.0196219>.

417 Chandel, R. S., V. K. Chandla, K. S. Verma, and Mandeep Pathania. 2022. "Chapter 21 - Insect Pests of Potato in India:  
 418 Biology and Management." Pp. 371–400 in, edited by A. Alyokhin, S. I. Rondon, and Y. B. T.-I. P. of P. (Second E.  
 419 Gao. Academic Press.

420 Clarke, Rebecca, Monica A. Kehoe, Sonya Broughton, and Roger A. C. Jones. 2020. "Host Plant Affiliations of Aphid  
 421 Vector Species Found in a Remote Tropical Environment." *Virus Research* 281(December 2019):197934. doi:  
 422 10.1016/j.virusres.2020.197934.

423 Döring, Thomas F. 2014. "How Aphids Find Their Host Plants, and How They Don't." *Annals of Applied Biology*  
 424 165(1):3–26. doi: <https://doi.org/10.1111/aab.12142>.

425 Ertunc, Filiz. 2020. "Chapter 46 - Emerging Plant Viruses." Pp. 1041–62 in, edited by M. M. B. T.-E. and R. V. P. Ennaji.  
 426 Academic Press.

427 Gadhve, Kiran R., Saurabh Gautam, David A. Rasmussen, and Rajagopalbabu Srinivasan. 2020. "Aphid Transmission of  
 428 Potyvirus: The Largest Plant-Infecting RNA Virus Genus." *Viruses* 12(7):773. doi: doi: 10.3390/v12070773.

429 Giannetti, Daniele, Mauro Mandrioli, Enrico Schifani, Cristina Castracani, Fiorenza A. Spotti, Alessandra Mori, and  
 430 Donato A. Grasso. 2021. "First Report on the Acrobat Ant *Crematogaster Scutellaris* Storing Live Aphids in Its Oak-  
 431 Gall Nests." *Insects* 12(2):108. doi: <https://doi.org/10.3390/insects12020108>.

432 Guo, Huijuan, Liyuan Gu, Fanqi Liu, Fajun Chen, Feng Ge, and Yucheng Sun. 2019. "Aphid-Borne Viral Spread Is  
 433 Enhanced by Virus-Induced Accumulation of Plant Reactive Oxygen Species 1." *Plant Physiol* 179(January):143–  
 434 55. doi: 10.1104/pp.18.00437.

435 Jakubczyk, Karolina, Klaudia Koprowska, Aleksandra Gottschling, and Katarzyna Janda-Milczarek. 2022. "Edible  
 436 Flowers as a Source of Dietary Fibre (Total, Insoluble and Soluble) as a Potential Athlete's Dietary Supplement."  
 437 *Nutrients* 14(12). doi: 10.3390/nu14122470.

438 Jaouannet, Maëlle, Patricia A. Rodriguez, Peter Thorpe, Camille J. G. Lenoir, and Ruari Macleod. 2014. "Plant Immunity  
 439 in Plant – Aphid Interactions." *Front Plant Sci.* 5(December):1–10. doi: 10.3389/fpls.2014.00663.

440 Jayaswal, Deepanshu, Pawan Mainkar, Kuldeep Kumar, Yamini Agarwal, and Ratna Prabha. 2022. "Pyramiding and  
 441 Evaluation of Segregating Lines Containing Lectin and Protease Inhibitor Genes for Aphid Resistance in Brassica  
 442 Juncea." *Indian Journal of Biochemistry & Biophysics* 59(August):800–807. doi: 10.56042/ijbb.v59i8.62319.

443 Jones, Roger A. C. 2022. "Alteration of Plant Species Mixtures by Virus Infection: Managed Pastures the Forgotten  
 444 Dimension." *Plant Pathology* 71(6):1255–81. doi: DOI: 10.1111/ppa.13571.

445 Jousset, Emmanuelle, Genson Gwenaelle, and Coeur D. Acie. Armelle. 2010. "Evolutionary Lability of a Complex Life  
 446 Cycle in the Aphid Genus *Brachycaudus*." *BMC Evolutionary Biology* 10(1). doi: 10.1186/1471-2148-10-295.

447 Kallas, John. 2010. *Edible Wild Plants*. Gibbs Smith.

448 Karami-jamou, Tahereh, Alinaghi Mirmoayedi, Abbasali Zamani, and Yadolah Khajehzadeh. 2018. "The Impact of Ant  
 449 Attendance on Protecting *Aphis Gossypii* against Two Aphidophagous Predators and It's Role on the Intraguild  
 450 Predation between Them." *Journal of Insect Behavior* 31:222–39. doi: DOI: 10.1007/s10905-018-9670-4.

451 Kennedy, J. S., and H. L. G. Stroyan. 1959. "Biology of Aphids." *Annual Review of Entomology* 4(1):139–60.

452 Kinley, Chimi, A. Najitha Banu, A. M. Raut, Johnson Wahengbam, and Thinley Jamtsho. 2021. "A Review on Past,  
 453 Present and Future Approaches for Aphids Management." *Journal of Entomological Research* 45(2):336–46. doi:  
 454 10.5958/0974-4576.2021.00053.0.

455 Kumar, Sarwan. 2019. "Aphid-Plant Interactions: Implications for Pest Management." P. Ch. 7 in *Plant Communities and*  
 456 *Their Environment*, edited by M. T. Oliveira, F. Candan, and A. Fernandes-Silva. Rijeka: IntechOpen.

457 Kumar, Sushil, Malay K. Bhowmick, and Puja Ray. 2021. "Weeds as Alternate and Alternative Hosts of Crop Pests."  
 458 *Indian Journal of Weed Science* 53(1):14–29. doi: 10.5958/0974-8164.2021.00002.2.

459 Liu, Xiang Dong, Ting Ting Xu, and Hai Xia Lei. 2017. "Refuges and Host Shift Pathways of Host-Specialized Aphids  
 460 *Aphis Gossypii*." *Scientific Reports* 7(1):1–9. doi: 10.1038/s41598-017-02248-4.

461 Maharani, Yani, Purnama Hidayat, Aunu Rauf, and Nina Maryana. 2018. "Short Communication: New Records of Aphid  
 462 Species Subfamily Aphidinae (Hemiptera: Aphididae) in West Java, Indonesia." *Biodiversitas* 19(2):460–65. doi:  
 463 10.13057/biodiv/d190219.

- Margaritopoulos, J. T., M. Tzortzi, K. D. Zarpas, J. A. Tsitsipis, and R. L. Blackman. 2006. "Morphological Discrimination of *Aphis Gossypii* (Hemiptera: Aphididae) Populations Feeding on Compositae." *Bulletin of Entomological Research* 96(2):153–65. doi: 10.1079/ber2005410.
- Meuninck, Jim. 2023. *Basic Illustrated Edible Wild Plants and Useful Herbs*. Rowman & Littlefield.
- Mo, Carmen, and Angela M. Smilanich. 2023. "Feeding on an Exotic Host Plant Enhances Plasma Levels of Phenoloxidase by Modulating Feeding Efficiency in a Specialist Insect Herbivore." *Frontiers in Physiology* 14(February):1–10. doi: 10.3389/fphys.2023.1127670.
- Naidu, VSGR. 2012. "Hand Book on Weed Identification."
- Nelson, Annika S., and Kailen A. Mooney. 2022. "The Evolution and Ecology of Interactions between Ants and Honeydew-Producing Hemipteran Insects." *Annual Review of Ecology, Evolution, and Systematics* 53:379–402. doi: <https://doi.org/10.1146/annurev-ecolsys-102220-014840>.
- Pettersson, Jan, W. Fred Tjallingii, and Jim Hardie. 2017. "Host-Plant Selection and Feeding." Pp. 173–95 in *Aphids as crop pests*. CABI Wallingford UK.
- Piron, PGM, MC de Haas, and MAHM Sonnemans. 2019. "The Presence of *Aphis* (Toxoptera) *Aurantii* (Homoptera: Aphididae) in the Netherlands." *Entomologische Berichten* 79(5):162–64.
- Santiago, Grazielle Silva, Luana Fonseca, Ernesto Oliveira Canedo-ju, Carla Rodrigues Ribas, Rafaela Pereira, and Guilherme Pereira Alves. 2017. "Isolated and Community Contexts Produce Distinct Responses by Host Plants to the Presence of Ant-Aphid Interaction: Plant Productivity and Seed Viability." *Plos One* 12(1):1–12. doi: 10.1371/journal.pone.0170915.
- Singh, Rajendra, and Garima Singh. 2021. "Aphids." Pp. 105–82 in *Polyphagous Pests of Crops*. Springer.
- Tegelaar, Karolina, Mattias Hagman, Robert Glinwood, Jan Pettersson, and Olof Leimar. 2012. "Ant–Aphid Mutualism: The Influence of Ants on the Aphid Summer Cycle." *Oikos* 121(1):61–66. doi: <https://doi.org/10.1111/j.1600-0706.2011.19387.x>.
- Yamamoto, Tetsuya, Mitsuru Hattori, and Takao Itino. 2020. "Seasonal Migration in the Aphid Genus *Stomaphis* (Hemiptera : Aphididae): Discovery of Host Alternation Between Woody Plants in Subfamily Lachninae." *Journal of Insect Science* 20(5):1–10. doi: 10.1093/jisesa/ieaa103.
- Zheng, Zhou, Mengqin Zhao, Zhijun Zhang, Xin Hu, Yang Xu, and Cong Wei. 2022. "Lactic Acid Bacteria Are Prevalent in the Infrabuccal Pockets and Crops of Ants That Prefer Aphid Honeydew." *Front. Microbiol.* 12(January):1–17. doi: 10.3389/fmicb.2021.785016.