

## 6. Bukti konfirmasi review kelima dan hasil revisi kelima

15 Desember 2023

The image shows two screenshots of an email interface. The top screenshot displays the header of an email from "Smujo Editors via SMUJO" with the subject "[biodiv] Editor Decision". The email body states that a decision has been reached regarding a submission to the Biodiversitas Journal of Biological Diversity, and that the decision is "Revisions Required". The bottom screenshot shows the content of "Reviewer A", which lists several specific comments and recommendations for revision, such as checking plagiarism, sentence shading, and the presence of aphids in monocot plants. The email interface includes a search bar, navigation icons, and a sidebar with various app icons.

smujo

[biodiv] Editor Decision Eksternal Kotak Masuk x

Smujo Editors via SMUJO <support@smujo.com> kepada saya, Erise

Jum, 15 Des 2023, 16.59

Terjemahkan ke Indonesia

Chandra Irsan, Erise Anggraini:

We have reached a decision regarding your submission to Biodiversitas Journal of Biological Diversity, "Species of aphids found in ornamental and wild Plants in Highland, Pagar Alam, South Sumatra". Complete your revision with a Table of Responses containing your answers to reviewer comments (for multiple comments) or enable Track Changes.

Our decision is: Revisions Required

Reviewer A:

Reviewer A:

Plagiarism is under 5% without references.

The sentences in shading need more attention to accomplish.

The strike-through sentence could be deleted without any reduced information.

Please check this Adiantum predatum plant was not found on any Tables/Figures and observation results.

Rhopalosiphum rice differs in Tables 3 and 4, as Rhopalosiphum padi. Please check.

The presence of aphids in monocot plants or plants. Please check whether this sentence (plant) is redundant or a monocot and dicot plant. Please clarify

macrocyini subfamily was not available online. Please clarify

Please clarify the wording Pagar Alam (as stated in the title); it is not a continuous pattern as Pagaralam.

This conclusion could be more deeply investigated by presenting (in brief) the diversity of aphid species found in ornamental and wild plants (this study's purposes), what kind of aphid species are preferred by ants, why aphids prefer the weed species, why *aphis gossypii* species could colonize 12 plants while *aphis citricola* only one, etc.

It is better to state that further research on the diversity of aphid species found in the area's ornamental and wild plants.

Recommendation: Revisions Required

# Species of aphids found in ornamental and wild Plants in Highland, Pagar Alam, South Sumatra

**Abstract.** Aphids are one of the crucial pests in the tropical and sub-tropical regions. The presence of aphids in a plant can be very detrimental due to their role as vectors. Aphids exhibit species diversity, but not much information has been reported about the species diversity of aphids associated with essential crops such as ornamental plants. Furthermore, many aphid species were found on plants that were not actually hosts such as wild plants. Therefore, this study reported the species of aphids found in ornamental plants and the wild plants. The field research employed a purposive and direct observation approach to inventory cultivated or wild plants hosting aphids and collecting aphids. The plant selection process included cultivated plants encompassing ornamental plants, as well as wild plants or weeds. The collection and identification of host plants, and aphids, involved systematic searches for the selected plants and subsequent examination for the presence of aphids. Observations were made to all existing plant species to find those colonized by aphids. This study revealed that 15 species of aphids were found in Pagaralam, namely *Aphis gossypii*, *Uroleucon* sp., *Toxoptera odinae*, *Macrosiphum rosae*, *Aphis citricola*, *Aphis craccivora*, *Toxoptera aurantii*, *Pentalonia nigronervosa*, *Hystenura* sp., *Aphis glycine*, *Greenidae* sp., *Rhopalosiphum padi*, *Rhopalosiphum maidis*, *Hyperomyzus* sp. *Lipaphis erysimi*.

**Keywords:** Aphids, ornamental plants, wild plants

**Running title:** Aphids found in ornamental and wild plants

## INTRODUCTION

Aphids are one of the crucial pests in the tropics and sub-tropics, exhibiting various polyphagous, oligophagous, and monophagous characteristics (Kennedy and Stroyan 1959). One species of aphids can host more than 400 species from 40 families (Bass et al. 2014). In addition to pests, aphids can also be vectors of plant viral diseases (Gadhawe et al. 2020). Aphids can transmit 275 viruses (Ertunc 2020). In tropical areas, aphids can ~~always~~ be found throughout the year due to their parthenogenetic nature of reproduction (Blackman and Eastop 2017). Aphids ~~suck phloem sap of tender plant parts~~ ~~consume young leaves sap~~, which can deplete essential nutrients for healthy growth (Cao et al. 2018). Moreover, ~~vector species when aphids transmit viral diseases from one plant to another, this~~ can further weaken and stunt the growth of infected plants (Jones 2022). ~~According to Kinley et al. (2021), aphids cause yield losses directly (35–40%) by sucking the plant sap or indirectly (20–80%) through viral transmission. Therefore, aphid infestations can have adverse effects on crop yields and overall plant health (Kumar 2019).~~

Due to their function as vectors, the presence of aphids on a plant can be highly damaging (Jaouannet et al. 2014). ~~They feed by piercing the plant's tissues and consuming its sap, which can reduce the plant's growth and productivity, ultimately leading to weakness and possible death (Chandel et al. 2022).~~ Additionally, as vectors, aphids can transmit a variety of plant diseases. They are as carriers for various plant viruses, and when they move from infected to healthy plants, these viruses can rapidly spread and cause extensive damage (Guo et al. 2019). ~~In addition, the honeydew that aphids secrete can lead to the growth of sooty mold, a black fungus that can prevent sunlight from reaching the plant's leaves, thereby impairing photosynthesis, the process by which plants produce food (Singh and Singh 2021).~~ Therefore, it is crucial to control aphid populations in gardens and crops.

Many aphid species ~~are~~ were found on plants that ~~are~~ were not their actual hosts (Maharani et al. 2018). Aphids have one or more secondary, or "alternative," host plants in addition to their primary host plants, which are the types of plants they feed on most frequently (Clarke et al. 2020). ~~Alternative plants An alternative host can also be a collateral host belonging to the same plant family as the primary host, helping crop pests to survive when the primary hosts are unavailable (Kumar et al., 2021). These secondary hosts may offer less adequate nutrition for insects (Mo and Smitanich 2023). However, they may provide a means of survival when primary hosts are unavailable, during certain seasons, or~~

under certain environmental conditions (Kumar et al., 2021). According to Liu et al. (2017), since hibiscus serves as an overwintering host for cotton-specialized aphids but not for cucurbit-specialized aphids, it is evident that host-specialized aphids have refuges during times of food shortage. The life cycles of numerous aphid species exhibit such complexity (Jousselin, Gwenaëlle, and Armelle 2010). They maintain a cycle of host alternation, shifting between their primary hosts (typically a woody plant) and secondary hosts (often herbaceous plants) (Yamamoto, Hattori, and Itino 2020). Weeds pose a continuous threat in both cropped and non-crop areas, providing food, shelter and reproductive sites for various pest organisms (Kumar et al., 2021). This indicates that weeds can serve as alternative hosts for aphids.

A study of aphid species on horticultural plants has been conducted (Maharani et al. 2018). However, information about aphid species on ornamental and wild plants has not received as much attention and remains largely unexplored. In South Sumatra, particularly in the highland areas like Pagar Alam, there are numerous ornamental and native plants. The research on the diversity of aphid species in ornamental and wild plants has received little attention. Therefore, this study reports was conducted in Pagar Alam, a highland region of South Sumatra, with the aim of obtaining information on the diversity of aphid species found in ornamental and wild plants found in this area. The findings from this study can serve as a valuable resource for aphid management.

MATERIALS AND METHODS

The field research employed a purposive and direct observation approach to inventory of cultivated or wild plants hosting aphids and collecting aphids. The plant selection process included cultivated plants encompassing ornamental plants, as well as wild plants or weeds. The collection and identification of host plants, and aphids, and natural enemies where available, involved systematic searches of for the selected plants and subsequent examination for the presence of aphids. Observations were made to all existing plant species to find those colonized by aphids. Any plants colonized by aphids were documented as aphid hosts. Aphids, along with their natural enemies within the aphid colonies, were systematically collected. All components of the collected observations were then identified.

Aphid identification was done using identification keys (Blackman and Eastop 2008). Identification of aphid species took place in the Laboratory of Entomology, Faculty of Agriculture, Universitas Sriwijaya. Identification relied on morphological characteristics. The host plants were identified using weed identification hand book (Kallas, 2010; Meuninck, 2023; Naidu, 2012). The location and size of aphid colonies, morphology of aphids including their life shape and color, as well as any symptoms observed in the host plants were recorded, and photographs of the aphid colonies and their host plants were recorded taken.

RESULTS AND DISCUSSION

Result

The results showed that 15 aphid species were found in Pagar Alam, namely (Tables 1, 2). Aphis gossypii, Uroleucon sp., Toxoptera odinae, Macrosiphum rosae, Aphis citricola, Aphis craccivora, Toxoptera aurantii, Pentalonía nigronervosa, Hystenura sp., Aphis glycine, Greenidae sp., Rhopalosiphum padi, Rhopalosiphum maidis, Hyperomyzus sp., Lipapis erysimi. Based on the observation, these aphids mostly colonised flowers of were found on various ornamental plants (Table 1, Figure 1). The primary colony locations were generally in flowers, and this study documented these colony locations in ornamental plants (Figure 1).

Table 1. Aphid species recorded found in ornamental plants and their colony locations

No	Host Plant	Aphid Species	Colony location
1	Aster alpinus	Sitobion luteum	flower
2	Brugmansia suaveolens	Aulacorthum solani	flower
		Neomyzus circumflexus	
		Myzus persicae	
3	Caladium sp.	Pentalonia sp	flower
4	Cananga odoratum	Aphis gossypii	flower
5	Canna indica	Pentalonia nigronervosa	flower
6	Catharanthus roseus	Aphis citricola	flower
7	Cestrum sp.	Aphis gossypii	flower
		Neomyzus circumflexus	
8	Clitoria ternatea	Aphis craccivora	flower
9	Cosmos caudatus	Uroleucon sp.	flower
10	Dahlia 'Kelvin'	Aphis gossypii	flower
11	Dendrobium sp.	Sinemogoura citricola	flower
12	Duranta sp.	Aphis gossypii	flower

Formatted: Highlight

Formatted: Highlight

Formatted: Highlight

Formatted: Highlight

Formatted: Font: Not Italic

Formatted: English (United States)

13	<i>Helianthus</i> sp.	<i>Aphis glycines</i> <i>Hyperomyzus</i> sp.	flower
14	<i>Hibiscus rosinensis</i>	<i>Aphis gossypii</i>	flower
15	<i>Ixora paludosa</i>	<i>Aphis gossypii</i> , <i>Toxoptera aurantii</i>	flower
16	<i>Ixora</i> sp.	<i>Aphis citricola</i> <i>Aphis gossypii</i> <i>Toxoptera aurantii</i>	flower
17	<i>Murraya paniculata</i>	<i>Aphis craccivora</i> <i>Toxoptera citricidus</i>	flower
18	<i>Mussaenda frondosa</i>	<i>Aphis citricola</i> <i>Toxoptera odinae</i>	flower
19	<i>Rosa indica</i>	<i>Macrosiphum rosae</i>	flower
20	<i>Spondias dulcissolana</i>	<i>Aphis citricola</i> <i>Hysteroneura setariae</i>	flower



**Fig 1. Photos showing colonies of different aphid species in ornamental plants: The location of aphid colonization on various plant parts.** a) *A. gossypii* in *D. kelvin* flower b) *A. gossypii* in *H. rosinensis* flower c) *A. gossypii* in tuberose flower, d) *A. craccivora* in *Clitoria ternatea* flower, e) *A. citricola* in *Helianthus* sp., f) *A. aurantii* on the *M. paniculata* flower, g) *T. odinae* in the *S. dulcissolana*, h) *Uroleucon* sp. in *chrysanthemums*, i) *Macrosiphum rosae* in *R. indica* flower, j) *Pentalonia nigronervosa* in *C. indica* leaves

98

99

100

101

102

103

104

105

106

107

108

109

110

110

In addition, this study documented the presence of weeds, which might serve as alternative hosts for aphids (Table 2). The location of aphid colonies also varied, namely on flowers, stalks, plant tops, young leaves and old leaves of wild plants (Table 2, Figure 2). The presence of specific plants or host plants within a habitat influenced the types of aphids found. Many aphid species are found on a broad range of plants or host plants, while others are highly specialized and are only found on specific plants or host plants. This is closely related to the polyphagous, oligophagous or monophagous nature of aphids (Blackman & Eastop 2000).

**Table 2.** Species of aphids found in wild or weed plants and their colony locations

No	Host Plant	Aphid species	Colony location
1	<i>Ageratum conyzoides</i>	<i>Aphis gossypii</i>	Shoots, young leaves, old leaves, flowers
2	<i>Alternanthera philoxeroides</i>	<i>Aphis gossypii</i>	Shoots, buds
3	<i>Alternanthera sessilis</i>	<i>Aphis gossypii</i>	Shoots, buds
4	<i>Amaranthus gracilis</i>	<i>Aphis craccivora</i>	Flowers, shoots, young leaves, old leaves
5	<i>Blumea lacera</i>	<i>Lipaphis erysimi</i>	Flowers, shoots, and buds
6	<i>Croton hirtus</i>	<i>Aphis gossypii</i>	Flowers, shoots, young leaves, old leaves, young twigs
7	<i>Cynodon dactylon</i>	<i>Schizaphis rotundiventris</i>	Flower, flower stalks
8	<i>Cyperus rotundus</i>	<i>Schizaphis rotundiventris</i>	Flower, flower stalks, leaf axils
9	<i>Cyperus compressus</i>	<i>Schizaphis rotundiventris</i>	Flower, flower stalks, leaf axils
10	<i>Digitaria ciliaris</i>	<i>Hysteronura setariae</i>	Flower, flower stalks
11	<i>Echinocloa crusgali</i>	<i>Hiperomyzus</i> sp.	Young leaves, old leaves
12	<i>Ecliptica prostrata</i>	<i>Aphis gossypii</i>	Shoots, young leaves
13	<i>Eleusin indica</i>	<i>Hysteronura setariae</i>	Flower, flower stalks, leaf axils
		<i>Rhopalosiphum maidis</i>	
14	<i>Emilia sonchifolia</i>	<i>Aphis gossypii</i>	Flower, flower stalks, shoots
15	<i>Eragrostis tenella</i>	<i>Hysteronura setariae</i>	Flower, flower stalks, seeds
16	<i>Euphorbia hirta</i>	<i>Aphis gossypii</i>	Young leaves, old leaves
17	<i>Eupotarium odoratum</i>	<i>Aphis gossypii</i> , <i>Aphis glycine</i>	Young leaves, old leaves, young twigs
18	<i>Hymenochera acutigluma</i>	<i>Hysteronura setariae</i>	Flowers, flower stalks, leaf axils
19	<i>Lagerstromia</i> Sp.	<i>Greenidea</i> sp.	Young leaves
20	<i>Lophatherum gracile</i>	<i>Hysteronura setariae</i>	Young leaves, old leaves, leaf axils
		<i>Rhopalosiphum maidis</i>	
21	<i>Melastoma affine</i>	<i>Aphis gossypii</i>	Shoots, young leaves
22	<i>Mikania mikranta</i>	<i>Aphis gossypii</i> <i>Aphis glycine</i>	Shoots, young leaves, old leaves
23	<i>Mimosa invisa</i>	<i>Aphis craccivora</i>	Shoots, pods
24	<i>Mimosa pudica</i>	<i>Aphis craccivora</i>	Shoots, pods, flowers
25	<i>Mimosa vigra</i>	<i>Aphis craccivora</i>	Shoots, pods
26	<i>Oryza rufipogon</i>	<i>Rhopalosiphum padi</i> , <i>Rhopalosiphum maidis</i>	Old leaves, young leaves (pupus), leaf axils
27	<i>Oxonopus compressus</i>	<i>Hysteronura setariae</i>	Flower, flower stalk, leaf axils
28	<i>Paspalum conjugatum</i>	<i>Hysteronura setariae</i>	Flower, flower stalk, seeds
29	<i>Phyllanthus neruri</i>	<i>Aphis citricola</i>	Shoot, young leaves, old leaves, young twigs, petioles
30	<i>Portulaca oleraceae</i>	<i>Aphis craccivora</i>	Shoots, young leaves, flower
31	<i>Physalis angulata</i>	<i>Aphis craccivora</i> , <i>A. gossypii</i>	Shoots, young leaves, old leaves
32	<i>Rorippa indica</i>	<i>Lipapis erysimi</i>	Flower, fruit, shoots, young leaves
33	<i>Sida rhombifolia</i>	<i>Aphis gossypii</i>	Shoots, young leaves, old leaves, fruit/seeds
34	<i>Sonchus arventris</i>	<i>Lipapis erysimi</i>	Young leaves, fruit stalks, flower, fruit

Formatted: Highlight

Formatted: Highlight





**Figure 2.** Aphids found infesting wild plants a) *A. gossypii* on *Ageratum conyzoides*, b) *A. gossypii* on *Croton* weed *hirtus* c) *A. gossypii* on *Eupatorium odoratum*, d) *A. gossypii* on *Pachystochys* sp., e) *A. gossypii* on *Caladium* sp., f) *A. gossypii* on *Alternanthera sessilis*, g) *A. gossypii* on *Portulaca oleraceae* weeds, h) *A. gossypii* on *Euphorbia hirta*, i) *A. citricola* on *Phyllanthus nerruri*, j) *A. citricola* on *Sida rhombifolia* weed, k) *A. citricola* on *Annona muricata*, l) *A. citricola* on *Ludwigia peruviana*, m) *A. craccivora* on *Mimosa pudica* weed, n) *A. craccivora* on

Formatted: Font: Not Italic

Formatted: Font: Italic

Formatted: Font: Not Italic

Formatted: Font: Not Italic

Formatted: Font: Not Italic

Formatted: Font: Not Italic

Formatted: Font: Not Italic

118 weeds-*Amaranthus gracilis*, o) *A. glycine* in *Mikania micranta*-weed, p) *Hysteneura* sp. in *Eleusin*-weeds, q) *Greenidae* sp. in kenidai  
119 trees (shrubs) *indica*, r) *Hyperomyzus* sp. in *Echinocloa crusgali*-Weed, s) *L. erysimi* ion weed-sonchus arventris, t) *Rhopalosiphum* rice  
120 ion the-weed-*Oryza rufipogon*, u) *Rhopalosiphum* Maidis ion the-weed-*Oryza rufipogon*.  
121

Formatted: Highlight

Formatted: Highlight

Formatted: Font: Not Italic

122 Discussion

123  
124 The plant species or host plant influences the distribution of aphids. There are aphid species that can be found on a  
125 wide range of host plants, which is closely related to the polyphagous nature of aphids, allowing them to colonize many  
126 different species of host plants. Host plants can also affect the distribution of aphids, as evidenced by the presence of aphid  
127 species exclusively found on certain host plants (Santiago et al. 2017). But there are some species of aphids found only on  
128 one particular host and are not found on other host plants (Döring 2014). *A. gossypii*, and *Aphis aurantii* have been found  
129 on many host plants because both aphids are classified as polyphagous aphids (Alotaibi et al., 2023).

130 Many aphid species can commonly be found infesting a variety of ornamental plants because these insects are  
131 attracted to such these plants due to the rich nutrient content in the plant sap (Braham et al. 2023). In the present study,  
132 some aphid species were found on some ornamental plants in Pagaram. The location of aphid colonization on the plants  
133 varied. On *Adiantum predatum* plants, aphids formed colonies on young leaf stalks and on newly emerging leaves. The  
134 aphids displayed brown and black coloration. The aphid colonies found were small, and the colonized plant parts showed  
135 no signs of disease. The identification results showed that the aphids were *Neotoxoptera* sp., and notably, they were not  
136 associated with ants. On *Aster alpinus*, aphids were found to form colonies on the stems or young leaf shoots, and the  
137 colonies were relatively large. The color of the aphids was dark brown to black. The colonized plant parts showed  
138 symptoms of stunting. The identification results showed that the aphids were *Uroleucon* sp., and they were associated with  
139 ants.

Formatted: Font color: Red, Highlight

Formatted: Font color: Red

Formatted: Highlight

140 On the *Brugmansia suaveolens*, *M. persicae* were found on the undersides of old leaves or leaves that have started to  
141 turn yellow. The colonies were relatively small. The aphids found were green and large bodies. The colonized plant parts  
142 did not show any signs of disease. On *Caladium* sp. (taro) was found one species of aphids: *A. gossypii*. The aphids formed  
143 colonies under the surface of young and older leaves. The occupied leaf areas did not display severe symptoms. The aphids  
144 were yellow green to dark green. The wingless adult aphids often had a white, flour-like appearance on their bodies. On  
145 the *Cananga odoratum* (ylang-ylang), colonies of *T. aurantii* were found on the undersides of the leaves, the shoots, buds,  
146 and unopened flower petals. The *T. aurantii* colonies found were relatively large. Colonized parts, especially shoots,  
147 showed signs of stunting. The aphids found were brown to black in color. The colonies of *T. aurantii* were found to be  
148 associated with black ants.

149 Aphids on *C. indica* (Indian shot, African arrowroot) were found to form colonies in the leaf axils and under the leaf  
150 surface near the leaf base. The colonies were quite large. The aphids were dark brown to dark red coloring with a medium-  
151 sized body. The identification results showed that the aphids were *P. nigronervosa*. The colonies of *P. nigronervosa* were  
152 found to be associated with ants. In the *Catharanthus roseus* (periwinkle), *A. citricola* aphids were found. The aphids were  
153 yellow-green, sifunculi, and black cauda. The aphids formed colonies on flowers and shoots, and the colonized plant parts  
154 did not show any symptoms of disease. On *Cestrum* sp. (Bastard jasmine), aphids formed colonies on the undersides of  
155 young leaves, shoots, and within flower parts, especially between petals or flower stalks that had not fully bloomed. The  
156 colonies were quite large. The body color of aphids was green to dark green with small to medium-sized bodies. The  
157 colonized plant parts, especially leaves, showed stunting symptoms. The identification results showed that the aphids were  
158 *A. gossypii*. The aphid colonies found were consistently associated with ants.

159 Aphids on *Clitoria ternatea* were found to form colonies on flower parts, flower crowns, stems and young leaves. The  
160 aphids were brown to black in color. Colonized plant parts, especially shoots and young leaves, showed stunting  
161 symptoms. The identification results showed that the aphids were *A. craccivora*. These colonies were consistently  
162 associated with ants. On the plant *Cosmos caudatus*, aphids were found on the flower petals. The colonies were not very  
163 large. The body color was green and light green. The identification results showed that the aphids were *A. gossypii*, and  
164 they were also associated with ants. The aphids on the *Dahlia kellyi* plant formed colonies on unopened flower buds, with  
165 a significant population among the blooming petals. The body color was green to dark green. The identification results  
166 showed that the aphids were *A. gossypii*. Aphids on *Datura metel* (amethyst) were found to form colonies on the  
167 undersides of old leaves. The aphids were medium sized with a green body color. The colonized plant parts did not show  
168 any symptoms of disease. The identification results showed that the aphids were *Myzus ornatus*. The aphid colonies were  
169 not associated with ants. Within *Dendrobium* sp., aphid colonies were found on the young leaves. The aphids were yellow,  
170 green to dark green. The colonized plants did not show any disease symptoms. The identification results showed that the  
171 aphids were *A. gossypii*, and they were associated with ants. On *Duranta* sp. (bonsai), colonies of aphids were located on  
172 the undersides of young leaves. The colonized plant parts showed stunting symptoms. The colonies were very large. The  
173 aphids were green in color. The identification results showed that the aphids were *A. gossypii*. The aphid colonies were  
174 consistently associated with ants.

175 On the *Helianthus annuus* (sunflower) plants, aphid colonies were found between the flower petals. The colonized  
176 flowers, especially the crowns, exhibited a tendency to fall off easily. The aphids were green and yellow in color. The  
177 colonies were small. The identification results showed that the aphids were *A. gossypii*. These aphid colonies were

associated with ants. Aphid colonies on *Helianthus* sp. were found on the undersides of old leaves. These colonies were small in size. The aphids were green with a medium body size. The colonized plant parts did not show any disease symptoms. The identification results showed that the aphids were *M. ornatus*. The aphid colonies were not associated with ants. Within the colonies, mummified aphids that were parasitized by Aphidiidae were found.

On the *Hibiscus rosa-sinensis*, aphids ranging in color from yellow to dark green were found. The aphids formed colonies on flower buds, unopened flower crowns, and the undersides of aging leaves. The colonies grew to be very large. The identification results showed that the aphids were *A. gossypii*. The aphid colonies were consistently associated with ants. Two types of aphids were found on the flowering plant *Ixora paludosa*. First, the aphids formed colonies on the undersides of young leaves that were still red or light green and sometimes on flower stalks that had not yet bloomed. The occupied plant parts showed symptoms such as stunted leaf growth, leaf shrinkage, necrotic spots on the leaf surface, and slightly downward-curved leaf edges. The upper leaf surface looked wet and sticky, similar to sugar. The aphids had yellow, green, or slightly dark green bodies, with some wingless adults having a powdery white upper surface. The identification results showed that the aphids were *A. gossypii*, and they were almost always associated with ants. The second type of aphids on *Ixora paludosa* formed colonies under the surface of young and older leaves. The colonies could also be found on newly emerging flowers and leaves. The plant parts occupied by these aphids did not show obvious signs of illness. These aphids were dark red to black, with once-branched stigma and venation in their black wings. The identification results showed that the aphids were *T. aurantii*. These aphids were also associated with ants.

In *Ixora* sp. flower plants, two forms of aphids were discovered. These aphids occupied the shoots, young leaves and unopened flowers. The affected plant parts did not show obvious symptoms. The aphids exhibited colors ranging from yellow and green to a slightly darker green. Sometimes the upper surface of the wingless imago's body appeared white, resembling flour. The identification results showed that these aphids were *A. gossypii*. These aphid colonies were almost always associated with ants. Another species of aphids was founded and formed colonies on flower stalks that had not yet bloomed and on newly emerging shoots or leaves. The presence of these aphids on the plant did not induce any symptoms of plant disease. The aphids were yellow or yellow green, with black cauda and siphunculi. Their bodies were very small to small. The identification results showed that the aphids were *A. citricola*. The colonies of *A. citricola* were also frequently found in association with ants.

Two types of aphids were found on *Mussaenda frondosa*, each forming colonies in different locations. The first type formed colonies on young leaves, shoots, and flowers. The plant parts they occupied showed no obvious disease symptoms. The aphids were yellow, green, and some with dark green. The identification results showed that the aphids were *A. gossypii*. The second type of aphids formed colonies on the stems or young twigs, appearing densely clustered as if piled up. The aphid colonies could also be found on young leaves, shoots and within flower parts. The plant parts they infested showed no signs of diseases. The aphids were yellow or yellow green, with black cauda and siphunculi. They had tiny to small bodies. The identification results showed that the aphids were *A. citricola*.

The results showed that 34 species of wild plants, including weeds, were growing in the yard colonized by aphids. This indicated that multiple species of aphids colonized various host plants. The aphid species colonizing these plants were generally consistent within the same taxon. *Ageratum conyzoides* was infested by *Aphis gossypii*. These aphids formed colonies on the flower sections, shoots, lower surfaces of young leaves, or leaves turning yellow. The aphids were green, yellow-green to dark green, often forming large colonies. *Alternanthera philoxeroides* or alligator grass was also colonized by *Aphis gossypii*. Small colonies were found on shoots or stems. These aphids had small bodies and were green, ranging from yellow-green to dark green. *Alternanthera sessilis* was colonized by *Aphis gossypii*, forming colonies on shoots, flowers, and fruit. The colonies were typically large, and they were often associated with tiny brown ants. *Amaranthus gracilis* was infested by *Aphis craccivora*. These aphids established colonies on shoots, flowers and young and old leaves. They were dark brown to black in color, with shiny black wingless imagoes. Colonies of these aphids were associated with both black and red ants. *Blumea lacera* was colonized by *Lipaphis erysimi* aphids. These aphids were bright green, and of medium size. The colonies formed on flowers, flower stalks and the undersides of the leaves at the top. The aphid colonies were not associated with ants. *Croton hirtus* or fire grass was infested by *Aphis gossypii*. The aphids were yellow-green to dark green. The colonies were found on the stems, leaves, buds and flowers, often forming large colonies. *Cynodon dactylon* or Bermuda grass was colonized by *Schizaphis rotundiventris*. The aphids colonized the flowers, flower stalks and sometimes in the leaf axils of the plant. Small colonies were formed. The aphids were brown to red-brown. They were associated with ants. *Cyperus rotundus* or nut grass was infested by *Schizaphis rotundiventris* aphids. The colonies were found on flower stalks, flowers, and leaf axils. The colonies were quite large and associated with both black and red ants. The aphids were dark brown in color. *Cyperus compressus* or grass puzzle was colonized by *Schizaphis rotundiventris* aphids, forming colonies in the flowers, flower stalks and sometimes in the axils and leaves of the shoots or buds. Small colonies were observed. *Digitaria ciliaris* was infested by *Hysteroneura setariae* aphids, with small colonies scattered on the flowers and flower stalks. These aphids were light brown to brown in color. *Echinocloa crussgali* or water hyacinth plants were colonized by *Hiperomyzus* sp. aphids. These aphids were dark brown to black and formed large colonies on the undersides of both young and old leaves. The aphid colonies were never found in association with ants. *Ecliptica prostrata* or urang aring was colonized by *Aphis gossypii*, forming small colonies on the shoots and flowers. The aphids were bright green to blackish green. The aphid colonies were also consistently associated with ants.



237 *Eleusine indica* was colonized by two species of aphids: *Hysteroneura setariae* and *Rhopalosiphum maidis*. *H. setariae*  
238 formed colonies in flower parts, flower stalks and leaf axils resulting in quite large colonies. *H. setariae* body color ranged  
239 from red brown to dark brown. The colonies were consistently associated with ants. The aphids of *R. maidis* formed  
240 colonies in the leaf axils and undersides of leaves and on leaf shoots that had not yet opened. The colonies were not  
241 densely packed. The leaf aphids of *R. maidis* were green in color, with distinct black siphunculi and cauda. These aphids  
242 had relatively large bodies with a slightly elongated shape. *R. maidis* colonies were always associated with ants. The plant  
243 *Emilia sonchifolia*, characterized by its purple flowers, was colonized by *Aphis gossypii*. The aphids were yellow to green  
244 in colour. The colonies formed near flowers, flower stalks, and shoot leaves.

245 *Eragrostis tenella* was infested by *Hysteroneura setariae* aphids. The aphids were brown to red brown. Small colonies  
246 formed on flowers near the seeds, with groups of aphids surrounding the plant's seeds. The aphids of *H. setariae* were  
247 consistently associated with ants. *Euphorbia hirta* or wart grass was colonized by *Aphis gossypii*. The aphids formed  
248 colonies on the undersides of leaves, resulting in stunted growth of the leaves. The aphids were yellow to dark green in  
249 color. *A. gossypii* colonies on *E. hirta* plants were consistently associated with ants. *Eupatorium odoratum* was colonized  
250 by both *Aphis gossypii* and *Aphis citricola*. *A. gossypii* formed colonies in the buds, young leaves, old leaves, and young  
251 twigs. Young leaves that were colonized by *A. gossypii* became stunted with an irregular shape. *A. gossypii* found in this  
252 plant showed yellow-green to dark green in body colour. The colonies of *A. citricola* formed on the young twigs near the  
253 shoots, with these aphids displaying yellow-green coloration and having black siphunculi and cauda. Aphid colonies of  
254 both *A. gossypii* and *A. citricola* on *E. odoratum* plants were associated with either black or red ants.

255 *Hymenochera acutigluma* or hair axis was colonized by *Hysteroneura setariae*, which formed colonies on the flower  
256 stalks and flowers. The colonized parts of the plants did not display any noticeable symptoms. *Lagerstromia* sp. or  
257 *kenidai*, was infested by *Greenidae* sp. These aphids had bright green bodies and distinctive elongated siphunculi with  
258 thorns. The aphids formed colonies on the undersides of leaves, especially on young leaves. The colonized leaves did not  
259 show any disease symptoms. *Lophatherum gracile* or bamboo grass plants were colonized by two species of aphids:  
260 *hysteroneura setariae* and *Rhopalosiphum maidis*. The aphids of *H. setariae* formed colonies on the undersides of leaves,  
261 leaf shoots, and leaf axils. The colonized leaves did not show any disease symptoms. *H. setariae* aphids were brown to  
262 red-brown. *R. maidis* aphids also formed colonies on the undersides of leaves, but the colonies were small. *R. maidis*  
263 aphids were green to bright green in color, with black siphunculi and cauda. It was possible for colonies of the two species  
264 of aphids on *L. gracile* to mix.

265 *Melastoma affine* was colonized by *Aphis gossypii*. The colonies formed on shoots, particularly near newly emerging  
266 shoots and on newly emerging fruits and flowers. The body colour of aphids ranged from yellow to green. The colonized  
267 plant parts did not show any disease symptoms. *Mikania miranta* was colonized by *Aphis gossypii* and *Aphis glycine*. *A.*  
268 *gossypii* formed colonies on the shoots, especially on the undersides of the leaves, resulting in stunted and curled leaves. *A.*  
269 *glycine* formed colonies on the branches. The colonies were densely populated. *A. Glycine* aphids were light green to green  
270 in color. The colonized plant parts became distorted. The two species of aphids could mix to form a single colony.

271 *Mimosa invisa* (cater-grass) was colonized by *Aphis craccivora*. The aphids of *A. craccivora* on *M. invisa* plants  
272 formed colonies only on the shoots with small colonies. The aphids appeared dark black with wingless imagoes. *Mimosa*  
273 *pudica* was observed to be colonized by *Aphis craccivora*. The aphids formed colonies on shoots, especially young shoots,  
274 and occasionally on flowers and pods. The aphids were black and of medium size, resulting in stunted growth of the  
275 colonized plant parts. The colonies were quite large. *Mimosa vigra* was colonized by *Aphis craccivora*. The colonies of  
276 aphids occupied the pods and shoots with small colonies. The nymphs of aphids were black, and wingless adults were  
277 shiny black. The colonized plant parts did not show any disease symptoms.

278 *Oryza rufipogon* was colonized by two species of aphids: *Rhopalosiphum rice* and *Rhopalosiphum maidis*. Both aphids  
279 colonized the same plant parts, namely the unopened leaves and the leaf axils with large colonies. The two species could  
280 be distinguished by their body color. *R. maidis* appeared green with black siphunculi and cauda, while *R. rice*  
281 appeared white. The colonies of *R. maidis* and *R. rice* in *O. rufipogon* plants were associated with the presence of red ants.  
282 *Oxonopus compressus* or pait grass was colonized by *Hysteroneura setariae* aphids. The colonies occupied flowers, flower  
283 stalks, seeds, and sometimes in the leaf axils. The aphids were brown to dark brown in color. Small colonies were formed,  
284 and they were also consistently associated with ants.

285 *Paspalum conjugatum* was colonized by *H. setariae* aphids. The colonies occupied flower parts, especially the seeds  
286 and flower stalks. Aphids had brown to dark brown bodies. *Phyllanthus niruri* was colonized by *Aphis citricola*. The  
287 colonies formed on the shoots and the undersides of leaves and petioles. The colonized parts became distorted, stunted,  
288 and wrinkled. The aphids had yellow bodies with black siphunculi and cauda, and the colonies formed were quite large.  
289 *Portulaca oleraceae* plants were colonized by *Aphis craccivora*. The aphids of *A. craccivora* in *P. oleraceae* plants  
290 formed colonies on the undersides of leaves, especially young leaves, shoots and in flowers. The colonized plant parts  
291 became stunted, and leaf edges curled downward. The aphids had dark brown to black bodies, with wingless imagoes that  
292 appeared glossy black.

293 *Physalis angulata* plants were colonized by *Aphis craccivora*. The aphids had dark green to black bodies, with glossy  
294 black wingless imagoes. *A. craccivora* formed colonies on the shoots or near the leaf buds. The colonized plant parts did  
295 not show any symptoms of disease. *Rorippa indica* or mustard land was colonized by *Lipaphis erysimi*. The colonies  
296 formed on the flowers, fruits, flower stalks and the lower surface of leaves. The colonized plant parts showed symptoms

such as curling and stunting. *Sida rhombifolia* or cacabean was colonized by *Aphis gossypii*. The aphids had green-yellow to green body colors. The colonies formed on the surface of lower leaves, stalks and flower petals. The colonized plant parts, especially the shoots, showed curling, and the leaf edges curled downward. *Sonchus arvensis* plants were colonized by *L. erysimi*. The aphids had green to whitish green body colours, and the colonies formed on flower stalks, under petals, and on young shoots or leaves. The colonized plant parts became stunted over time.

In general, aphids observed on ornamental and wild plants formed colonies. The colonized plant parts typically displayed typical symptoms of damage, but some did not show any symptoms. Generally, the symptoms of the plants caused by aphid colonies were relatively the same, such as stunted growth, abnormal shape, and stunted or curly leaves. These characteristic symptoms serve as indicators of aphid infestations. However, some plants or plant parts did not show symptoms when colonized by aphids. This condition happened because the colonized parts had reached their maximum growth or development. It indicated that the colonized part was not currently undergoing a growth phase. The colonies that did not induce symptoms typically occurred when the colonized leaves had reached their maximum growth or when the leaves and plant parts were old. The old leaves or twigs might not show the typical symptoms associated with aphid infestations.

The part of the plant exhibiting characteristic symptoms when colonized by aphids also often experienced a cessation in growth due to the piercing by the aphids. In contrast, the areas surrounding the puncture site continued to grow, resulting in some parts developing normally while others become stunted (Pettersson, Tjallingii, and Hardie 2017). This condition could lead to the bending of shoots or young stems, curling of leaves, downward curling of leaf edges, or stunted leaf growth. In this observation, monocot plants or groups of grasses with narrow leaves generally did not display any distinctive symptoms when colonized by aphids. This might be because the growth or development of their leaves differed from that of dicot plants. Therefore, the presence of aphids in monocot plants or plants was often easier to recognize through the presence of ants. If a plant was found to have a significant number of ants, there was a possibility that aphids had colonized the plant (Tegelaar et al. 2012). Therefore, the presence of ants could serve as an indicator of the presence of aphid colonies.

Throughout their life cycle, aphids exhibited host alternation by switching between two distinct host plants (Yamamoto et al. 2020). They overwintered on woody plants, reproduced in the spring, and migrated to herbaceous plants during the summer before returning to their primary host in the autumn (Yamamoto et al. 2020). This allowed aphids to maximize resource utilization, avoid congestion and competition, evade predators and parasites, circumvent plant defenses, and colonize new areas. Aphids could distribute their population efficiently, thereby avoiding overcrowding, predators and parasites, and plant defenses developing over time through host switching (Yamamoto et al. 2020). This behavior was essential for the survival and environmental adaptation of aphids.

Aphids colonized flowers because they may offer an accessible and rich food source, sugary plant sap found in new growth or reproductive parts of plants. Flowers contain a nutrient-rich nature and easy access to sap, therefore aphids were attractive to sap the flowers. Some aphid species were drawn to certain colors (Jakubczyk et al. 2022). Herbs served as an alternative host for aphids in this present study. Aphids consume sugar-rich liquid in plants, known as "sap". Aphids considered herbs and other green vegetation as abundant food sources. Aphids utilize needle-like mouthparts to penetrate plant tissues and access this fluid (Brożek et al. 2015). Several aphids colonized herbs such as Indian mustards, *Lipaphis erysimi*, and *Myzus persicae* are the most devastating insects, infesting leaves, stems, and floral parts (Jayaswal et al. 2022).

Due to a symbiotic relationship, the prevalence of aphids and ants was frequently correlated. Aphids produced a delicious substance known as honeydew as a waste product, which ants found highly attractive as a food source (Nelson and Mooney 2022). The honeydew contained an abundance of sugars, extracted by aphids from the plant juice (Zheng et al. 2022). Ants were drawn to this nutrient-rich food source and would often 'farm' aphids for it. In exchange for honeydew, ants provided aphids with protection from other insects and predators, such as ladybugs, lacewing larvae, and parasitic wasps (Karami-jamouri et al. 2018). Certain species of ants would transport aphids to new host plants for improved foraging opportunities, ensuring that aphids had a continuous food source (Giannetti et al. 2021). Honeydew not only nourished the ant colony, but its high sugar content also supported the development of their fungus farms (in certain species) and provided energy for the growth of their own progeny (Biedermann and Vega 2020).

## CONCLUSION

15 species of aphids were found in ornamental and wild plants in Pagaralam, namely *Aphis gossypii*, *Uroleucon* sp., *Toxoptera odinae*, *Macrosiphum rosae*, *Aphis citricola*, *Aphis craccivora*, *Toxoptera aurantii*, *Pentalonia nigronervosa*, *Hystenura* sp., *Aphis glycine*, *Greenidea* sp., *Rhopalosiphum padi*, *Rhopalosiphum maidis*, *Hyperomyzus* sp. *Lipaphis erysimi*.

## ACKNOWLEDGMENTS

The authors thank Universitas Sriwijaya, that supported this research.

## REFERENCES

- Alotaibi, N. J., T. Alsufyani, N. H. M'sakni, M. A. Almalki, E. M. Alghamdi, and D. Spiteller. 2023. "Rapid Identification of Aphid Species by Headspace GC-MS and Discriminant Analysis." *Insects* 14(589). doi: <https://doi.org/10.3390/insects14070589>.
- Bass, Chris, Alin M. Puinean, Christoph T. Zimmer, Ian Denholm, Linda M. Field, Stephen P. Foster, Oliver Gutbrod, Ralf Nauen, Russell Slater, and Martin S. Williamson. 2014. "The Evolution of Insecticide Resistance in the Peach Potato Aphid, *Myzus persicae*." *Insect Biochemistry and Molecular Biology* 51:41–51. doi: [10.1016/j.ibmb.2014.05.003](https://doi.org/10.1016/j.ibmb.2014.05.003).
- Biedermann, Peter H. W., and Fernando E. Vega. 2020. "Ecology and Evolution of Insect–Fungus Mutualisms." *Annual Review of Entomology* 65:431–55. doi: <https://doi.org/10.1146/annurev-ento-011019-024910>.
- Blackman, Roger L., and Victor F. Eastop. 2008. *Aphids on the World's Herbaceous Plants and Shrubs, 2 Volume Set*. John Wiley & Sons.
- Blackman, Roger L., and Victor F. Eastop. 2017. "Taxonomic Issues." Pp. 1–36 in *Aphids as crop pests*. CABI Wallingford UK.
- Braham, Mohamed, Synda Boulahia-kheder, Mouna Kahia, and Siwar Nouri. 2023. "Aphids and Citrus Responses to Nitrogen Fertilization." *Journal of the Saudi Society of Agricultural Sciences* 22(6):374–83. doi: [10.1016/j.jssas.2023.03.003](https://doi.org/10.1016/j.jssas.2023.03.003).
- Brożek, Jolanta, Ewa Mróz, Dominika Wylężek, Łukasz Depa, and Piotr Węgierek. 2015. "The Structure of Extremely Long Mouthparts in the Aphid Genus *Stomaphis* Walker (Hemiptera: Sternorrhyncha: Aphididae)." *Zoomorphology* 134:431–45. doi: <https://doi.org/10.1007/s00435-015-0266-7>.
- Cao, He-he, Zhan-feng Zhang, Xiao-feng Wang, and Tong-xian Liu. 2018. "Nutrition versus Defense: Why *Myzus persicae* (Green Peach Aphid) Prefers and Performs Better on Young Leaves of Cabbage." *PloS One* 13(4):1–16. doi: <https://doi.org/10.1371/journal.pone.0196219>.
- Chandel, R. S., V. K. Chandra, K. S. Verma, and Mandeep Pathania. 2022. "Chapter 21 - Insect Pests of Potato in India: Biology and Management." Pp. 371–400 in, edited by A. Alyokhin, S. I. Rondon, and Y. B. T.-I. P. of P. (Second E. Gao. Academic Press.
- Clarke, Rebecca, Monica A. Kehoe, Sonya Broughton, and Roger A. C. Jones. 2020. "Host Plant Affiliations of Aphid Vector Species Found in a Remote Tropical Environment." *Virus Research* 281(December 2019):197934. doi: [10.1016/j.virusres.2020.197934](https://doi.org/10.1016/j.virusres.2020.197934).
- Döring, Thomas F. 2014. "How Aphids Find Their Host Plants, and How They Don't." *Annals of Applied Biology* 165(1):3–26. doi: <https://doi.org/10.1111/aab.12142>.
- Ertunc, Filiz. 2020. "Chapter 46 - Emerging Plant Viruses." Pp. 1041–62 in, edited by M. M. B. T.-E. and R. V. P. Ennaji. Academic Press.
- Gadhavi, Kiran R., Saurabh Gautam, David A. Rasmussen, and Rajagopalbabu Srinivasan. 2020. "Aphid Transmission of Potyvirus: The Largest Plant-Infecting RNA Virus Genus." *Viruses* 12(7):773. doi: [10.3390/v12070773](https://doi.org/10.3390/v12070773).
- Giannetti, Daniele, Mauro Mandrioli, Enrico Schifani, Cristina Castracani, Fiorenza A. Spotti, Alessandra Mori, and Donato A. Grasso. 2021. "First Report on the Acrobasis Ant Crematogaster Scutellaris Storing Live Aphids in Its Oak-Gall Nests." *Insects* 12(2):108. doi: <https://doi.org/10.3390/insects12020108>.
- Guo, Huijuan, Liyuan Gu, Fanqi Liu, Fajun Chen, Feng Ge, and Yucheng Sun. 2019. "Aphid-Borne Viral Spread Is Enhanced by Virus-Induced Accumulation of Plant Reactive Oxygen Species 1." *Plant Physiology* 179(January):143–55. doi: [10.1104/pp.18.00437](https://doi.org/10.1104/pp.18.00437).
- Jakubczyk, Karolina, Klaudia Koprowska, Aleksandra Gottschling, and Katarzyna Janda-Milczarek. 2022. "Edible Flowers as a Source of Dietary Fibre (Total, Insoluble and Soluble) as a Potential Athlete's Dietary Supplement." *Nutrients* 14(12). doi: [10.3390/nu14122470](https://doi.org/10.3390/nu14122470).
- Jaouannet, Maëlle, Patricia A. Rodriguez, Peter Thorpe, Camille J. G. Lenoir, and Ruari Macleod. 2014. "Plant Immunity in Plant – Aphid Interactions." *Front Plant Sci.* 5(December):1–10. doi: [10.3389/fpls.2014.00663](https://doi.org/10.3389/fpls.2014.00663).
- Jayaswal, Deepanshu, Pawan Mainkar, Kuldeep Kumar, Yamini Agarwal, and Ratna Prabha. 2022. "Pyramiding and Evaluation of Segregating Lines Containing Lectin and Protease Inhibitor Genes for Aphid Resistance in Brassica Juncea." *Indian Journal of Biochemistry & Biophysics* 59(August):800–807. doi: [10.56042/ijbb.v59i8.62319](https://doi.org/10.56042/ijbb.v59i8.62319).
- Jones, Roger A. C. 2022. "Alteration of Plant Species Mixtures by Virus Infection: Managed Pastures the Forgotten Dimension." *Plant Pathology* 71(6):1255–81. doi: [10.1111/ppa.13571](https://doi.org/10.1111/ppa.13571).
- Jousset, Emmanuelle, Genson Gwenaëlle, and Coeur D. Acie. Armelle. 2010. "Evolutionary Lability of a Complex Life Cycle in the Aphid Genus *Brachycaudus*." *BMC Evolutionary Biology* 10(1). doi: [10.1186/1471-2148-10-295](https://doi.org/10.1186/1471-2148-10-295).
- Kallas, John. 2010. *Edible Wild Plants*. Gibbs Smith.
- Karami-jamou, Tahereh, Alinaghi Mirmoayedi, Abbasali Zamani, and Yadolah Khajehzadeh. 2018. "The Impact of Ant Attendance on Protecting Aphis Gossypii against Two Aphidophagous Predators and Its Role on the Intraguild Predation between Them." *Journal of Insect Behavior* 31:222–39. doi: [10.1007/s10905-018-9670-4](https://doi.org/10.1007/s10905-018-9670-4).
- Kennedy, J. S., and H. L. G. Stroyan. 1959. "Biology of Aphids." *Annual Review of Entomology* 4(1):139–60.
- Kinley, Chimi, A. Najitha Banu, A. M. Raut, Johnson Wahengbam, and Thinley Jamtsho. 2021. "A Review on Past, Present and Future Approaches for Aphids Management." *Journal of Entomological Research* 45(2):336–46. doi: [10.5958/0974-4576.2021.00053.0](https://doi.org/10.5958/0974-4576.2021.00053.0).
- Kumar, Sarwan. 2019. "Aphid-Plant Interactions: Implications for Pest Management." P. Ch. 7 in *Plant Communities and Their Environment*, edited by M. T. Oliveira, F. Candan, and A. Fernandes-Silva. Rijeka: IntechOpen.
- Kumar, Sushil, Malay K. Bhowmick, and Puja Ray. 2021. "Weeds as Alternate and Alternative Hosts of Crop Pests." *Indian Journal of Weed Science* 53(1):14–29. doi: [10.5958/0974-8164.2021.00002.2](https://doi.org/10.5958/0974-8164.2021.00002.2).
- Liu, Xiang Dong, Ting Ting Xu, and Hai Xia Lei. 2017. "Refuges and Host Shift Pathways of Host-Specialized Aphids Aphis Gossypii." *Scientific Reports* 7(1):1–9. doi: [10.1038/s41598-017-02248-4](https://doi.org/10.1038/s41598-017-02248-4).
- Maharani, Yani, Purnama Hidayat, Aunu Rauf, and Nina Maryana. 2018. "Short Communication: New Records of Aphid Species Subfamily Aphidinae (Hemiptera: Aphididae) in West Java, Indonesia." *Biodiversitas* 19(2):460–65. doi: [10.13057/biodiv/d190219](https://doi.org/10.13057/biodiv/d190219).
- Margaritopoulos, J. T., M. Tzortzi, K. D. Zarpas, J. A. Tsitsipis, and R. L. Blackman. 2006. "Morphological Discrimination of Aphis Gossypii (Hemiptera: Aphididae) Populations Feeding on Compositae." *Bulletin of Entomological Research* 96(2):153–65. doi: [10.1079/ber.2005410](https://doi.org/10.1079/ber.2005410).
- Meuninck, Jim. 2023. *Basic Illustrated Edible Wild Plants and Useful Herbs*. Rowman & Littlefield.
- Mo, Carmen, and Angela M. Smilanich. 2023. "Feeding on an Exotic Host Plant Enhances Plasma Levels of Phenoloxidase by Modulating Feeding Efficiency in a Specialist Insect Herbivore." *Frontiers in Physiology* 14(February):1–10. doi: [10.3389/fphys.2023.1127670](https://doi.org/10.3389/fphys.2023.1127670).
- Naidu, VSGR. 2012. "Hand Book on Weed Identification."
- Nelson, Annika S., and Kailen A. Mooney. 2022. "The Evolution and Ecology of Interactions between Ants and Honeydew-Producing Hemipteran Insects." *Annual Review of Ecology, Evolution, and Systematics* 53:379–402. doi: <https://doi.org/10.1146/annurev-ecolsys-102220-014840>.
- Petterson, Jan, W. Fred Tjallingii, and Jim Hardie. 2017. "Host-Plant Selection and Feeding." Pp. 173–95 in *Aphids as crop pests*. CABI Wallingford UK.
- Piron, PGM, MC de Haas, and MAHM Sonnemans. 2019. "The Presence of Aphis (Toxoptera) Aurantii (Homoptera: Aphididae) in the Netherlands." *Entomologische Berichten* 79(5):162–64.

420 Santiago, Grazielle Silva, Luana Fonseca, Ernesto Oliveira Canedo-ju, Carla Rodrigues Ribas, Rafaela Pereira, and Guilherme Pereira Alves. 2017.  
421 "Isolated and Community Contexts Produce Distinct Responses by Host Plants to the Presence of Ant-Aphid Interaction : Plant Productivity and  
422 Seed Viability." Plos One 12(1):1–12. doi: 10.1371/journal.pone.0170915.  
423 Singh, Rajendra, and Garima Singh. 2021. "Aphids." Pp. 105–82 in *Polyphagous Pests of Crops*. Springer.  
424 Tegelaar, Karolina, Mattias Hagman, Robert Glinwood, Jan Pettersson, and Olof Leimar. 2012. "Ant–Aphid Mutualism: The Influence of Ants on the  
425 Aphid Summer Cycle." Oikos 121(1):61–66. doi: <https://doi.org/10.1111/j.1600-0706.2011.19387.x>.  
426 Yamamoto, Tetsuya, Mitsuru Hattori, and Takao Itino. 2020. "Seasonal Migration in the Aphid Genus Stomaphis (Hemiptera : Aphididae): Discovery of  
427 Host Alternation Between Woody Plants in Subfamily Lachninae." Journal of Insect Science 20(5):1–10. doi: 10.1093/jisesa/ieaa103.  
428 Zheng, Zhou, Mengqin Zhao, Zhijun Zhang, Xin Hu, Yang Xu, and Cong Wei. 2022. "Lactic Acid Bacteria Are Prevalent in the Infrabuccal Pockets and  
429 Crops of Ants That Prefer Aphid Honeydew." Front. Microbiol. 12(January):1–17. doi: 10.3389/fmicb.2021.785016.  
430

# Species of aphids found in ornamental and wild plants in Pagar Alam District, South Sumatra, Indonesia

CHANDRA IRSAN<sup>1,\*</sup>, ERISE ANGGRAINI<sup>1,2</sup>, WENNY RAMADHANI<sup>3</sup>

<sup>1</sup>Department of Plant Pests and Diseases, Faculty of Agriculture, Universitas Sriwijaya. Kampus Indralaya, Jl. Palembang-Prabumulih, KM 32 Inderalaya Ogan Ilir 30662, South Sumatra, Indonesia. Tel.: +62-711-580059, \*email: chandrairsan@fp.unsri.ac.id

<sup>2</sup>Prograsm Study of Agroecotechnology, Faculty of Agriculture, Universitas Sriwijaya. Kampus Indralaya, Jl. Palembang-Prabumulih, KM 32 Inderalaya Ogan Ilir 30662, South Sumatra, Indonesia

<sup>3</sup>Plant Quarantine. Jl. Kol. H. Barlian Km. 6 No. 78, Srijaya, Alang-Alang Lebar, Palembang 30153, South Sumatera, Indonesia

Manuscript received: 8 September 2023. Revision accepted: xxx December 2023.

**Abstract.** Irsan C, Anggraini E, Ramadhani W. 2023. Species of aphids found in ornamental and wild plants in Pagar Alam District, South Sumatra, Indonesia. Biodiversitas 24: xxx-xxx. Aphids are one of the crucial pests in tropical and sub-tropical regions. The presence of aphids in a plant can be very detrimental due to their role as vectors. Aphids exhibit species diversity, but not much information has been reported about the species diversity of aphids associated with essential crops such as ornamental plants. Furthermore, many aphid species, such as wild plants, were found on plants that were not hosts. Therefore, this study reported the species of aphids found in ornamental and wild plants. The field research employed purposive and direct observation to inventory cultivated or wild plants hosting and collecting aphids. The plant selection process included cultivated plants encompassing ornamental plants and wild plants or weeds. The collection and identification of host plants and aphids involved systematic searches for the selected plants and subsequent examination for the presence of aphids. Observations were made to all existing plant species to find those colonized by aphids. This study revealed that the total of 15 species aphids found in Ornamental plants, *Aphis craccivora* Koch, 1854, *Aphis spiraecola* Patch, 1914, *Aphis glycines* Matsumura, 1917, *Aphis gossypii* Glover, 1877, *Aulacorthum solani* Kaltenbach, 1843, *Macrosiphoniella sanborni* Gillette, 1908, *Macrosiphum rosae* Linnaeus, 1758, *Myzus persicae* Sulzer, 1776, *Neomyzus circumflexus* Buckton, 1876, *Pentalonia caladii* van der Goot, 1917, *Rhopalosiphum nymphaeae* Linnaeus, 1761, *Sinemegoura citricola* van der Goot, 1917, *Toxoptera aurantii* Boyer de Fonscolombe, 1841, *Toxoptera citricidus* Kirkaldy, 1907, *Toxoptera odinae* van der Goot, 1917 and the total of 11 species aphids found in weeds, *A. gossypii*, *A. craccivora*, *A. glycines*, *A. citricola*, *Greenidea* sp., *Hystroeneura setariae* Thomas, 1878, *Hiperomyzus* sp., *Lipaphis erysimi* Kaltenbach, 1843, *Rhopalosiphum maidis* Fitch, 1856, *Rhopalosiphum padi* Linnaeus, 1758, *Schizaphis rotundiventris* Signoret, 1860.

**Keywords:** Aphids, ornamental plants, wild plants

## INTRODUCTION

Aphids are crucial pests in the tropics and sub-tropics, exhibiting various polyphagous, oligophagous, and monophagous characteristics (Kennedy and Stroyan 1959). One species of aphids can host more than 400 species from 40 families (Bass et al. 2014). In addition to pests, aphids can also be vectors of plant viral diseases (Gadhawe et al. 2020); aphids can transmit 275 viruses (Ertunc 2020). In tropical areas, aphids can be found throughout the year due to their parthenogenetic nature of reproduction (Blackman and Eastop 2017). Aphids suck phloem sap of tender plant parts, which can deplete essential nutrients for healthy growth (Cao et al. 2018). Moreover, vector species can further weaken and stunt the growth of infected plants (Jones 2022). In addition, the honeydew that aphids secrete can lead to the growth of sooty mold, a black fungus that can prevent sunlight from reaching the plant's leaves, thereby impairing photosynthesis, the process by which

plants produce food (Singh and Singh 2021). Therefore, it is crucial to control aphid populations in gardens and crops.

Many aphid species are found on plants that are not their actual hosts (Maharani et al. 2018). Aphids have one or more secondary, or alternative, host plants in addition to their primary host plants, which are the types of plants they feed on most frequently (Clarke et al. 2020). Alternative plants provide a means of survival when primary hosts are unavailable, during certain seasons, or under certain environmental conditions (Kumar et al. 2021). These secondary hosts may offer less adequate nutrition for insects (Mo and Smilanich 2023). However, they may provide a means of survival when primary hosts are unavailable, during certain seasons, or under certain environmental conditions (Kumar et al., 2021). According to Liu et al. (2017), hibiscus serves as an overwintering host for cotton-specialized aphids but not for cucurbit-specialized aphids, it is evident that host-specialized aphids



have refuges during times of food shortage. The life cycles of numerous aphid species exhibit such complexity (Jousselin et al. 2010). They maintain a cycle of host alternation, shifting between their primary hosts (typically woody plants) and secondary hosts (often herbaceous plants) (Yamamoto et al. 2020). Weeds pose a continuous threat in both cropped and non-crop areas, providing food, shelter, and reproductive sites for various pest organisms (Kumar et al., 2021). This indicates that weeds can serve as alternative hosts for aphids.

In South Sumatra, particularly in the highland areas like Pagar Alam, there are numerous ornamental and wild plants. Research on the inventory of aphid species in ornamental and wild plants has less noticed. This study reports the diversity of aphid species found in ornamental and wild plants found in this area. The findings from this study can serve as a valuable resource for aphid management.

## MATERIALS AND METHODS

The field research employed a purposive and direct observation to inventory cultivated or wild plants hosting and collecting aphids. The plant selection included cultivated plants encompassing ornamental plants, as well as wild plants or weeds. Where available, collecting and identifying host plants, aphids, and natural enemies involved systematic searches of all existing plant species to find those colonized by aphids. Any plants colonized by aphids are documented as aphid hosts. Aphid identification was done using identification keys (Blackman and Eastop 2008) in the Laboratory of Entomology, Faculty of Agriculture, Universitas Sriwijaya. Identification relied on morphological characteristics. The host plants were identified using the weed identification handbook (Kallas

2010; Naidu 2012; Meuninck 2023). The location and aphid colony sizes, including their life color, and photographs of the aphid colonies and their host plants were recorded.

## RESULTS AND DISCUSSION

### Result

#### **Aphids infesting in ornamental plants**


The results showed that 15 aphid species were found in Pagar Alam (Tables 1 and 2). These aphids mostly colonized flowers of various ornamental plants (Table 1, Figure 1).

The relationship between aphids and ants was also recorded. Aphids produce a sweet, sticky substance called honeydew; ants are attracted to this honey because it serves as a food source. When aphids are present, they secrete honeydew, which attracts ants. This research recorded the presence of ants on plant parts colonized by aphids (Table 2).

#### **Aphids infesting in wild plants (weed or non-weed plants)**

In addition, this study documented aphid colonies on flowers, stalks, plant tops, young leaves, and old leaves of wild plants (Table 3, Figure 2).

The presence of ants in aphid colonization symbolizes a mutually beneficial relationship where the ants receive food from the aphids while protecting them. This study recorded the ant attendance in aphids colonization (Table 4).



**Figure 1.** Photos showing colonies of different aphid species in ornamental plants: A. *Aphis gossypii* in *Dahlia* sp. flower; B. *Aphis gossypii* in *Hibiscus rosinensis* flower; C. *Aphis gossypii* in *Cestrum* twig and flower; D. *Aphis craccivora* in *Clitoria ternatea* flower; E. *Aphis glycines* in *Helianthus giganteus* flower; F. *Aphis craccivora* on the *Murayya paniculata* flower; G. *Toxoptera odinae* in the *Mussaenda frondosa*; H. *Macrosiphoniella sanborni*. in *Chrysanthemum* sp. Leaves; I. *Macrosiphum rosae* in *Rosa indica* flower; J. *Rhopalosiphum nymphaeae* in *Canna indica* leaves. Chandra Irsan captured all the photos.

**Table 1.** Aphid species recorded in ornamental plants and their colony locations

Host plant	Aphid species	Colony location
<i>Aster alpinus</i>	<i>Macrosiphoniella sanborni</i>	Leaves, young twig, flower
<i>Brugmansia suaveolens</i>	<i>Aulacorthum solani</i>	Leaves, flower
	<i>Neomyzus circumflexus</i>	Leaves
	<i>Myzus persicae</i>	Leaves, flower
<i>Caladium</i> sp.	<i>Pentalonia caladii</i>	Leaves,
<i>Cananga odoratum</i>	<i>Aphis gossypii</i>	Leaves, flower
<i>Canna indica</i>	<i>Rhopalosiphum nymphaeae</i>	Leaf
<i>Catharanthus roseus</i>	<i>Aphis spiraeicola</i>	Shoot, young leaves, flower


<i>Cestrum</i> sp.	<i>Aphis gossypii</i>	Shoot, flower
	<i>Neomyzus circumflexus</i>	Young leaves
<i>Clitoria ternatea</i>	<i>Aphis craccivora</i>	Flower
<i>Chrysanthemum</i> sp.	<i>Macrosiphoniella sanborni</i>	Shoot, twig
<i>Dahlia</i> sp.	<i>Aphis gossypii</i>	Flower
<i>Dendrobium</i> sp.	<i>Sinemogoura citricola</i>	Flower
<i>Duranta</i> sp.	<i>Aphis gossypii</i>	Shoot, flower
<i>Helianthus giganteus</i> .	<i>Aphis glycines</i>	Flower
<i>Hibiscus rosasinensis</i>	<i>Aphis gossypii</i>	Flower
<i>Ixora paludosa</i>	<i>Aphis gossypii</i> ,	Flower
	<i>Toxoptera aurantii</i>	Shoot, young leaves
<i>Ixora</i> sp.	<i>Aphis spiraecola</i>	Flower
	<i>Aphis gossypii</i>	Flower
	<i>Toxoptera aurantii</i>	Shoot, flower
<i>Murraya paniculata</i>	<i>Aphis craccivora</i>	Young Twig
	<i>Toxoptera citricidus</i>	Shoot, flower
<i>Mussaenda frondosa</i>	<i>Aphis spiraecola</i>	Shoot, flower
	<i>Toxoptera odinae</i>	Shoot, flower
<i>Rosa indica</i>	<i>Macrosiphum rosae</i>	Flower
<i>Spondias dulcis</i>	<i>Aphis spiraecola</i>	Flower

**Table 2.** Aphid species recorded in ornamental plants and the presence of the ants in the plant parts colonized

Aphid species	Ornamental plants	Aphids life color	Plant parts colonized	Ant attendance	Total individual of ant
<i>Aphis craccivora</i>	<i>Clitoria ternatea</i>	Black	Flowers	+	3
	<i>Murraya paniculata</i>	Black	Flowers	+	2
<i>Aphis spiraecola</i>	<i>Catharanthus roseus</i>	Greenish yellow	Flowers	+	2
	<i>Ixora</i> sp.	greenish yellow	Flowers	+	3
	<i>Mussaenda frondosa</i>	greenish yellow	Shoots, flowers	+	7
	<i>Spondias dulcis</i>	greenish yellow	Flowers	+	8
<i>Aphis glycines</i>	<i>Helianthus giganteus</i>	Greenish yellow	Flowers	+	3
<i>Aphis gossypii</i>	<i>Cestrum</i> sp.	Green	Shoots, flowers	+	4
	<i>Cananga odoratum</i>	Light green	Shoots, flowers	+	1
	<i>Dahlia</i> sp.	Green dark	Flowers	+	2
	<i>Duranta</i> sp.	Light green	Shoots, flowers	+	5
	<i>Hibiscus rosasinensis</i>	Dark green	Flowers	+	6
	<i>Ixora paludosa</i>	Light green	Flowers	+	2
	<i>Ixora</i> sp.	Light green	Flowers	+	7
<i>Aulacorthum solani</i>	<i>Brugmansia suaveolens</i>	Greenish yellow	Leaves, flowers	-	0
<i>Macrosiphoniella sanborni</i>	<i>Aster alpinus</i>	Brown black	Leaves, twigs, flowers	+	5
	<i>Chrysanthemum</i> sp.	Reddish brown	Leaves, twigs	+	5
<i>Macrosiphum rosae</i>	<i>Rosa indica</i>	Green	Flowers	-	0
<i>Myzus persicae</i>	<i>Brugmansia suaveolens</i>	Greenish yellow	Leaves, flowers	-	0
<i>Neomyzus circumflexus</i>	<i>Cestrum</i> sp.	Light green	Young leaves, flowers	-	0
	<i>Brugmansia suaveolens</i>	Light green	Flowers	-	0
<i>Pentalonia caladii</i>	<i>Caladium</i> sp.	Brown-black	Leaves	+	7
<i>Rhopalosiphum nymphaeae</i>	<i>Canna indica</i>	Green black	Leaves	+	1
<i>Sinemogoura citricola</i>	<i>Dendrobium</i> sp.	Brown	Flowers	-	0
<i>Toxoptera aurantii</i>	<i>Ixora paludosa</i>	Brown black	Flowers	+	5
	<i>Ixora</i> sp.	Brown black	Flowers	+	4
<i>Toxoptera citricidus</i>	<i>Murraya paniculata</i>	Black	Stems	+	6
<i>Toxoptera odinae</i>	<i>Mussaenda frondosa</i>	Reddish-brown	Flowers	+	4

Notes: (+) = present; (-) = absent





**Figure 2.** Aphids found infesting wild plants A. *Aphis gossypii* in *Ageratum conyzoides*; B. *Aphis gossypii* in *Croton hirtus*; C. *Aphis gossypii* in *Eupatorium odoratum*; D. *Aphis gossypii* in *Pachystochys* sp.; E. *Pentalonia caladii* in *Caladium* sp.; F. *Aphis gossypii* in *Alternanthera sessilis*; G. *Aphis gossypii* in *Portulaca oleraceae*; H. *Aphis gossypii* in *Euphorbia hirta*; I. *Aphis spiraecola* in *Phyllanthus nerruri*; J. *Aphis spiraecola* in *Sida rhombifolia*; K. *Aphis spiraecola* in *Bridelia tomentosa*; L. *Aphis spiraecola* in *Ludwigia peruviana*; M. *A. craccivora* in *Mimosa pudica*; N. *Aphis craccivora* in *Amaranthus viridis*; O. *Aphis glycine* in *Mikania micrantha*; P. *Hysteneura* sp. in *Eleusine* sp.; Q. *Greenidae* sp. in *Bridelia tomentosa*; R. *Hyperomyzus* sp. in *Echinochloa crusgali*; S. *Lipaphis erysimi* in *sonchus arvensis*; T. *Rhopalosiphum padi* in *Oryza rufipogon*; U. *Rhopalosiphum maidis* in *Oryza rufipogon*. All the photos were captured by Chandra Irsan.

**Table 3.** Species of aphids found in wild (weed or non-weed) plants and their colony locations

Host plant	Weeds or non-weed plants	Aphid species	Colony location
<i>Ageratum conyzoides</i>	Weed	<i>Aphis gossypii</i>	Shoots, young leaves, old leaves, flowers
<i>Alternanthera philoxeroides</i>	Weed	<i>Aphis gossypii</i>	Shoots, buds
<i>Alternanthera sessilis</i>	Weed	<i>Aphis gossypii</i>	Shoots, buds
<i>Amaranthus viridis</i>	Weed	<i>Aphis craccivora</i>	Flowers, shoots, young leaves, old leaves
<i>Blumea lacera</i>	Weed	<i>Lipaphis erysimi</i>	Flowers, shoots, and buds
<i>Bridelia tomentosa</i>	Non-weed	<i>Greenidea</i> sp.	Young leaves
		<i>Aphis spiraeicola</i>	Shoot, young leaves
<i>Croton hirtus</i>	Weed	<i>Aphis gossypii</i>	Flowers, shoots, young leaves, old leaves, young twigs
<i>Cynodon dactylon</i>	Weed	<i>Schizaphis rotundiventris</i>	Flower, flower stalks
<i>Cyperus rotundus</i>	Weed	<i>Schizaphis rotundiventris</i>	Flower, flower stalks, leaf axils
<i>Cyperus compressus</i>	Weed	<i>Schizaphis rotundiventris</i>	Flower, flower stalks, leaf axils
<i>Digitaria ciliaris</i>	Weed	<i>Hystroneura setariae</i>	Flower, flower stalks
<i>Echinocloa crusgali</i>	Weed	<i>Hyperomyzus</i> sp.	Young leaves, old leaves
<i>Ecliptica prostrata</i>	Weed	<i>Aphis gossypii</i>	Shoots, young leaves
<i>Eleusin indica</i>	Weed	<i>Hysteroneura setariae</i>	Flower, flower stalks, leaf axils
		<i>Rhopalosiphum maidis</i>	Flower, flower stalks, leaf axils
<i>Emilia sonchifolia</i>	Weed	<i>Aphis gossypii</i>	Flower, flower stalks, shoots
<i>Eragrostis tenella</i>	Weed	<i>Hysteroneura setariae</i>	Flower, flower stalks, seeds
<i>Euphorbia hirta</i>	Weed	<i>Aphis gossypii</i>	Young leaves, old leaves
<i>Eupatorium odoratum</i>	Weed	<i>Aphis gossypii</i>	Young leaves, old leaves,
		<i>Aphis glycines</i>	Shoot, young twigs
<i>Hymenochera acutigluma</i>	Weed	<i>Hysteroneura setariae</i>	Flowers, flower stalks, leaf axils
<i>Lophatherum gracile</i>	Weed	<i>Hysteroneura setariae</i>	Young leaves, old leaves, leaf axils
		<i>Rhopalosiphum maidis</i>	Young leaves, old leaves, leaf axils
<i>Melastoma affine</i>	Non-weed	<i>Aphis gossypii</i>	Shoots, young leaves
<i>Mikania micrantha</i>	Weed	<i>Aphis gossypii</i>	Shoots, young leaves, old leaves
		<i>Aphis glycines</i>	Shoot, young twig
<i>Mimosa invisa</i>	Weed	<i>Aphis craccivora</i>	Shoots, pods
<i>Mimosa pudica</i>	Weed	<i>Aphis craccivora</i>	Shoots, pods, flowers
<i>Mimosa vigra</i>	Non-weed	<i>Aphis craccivora</i>	Shoots, pods
<i>Oryza rufipogon</i>	Weed	<i>Rhopalosiphum padi</i> ,	Old leaves, young leaves (shoot), leaf axils
		<i>Rhopalosiphum maidis</i>	Old leaves, young leaves (shoot), leaf axils
<i>Oxonopus compressus</i>	Weed	<i>Hysteroneura setariae</i>	Flowers, flower stalks, leaf axils
<i>Paspalum conjugatum</i>	Weed	<i>Hysteroneura setariae</i>	Flowers, flower stalks, seeds
<i>Phyllanthus neruri</i>	Weed	<i>Aphis spiraeicola</i>	Shoot, young leaves, old leaves, young twigs, petioles
<i>Portulaca oleraceae</i>	Weed	<i>Aphis craccivora</i>	Shoots, young leaves, flowers
<i>Physalis angulata</i>	Weed	<i>Aphis craccivora</i>	Shoots, young leaves, old leaves
		<i>Aphis gossypii</i>	Shoots, young leaves, old leaves
<i>Rorippa indica</i>	Weed	<i>Lipaphis erysimi</i>	Flowers, fruits, shoots, young leaves
<i>Sida rhombifolia</i>	Weed	<i>Aphis gossypii</i>	Shoots, young leaves, old leaves, fruit/seeds
<i>Sonchus arvensis</i>	Weed	<i>Lipaphis erysimi</i>	Young leaves, fruit stalks, flowers, fruits

**Table 4.** Aphid species were recorded in wild plants, and the presence of ants in the plant parts colonized

Aphid species	Wild plants	Aphids life color	Plant parts colonized	Ant attendance	Total individual of ant
<i>Aphis gossypii</i>	<i>Ageratum conyzoides</i>	Light green	Shoots, young leaves, old leaves, flowers	+	5
	<i>Alternanthera philoxeroides</i>	Light green	Shoots, buds	+	3
	<i>Alternanthera sessilis</i>	Light green	Shoots, buds	-	0
	<i>Croton hirtus</i>	Dark green	Flowers, shoots, young leaves, old leaves,	+	7
	<i>Ecliptica prostrata</i>	Green	young twigs	+	5
	<i>Emilia sonchifolia</i>	Green	Shoots, young leaves	+	6
	<i>Euphorbia hirta</i>	Light green	Flower, flower stalks, shoots	+	7
	<i>Eupatorium odoratum</i>	Light green	Young leaves, old leaves	+	8
	<i>Melastoma affine</i>	Light green	Young leaves, old leaves, young twigs	+	8
	<i>Mikania micrantha</i>	Light green	Shoots, young leaves	+	9
	<i>Physalis angulata</i>	Yellowish green	Shoots, young leaves, old leaves	+	10
	<i>Sida rhombifolia</i>	Yellowish green	Shoots, young leaves, old leaves, fruit/seeds	-	0
	<i>Amaranthus viridis</i>	Black	Flowers, shoots, young leaves, old leaves	+	3
	<i>Mimosa invisa</i>	Black	Shoots, pods	+	2
<i>Aphis craccivora</i>	<i>Mimosa pudica</i>	Black	Shoots, pods, flowers	+	3
	<i>Mimosa vigra</i>	Black	Shoots, pods	+	4
	<i>Portulaca oleraceae</i>	Black	Shoots, young leaves, flowers	+	7
	<i>Physalis angulata</i>	Black	Shoots, young leaves, old leaves	+	4
<i>Aphis glycines</i>	<i>Eupatorium odoratum</i>	Greenish yellow	Young leaves, old leaves, young twigs	+	6



<i>Aphis spiraecola</i>	<i>Mikania micrantha</i>	Light green	Shoots, young leaves, old leaves	+	4
	<i>Phyllanthus neruri</i>	Greenish yellow	Shoot, young leaves, young twigs, petioles	+	5
<i>Greenidea</i> sp.	<i>Bridelia Tomentosa</i>	Greenish yellow	Shoot, young leaves	+	2
	<i>Bridelia Tomentosa</i>	Greenish yellow	Young leaves	-	0
<i>Hystroneura setariae</i>	<i>Digitaria ciliaris</i>	Reddish-brown	Flower, flower stalks	+	3
	<i>Eleusin indica</i>	Reddish-brown	Flower, flower stalks, leaf axils	+	4
	<i>Eragrostis tenella</i>	Reddish-brown	Flower, flower stalks, seeds	+	4
	<i>Hymenochera acutigluma</i>	Reddish-brown	Flowers, flower stalks, leaf axils	+	3
	<i>Lophatherum gracile</i>	Reddish-brown	Young leaves, old leaves, leaf axils	+	6
	<i>Oxonopus compressus</i>	Reddish-brown	Flower, flower stalk, leaf axils	+	3
	<i>Paspalum conjugatum</i>	Reddish-brown	Flower, flower stalk, seeds	+	6
	<i>Echinocloa crussgali</i>	Black	Young leaves, old leaves	-	0
	<i>Blumea lacera</i>	Whitish green	Flowers, shoots, and buds	+	4
	<i>Rorippa indica</i>	Whitish green	Flower, fruit, shoots, young leaves	+	4
<i>Rhopalosiphum maidis</i>	<i>Sonchus arvensis</i>	Whitish green	Young leaves, fruit stalks, flowers, fruit	+	5
	<i>Eleusin indica</i>	Green	Flower, flower stalks, leaf axils	+	3
	<i>Lophatherum gracile</i>	Green	Young leaves, old leaves, leaf axils	+	4
<i>Rhopalosiphum padi</i>	<i>Oryza rufipogon</i>	Green	Old leaves, young leaves (shoot), leaf axils	-	0
	<i>Oryza rufipogon</i>	Whitish green	Old leaves, young leaves (shoot), leaf axils	+	4
<i>Schizaphis rotundiventris</i>	<i>Cynodon dactylon</i>	Green	Flowers, flower stalks	+	6
	<i>Cyperus compressus</i>	Green	Flowers, flower stalks, leaf axils	+	4
	<i>Cyperus rotundus</i>	Green	Flowers, flower stalks, leaf axils	+	4

Notes: (+): present, (-): absent

## Discussion

In the present study, some aphid species were found on several ornamental plants in Pagar Alam, the location of aphid colonization on the plants varied. On *Aster alpinus* L., aphids were found to form colonies on the stems or young leaf shoots, and the colonies were relatively large. The color of the aphids was dark brown to black. The colonized plant parts showed symptoms of stunting. The identification results showed that the aphids were *Macrosiphoniella sanborni* Gillette, 1908 associated with ants. On the *Brugmansia suaveolens* (Humb. & Bonpl. ex Willd.) Bercht. & J.Presl, *Myzus persicae* Sulzer, 1776 were found on the undersides of old leaves or leaves that have turned yellow. The colonies were relatively small. The aphids found were green and large bodies. The colonized plant parts did not show any signs of disease. On *Caladium* sp. was found one species of aphids: *Pentalonia caladii* van der Goot, 1917. *P. caladii* was known and found in taro plants; the aphids formed colonies under the surface of young and older leaves (Bhadra and Agarwala 2014). This study found that the occupied leaf areas did not display severe symptoms; the aphids were yellow-green to dark green. The wingless adult aphids often had a white, flour-like appearance on their bodies. On the *Cananga odorata* (Lam.) Hook.f. & Thomson (ylang-ylang), colonies of *Toxoptera aurantii* Boyer de Fonscolombe, 1841 were found on the undersides of the leaves, the shoots, buds, and unopened flower petals. The *T. aurantii* colonies found were relatively large. Colonized parts, especially shoots, showed signs of stunting. The aphids found were brown to black. The colonies of *T. aurantii* were found to be associated with black ants. Aphids on *Canna indica* L. (Indian shot, African arrowroot) were found to form colonies in the leaf axils and under the leaf surface near the leaf base. The colonies were quite large. The aphids were

dark brown to dark red coloring with a medium-sized body and the identification results showed that the aphids were *Rhopalosiphum nymphaeae* Linnaeus, 1761 (Ghosh and Singh 2004). The colonies of *R. nymphaeae* were found to be associated with ants. In the *Catharanthus roseus* (L.) G.Don (periwinkle), *Aphis spiraecola* Patch, 1914 aphids were found. The aphids were yellow-green, sifunculi, and black cauda. The aphids formed colonies on flowers and shoots, and the colonized plant parts showed no disease symptoms. On *Cestrum* sp. (Bastard jasmine), aphids formed colonies on the undersides of young leaves, shoots, and within flower parts, especially between petals or stalks that had not fully bloomed; the colonies were quite large. The body color of aphids was green to dark green, with small to medium-sized bodies. The colonized plant parts, especially leaves, showed stunting symptoms. The identification results showed that the aphids were *Aphis gossypii* Glover, 1877. The aphid colonies found were consistently associated with ants. Aphids on *Clitoria ternatea* L. were found to form colonies on flower parts, flower crowns, stems, and young leaves. The aphids were brown to black. Colonized plant parts, especially shoots and young leaves, showed stunting symptoms. The identification results showed that the aphids were *Aphis craccivora* Koch, 1854. These colonies were consistently associated with ants. The aphids on the *Dahlia* sp. formed colonies on unopened flower buds, with a significant population among the blooming petals. The body color was green to dark green. The identification results showed that the aphids were *A. gossypii*. According to this present study, *Sinemegoura citricola* van der Goot, 1917 colonies were found on the young leaves of *Dendrobium* sp., with the color body of the *S. citricola* aphids were yellow, green to dark green, and the colonized plants showing no disease

symptoms, and were associated with ants. On *Duranta* sp., colonies of aphids on the undersides of young leaves, and the colonized plant parts showed stunting symptoms. The colonies were very large. The aphids were green. The identification results showed that the aphids were *A. gossypii*. The aphid colonies were consistently associated with ants. Furthermore, on the *Helianthus annuus* L., aphid colonies were found between the flower petals. The colonized flowers, especially the crowns, tended to fall off easily. The aphids were green and yellow in color. The colonies were small. The identification results showed that the aphids were *A. gossypii*. These aphid colonies were associated with ants. Aphid colonies on *Helianthus* sp. were found on the undersides of old leaves. These colonies were small in size. The aphids were green with a medium body size. The colonized plant parts did not show any disease symptoms. The identification results showed that the aphids were *Aphis glycines* Matsumura, 1917. The aphid colonies were not associated with ants. Within the colonies, mummified aphids that Aphidiidae parasitized were found. On the *Hibiscus rosa-sinensis* L., aphids ranging from yellow to dark green were found. The aphids formed colonies on flower buds, unopened flower crowns, and the undersides of aging leaves. The colonies grew to be very large. The identification results showed that the aphids were *A. gossypii*. The aphid colonies were consistently associated with ants. Two types of aphids were found on the flowering plant *Ixora paludosa* (Blume) Kurz. First, the aphids formed colonies on the undersides of young leaves that were still red or light green and sometimes on flower stalks that had not yet bloomed. The occupied plant parts showed symptoms such as stunted leaf growth, leaf shrinkage, necrotic spots on the leaf surface, and slightly downward-curved leaf edges. The upper leaf surface looked wet and sticky, like sugar. The aphids had yellow, green, or slightly dark green bodies, with some wingless adults having a powdery white upper surface. The identification results showed that the aphids were *A. gossypii* almost always associated with ants. The second type of aphids on *Ixora paludosa* formed colonies under the surface of young and older leaves. The colonies could also be found on newly emerging flowers and leaves. The plant parts occupied by these aphids did not show obvious signs of illness. These aphids were dark red to black, with once-branched stigma and venation in their black wings. The identification results showed that the aphids were *T. aurantii*. These aphids were also associated with ants. Moreover, two forms of aphids were discovered in *Ixora* sp. flower plants. These aphids occupied the shoots, young leaves, and unopened flowers; the affected plant parts did not show obvious symptoms. The aphids exhibited colors ranging from yellow and green to a slightly darker green.

Sometimes, the upper surface of the wingless imago's body appeared white, resembling flour. The identification results showed that these aphids were *A. gossypii*. These aphid colonies were almost always associated with ants. Another species of aphids formed colonies on flower stalks that had not yet bloomed and on newly emerging shoots or leaves. The presence of these aphids on the plant did not induce plant disease symptoms. The aphids were yellow or yellowish green, with black cauda and siphunculi. Their bodies were very small to small. The identification results showed that the aphids were *A. citricola*. The colonies of *A. citricola* were also frequently found in association with ants. Two types of aphids were found on *Mussaenda frondosa* L., each forming colonies in different locations. The first type formed colonies on young leaves, shoots, and flowers. The plant parts they occupied showed no obvious disease symptoms. The identification results showed that the aphids were *Toxoptera odinae* van der Goot, 1917. The aphids were yellow, green, and dark green (Blackman et al. 2011). The second type of aphids formed colonies on the stems or young twigs, appearing densely clustered as if piled up. The aphid colonies could also be found on young leaves, shoots, and within flower parts. The plant parts they infested showed no signs of diseases. The aphids were yellow or yellow-green, with black cauda and siphunculi. They had tiny to small bodies. The identification results showed that the aphids were *A. citricola*. Many aphid species infest various ornamental plants because these insects are attracted to such plants due to the rich nutrient content in the plant sap (Braham et al. 2023).

The results showed that 34 species of wild plants, including weeds, were growing in the yard colonized by aphids. This indicated that multiple species of aphids colonized various host plants. The aphid species colonizing these plants were generally consistent within the same taxon. *Ageratum conyzoides* L. was infested by *A. gossypii*. These aphids formed colonies on the flower sections, shoots, lower surfaces of young leaves, or leaves turning yellow. The aphids were green, yellow-green to dark green, often forming large colonies. *Alternanthera philoxeroides* (Mart.) Griseb., or alligator grass, was also colonized by *A. gossypii*. Small colonies were found on shoots or stems. These aphids had small bodies and were green, ranging from yellow-green to dark green. *Alternanthera sessilis* (L.) R.Br. ex DC. was colonized by *A. gossypii*, forming colonies on shoots, flowers, and fruit. The colonies were typically large, and often associated with tiny brown ants. *Amaranthus viridis* Linnaeus was infested by *A. craccivora*. These aphids established colonies on shoots, flowers, and young and old leaves. They were dark brown to black, with shiny black wingless imagoes. Colonies of these aphids were associated with both black and red ants.

*Blumea lacera* (Burm.fil.) DC. was colonized by *Lipaphis erysimi* Kaltenbach, 1843. These aphids were bright green and of medium size. The colonies formed on flowers, flower stalks, and the undersides of the leaves at the top. The aphid colonies were not associated with ants. *Croton hirtus* L'Hér., or fire grass, was infested by *A. gossypii*; the aphids were yellow-green to dark green. The colonies were found on the stems, leaves, buds, and flowers, often forming large colonies. *Cynodon dactylon* (L.) Pers. or Bermuda grass was colonized by *Schizaphis rotundiventris* Signoret, 1860. The aphids colonized the flowers, flower stalks, and sometimes the plant leaf axils. Small colonies were formed. The aphids were brown to reddish brown. They were associated with ants. *Cyperus rotundus* L., or nut grass, was infested by *S. rotundiventris* aphids. The colonies were found on flower stalks, flowers, and leaf axils. The colonies were quite large and associated with both black and red ants. The aphids were dark brown in color. *Cyperus compressus* L., or grass puzzle, was colonized by *S. rotundiventris* aphids, forming colonies in the flowers, flower stalks, and sometimes in the axils and leaves of the shoots or buds. Small colonies were observed. *Digitaria ciliaris* (Retz.) Koeler was infested by *Hysteroneura setariae* Thomas 1878 aphids, with small colonies scattered on the flowers and flower stalks. These aphids were light brown to brown in color. *Echinochloa crus-galli* (L.) P.Beauv., or water hyacinth plants, were colonized by *Hiperomyzus* sp. aphids. These aphids were dark brown to black and formed large colonies on the undersides of both young and old leaves. The aphid colonies were never found in association with ants. *Eclipta prostrata* (L.) L., or urang-aring, was colonized by *A. gossypii*, forming small colonies on the shoots and flowers. The aphids were bright green to blackish green. The aphid colonies were also consistently associated with ants. *Eleusin indica* (L.) Gaertn. was colonized by two species of aphids: *Hysteroneura setariae* Thomas, 1878 and *Rhopalosiphum maidis* Fitch, 1856. *H. setariae* formed colonies in flower parts, flower stalks, and leaf axils, resulting in large colonies. *H. setariae*'s body color ranged from red-brown to dark brown. The colonies were consistently associated with ants. The aphids of *R. maidis* formed colonies in the leaf axils and undersides of leaves and leaf shoots that had not yet opened. The colonies were not densely packed. The leaf aphids of *R. maidis* were green in color, with distinct black siphunculi and cauda. These aphids had relatively large bodies with a slightly elongated shape. *R. maidis* colonies were always associated with ants. The plant *Emilia sonchifolia* (L.) DC. ex Wight, characterized by its purple flowers, was colonized by *A. gossypii*; the aphids were yellow to green in color. The colonies formed near flowers, flower stalks, and shoot

leaves. *Eragrostis tenella* was infested by *H. setariae* aphids. The aphids were brown to red-brown. Small colonies formed on flowers near the seeds, with groups of aphids surrounding the plant's seeds. The aphids of *H. setariae* were consistently associated with ants. *Euphorbia hirta* L., or wart grass, was colonized by *A. gossypii*. The aphids formed colonies on the undersides of leaves, resulting in stunted growth of the leaves. The aphids were yellow to dark green in color. *A. gossypii* colonies on *E. hirta* plants were consistently associated with ants. *Eupatorium odoratum* L. was colonized by *A. gossypii* and *A. citricola*. *A. gossypii* formed colonies in the buds, young leaves, old leaves, and young twigs. Young leaves colonized by *A. gossypii* became stunted with an irregular shape. *A. gossypii* found in this plant showed yellow-green to dark-green body color. The colonies of *A. citricola* formed on the young twigs near the shoots, with these aphids displaying yellow-green coloration and having black siphunculi and cauda. Aphid colonies of *A. gossypii* and *A. citricola* on *E. odoratum* plants were associated with either black or red ants. *Hymenachne acutigluma* (Steud.) Gilliland, or hair axis, was colonized by *H. setariae*, which formed colonies on the flower stalks and flowers. The colonized parts of the plants did not display any noticeable symptoms. *Lagerstromia* sp., or *kenidai*, was infested by *Greenidae* sp. These aphids had bright green bodies and distinctive elongated siphunculi with thorns. The aphids formed colonies on the undersides of leaves, especially on young leaves. The colonized leaves did not show any disease symptoms. *Lophatherum gracile* Brongn. or bamboo grass plants, were colonized by two species of aphids: *H. setariae* and *R. maidis*. The aphids of *H. setariae* formed colonies on the undersides of leaves, leaf shoots, and leaf axils. The colonized leaves did not show any disease symptoms. *H. setariae* aphids were brown to red-brown. *R. maidis* aphids also formed colonies on the undersides of leaves, but the colonies were small. *R. maidis* aphids were green to bright green, with black siphunculi and cauda. It was possible for colonies of the two species of aphids on *L. gracile* to mix. In addition, *Melastoma affine* D.Don was colonized by *A. gossypii*. The colonies formed on shoots, particularly near newly emerging shoots and newly emerging fruits and flowers. The body color of aphids ranged from yellow to green. The colonized plant parts did not show any disease symptoms. *Mikania micrantha* Kunth was colonized by *A. gossypii* and *Aphis glycines* Matsumura, 1917. *A. gossypii* formed colonies on the shoots, especially on the undersides of the leaves, resulting in stunted and curled leaves. *A. glycines* formed colonies on the branches. The colonies were densely populated. *A. glycines* aphids were light green to green in color. The colonized plant parts became distorted. The two

species of aphids could mix to form a single colony. *Mimosa invisa* Mart. ex Colla (cater-grass) was colonized by *A. craccivora*. The aphids of *A. craccivora* on *M. invisa* plants formed colonies only on the shoots with small colonies. The aphids appeared dark black with wingless imagoes. *Mimosa pudica* L. was observed to be colonized by *A. craccivora*. The aphids formed colonies on shoots, especially young shoots, and occasionally on flowers and pods. The aphids were black and of medium size, resulting in stunted growth of the colonized plant parts. The colonies were quite large. *Mimosa pigra* L. was colonized by *A. craccivora*. The colonies of aphids occupied the pods and shoots with small colonies. The nymphs of aphids were black, and wingless adults were shiny black. The colonized plant parts did not show any disease symptoms. *Oryza rufipogon* Griff. was colonized by two species of aphids: *Rhopalosiphum padi* and *R. maidis*. Both aphids colonized the same plant parts, namely the unopened leaves and the leaf axils with large colonies. The two species could be distinguished by their body color. *R. maidis* appeared green with black siphunculi and cauda, while *Rhopalosiphum padi* Linnaeus, 1758 appeared white. The colonies of *R. maidis* and *R. padi* in *O. rufipogon* plants were associated with the presence of red ants. *Axonopus compressus* (Sw.) P.Beauv., or *pait* grass, was colonized by *H. setariae* aphids. The colonies occupied flowers, flower stalks, seeds, and sometimes the leaf axils. The aphids were brown to dark brown. Small colonies were formed, and they were also consistently associated with ants. *Paspalum conjugatum* was colonized by *H. setariae* aphids. The colonies occupied flower parts, especially the seeds and flower stalks. Aphids had brown to dark brown bodies. *Phyllanthus neruri* L. was colonized by *A. citricola*. The colonies formed on the shoots and the undersides of leaves and petioles. The colonized parts became distorted, stunted, and wrinkled. The aphids had yellow bodies with black siphunculi and cauda; the colonies formed were large. *Portulaca oleracea* L. plants were colonized by *A. craccivora*. The aphids of *A. craccivora* in *P. oleraceae* plants formed colonies on the undersides of leaves, especially young leaves, shoots, and flowers. The colonized plant parts became stunted, and leaf edges curled downward. The aphids had dark brown to black bodies, with wingless imagoes that appeared glossy black. *Physalis angulata* plants were colonized by *Aphis craccivora*. The aphids had dark green to black bodies, with glossy black wingless imagoes. *A. craccivora* formed colonies on the shoots or near the leaf buds. The colonized plant parts did not show disease symptoms. *Rorippa indica* (L.) Hiern, or mustard land, was colonized by *L. erysimi*. The colonies formed on the flowers, fruits, flower stalks, and the lower leaf's surface. The colonized plant parts showed symptoms

such as curling and stunting. *Sida rhombifolia* L., or cacabea, was colonized by *A. gossypii*. The aphids had green-yellow to green body colors. The colonies formed on the surface of lower leaves, stalks, and flower petals. The colonized plant parts, especially the shoots, showed curling, and the leaf edges curled downward. *Sonchus arvensis* L. plants were colonized by *L. erysimi*. The aphids had green to whitish green body colors, and the colonies formed on flower stalks, under petals, and on young shoots or leaves. The colonized plant parts became stunted over time.

In general, aphids observed on ornamental and wild plants formed colonies. The colonized plant parts typically displayed typical damage symptoms, but some did not show any symptoms at all. Generally, the plants' symptoms due to aphid colonies were relatively the same, such as stunted growth, abnormal shape, and stunted or curly leaves. These characteristic symptoms serve as indicators of aphid infestations. However, some plants or plant parts did not show symptoms when colonized by aphids. This condition occurred because the colonized parts had reached maximum growth or development. It indicated that the colonized part was not currently undergoing a growth phase. The colonies that did not induce symptoms typically occurred when the colonized leaves had reached their maximum growth or when the leaves and plant parts were old. Furthermore, the old leaves or twigs might not show the typical symptoms associated with aphid infestations. The plant parts exhibiting characteristic symptoms when colonized by aphids also often experienced a cessation in growth due to the piercing by the aphids. In contrast, the areas surrounding the puncture site continued growing, resulting in some parts developing ordinary while others became stunted (Pettersson et al. 2017). This condition could lead to bending shoots or young stems, curling leaves, downward curling of leaf edges, or stunted leaf growth. In this observation, monocot plants or groups of grasses with narrow leaves generally did not display any distinctive symptoms when colonized by aphids. This might be because the growth or development of their leaves differed from that of dicot plants. Therefore, the presence of aphids in monocot plants was often easier to recognize through the presence of ants. If a plant was found to have a significant number of ants, there was a possibility that aphids had colonized the plant (Tegelaar et al. 2012). Therefore, the presence of ants could serve as an indicator of the aphid colonies. According to this study, ants were present in some aphid colonies from the subfamily Aphidinae, while the ants were absent in some aphid colonies from the macrosiphini subfamily. The bodies of aphids from the subfamily Aphidinae are relatively small and have short siphunculi. On the other hand, aphids, which

have large bodies and relatively long sifunculi, are never visited by ants. This happens because long sifunculi are reported to disturb ants, so the ants don't like to come close. Additionally, large aphids and long sifunculi generally do not produce honeydew, so ants do not want to come close.

The absence of ants in aphid colonies could be because the colonies have just formed or the population is still low (Kummel et al. 2013). Aphids colonized flowers because they may offer an accessible and rich food source, sugary plant sap found in new growth or reproductive plant parts. Flowers contain a nutrient-rich nature and easy access to sap; therefore, aphids were attracted to flower saps. In addition, some aphid species were drawn to certain colors (Jakubczyk et al. 2022). Herbs served as an alternative host for aphids in this present study. Aphids consume sugar-rich liquid in plants, known as sap. Aphids considered herbs and other green vegetation as abundant food sources. Aphids utilize needle-like mouthparts to penetrate plant tissues and access this fluid (Brożek et al. 2015). Several aphids colonized herbs such as Indian mustards, *L. erysimi*, and *M. persicae*, the most devastating insects, infesting leaves, stems, and floral parts (Jayaswal et al. 2022). Due to a symbiotic relationship, the prevalence of aphids and ants was frequently correlated. Aphids produced a delicious substance known as honeydew as a waste product, which ants found highly attractive food sources (Nelson and Mooney 2022). The honeydew contained abundant sugars extracted by aphids from the plant juice (Zheng et al. 2022). Ants were drawn to this nutrient-rich food source and would often farm aphids for it. In exchange for honeydew, ants protected aphids from other insects and predators, such as ladybugs, lacewing larvae, and parasitic wasps (Karami-Jamour et al. 2018). Certain ant species would transport aphids to new host plants for improved foraging opportunities, ensuring that aphids had a continuous food source (Giannetti et al. 2021). Honeydew not only nourished the ant colony, but its high sugar content also supported the development of their fungus farms (in certain species) and provided energy for the growth of their progeny (Biedermann and Vega 2020). Ornamental plants, and also weeds are generally grown with simple maintenance and usually free pesticides. The ecological habitat of ornamental plants and weeds is assumed to be the same. Therefore, many species of aphids found on ornamental plants were also found on weeds.

In conclusion total of 15 species aphids found in ornamental plants, *A. craccivora*, *A. citricola*, *A. glycines*, *A. gossypii*, *A. solani*, *M. sanborni*, *M. rosae*, *M. persicae*, *N. circumflexus*, *P. caladii*, *R. nymphaeae*, *S. citricola*, *T. aurantii*, *T. citricidus*, *T. odinae*. The total of 11 species aphids found in weeds, *A. gossypii*, *A. craccivora*, *A.*

*glycines*, *A. citricola*, *Greenidea* sp., *H. setariae*, *Hiperomyzus* sp., *L. erysimi*, *R. maidis*, *R. padi*, *S. rotundiventris*.

## ACKNOWLEDGMENTS

The authors thank Universitas Sriwijaya, who supported this research. This research is a part of contract number 0188/UN9.3.1/SK/2023, 18 April 2023, with the chairman Chandra Irsan.

## REFERENCES

- Ghosh S, Singh R. 2004. Aphids on medicinal plants in North East India (Insecta : Homoptera : Aphididae). Rec Zool Surv India 102 (1-2): 169-186. DOI: 10.26515/rzsi/v103/i1-2/2004/159495.
- Bass C, Puinean AM, Zimmer CT, Denholm I, Field LM, Foster SP, Gutbrod O, Nauen R, Slater R, Williamson MS. 2014. The evolution of insecticide resistance in the peach potato aphid, *Myzus persicae*. Insect Biochem Mol Biol 51: 41-51. DOI: 10.1016/j.ibmb.2014.05.003.
- Bhadra P, Agarwala BK. 2014. On the morphological and genotypic variations of two congeneric species of banana aphid *Pentalonia* (Homoptera : Aphididae) from India. Aadv Life Sci 2 (3): 75-81. DOI: 10.5932/j.als.20120203.06.
- Biedermann PHW, Vega FE. 2020. Ecology and evolution of insect-fungus mutualisms. Ann Rev Entomol 65: 431-55. DOI: 10.1146/annurev-ento-011019-024910.
- Blackman RL, Eastop VF. 2008. Aphids On The World's Herbaceous Plants And Shrubs 2 Volume Set. John Wiley & Sons, Hoboken, New Jersey.
- Blackman RL, Eastop VF. 2017. Taxonomic Issues. CABI, Wallingford, UK. DOI: 10.1079/9781780647098.0001.
- Blackman RL, Sorin M, Miyazaki M. 2011. Sexual morphs and colour variants of *Aphis* (Formerly *Toxoptera*) *Odinae* (Homoptera, Aphididae) in Japan. Zootaxa 3110 (1): 53-60. DOI: 10.11646/zootaxa.3110.1.5.
- Braham M, Boulahia-kheder S, Kahia M, Noura S. 2023. Aphids and Citrus responses to nitrogen fertilization. J Saudi Soc Agric Sci 22 (6):374-83. DOI: 10.1016/j.jssas.2023.03.003.
- Brożek J, Mróz E, Wylęzek D, Depa Ł, Węgierek P. 2015. The structure of extremely long mouthparts in the aphid genus *Stomaphis* Walker (Homoptera: Sternorrhyncha: Aphididae). Zoomorphology 134:431-45. DOI: 10.1007/s00435-015-0266-7.
- Cao HH, Zhang ZF, Wang XF, Liu TX. 2018. Nutrition versus defense : Why *Myzus persicae* (Green peach aphid) prefers and performs better on young leaves of cabbage PloS One 13 (4):1-16. DOI: 10.1371/journal.pone.0196219.
- Clarke R, Kehoe MA, Broughton S, Jones RAC. 2020. Host plant affiliations of aphid vector species found in a remote tropical environment. Virus Res 281: 197934. DOI: 10.1016/j.virusres.2020.197934.
- Ertunc F. 2020. Emerging plant viruses. In: Emerging and Reemerging Viral Pathogens. Academic Press, Cambridge, Massachusetts.
- Gadhawe KR, Gautam S, Rasmussen DA, Srinivasan R. 2020. Aphid transmission of potyvirus: The largest plant-infecting RNA virus genus. Viruses 12 (7): 773. DOI: 10.3390/v12070773.
- Giannetti D, Mandrioli M, Schifani E, Castracani C, Spotti FA, Mori A, Grasso DA. 2021. First report on the acrobat ant *crematogaster* *scutellaris* storing live aphids in its oak-gall nests. Insects 12 (2): 108. DOI: 10.3390/insects12020108.
- Jakubczyk K, Koprowska K, Gottschling A, Janda-Milczarek K. 2022. Edible flowers as a source of dietary fibre (Total, Insoluble and Soluble) as a potential athlete's dietary supplement. Nutrients 14 (12): 2470. DOI: 10.3390/nu14122470.
- Jayaswal D, Mainkar P, Kumar K, Agarwal Y, Prabha R. 2022. Pyramiding and evaluation of segregating lines containing lectin and protease inhibitor genes for aphid resistance in *Brassica juncea*.



- Indian J Biochem Biophys 59 (8): 800-807. DOI: 10.56042/ijbb.v59i8.62319.
- Jones RAC. 2022. Alteration of plant species mixtures by virus infection: Managed pastures the forgotten dimension. Plant Pathol 71 (6): 1255-81. DOI: DOI: 10.1111/ppa.13571.
- Jousselin, Emmanuelle, Genson Gwenaelle, and Coeur D. Acie. Armelle. 2010. "Evolutionary Lability of a Complex Life Cycle in the Aphid Genus *Brachycaudus*." BMC Evolutionary Biology 10(1). doi: 10.1186/1471-2148-10-295.
- Kallas J. 2010. Edible Wild Plants. Gibbs Smith, Layton, Utah.
- Karami-Jamour T, Mirmoayedi A, Zamani A, Khajehzadeh Y. 2018. The impact of ant attendance on protecting *Aphis gossypii* against two aphidophagous predators and it's role on the intraguild predation between them. J Insect Behav 31: 222-39. DOI: 10.1007/s10905-018-9670-4.
- Kennedy JS, Stroyan HLG. 1959. Biology of aphids. Ann Rev Entomol 4 (1): 139-60. DOI: 10.1146/annurev.en.04.010159.001035.
- Kumar S, Bhowmick MK, Ray P. 2021. Weeds as alternate and alternative hosts of crop pests. Indian J Weed Sci 53 (1): 14-29. DOI: 10.5958/0974-8164.2021.00002.2.
- Kummel M, Brown D, Bruder A. 2013. How the aphids got their spots: predation drives self-organization of aphid colonies in a patchy habitat. Oikos 122 (6): 896-906. DOI: 10.1111/j.1600-0706.2012.20805.x.
- Liu, Xiang Dong, Ting Ting Xu, and Hai Xia Lei. 2017. "Refuges and Host Shift Pathways of Host-Specialized Aphids *Aphis Gossypii*." Scientific Reports 7(1):1–9. doi: 10.1038/s41598-017-02248-4.
- Maharani Y, Hidayat P, Rauf A, Maryana N. 2018. Short Communication: New records of aphid species subfamily Aphidinae (Hemiptera: Aphididae) in West Java, Indonesia. Biodiversitas 19 (2):460-65. DOI: 10.13057/biodiv/d190219.
- Meuninck J. 2023. Basic Illustrated Edible Wild Plants and Useful Herbs. Rowman & Littlefield, Lanham, Maryland, US.
- Mo, Carmen, and Angela M. Smilanich. 2023. "Feeding on an Exotic Host Plant Enhances Plasma Levels of Phenoloxidase by Modulating Feeding Ef Fi Cieny in a Specialist Insect Herbivore." Frontiers in Physiology 14(February):1–10. doi: 10.3389/fphys.2023.1127670.
- Naidu VSGR. 2012. Hand Book on Weed Identification. Directorate of Weed Science Research, Jabalpur, India.
- Nelson AS, Mooney KA. 2022. The evolution and ecology of interactions between ants and honeydew-producing hemipteran insects. Ann Rev Ecol Evol Syst 53: 379-402. DOI: 10.1146/annurev-ecolsys-102220-014840.
- Pettersson J, Tjallingii WF, Hardie J. 2017. Host-plant selection and feeding. In: Aphids As Crop Pests. CABI, Wallingford, UK. DOI: 10.1079/9781780647098.0173.
- Singh, Rajendra, and Garima Singh. 2021. "Aphids." Pp. 105–82 in Polyphagous Pests of Crops. Springer. Singapore. [https://doi.org/10.1007/978-981-15-8075-8\\_3](https://doi.org/10.1007/978-981-15-8075-8_3).
- Tegelaar K, Hagman M, Glinwood R, Pettersson J, Leimar O. 2012. Ant-aphid mutualism: The influence of ants on the aphid summer cycle." Oikos 121 (1): 61-66. DOI: 10.1111/j.1600-0706.2011.19387.x.
- Yamamoto, Tetsuya, Mitsuru Hattori, and Takao Itino. 2020. "Seasonal Migration in the Aphid Genus *Stomaphis* (Hemiptera : Aphididae): Discovery of Host Alternation Between Woody Plants in Subfamily Lachninae." Journal of Insect Science 20(5):1–10. doi: 10.1093/jisesa/ieaa103.
- Zheng Z, Zhao M, Zhang Z, Hu X, Xu Y, Wei C. 2022. Lactic acid bacteria are prevalent in the infrabuccal pockets and crops of ants that prefer aphid honeydew. Front Microbiol 12 (1): 1-17. DOI: 10.3389/fmicb.2021.785016.

