

Antibacterial Activity of Methanol Extract from Seagrass of Halodule uninervis in the Coastal of Lampung

by Ace Baehaki

Submission date: 18-Apr-2020 05:11PM (UTC+0700)

Submission ID: 1300878236

File name: rom_Seagrass_of_Halodule_Uninervis_in_the_Coastal_of_Lampung.pdf (103.95K)

Word count: 1379

Character count: 8030



Scholars Research Library

Der Pharmacia Lettre, 2016, 8 (4):77-79
(<http://scholarsresearchlibrary.com/archive.html>)



Antibacterial Activity of Methanol Extract from Seagrass of *Halodule Uninervis* in the Coastal of Lampung

Agus Supriadi*, Ace Baehaki and Muhammad Cahya Pratama

Department of Fisheries Product Technology, Faculty of Agriculture, Sriwijaya University, Indralaya, South Sumatera, Indonesia

ABSTRACT

The purpose of the research was to analyze bacterial activity from methanol extract of *Halodule Uninervis* against Gram-negative bacteria (*Aeromonas hydrophila* and *Vibrio harveyi*) and Gram-positive bacteria (*Bacillus subtilis* and *Listeria monocytogenes*). The result showed content of phenolic was 20.17 ppm and content of tannin was 1,223 ppm. Antibacterial activity of an extract showed that seagrass can inhibit several types of Gram-negative bacteria (*Aeromonas hydrophila* and *Vibrio harveyi*) and Gram-positive bacteria (*Bacillus subtilis* and *Listeria monocytogenes*). As for the highest inhibitory activity present in the type of Gram-positive bacteria (*Bacillus subtilis* and *Listeria monocytogenes*) between 7 mm to 9 mm.

Keywords : Seagrass, *Halodule uninervis*, antibacterial

INTRODUCTION

Seagrasses are the marine flowering plants. They are the only angiosperms that successfully growth in tidal and sub tidal marine environment [1]. The seagrasses was found to show prominent effect against the human bacterial pathogens [2-5] and hence they were tested for the effect against the pathogenic bacteria by using their crude extract of *Halodule Uninervis* seagrass.

However reports on the antibacterial of seagrasses of the coastal of Lampung, Indonesia are limited with the exception of few studies. Therefore the present study was undertaken to evaluate the antibacterial, and phytochemical constituents of methanol extract of *Halodule Uninervis* seagrasses.

MATERIALS AND METHODS

Leaves of *Halodule Uninervis* were collected from Coastal of Lampung and immediately brought to the laboratory in sterile plastic bags containing water to prevent evaporation. Sea grasses were washed thoroughly with distilled water to remove extraneous materials and shade-dried for 10 days at room temperature until constant weight obtained. The dried Sea grasses were powdered and stored in refrigerator for future use.

Preparation of Sea grass extract

Sea grass powder was soaked in 2L organic solvents with methanol (1:4 w/v), and kept for 10 days in a shaker. The extraction was repeated thrice and pooled. Each filtrate was concentrated to dryness under reduced pressure using a rotary flash evaporator. The dry aqueous extracts were stored in a refrigerator until further analysis.

The total phenolic content

The total phenolic content was determined by the Folin-Ciocalteu method [6]. The methanol extracts 250 μ L was mixed with 125 ml of Folin-Ciocalteu's phenol reagent. After 5 min, 250 ml of a 7% Na_2CO_3 solution was added to

the mixture followed by the addition of 125 ⁶ ml of deionized distilled water and mixed thoroughly. The mixture was kept in the dark at room temperature for 1 h. Absorbance was measured at 725 nm. The content of phenolic compounds was standardized with gallic acid and defined as mg of gallic acid equivalents per 1 g of sample

The total tannin content:

A total of 0.2 g of sample is introduced into the erlenmeyer containing 10 ¹⁰ ml of methanol, and then stirred using a mechanic ¹⁷ shaker for 1 hour. Taken 1.0 ml of the supernatant ¹³ then mix them with distilled water in a test tube. Added 0.3 ml of 0.1 M FeCl₃ and shake. Added 0.3 ml of 0.008 M K₃Fe(CN)₆. Then the mixture was allowed to stand for 10 minutes, the sample absorbance is read at a wavelength of 720 nm. Documents used is the whole solution, but without the sample. Tannin content is calculated by the following formula:

$$\text{Tanin (ppm)} = \frac{\text{absorbance sample}}{\text{absorbance blanko}} \times \frac{100}{\text{Weight of sample(g)}} \times 100$$

Antibacterial test

Pathogens bacterial (*Aeromonas hydrophila*, *Vibrio harveyi*, *Bacillus subtilis* and *Listeria monocytogenes*) used in this study. Antibacterial activity was evaluated using diffusion method [7]. Actively growing lag phase cultures of bacteria were mixed in Nutrien agar (Nutrien broth with 1.5% agar) and plate ¹⁰ The various extract (500 ppm, 1,000 ppm, 1,500 ppm and 2,000 ppm) were loaded onto different ¹⁹ per discs (Whatman no. 1 filter paper). The discs were placed on the agar medium containing cultures incubated for 24 h at 37°C. Zone of inhibition was recorded in millimeters and mean values were reported.

7 RESULTS AND DISCUSSION

The total phenolic and tannin content

Table 1. The total phenolic and tannin content of *Halodule uninervis*

Extract of	Total Phenolic (ppm)	Total tannin (ppm)
<i>Halodule uninervis</i>	20.17	1,223

Our study shows a small content of phenolic compared to total tannin which is around 20.17 ppm (20.17 ppm to 1,223 ppm, respectively). The phytochemical compounds detected such as tannins and phenolic have previously been reported to have antimicrobial and antioxidant activity [8].

Antibacterial activity from *H. uninervis* against Gram-positive bacteria

Antibacterial activity from *H. uninervis* with different concentration of extract against Gram-positive bacteria are depicted in Figure 1. The Antibacterial activity from *Halodule uninervis* toward *Bacillus subtilis* and *Listeria monocytogenes* were peaked after 2,000 ppm of extract concentration.




Figure 1. Bacterial activity of *H. uninervis* against Gram-positive bacteria (A=*Bacillus subtilis* and B=*Listeria monocytogenes*)

In antimicrobial assay the *H. uninervis* extracts have shown good antimicrobial activities against the *Bacillus subtilis* and *Listeria monocytogenes* with a maximum inhibitory effect at 9mm and minimum inhibitory effect at 6 mm.

Antibacterial activity from *H. uninervis* against Gram-negative bacteria

Antibacterial activity from *H. uninervis* with different concentration of extract against Gram-negative bacteria are depicted in Fig. 2.




Figure 2. Bacterial activity of *Halodule uninervis* against Gram-negative bacteria (A= *Aeromonas hydrophila* and B= *Vibrio harveyi*)

Bacterial activity of *Halodule uninervis* against *Bacillus subtilis* was from 6 mm to 8 mm and the extract of *Halodule uninervis* had the highest bacterial activity, which was 8 mm at the concentration was 2,000 ppm. While bacterial activity of *Halodule uninervis* against *Listeria monocytogenes* was from 6 mm to 9 mm and the extract had the highest bacterial activity which 9 mm at concentration was 2,000 ppm.

These findings suggest that antimicrobial activity of *H. uninervis* extract may be primarily due to the presence of tannins and phenolic compound. The inhibition of microorganisms by phenolic compounds may also be due to iron deprivation or hydrogen binding with vital proteins such as microbial enzymes [9]. Tannins have been traditionally used for treatment of catarrh, hemorrhoids and diarrhea [10].

The bacterial activity of *H. uninervis* extract against Gram-positive bacteria higher than that of these Gram-negative bacteria. A small activity of *uninervis* extract against Gram-negative bacteria due to Gram-negative bacteria contain three layer : liposaccharide, peptidoglycan and protein so that compounds of seagrasses can not damage the cell of bacteria.

CONCLUSION

From the present study the extract of seagrass (*Halodule uninervis*) has anti-bacterial effect and effective on gram-positive (*Bacillus subtilis* and *Listeria monocytogenes*).

REFERENCES

- [1] Gravatt, G. L.; Baguley, B. C.; Wilson, W. R. *J. Med. Chem.*, 1994, 37, 4338.
- [1] Vijayakumar, S.; Manakha, M. I.; Anderson, A.; Lingakumar, K. *Int J Microbiol & Parasit*, 2014, 1(1), 1-10
- [2] Umamaheswari, R.; Thirumaran, G.; Anantharaman, P. *Adv. Bio J Res*, 2009, 3, 140-143.
- [3] Kannan, R. R. R.; Arumugam, R.; Anantharaman, P. *Asian Pacific J. Trop Med.*, 2010, 890-893
- [4] Kannan, R. R. R.; Arumugam, R.; Iyapparaj, P.; Thangaradjou, T.; Anantharaman, P. *Food Chem.*, 2013, 136, 1484-1489
- [5] Sangeetha, J.; Asokan, S. *World J. Pharmacy and Pharma. Sci.*, 2015, 4(12), 677-683.
- [6] Chandler, S. F.; Dodds, J. H. *Plant cell reports*, 1983, 2(4), 205-208.
- [7] Elgayyar, M.; Draughon, F. A.; Golden, D. A.; Mount, J. R. *J. Food Prot.*, 2001, 64(7), 1019-1024
- [8] Leven, M.; Vannenberghe, D. A.; Mertens, F. *J. Planta Med.*, 1979, 36, 311-321.
- [9] Scalbert, A. *Phytochemistry*, 1991, 30, 3875-3883.
- [10] Senthilkumar, M. *Int. J. Sci Res Pub*, 2013, 3(1), 1-5

Antibacterial Activity of Methanol Extract from Seagrass of Halodule Uninervis in the Coastal of Lampung

ORIGINALITY REPORT

47%

SIMILARITY INDEX

%

INTERNET SOURCES

31%

PUBLICATIONS

41%

STUDENT PAPERS

PRIMARY SOURCES

1	Rengasamy Ragupathi Raja Kannan, Radjassegarin Arumugam, Palanisamy Iyapparaj, Thirunavukarasu Thangaradjou et al. "In vitro antibacterial, cytotoxicity and haemolytic activities and phytochemical analysis of seagrasses from the Gulf of Mannar, South India", Food Chemistry, 2013 Publication	8%
2	Submitted to Universitas Diponegoro Student Paper	7%
3	Submitted to Kingston University Student Paper	6%
4	Submitted to Universiti Sains Malaysia Student Paper	5%
5	Submitted to Universitas Andalas Student Paper	4%
6	Submitted to Higher Education Commission Pakistan Student Paper	2%

7	Dutsadee Chinnapun. " Antioxidant activity and DNA protection against oxidative damage of bambara groundnut seeds ((L.) Verdc.) as affected by processing methods ", International Journal of Food Properties, 2018 Publication	2%
8	E. Warsiki, J.T. Bawardi. "ASSESSING MECHANICAL PROPERTIES AND ANTIMICROBIAL ACTIVITY OF ZINC OXIDE-STARCH BIOFILM", IOP Conference Series: Earth and Environmental Science, 2018 Publication	2%
9	Rengasamy Ragupathi Raja Kannan, Rajasekaran Arumugam, Perumal Anantharaman. "Antibacterial potential of three seagrasses against human pathogens", Asian Pacific Journal of Tropical Medicine, 2010 Publication	2%
10	Submitted to Surabaya University Student Paper	2%
11	Submitted to Southern Cross University Student Paper	2%
12	Sigrun Eick, Kevin Hofpeter, Anton Sculean, Claudia Ender, Susann Klimas, Sebastian Vogt, Sandor Nietzsche. "Activity of Fosfomycin- and Daptomycin-Containing Bone Cement on	1%

Selected Bacterial Species Being Associated with Orthopedic Infections", BioMed Research International, 2017

Publication

13

H.-S. Chiang. "Comparative constitutive resistance in soybean lines to Mexican bean beetle", *Entomologia Experimentalis et Applicata*, 09/1986

Publication

1%

14

Submitted to Sheffield Hallam University

Student Paper

1%

15

Malairaj Sathuvan, Anadhan Vignesh, Ramar Thangam, Perumal Palani, Ramasamy Rengasamy, Kandasamy Murugesan. "In Vitro Antioxidant and Anticancer potential of Bark of *Costus pictus* D.DON", *Asian Pacific Journal of Tropical Biomedicine*, 2012

Publication

1%

16

Xosé Anxelu G. Morán, Federico Baltar, Cátia Carreira, Christian Lønborg. "Responses of physiological groups of tropical heterotrophic bacteria to temperature and DOM additions: food matters more than warming", *Environmental Microbiology*, 2020

Publication

1%

17

Salah A Al-Maiman, Dilshad Ahmad. "Changes in physical and chemical properties during

1%

pomegranate (Punica granatum L.) fruit
maturation", Food Chemistry, 2002

Publication

Exclude quotes	Off	Exclude matches	Off
Exclude bibliography	On		