

🔨 10.22146/ijc.21456 í Abstract views : 266 🔤 views : 717	
SORPTION OF Au(III) BY Saccharomyces cerevisiae BIOMASS	177-183
Amaria Amaria, Suyono Suyono, Eko Sugiharto, Anis N Rohmah	
🔨 10.22146/ijc.21457 í Abstract views : 280 🔤 views : 290	
EFFECT OF SCANDIUM ON HIDROGEN DISSOCIATION ENERGY AT MAGNESIUM SURFACE: AB INITIO DFT STUDY	184-188
l Wayan Sutapa, Ria Armunanto, Karna Wijaya	
🔨 <i>10.22146/ijc.21458 🛍</i> ll Abstract views : 438 🔤 views : 372	
CYCLISATION-ACETYLATION KINETIC OF (R)-(+)-CITRONELLAL BY Zn ²⁺ -NATURAL ZEOLITE AS SOLID SOLVENT CATALYST	189-194
U10.22146/ijc.21459 MIII Abstract views : 301 🔤 views : 456	
STRUCTURE ELUCIDATION OF ALKALOIDS FROM LEAVES OF <i>Voacanga foetida</i> (BI.) Rolfe OF LOMBOK ISLAND	195-200
Surya Hadi, Dina Asnawati, Novi Febrianti	
10.22146/ijc.21460 iiiiii Abstract views: 279 22 views: 330	
CYCLIZATION AND HYDROGENATION OF (+)-CITRONELLAL TO MENTHOLS OVER $ZnBr_2$ AND Ni CATALYSTS SUPPORTED ON γ -Al_2O_3	201-206
Elvina Dhiaul Iftitah, M. Muchalal, Wega Trisunaryanti, Ria Armunanto	
🔨 10.22146/ijc.21461 í Abstract views : 267 🔤 views : 283	
THE ADDITION OF PETROLEUM BENZIN IN MIXING OF CRUDE PALM OIL AND METHANOL ON BIODIESEL PRODUCTION	207-204
Abdullah Abdullah, F Fadhillah, Supranto Supranto, A. Prasetya	
[©] 10.22146/ijc.21462 前側 Abstract views : 298 views : 308	
6,6'-DIMETHOXY-4,4'-DIHYDROXY-3',2'-FURANO-ISOFLAVANE, A NEW COMPOUND FROM <i>Melochia</i> umbellata (Houtt.) Stapf var. Degrabrata K. (<i>Paliasa</i>)	215-218
\bigvee 10.22146/IJC.21463 IIIIII Abstract views : 533 $\stackrel{\text{\tiny IM}}{=}$ views : 381	
ANTIBIOTIC COMPOUND FROM MARINE ACTINOMYCETES (<i>Streptomyces sp A11</i>): ISOLATION AND STRUCTURE ELUCIDATON	219-225
Rofiq Sunaryanto, Bambang Marwoto, Tun Tedja Irawadi, Zainal Alim Mas'ud, Liesbetini Hartoto	
🔮 10.22146/ijc.21464 📶 Abstract views : 327 📟 views : 338	
UGONIN J FLAVONOID FROM TUNJUK LANGIT (Helminthostachys zeylanica Linn.) ROOT EXTRACT	226-231
Fitrya Fitrya, Lenny Anwar, Eliza Eliza, Muharni Muharni	
🔨 10.22146/ijc.21465 í Abstract views : 379 🔤 views : 345	
BIOACTIVITY AND CLONING OF A NEW ANTIBACTERIAL LECTIN PROTEIN IN SPONGE <i>Gelliodes</i> sp. FROM BARANGLOMPO ISLAND IN SOUTH SULAWESI INDONESIA TERRESTRIAL	232-239
Hanapi Usman, Prastawa Budi, Ahyar Ahmad	
🔨 10.22146/ijc.21466 🏭 Abstract views : 348 🔤 views : 256	
ANTICANCER ACTIVITY OF CALANONE ON HeLa CELL LINE Heny Ekowati, Indwiani Astuti, Mustofa Mustofa	240-244
🔨 10.22146/ijc.21467 í Abstract views : 353 🔤 views : 1210	
MUTATION ON WD DIPEPTIDE MOTIFS OF THE p48 SUBUNIT OF CHROMATIN ASSEMBLY FACTOR-1 CAUSING VIABILITY AND GROWTH OF DT40 CHICKEN B CELL LINE	245-250
Ahyar Ahmad, Harningsih Karim	
₩ 10.22146/ijc.21468 milli Abstract views : 259 ₩ views : 305	
EFFECTS OF DIETARY ADENINE ON GROWTH, LIPID SECRETION, AND LIVER LIPID CONTENT Yohanes Buang	251-255
🔨 10.22146/ijc.21469 í Abstract views : 280 🔤 views : 287	
PRODUCTION AND CHARACTERIZATION OF CHITINASE ENZYMES FROM SULILI HOT SPRING IN	256-260

adsorption

antioxidant catalyst characterization chitosan cytotoxic eugenol extraction ash heavy metals humic acid immobilization kinetics molecular docking silica synthesis transesterificatio zeolite

Indones. J. Cherr indexed by:

Scopus°











CURRENT ISSUE

ATOM	1.0
RSS	2.0
RSS	1.0

233

UGONIN J FLAVONOID FROM TUNJUK LANGIT (*Helminthostachys zeylanica* Linn.) ROOT EXTRACT

Fitrya^{1.}*, Lenny Anwar², Eliza¹, and Muharni¹

¹Department of Chemistry, University of Sriwijaya, JI. Raya Palembang-Prabumulih Km 32, Indralaya, OI, Sumsel ²Department of Chemistry, University of Riau, JI. Prof. Dr. Muchtar Lutfi Simpang Baru Pekanbaru

Received February 18, 2010; Accepted April 5, 2010

ABSTRACT

A flavonoid compound was isolated from dried rihzomes of tunjuk langit (Helminthostachis zaylanica Lin), a traditional medicine from South Sumatera, Indonesia. Extraction was done by maceration methode and separation of isolated compound was conducted by chromatographic technique. The structure of this compound was determined base on spectroscopic data such as including UV, IR, 1-D, 2-D NMR, and comparison with the reported data. Based on spectral data analysis, concluded that isolated compound was Ugonin J (5,7,3',4'-tetrahydroxy-6-(6,6-dimethyil-2-methylenecyclo-hexylmethyl)flavone).

Keywords: Flavonoid, Ugonin J, Helmintohostacys zaylanica

INTRODUCTION

Tunjuk langit (Helminthostacys zaylanica Linn.) (Ophiolglosaceae) is pteridophyte show medicial utility. The rhizome of tunjuk langit traditionally used as cytotoxic, antinflammatory and pulmonary disease. The rhizome of tunjuk langit is Chinese herbal medicine used as antypiretic and antiphlogistic agent. In India, the rhizome of tunjuk langit used for curing impotency [1]. In Malaysia, the rhizome used as antidiarrheal agent and chewed with areca for whooping cough relief [2]. Suja reported that ethanol extract of rhizome of H. zeylanica showed aphrodiciac properties [4]. Study of ethanol rhizome extract of also showed significant hepatoprotective effect [5]. Huang yielded eight flavonoids, ugonin E-L from the rhizome of tunjuk langit Ugonin J, K and L showed significant antioxidant activity [6]. Yi Chen reported that ugonin K, a flavonoid from rhizome of tunjuk langit has neuroprotective activity in neuroblastoma SH-SY5Y cells [7] and ugonin L showed antiinflammatory effect [8].

H. *zaylanica* is pteridophytes, widely distributed in tropical Asia and Australia. This plant is locally named, paku payung, pancar bumi (South Sumatra), tapak jalak (Sunda), bute-bute (Makasar), pakis urang, pakis kaler, ceker ayam (Java) [9]. Three new cyclized stilbenes, ugonstilben A, B and C and 3-hidroxy asetophenon compound were isolated from the rhizome of *Helminthostachys zeylanica* [10]. Four flavonoid compounds ugonin A-D were isolated too [11].

This plant widely growth in Lahat, South Sumatra. In Indralaya, traditionally used as cytotoxic, antiinflammatory and pulmonary disease treatment. In this paper we described the isolation and structure

* Corresponding author. Tel/Fax : +62-85758666119 Email address : fitrya_apt@yahoo.com flavonoid compound from ethylacetate extract from rhizome of H. *zeylanica*.

EXPERIMENTAL SECTION

Materials

The rhizome of H. *zeylanica* were collected from Jati village and Kuba village, Kecamatan Pulau Pinang Kabupaten Lahat South Sumatra. The plant was also identified by comparison with a voucher specimen already deposited at the Herbarium ANDA, Andalas University. Vacuum liquid chromatography (VLC) was carried out using Merck Si gel 60 GF₂₅₄ (230-400 Mesh) and column chromatography using Si gel Merck G 60 (70-230 Mesh), thin layer chromatography (TLC) analysis was performed on precoated Si Gel plates (Merck Kiesel gel 60 GF₂₅₄, 0.25 mm 20 x 20 cm).

Instrumentation

UV and IR spectra were measured with spektrofotometer Beckman DU-700 and Shimadzu FTIR 8400. ¹H and ¹³C NMR spectra was recorded JEOL JNM ECA-500 500 MHz (¹H) and 125 MHz (¹³C) using internal standard TMS.

Procedure

Extraction and isolation

The powdered of rhizome of *H zeylanica* (5 kg) was extracted with n-hexane, ethylacetate and methanol respectively (8 Lx 3). The extract was filtered

and the filtrate concentrated under reduced pressure. The EtOAc extract was chromatographed on column.

EtOAc extract (50 g), was subjected to vacuum liquid chromatography eluted with a gradient system (n-hexana, n-hexane: EtOAc = 9:1; 8:2; 7:3; 6:4 and EtOAc) to afford 5 fractions F1–F5. Fraction F3 was further purified column chromatography (eluted with MTC-MeOH 9.5:0.5) to afford 4 subfraction F3.1-F3.4. Subfraction F3.3 was repeatedly separated over silica gel colomns to afford F3.3.1-F3.3.1. F3.3.4 after purification with recrystalization gave a pure compound as yellow powder (50 mg).

Structure elucidation

The structure was elucidating using UV, IR, 1D and 2D NMR spectroscopic data. The ¹³C NMR signals were assigned from DEPT, HMQC, HMBC and COSY spectra. The ¹H and ¹³C spectra data also compared with those reported in the literature. In the UV spectra, we can obtain information about chromophore, information from IR spectra was specific functional group. ¹H NMR we were obtain information about quantity of proton, chemical shift, coupling constant, and ¹³C NMR used to implied total carbon and DEPT (Distortionless Enhancement by Polarization Transfer) spectrum showed kinds of C, CH, CH₂ and CH₃.HMQC (Heteronuclear Multiple Quantum Coherence) used to identified signals of carbon which attached proton, HMBC (Heteronuclear Multiple Bond Connectivity) showed long range correlation ²J and ³J of proton to carbon, and COSY (Correlated Spectroscopy) showed proton adjacent. [12-13].

RESULT AND DISCUSSION

Extraction and Isolation

Dried rhizome of H. *zeylanica* (5 kg) extracted with n-hexane, EtOAc and MeOH respectively and resulted n-hexane extract (30 g), EtOAc extract (51 g) and MeOH extract (30 g). EtOAc extract separated over silica gel column to afford a pure compound.

Structure Elucidation

The pure compound as yellow powder has melting point 229-230 °C and $[\alpha]_D^{20}$ -49° (c 1,0, MeOH). The structure was elucidating using 1D and 2D NMR spectroscopic data. The ¹³C NMR signals were assigned from DEPT, HMQC, HMBC and COSY spectra. The ¹H and ¹³C spectra data also compared with those reported in the literature. The UV spectrum (MeOH) showed λ_{max} absorption at 347(4.27), 275 (4.16) and 215 (4.5) nm characteristic to flavone nucleus [14]. The UV spectrum in the presence of NaOH showed bathochromic shift λ_{max}

(nm) 405, 278 and 215 indicated there was phenolic chromophore. The IR spectrum (KBr) showed absorptions typical of hydroxyl (3440 cm⁻¹), C-H aliphatic (2920, 2962 cm⁻¹), carbonyl (1651), aromatic (1651, 1612 and 1465 cm⁻¹) and alcohol (1172 cm⁻¹). The ¹H NMR and ¹³C HNMR (methanol-*d*4)

The ¹H NMR and ¹³C HNMR (methanol-*d4*) spectra of pure compound (Table 1) showed characteristic signals for flavone. ¹H NMR spectrum showed signal at 7.36 (1H, *dd*, *J* = 1.8 and 8.5 Hz), can be attribute to aromatic proton which *orto* and *meta* coupling with 6.89 (1H, *d*, *J* = 8.5 Hz) and $\delta_{\rm H}$ 7.35 (1H, *d*, *J* = 1.8 Hz) proton from trisubstituted benzene ring. Signals at $\delta_{\rm H}$ 6.51 (1H, *s*) and 6.42 (1H, *s*) assigned two aromatic which uncoupled. Five specific protons indicated that this compound was flavone.

Furthermore, signals at $\delta_{\rm H}$ 2.73 (H-9a) (1H, *dd*, J = 3.7 and 12.9 Hz), and 2.94 (H-9b) (1H, *t*, J = 12.9) attribute to methylene group which *orto*-coupling with proton at $\delta_{\rm H}$ 2.41 (1H, *dd*, J = 3.7) and coupling gemynal. Signal at $\delta_{\rm H}$ 4.42 and 4.19 (1H, *brs*) was for methine proton from C-18, suggesting gemynal coupling. ¹H NMR spectrum showed three signals for methylene SP3 at 2.56 (H-12a), 1.95 (12b), 1.59 (13-a), 1.48 (13-b), 1.70 (14a), and 1.28 (14-b) respectively (1H, *m*). Signals at $\delta_{\rm H}$ 1.06 (3H, *s*) and 0.94 (3H, *s*) attributed to methyl proton from prenyl unit. ¹H NMR spectrum showed at Fig. 1. Based on signals at under 3 ppm suggested that the pure compound contain syclic geranyl unit. That was flavone which siclic geranyl substituted.

¹³C NMR spectrum showed 25 signals and 15 of these were SP2 signals for aromatic carbon and one of these for carbonyl carbon at 184.0 ppm, can see at Fig. 2. These signals specific for carbon or flavonoid typical flavone. Nine remaining signals were signals typical resulted from geranyl unit which has one carbon SP2. That was strengthen two signals for CH₃ only, revealed the presence two signals CH₃. Therefore, these data supporting isolated pure compound to be a flavone which geranyl substituted.

At DEPT Spectrum (Fig. 2) showed two signals methyl at $\delta_{\rm C}$ 28.6 (C-16) and 28.8 (C-17), five methylene signals (CH₂) at $\delta_{\rm C}$ 22,4 (C-9); 24,6 (C-13); 32.3 (C-12); 35.9 (C-14); and 109.8 (C-18), six methine signals (CH) at $\delta_{\rm C}$ 53.6 (C-10); 93.9 (C-8); 103.8 (C-3); 114.2 (C-2'); 116.5 (C-5') and 120.3 (C-6'), and twelve signals kuaterner carbon at $\delta_{\rm C}$ 35.5 (C-15); 104.9 (C-C-4^a); 113.5 (C-6); 123.9 (C-1'); 147.1 (C-3); 150.9 (C-11); 151.2 (C-4); and 157.2. Furthermore, isolated compound structure predicted supporting by NMR 2-D data analysis. HMQC spectrum showed correlation between proton at $\delta_{\rm H}$ 6.51 (H-3) with carbon at $\delta_{\rm C}$ 103.8 and HMBC spectrum showed there were long range correlation with C-4a at $\delta_{\rm C}$ 104.9; 123.9 (C-1'); and displayed correlation (²J) with carbon at $\delta_{\rm C}$ (C-2)





Fig 4. HMBC at δ_C 6.8 – 7.4 ppm spectrum

and 184.0 (C-4), so this proton deduced attached to C-2. The HMBC spectrum also displayed long range correlation 3J between proton at $\delta_{\rm H}$ 6.42 (H-8) with carbon at $\delta_{\rm C}$ 104.9 (C-4a); 113.5 (C-6) and 2J coupling with carbon at $\delta_{\rm C}$ 157.2 (C-8a) and 164.3 (C-7) so this proton attributed to C-8. The HMQC spectrum displayed this proton correlated to carbon at $\delta_{\rm C}$ 93.9.

Furthermore, the HMQC spectrum (Fig. 3) showed proton at 6.89 attached at $\delta_{\rm C}$ 116.8 ppm (C-5'), and HMBC spectrum at Fig. 4 showed correlation with $\delta_{\rm C}$ 147.1 (C-3') and 123.9 (C-1') so this proton attributed at C-5'. Proton at 7.35 (H-2') and 7.36 (C-6') at spectrum HMQC, attached to carbon at $\delta_{\rm C}$ 114.2 and 120.3 showed ³J coupling with carbon at $\delta_{\rm C}$ 120.3 (C-6') and 114.2 (C-2') and ²J with carbon at $\delta_{\rm C}$ 147.1 (C-3'), so deduced this proton attached at C-2' and C-6'.

HMQC spectrum also showed proton at δ_{H} 2.73 (H-9a) and 2.94 (H-9b) attached to carbon at 22.4 ppm and HMBC spectrum displayed correlation with carbon at 53.6 (C-10), and 113.5 (C-6) attributed to C-9. In the COSY spectrum (Fig. 5) displayed correlation between proton at δ_{H} 2.73 (H-9a) and δ_{H} 2.94 (H-9b) with proton



Fig 6. HMBC and COSY correlation of ugonin J (A)

at 2.41 (H-10), in HMBC spectrum displayed correlation with carbon at 35.5 (C-15) so this signal appeared as doublet-doublet. The COSY spectrum also displayed correlation between proton at δ_H 2.56 (H-12a) with proton at δ_H 1.59 (H-12b), proton at 1.59 (C-13a) and 1.48 (C-13b), and proton at δ_H 1.70 (H-14a) and 1.28 (H-14b) so this proton appeared as multiplets.

Correlation long range coupling ²J and ³J between proton with carbon from HMBC and correlation between protons with proton at COSY showed at Fig. 6.

The ¹³C NMR data of isolated compound (A) showed same with Ugonin J (A*) as reported data (Table 1), but there was difference because isolated compound measured in methanol-*d4* whereas (-)-Ugonin J measured in DMSO-*d6*. There was also same of melting point and optical density isolated compound 229-230 °C and $[\alpha]_D$ -49° (c 1,0; MeOH) respectively whereas ugonin J as reported data was $[\alpha]_D^{20}$ -50° (MeOH). Based on above data concluded the isolated compound was Ugonin J showed at Fig. 7.

Position	δ_{H} (ppm), integration, mult, <i>J</i> (Hz)	δC (ppm)		DEPT	HMBC	COSY
	A	А	A*	А	А	А
2		165.9	164.7	С		
3	6.51(1H, <i>s</i>)	103.8	104.2	CH	C-4a, C-1', C2, C4	
4		184.0	183.0	С		
4a		104.9	104.9	С		
5		160.5	160.3	С		
6		113.5	112.4	С		
7		164.3	162.7	С		
8	6.42 (1H, <i>s</i>)	93.9	93.7	СН	C-4a, C-6, C8a, C7	
8 ^a		157.2	156.3	С		
9	2.73 (1H,dd, 3,7; 12,9)	22.4	21.9	CH2	C-5, C-6, C-7, C-10	H-10
	2.94 (1H, t, 7,9; 12,9)					
10	2.41 (1H, dd, 3,7; 7,9)	53.6	53.0	СН	C-15	H-9a, H-9b
11		150.9	150.5	С		
12	2.56 (1H, <i>m</i>)	32.3	32.0	CH2		H-12b
	1.95 (1H. <i>m</i>)					H-12a
13	1.59 (1H, <i>m</i>)	24.6	24.1	CH2		H-12 ^a
	1.48 (1H, <i>m</i>)					H-12b,
						H-14 ^a
14	1.70 (1H, <i>m</i>)	35.9	35.2	CH2		H-14b
	1.28 (1H. <i>m</i>)					H-14a
15		35.5	35.2	С		
16	1.06 (3H. s)	28.6	28.1	CH3	C-10. C-15. C-17	
17	0.94 (3H, s)	28.8	28.5	CH3	C-10, C-16	
18	4.42 (1H, brs)	109.8	109.4	CH2	,	H-18b
	4.19 (1H.brs)					H-18a
1'	- ())	123.9	123.9	С		
2	7.35 (1H, <i>m</i> , 1,8)	114.2	113.9	ĊН	C-3', C-6'	
3		147.1	146.3	С	,	
4		151.2	149.8	C		
5	6.89 (1H, <i>d</i>)	116.8	116.5	ĊН	C-3', C-1', C-4'	H-6'
6	7.40 (1H, dd, 1,8))	120.3	120.0	СН	C-4', C-2	H-5'

Table 1. ¹H and ¹³C NMR 1-D and 2-D data (methanol-*d4*) for isolated compound (A) and Ugonin K (A*) as comparing data [6].



Fig 7. Structure of ugonin J

CONCLUSION

Ugonin J or 5,7,3',4'-tetrahidoksi-6-(6,6-di-metil-2metilen-sikloheksilmetil) flavone) had been isolated from ethylacetate extract of rhizome of *Helmynthostachys zeylanica*

ACKNOWLEDGEMENT

The authors are grateful to the Herbarium ANDA, Padang, Indonesia for identification of the plant specimen and LIPI Research Center for Chemistry, PUSPITEK, Serpong, Indonesia for spectrum measurement.

REFERENCES

- Singh, V., Ali, Z.A., Zaidi, S.T.H., and Siddqui, M.K., 1996, *Fitoterapia*, Department Botany Aligarh Muslim Universities Aligarh, India.
- Jalil, J., Bidin, A.A., and Chye, T,S., 1986, *Phytochemical Study of Ophyoglosaceae Family Species, Proc. Malays Biochemistry Society Conference*, Fakultas Sains Hayati Universitas Kebangsaan Malaysia, Bangi Malaysia, 12, 160-164.

237

- 3. Napralert, 2003, *Informasi Tumbuhan Helminthostachys zeylanica (L) Hook*, Faculty of Pharmacy, Andalas University.
- 4. Suja, S.R., Latha, P.G, Pushpangadan, P., and Raasekharan, S., 2002, J. Trop. Med. Plants, 3, 191-195.
- 5. Suja, S.R, Latha, P.G, Pushpangadan, P., and Raasekharan. S., 2004, J. Ethnopharmacol., 92, 1, 61-66.
- 6. Huang, Y-L., Yeh, P.Y., Shen, C.C., and Chen, C.C., 2003, *Phytochemistry*, 64, 7, 1277-1283.
- Lin, Y.C., Huang, Y.C., Chen, S.C., Liaw, C.C., Kuo, S.C., Huang, L.H., and Gean, P.W., 2009, *Neurochem. Res.*, 34, 923-930.
- Huang, Y.C., Hwang, T.L., Chang, C.S., Yang, Y.L., Shen, C.N., Liao, W.Y., Chen, S.C., and Liaw, C.C., 2009, *J. Nat. Prod.*, 72, 7, 1273-1278.

- 9. Heyne, K., 1987, *Tumbuhan berguna Indonesia*, Jilid III, Jakarta: Yayasan Sarana Wana Jaya, 1387
- 10. Chen, C.C., Huang, Y.L., Yeh, P.Y., and Ou, J.C., 2003, *Planta Med.*, 69, 10, 964-967.
- 11. Murakami, T., Hagiwara, M., Tanaka, K., and Chen, C.M., 1973, *Chem. Pharm. Bull.*, 21, 1851-1852.
- 12. Breitmaier, E., 1993, *Structure elucidation by NMR in Organic Chemistry, A Practical Guide*, England: John Wiley & Sons Ltd.
- 13. Nakanishi, K., 1990, One-Dimensional and Two-Dimensional NMR Spectra by Modern Pulse Technique, Tokyo: University Science Books Mill Valley.
- 14. Markham, K.R., 1988, *Cara Mengidentifikasi Flavonoid*, Penerbit ITB Bandung.