

File ini lengkap berisi :
a cek itenticate
b bukti korespondensi
c lembar peer review

jurnal4

by Mardiyanto Mardiyanto

Submission date: 21-May-2020 10:56AM (UTC+0700)

Submission ID: 1328868741

File name: jurnal4.pdf (1.36M)

Word count: 2563

Character count: 15210

⁴
**THE SCREENING OF POTENTIAL ANTIBIOTICS FROM HOSPITAL
WASTEWATER IN TROPICAL REGION (CASE STUDY AT
PALEMBANG, SOUTH SUMATRA, INDONESIA)**

I. KURNIAWAN^{1,2}, S. NASIR³, HERMANSYAH⁴ AND MARDIYANTO⁴

¹*Sriwijaya University, Palembang, Indonesia*

²*Faculty of Health Science, Muscharitas Catholic University, Palembang, Indonesia*

³*Faculty of Engineering, Sriwijaya University, Palembang, Indonesia*

⁴*Faculty of Mathematics and Science, Sriwijaya University, Palembang, Indonesia*

(Received 1 February, 2017; accepted 15 March, 2017)

ABSTRACT

Clean water is a primary one of the causes of water quality degradation is pollution through the residue of human activity. Hospital is a potential pollutant source with the characteristics of hazardous and toxic wastewater. The components of pollutants in the wastewater of hospital is still limited investigated is antibiotic. The regulation of hospital wastewater in Indonesia as tropical and developing countries do not regulate antibiotics, while in the world the issue of zero tolerance of antibiotics becomes an important concern. Antibiotics have a persistent and resistant impact for the environment. In this research, screening for antibiotics by using samples from one type B hospital in Palembang City, Indonesia is done. Hospital wastewater samples in the non permanent ¹ and are the residue of all process activities mixed in one pond. Screening using High-Resolution ¹ Mass Spectrometry (LC / Q-TOF / MS) 6500 Series for the Detection of Pharmaceuticals in Water is used to detect the antibiotics. The result of laboratory analysis shows that there are five types of antibiotics detected, namely Ciprofloxacin, Lincomycin, Metronidazole, Netilmicin, and Ofloxacin/Levofloxacin. The qualitative analysis concluded that Ciprofloxacin is one of a type potential antibiotics in the hospital wastewater. The result of this research recommends that the Indonesian Government should revise the regulation about antibiotics in the hospital wastewater.

KEY WORDS : Clean water, Environment, Antibiotics, Hospital wastewater, Ciprofloxacin

INTRODUCTION

World Health Statistics reported by WHO in 2015 shows that about 85% of the Indonesian population needs access to clean water for consumption. The community in the late decades concerns the importance to solve the environmental problems, especially about clean water conservation. International Water Institute predicts in 2025 Java and the others Islands in Indonesia will have a crisis of clean water. The water global scarcity has increased and is needed the other sources to make it balance with the water need because existing

water sources has not been used because of the pollution and wastewater. The problem of clean water sources because of containing components of polluters in high concentrations, such as heavy metals, toxic compounds, and microbial pathogens.

The data released by WWF world organization in its report explain that the data between 1930 to 2000 about the production of chemicals caused by human activities increased from 1 million to 400 million/year. The statistical data published by Euro Stat in 2013 explain that between 2002 and 2011 more than 50% of the total chemical production produced the dangerous compounds and 70% out of them has a

*Corresponding author's email: iankurniawan019@gmail.com

1. Doctoral Student of Environmental Science

significant impact on the environment. Besides, human activities have affected the contamination of water resources with micropollutants biological, such as viruses and bacteria that are toxic (Xie, 2012; Periasamy and Sundaram, 2013; Sui *et al.*, 2015).

Contaminants transportation in the aquatic environment is described in Figure 1 (the model source-line-receptors) describes the path of pollution of wastewater from various sources, including hospitals, which can be harmful to the environment especially for human health and natural ecosystems. The residual chemicals which go into the exhaust system will go towards the body of water, such as river, lake or sea. On the low speed, this waste flows into the groundwater if there is a highly enough concentration or if there are gaps and cracks on the ground surface. That fact is urgent and becomes a challenge in doing particular observation and investigation which must be done to anticipate micro pollutants contaminants entering the body of water through various research and studies (Jiang *et al.*, 2013).

Hospital is one of the pollutant sources which results from wastewater from drugs. The installation

of wastewater treatment owned by hospital generally is not be able to eliminate these compounds (Verlicchi *et al.*, 2012). The residue of the chemical and pharmaceuticals from hospital wastewater can not be eliminated by the installation system and generally, it becomes residue, virus, and bacteria and then transforms as a multiresistant agent in the environment (Yamina *et al.*, 2014; Yuan *et al.*, 2015). Several studies which have been done (Zorita *et al.*, 2009; Kosma *et al.*, 2010; Behera *et al.*, 2011; Deegan *et al.*, 2011; Jelic *et al.*, 2011; Gracia-Lor *et al.*, 2012) conclude that wastewater treatment installation owned by the hospital does not have the capability to reduce wastewater containing drugs.

The activities done in the hospital will produce the residue of drugs, one of them is antibiotics. Antibiotic is an important part of treatment because it helps to treat the humans' infection (Hirsch *et al.*, 1999). Wastewater from the health services, for example, from hospital containing chemicals and drugs which are carcinogenic and genotoxic which cause cancer and genetic abnormalities (Escher *et al.*, 2010; Orias and Perrodin, 2013; Bayer *et al.*, 2014; Sharma *et al.*, 2015).




Fig. 1. Pathways of Pollution in Water (Stuart *et al.*, 2012)

Yoon *et al.* (2010) found that 31 micropollutants (Atorvastatin, Octylphenol, Gemfibrozil, Triclosan, Fluoxetine, Musk ketone, Estradiol, Progesterone, Ibuprofen, Ethynylestradiol, BHA, Testosterone, Bisphenol A, Naproxen, Benzophenone, Estrone, the TCPP, Diazepam, Dilantin, Carbamazepine, Atrazine, DEET, Trimethoprim, Sulfa-methoxazole, TCEP, Primidone, Diclofenac, Meprobamate, Caffeine, Iopromide) which are compound of the endocrine system disorder and drugs in the surface water and in the river in Seoul, South Korea in the summer and winter condition. Hamjinda *et al.* (2015) also observe the character of wastewater hospital using antibiotic compound as the research subject in the seven hospitals in Bangkok city and found some of antibiotics compound (Levofloxacin, Ofloxacin, Quinolone, Ciprofloxacin, Norfloxacin, Sulfametoksazol, Norflokasin) which have an effect on the toxicity on several types of microorganisms (*Chlorella vulgaris* and *Scenedesmus quadricauda*), and Microcrustacean (*Moina macrocopa*).

Prayitno *et al.* (2012) conduct a research about the quality of the hospital wastewater in Malang city and some of the result of the studies about wastewater treatment quality in several hospitals in Indonesia (RSUD Kelet Jepara, RSUD Dr. Muhammad Zein Painan Sumbar, RS Permadi, RSU Prov. NTB, RSUP Dr. Wahidin Sudirohusodo Sulse, RS PKT Bontang, RSUP Persahabatan) conclude that facts and data are relatively similar (in aspect Physics, Biology, Chemistry, and Radioactivity) and they do not focus on researching the content of drugs in the wastewater because of the limitations of technology and the regulation done by Indonesia Government does not regulate specifically about the contaminant material, whereas in the world, zero tolerance of antibiotics effect is being socialized.

The detection of micropollutants content in wastewater with specific containing drugs can be done using High-Performance Liquid Chromatography (HPLC) (Ren *et al.*, 2008; Dolar *et al.*, 2013; Mendoza *et al.*, 2015; Sabry *et al.*, 2015). According to Ferrer and Thurman (2012), the use of High-Resolution Mass Spectrometry (LC/Q-TOF/MS) with the principle of mass spectrometer enables the identification and characterization of more than 100 drugs and metabolite in the water source at small concentrations in a part per trillion (ppt). The examination method can provide a good sensitivity and selectivity in identifying the types of antibiotics which are unknown its group (untargeted) in low

concentrations of the water sample.

In this research, screening and identification of the type of antibiotics in the hospital wastewater has been done in Indonesia. The identification of antibiotic which is detected can be used as the recommendations to the Government of Indonesia. The recommendation explains about the new element polluters in the hospital wastewater which needs special regulation to keep the environmental sustainability.

MATERIALS AND METHODS

This research was conducted during August – December 2016 by taking wastewater hospital in one of the hospital type B in Palembang City, South Sumatra Province, Indonesia. Wastewater sample used is a mixed of all activities done in the hospital which has liquid characteristic and hazardous and toxic material (B3). The assumption which can be used for the same activities having the similarities in terms of the characteristic properties of the wastewater produced. A sample of the waste used is the wastewater hospital in reception pond before entering to the installation process of wastewater treatment with size 2.4 m x 1.9 m x 4 m and volume ± 18,2 m³. The selection of the sample research was done by using purposive sampling with the criteria the hospital which has technically wastewater treatment pond located in the underground so that the confounding variable, namely rainfall and extreme weather can be neglected.

The sampling instrumentation used vandorn sampler which can be used to a specific depth. Wastewater retrieval methods used grab method and composite. Wastewater sample mixed is taken from different points (basic, middle, and the surface of the pond) in different time span, with the same volume of 2 liters and homogenized. The sample is stored in the sample bottle of polyethylene as much as 2 liters and then sent it to the examination laboratory, Anglerbiochem Laboratory Surabaya, Indonesia.

The screening process will be done to predict qualitatively the types of antibiotics contained in the wastewater by using High-Resolution Mass Spectrometry (LC/Q-TOF/MS) 6500 Series for the Detection of Pharmaceuticals in Water.

Standard operating procedure using High-Resolution Mass Spectrophotometry (LC/Q-TOF/MS) 6500 Series to examine the hospital wastewater are as follows the analysis uses the Mobile Phase

and C18 column A (10% Methanol with 5 mM Ammonium acetate), Mobile Phase B (10% Methanol with 5 mM Ammonium acetate). The wastewater sample is prepared then done the evaporation at a temperature of 40°C until dry (residue), residue then is added to methanol solvent 50%. The process uses Vortec and Ultrasonic, Centrifuge 10,000 rpm for 5 minutes, Syringe with filter 0.22 µm, and injected to a High-Resolution Mass Spectrometry (LC/Q-TOF/MS) 6500 Series.

The use of the Agilent 6540 Q-TOF/LC/MS is an instrumentation system which is integrated with some supporting software, namely Molecular Feature Extractor, Molecular Formula Generator, and Accurate Mass Databases to detect, identify, and predict the concentration of antibiotics that exist in the water.

RESULTS AND DISCUSSION

The Qualitative Antibiotics Data Consumption in Hospital

The wastewater treatment system in Hospital X uses a combination of technology systems treatment processes in physics, chemistry and biology integratedly. The pool is made of cast concrete material and means residence time - approximately 30 minutes with a processing capacity of 1000 m³/day. The wastewater treatment process is not designed to reduce the level of medicines, especially antibiotics. Antibiotics is a pollutant compound that has characteristics and requires further processing (Amand *et al.*, 2014).

Each hospital provides services including inpatient, emergency, operations, radiology, laboratory, pharmacy, kitchen, laundry, administrative, education and training, and other supporting services. The potential discovery of residual antibiotics use that comes with wastewater from the process of hospital services is by performing sampling on the primary pond.

The activities of the Hospital X, known as a type B-Hospital, which are mainly related to medical activities will be using the medicines - either antibiotics or other medicines. Primary data collection technique of antibiotics use applies to time series method, that is a data-retrieval of antibiotic use in a period of a month or a certain period (July - September 2016), so that this data can be regarded as historical data or time series. The results of qualitative data of antibiotics consumption in

Hospital X are presented in Table 1.

Based on Table 1, the use of antibiotics in Hospital X can be classified based on the mechanism of action of the usefulness of these antibiotics, such as groups of Beta-lactam, Cephalosporins, Lincosamide, Nitroimidazole aim to inhibit the synthesis or to damage the cell wall of bacteria or protozoa. Groups of Aminoglycosides, Thiamphenicol, Tetracycline, Macrolide, Clindamycin, aim to modify or to inhibit protein synthesis. Group of Sulfonamides aims to inhibit essential enzymes in the metabolism of folate. Group of Quinolones affects the synthesis or nucleic acid metabolism.

Molecular Feature Extractor (MFE)

Molecular Feature Extractor (MFE) produces the first output from the results of wastewater samples in the form of the chromatogram of the antibiotics compound (Fig. 2, Fig. 3, Fig. 4, Fig. 5 and Fig. 6).

Molecular Formula Generator (MFG)

Molecular Formula Generator (MFG) will identify the form of the *chromatogram* in order to be matched with a particular compound formula to produce the spectrum of antibiotics (Fig. 7 Ciprofloxacin, Fig. 8 Lincomycin, Fig. 9 Metronidazole, Fig. 10 Netilmicin, and Fig. 11 Ofloxacin/Levofloxacin).

Accurate Mass Database (AMD)

The final step of the instrumentation system is Accurate Mass Database. In this step, the spectrum from MFG will be matched with the mass of molecular compounds and retention time in the library tools, then detected antibiotics types will be obtained. Five Antibiotics are detected from Hospital Wastewater in the tropical country (Table 2).


Ciprofloxacin Issue

The identification result of antibiotics type detected in waste water samples of the hospital produces 5 types of antibiotics detected from 126 types of antibiotics in which screening are conducted. The use of five types of antibiotics to patients through oral (tablets, capsules, and syrups), injection and infuse will remain the rest of antibiotics due to the improper use and disposal, hence these remains are mixed with wastewater of the hospital.

The research conducted on wastewater effluent of the hospital found that even though the wastewater treatment process had been done, but the content of

Table 1. Qualitative Data of Antibiotics Consumption in Hospital X

Type of Antibiotics	Brand of Products	Type of Use	Group
Amikacin	Amikacin, Amiosin	Injection	Aminoglycosides
Amoxicillin	Amoxicillin, Amoxsan	Syrup, pill	Beta-lactam
Amoxicillin, Clavulanat	Co Amoxiclav, Capsinat	Syrup, pill	Beta-lactam
Azitromycin	Azomax, Infimycin, Zibramax, Zistic	Syrup, Pill, Injection	Macrolide
Ampicillin, Sulbactam	Bactesyn	Injection dan Tablet	Beta-lactam
Cefadroxyl	Cefadroxyl, Cefat, Renasistin	Syrup, Pill	Cephalosporins
Cefixime	Cefixime, Fixacep, Nucef, Starcef	Syrup, Pill, drop	Cephalosporins
Cefuroxime	Anbacin	Injection	Cephalosporins
Cefotaxime	Cefotaxime, Biocef, Kalfoxim	Injection	Cephalosporins
Cefoperazone	Cefoperazone, Cefoject	Injection	Cephalosporins
Ceftriaxone	Broadced	Injection	Cephalosporins
Ceftazidime	Ceftazidime, Ceftum, Zibac	Injection	Cephalosporins
Ceftizoxime	Ceftizoxime	Injection	Cephalosporins
Ceftriaxone	Ceftriaxone, Elpicef	Injection	Cephalosporins
Clindamycin	Clindamycin	Pill	Clindamycin
Cotrimoxazol	Cotrimoxazol	Syrup, Tablet	Sulfonamides
Ciprofloxacin	Ciprofloxacin, Baquinor, Renator	Pill	Quinolones
Doxixyclin	Doxixyclin	Kapsul	Tetracycline
Erytromycin	Erytromycin, Erysanbe	Pill, Syrup	Macrolide
Kanamycin	Kanamycin	Injection	Aminoglycosides
Levofloxacin	Cravit, Volequin	Tablet, Infus	Quinolones
Lincomycin	Lincomycin	Pill, Syrup	Lincosamide
Metronidazole	Farnat	Infus	Nitroimidazole
Meropenem	Merem, Merofen, Meropenem	Injection	Beta-lactam
Netilmicin	Netromycin	Injection	Aminoglycosides
Ofloxacin	Ofloxacin, Pharflor	Pill	Quinolones
Primadex	Primadex	Pill, Syrup	Sulfonamides
Tetracycline	Tetracycline, Tetrasanbe	Pill	Tetracycline
Thiamphenicol	Thiamphenicol, Biothicol	Pill	Thiamphenicol

**Fig. 2.** Chromatogram of Ciprofloxacin

antibiotic Ciprofloxacin was still detected. Hamjinda *et al.* (2015) who conducted a research in Bangkok, Thailand found Kuinolon, Ofloxacin, Levofloxacin, Norfloksasin, Sulfametoksazol, Norfloksasin dan Ciprofloxacin. Mater *et al.* (2014) found Ciprofloxacin, Tamoxifen and Cyclo-phosphamide

in France. Furthermore, Varela *et al.* (2014) who investigated the sample from hospital in Portugal, detected Amoxicillin, Ciprofloxacin, Fluoroquinolones, Arsenic, Mercury, Metracyclines, Sulphonamides, and Penicillin G. Santos *et al.*, (2013) detected antibiotics such as Ciprofloxacin,




Fig. 3. Chromatogram of Lincomycin




Fig. 4. Chromatogram of Metronidazole




Fig. 5. Chromatogram of Netilmicin




Fig. 6. Chromatogram of Ofloxacin/Levofloxacin

ORIGINALITY REPORT

3%

SIMILARITY INDEX

0%

INTERNET SOURCES

2%

PUBLICATIONS

1%

STUDENT PAPERS

PRIMARY SOURCES

1

Submitted to University of Portsmouth

Student Paper

1%

2

Sumarto, Sylvana Novilia, Frans Van Galen, Zulkardi Zulkardi, and Darmawijoyo Darmawijoyo. "Proportional Reasoning: How do the 4th Graders Use Their Intuitive Understanding?", International Education Studies, 2013.

Publication

1%

3

Jun-ichi Yamaguchi, Yumiko Watanabe, Mari Ohmichi, Shigeji Jingu, Naoyoshi Ogawa, Jun Kokatsu, Kiyomi Fukushima, Junichi Goto. "Ultrasensitive determination of NE-100, a novel sigma ligand, in human plasma by liquid chromatography and electrospray ionization tandem mass spectrometry combined with a column-switching technique", Journal of Chromatography B: Biomedical Sciences and Applications, 1999

Publication

<1%

4

Aryansyah Batubara, Eddy Ibrahim, Subriyer Nasir, Muhammad Said, Ian Kurniawan, Adri Huda. "A New Prototype Design and Experimental Study for Assessing Spontaneous Coal Combustion", Journal of Ecological Engineering, 2019

Publication

<1%

5

iowamedicaidpdl.com

Internet Source

<1%

Exclude quotes On

Exclude matches Off

Exclude bibliography On



[Download Sample Paper](#)

ISSN: 0257-8050

Chief Editor

Prof. (Dr.) R.K.Trivedy, Pune, India

Associate Editors


Dr. S. Kaul, MOEF, New Delhi India

[Editorial Board](#)

[Download Sample Issue](#)

[Subscription Rates](#)

[Submit Article](#)



[Click here for scimagojr.com](http://www.scimagojr.com)

SCOPUS - H Index - 20

NAAS Rating - 4.97

Brief About Pollution Research

Published Quarterly Since 1982. Pollution Research is published in March, June, September and December every year.

POLLUTION RESEARCH is one of the leading enviromental journals in world and is widely subscribed in India and abroad by Institutions and Individuals in Industry, Research and Govt. Departments.

Pollution Research is abstracted/covered in:

- Chemical Abstracts, U.S.A.
- SCOPUS
- EBSCO Publishing, U.S.A.
- Cambridge Science Abstracts
- Ecology Abstracts
- Pollution Abstracts
- Eco-Disc CD Rom
- Geological Abstracts
- International Development Abstracts
- Oceanographic Literature Review
- Zoological Records
- Indian Science Abstracts, Nisclair, India
- Elsevier's Compendex
- Elsevier's Current Awareness in Biological Sciences
- Elsevier's Encompass
- Elsevier's Geobase

Pollution Research journal is accredited with National Academy of Agricultural sciences, NAAS, India.

Pollution Research journal is covered by SCOPUS.

Pollution Research journal also features in Uhlrich International Periodical Directory, U.K., Gale Directory, U.K. and SAARC directory of periodicals.

Journal Issues

- [Vol. 36, Issue 2, 2017](#)
- [Vol. 36, Issue 1, 2017](#)
- [Vol. 35, Issue 4, 2016](#)
- [Vol. 35, Issue 3, 2016](#)
- [Vol. 35, Issue 2, 2016](#)
- [Vol. 35, Issue 1, 2016](#)
- [Vol 34, Issue 4, 2015](#)
- [Vol 34, Issue 3, 2015](#)
- [Vol 34, Issue 2, 2015](#)
- [Vol 34, Issue 1, 2015](#)
- [Vol 33, Issue 04, 2014](#)
- [Vol 33, Issue 03, 2014](#)
- [Vol 33, Issue 02, 2014](#)
- [Vol.33, Issue 01, 2014](#)
- [Vol.32, Issue 04, 2013](#)
- [Vol.32, Issue 03, 2013](#)
- [Vol.32, Issue 02, 2013](#)
- [Vol.32, Issue 1, 2013](#)
- [Vol.31, Issue 04, 2012](#)
- [Vol.31, Issue 3, 2012](#)
- [Vol.31, Issue 2, 2012](#)
- [Vol.31, Issue 1, 2012](#)
- [Vol.30, Issue 4, 2011](#)
- [Vol.30, Issue 3, 2011](#)
- [Vol.30, Issue 2, 2011](#)
- [Vol.30, Issue 1, 2011](#)
- [Vol.29, Issue 4, 2010](#)
- [Vol.29, Issue 3, 2010](#)
- [Vol.29, Issue 2, 2010](#)
- [Vol.29, Issue 1, 2010](#)
- [Vol.26, Issue 3, 2007](#)
- [Vol.20, Issue 02, 2001](#)
- [Vol.20, Issue 01, 2001](#)
- [Vol.19, Issue 03, 2000](#)
- [Vol.19, Issue 01, 2000](#)
- [Vol.18, Issue 02, 1999](#)
- [Vol.17, Issue 04, 1998](#)
- [Vol.17, Issue 03, 1998](#)
- [Vol.17, Issue 01, 1998](#)



EM INTERNATIONAL

C-101, Prakriti, Balewadi, Baner,
Pune 411 045, Maharashtra, India

☎ 91-20-46745119, 9326712297

Email: rktem@pn3.vsnl.net.in

Website: www.envirobiotechjournals.com

INTERNATIONAL JOURNALS

POLLUTION RESEARCH	Quarterly	ISSN 0257 - 8050
JR.OF INDUSTRIAL POLLUTION CONTROL	Biannual	ISSN 0970 - 2083
ECOLOGY, ENVIRONMENT AND CONSERVATION	Quarterly ISSN	0971 - 765 X

I. Kurniawan^{1,2}, S. Nasir³, Hermansyah⁴ and Mardiyanto⁴

Date: 15.3.2017

¹ Doctoral Student of Environmental Science,

Ref.: PR-F-170

Sriwijaya University, Palembang, Indonesia

² Faculty of Health Science, Musi Charitas Catholic University,
Palembang, Indonesia

³ Faculty of Engineering, Sriwijaya University, Palembang, Indonesia

⁴ Faculty of Mathematics and Science, Sriwijaya University,
Palembang, Indonesia

Dear Authors,

The Editor thanks for your manuscript entitled –

- **The Screening of Potential Antibiotics from Hospital Wastewater in Tropical Region
(Case Study at Palembang, South Sumatra, Indonesia)**

Submitted for publication in **POLLUTION RESEARCH**. Pl. always quote **Ref. No.** –
PR - F - 170, while doing any correspondence with us.

Editor is pleased to inform you that your paper is accepted for publication in **POLLUTION RESEARCH** in the year **2017 Issue no. (2)**.

Kindly note the journal has **SCOPUS h index 18.00** and **NAAS India impact rating 4.97**
(www.envirobiotechjournals.com).

Thanking you & with regards.

Yours Sincerely

Publisher

THE SCREENING OF POTENTIAL ANTIBIOTICS FROM HOSPITAL WASTEWATER IN TROPICAL REGION (CASE STUDY AT PALEMBANG, SOUTH SUMATRA, INDONESIA)

I. KURNIAWAN, S. NASIR, HERMANSYAH AND MARDIYANTO

Abstract

Clean water is a primary one of the causes of water quality degradation is pollution through the residue of human activity. Hospital is a potential pollutant source with the characteristics of hazardous and toxic wastewater. The components of pollutants in the wastewater of hospital is still limited investigated is antibiotic. The regulation of hospital wastewater in Indonesia as tropical and developing countries do not regulate antibiotics, while in the world the issue of zero tolerance of antibiotics becomes an important concern. Antibiotics have a persistent and resistant impact for the environment. In this research, screening for antibiotics by using samples from one type B hospital in Palembang City, Indonesia is done. Hospital wastewater samples in the non permanent pond are the residue of all process activities mixed in one pond. Screening using High-Resolution Mass Spectrometry (LC / Q-TOF / MS) 6500 Series for the Detection of Pharmaceuticals in Water is used to detect the antibiotics. The result of laboratory analysis shows that there are five types of antibiotics detected, namely Ciprofloxacin, Lincomycin, Metronidazole, Netilmicin, and Ofloxacin/Levofloxacin. The qualitative analysis concluded that Ciprofloxacin is one of type potential antibiotics in the hospital wastewater. The result of this research recommends that the Indonesian Government should revise the regulation about antibiotics in the hospital wastewater.

Enter your contact information below to receive full paper.

Your Name :

Email:

Phone:

City:

Cost of Full Paper: Rs.75 for Indian Nationals or \$20 (USD) for international subscribers.

By clicking on Request Paper you Agree to pay the above mentioned cost per paper.

[Request Paper](#)

[Back](#)

Journal Issues

[Vol. 36, Issue 2, 2017](#)

[Vol. 36, Issue 1, 2017](#)

[Vol. 35, Issue 4, 2016](#)

[Vol. 35, Issue 3, 2016](#)

[Vol. 35, Issue 2, 2016](#)

[Vol. 35, Issue 1, 2016](#)

[Vol. 34, Issue 4, 2015](#)

[Vol. 34, Issue 3, 2015](#)

[Vol. 34, Issue 2, 2015](#)

[Vol. 34, Issue 1, 2015](#)

[Vol. 33, Issue 04, 2014](#)

[Vol. 33, Issue 03, 2014](#)

[Vol. 33, Issue 02, 2014](#)

[Vol. 33, Issue 01, 2014](#)

[Vol. 32, Issue 04, 2013](#)

[Vol. 32, Issue 03, 2013](#)

[Vol. 32, Issue 02, 2013](#)

[Vol. 32, Issue 1, 2013](#)

[Vol. 31, Issue 04, 2012](#)

[Vol. 31, Issue 3, 2012](#)

[Vol. 31, Issue 2, 2012](#)

[Vol. 31, Issue 1, 2012](#)

[Vol. 30, Issue 4, 2011](#)

[Vol. 30, Issue 3, 2011](#)

[Vol. 30, Issue 2, 2011](#)

[Vol. 30, Issue 1, 2011](#)

[Vol. 29, Issue 4, 2010](#)

[Vol. 29, Issue 3, 2010](#)

[Vol. 29, Issue 2, 2010](#)

[Vol. 29, Issue 1, 2010](#)

[Vol. 26, Issue 3, 2007](#)

[Vol. 20, Issue 02, 2001](#)

[Vol. 20, Issue 01, 2001](#)

[Vol. 19, Issue 03, 2000](#)

[Vol. 19, Issue 01, 2000](#)

[Vol. 18, Issue 02, 1999](#)

[Vol. 17, Issue 04, 1998](#)

[Vol. 17, Issue 03, 1998](#)

[Vol. 17, Issue 01, 1998](#)

Looking for Past Issues?

[Click here to get them!!](#)

POLLUTION RESEARCH

VOL. 36 (2) : 2017

CONTENTS

- 159–167 Anaerobic Digestion of Dewatered Primary Sludge (DPMS) from the Nagaon Paper Mill Morigaon Assam
—*Niro Akbary, Ajay Kalamdhad and Manfred Koch*
- 168–174 Heavy Metals in Wild Indonesian Shortfin Eel, *Anguilla Bicolor Bicolor* (McClelland 1844), and Giant Mottled Eel, *A. Marmorata* (Quoy & Gaimard 1824), in The Northwest of Peninsular Malaysia
—*N.A. Khalil, M.N.A. Amal, A. Ismail, S.Z. Zulkifli and F. Rahman*
- 175–184 Adsorption-assisted Electrocoagulation of Real Car Wash Wastewater with Equilibrium and Kinetic Studies
—*Humberto Rubí-Juárez, Carlos Barrera-Díaz and Fernando Ureña-Nuñez*
- 185–196 Adsorption Isotherm Models Test by Error Function Analysis in the Removal of Aromatic Compounds and Colour Using Zeliac
—*Nurazim Ibrahim, Hamidi Abdul Aziz and Mohd Suffian Yusoff*
- 197–204 Optimizing Conditions for Crude oil Degrading Bacterial Consortium Isolated from Aquatic Environment
—*M. Azab El-liethy, Bahaa A. Hemdan, Farag A. Samhan, Sahar S. Ali and Gamila E. El-taweel*
- 205–211 The Waste Management of Clothing Home Industries in Makassar City, Indonesia
—*Hamidah Suryani, Gufran Darma Dirawan, Muh. Yahya and Suradi Tahmir*
- 212–219 Spatial Linear Programming Model to Reduce a Pollution Emissions of Sugarcane Transportation in Northeastern Thailand
—*Patiwat Littidej*
- 220–229 Monthly Average Spatial Interpolation for Air Pollutants in Egypt
—*Sara Osama Ahmed, Reda Mazloum and Hala Abou-Ali*
- 230–240 Decolorization and Detoxification of Methylene Blue by A Lebanese Bacterial Strain; *Bacillus Licheniformis* He
—*HodaYusef, Salwa Abdel Rahman, Enas Dabbous and Hassan Hammud*
- 241–248 Air Pollution Assessments Using Lichen Biodiversity Index (LBI) in Kuala Lumpur, Malaysia
—*Azlan Abas and Azahan Awang*
- 249–252 Bioremediation of Formalin by *Ricinus Communis*
—*I. Jeevan Madhukar and C.V. Narasimha Murthy*
- 253–259 Assessment of Groundwater Quality and Mapping Using Gis- A Case Study
—*P. Eshanthini, Christina Mary. V and M. Kalaiivani*
- 260–263 Impact of Building Construction-dust on Foliar Biochemistry of *Delonix Regia* (Boj.ex.hook.) Raf. and *Peltophorum Pterocarpum* (Dc) Backer.
—*Vaishnavi V. More*
- 264–270 Investigation of Dominant Leaching Factors for Release of Cr from Indian Fly Ash
—*Manoj Kumar Tiwari, Samir Bajpai and Umesh Kumar Dewangan*
- 271–276 Relation of Groundwater Arsenic, Iron, and Redox-sensitive Manganese to Aquifer Type in Barak Basin, Assam, India
—*Tushar Deb Kanungo*
- 277–281 Organophosphates and Carbamates as Inhibitors of Acetylcholinesterase in *Eisenia fetida*
—*Jyoti Yadav, Dharambir Singh, Jayant Yadav and Deepak Kumar*
- 282–291 Environmental Analysis and Treatment of Effluents from Agro-based Paper Mills
—*Sangeeta Sharma, Sarabjit Singh and Harleen Kaur*
- 292–295 Effects of Cd and Zn Toxicity on Biomolecules of *Eisenia fetida*
—*Jyoti Yadav and R.K. Gupta*
- 296–305 Tourism Impacts on the Water Quality of Bhimtal Lake in Central Himalaya
—*Rajesh Kumar Lohchab, Devendra K. Agrawal and C.P. Kaushik*

(Continued on Inside Back Cover)

POLLUTION RESEARCH

VOL. 36 (2) : 2017

(Contents Continued from Back Cover)

- 306–311 Characterization and Optimization of Physicochemical Properties of Dairy Effluents
—Asha Rani Garg, Seema Garcha, Ashish Kumar Singh, Kamaldeep and Ankit Magotra
- 312–318 Evaluation of Antioxidant Enzyme Responses in Earthworms Exposed to Polystyrene Nano Particles and Pamam Dendrimers
—B. Siva Prasad, P. Sankar Ganesh, K. Lagan, T. Vishaka, S.S. Vutukuru and S.K. Sarkar
- 319–328 Assessment of Heavy Metal Contents in Vegetables Grown in the Vicinity of Sirsa and their Effects on Human Health
—Sangeeta Sharma, Davender Kumar and Jitender Pal
- 329–332 Mycorrhiza Assisted Approach for Bioremediation With Special Reference to Biosorption
—Shweta Pathak, Prasann Kumar, Pankaj Kumar Mishra and Mukul Kumar
- 333–339 Phytoremediation Potential of Some Native Plants Growing in the Vicinity of Sponge Iron Industry
—Sunanda Sahoo
- 340–342 Physico-chemical Status of Bhima, Bhama and Indrayani River Flowing Through Khed Taluka, Rajgurunagar, Pune District
—S.S. Kolekar
- 343–351 The Screening of Potential Antibiotics from Hospital Wastewater in Tropical Region (Case Study at Palembang, South Sumatra, Indonesia)
—I. Kurniawan, S. Nasir, Hermansyah and Mardiyanto
- 352–358 Adsorption Characteristics of Fe²⁺ on Mixed Micelles of Sodium Dodecyl Benzene Sulphonate Using Micellar Enhanced Ultrafiltration
—C.S. Nura, A. Chattree and R.P. Singh
- 359–365 Efficient Ecofriendly Bioremediation of Crude Oil Waste by Extremophilic Marine *Streptomyces Indiaensis* Act 7 and *Streptomyces Hygroscopicus* ACT 14
—B. Sathya Priya, T. Stalin and K. Selvam
- 366–372 Bioaccumulation and Health Risks Assessment of Lead (Pb) in Freshwater Asian Clams (*Corbicula fluminea*, Müller) from Laguna de Bay, Philippines
—Christian Paul P. de la Cruz, Nestor M. De Vera, Lustina P. Lapie, Ma. Neda A. Catalma and Rosalina V. Bunal
- 373–379 Production of Biodiesel from Lipase Mediated Trans-esterification by Reducing free Fatty Acid Content of Algal Oil Produced from *Botryococcus Braunii*
—Shivandappa, Pushpa Agrwal, V. Krishna and S. Vidyashree

THE SCREENING OF POTENTIAL ANTIBIOTICS FROM HOSPITAL WASTEWATER IN TROPICAL REGION (CASE STUDY AT PALEMBANG, SOUTH SUMATRA, INDONESIA)

I. KURNIAWAN^{1,2}, S. NASIR³, HERMANSYAH⁴ AND MARDIYANTO⁴

¹*Sriwijaya University, Palembang, Indonesia*

²*Faculty of Health Science, Musi Charitas Catholic University, Palembang, Indonesia*

³*Faculty of Engineering, Sriwijaya University, Palembang, Indonesia*

⁴*Faculty of Mathematics and Science, Sriwijaya University, Palembang, Indonesia*

(Received 1 February, 2017; accepted 15 March, 2017)

ABSTRACT

Clean water is a primary one of the causes of water quality degradation is pollution through the residue of human activity. Hospital is a potential pollutant source with the characteristics of hazardous and toxic wastewater. The components of pollutants in the wastewater of hospital is still limited investigated is antibiotic. The regulation of hospital wastewater in Indonesia as tropical and developing countries do not regulate antibiotics, while in the world the issue of zero tolerance of antibiotics becomes an important concern. Antibiotics have a persistent and resistant impact for the environment. In this research, screening for antibiotics by using samples from one type B hospital in Palembang City, Indonesia is done. Hospital wastewater samples in the non permanent pond are the residue of all process activities mixed in one pond. Screening using High-Resolution Mass Spectrometry (LC / Q-TOF / MS) 6500 Series for the Detection of Pharmaceuticals in Water is used to detect the antibiotics. The result of laboratory analysis shows that there are five types of antibiotics detected, namely Ciprofloxacin, Lincomycin, Metronidazole, Netilmicin, and Ofloxacin/Levofloxacin. The qualitative analysis concluded that Ciprofloxacin is one of a type potential antibiotics in the hospital wastewater. The result of this research recommends that the Indonesian Government should revise the regulation about antibiotics in the hospital wastewater.

KEY WORDS : Clean water, Environment, Antibiotics, Hospital wastewater, Ciprofloxacin

INTRODUCTION

World Health Statistics reported by WHO in 2015 shows that about 85% of the Indonesian population needs access to clean water for consumption. The community in the late decades concerns the importance to solve the environmental problems, especially about clean water conservation. International Water Institute predicts in 2025 Java and the others Islands in Indonesia will have a crisis of clean water. The water global scarcity has increased and is needed the other sources to make it balance with the water need because existing

water sources has not been used because of the pollution and wastewater. The problem of clean water sources because of containing components of polluters in high concentrations, such as heavy metals, toxic compounds, and microbial pathogens.

The data released by WWF world organization in its report explain that the data between 1930 to 2000 about the production of chemicals caused by human activities increased from 1 million to 400 million/year. The statistical data published by Euro Stat in 2013 explain that between 2002 and 2011 more than 50% of the total chemical production produced the dangerous compounds and 70% out of them has a

*Corresponding author's email: iankurniawan019@gmail.com

1. Doctoral Student of Environmental Science

significant impact on the environment. Besides, human activities have affected the contamination of water resources with micropollutants biological, such as viruses and bacteria that are toxic (Xie, 2012; Periasamy and Sundaram, 2013; Sui *et al.*, 2015).

Contaminants transportation in the aquatic environment is described in Figure 1 (the model source-line-receptors) describes the path of pollution of wastewater from various sources, including hospitals, which can be harmful to the environment especially for human health and natural ecosystems. The residual chemicals which go into the exhaust system will go towards the body of water, such as river, lake or sea. On the low speed, this waste flows into the groundwater if there is a highly enough concentration or if there are gaps and cracks on the ground surface. That fact is urgent and becomes a challenge in doing particular observation and investigation which must be done to anticipate micro pollutants contaminants entering the body of water through various research and studies (Jiang *et al.*, 2013).

Hospital is one of the pollutant sources which results from wastewater from drugs. The installation

of wastewater treatment owned by hospital generally is not be able to eliminate these compounds (Verlicchi *et al.*, 2012). The residue of the chemical and pharmaceuticals from hospital wastewater can not be eliminated by the installation system and generally, it becomes residue, virus, and bacteria and then transforms as a multiresistant agent in the environment (Yamina *et al.*, 2014; Yuan *et al.*, 2015). Several studies which have been done (Zorita *et al.*, 2009; Kosma *et al.*, 2010; Behera *et al.*, 2011; Deegan *et al.*, 2011; Jelic *et al.*, 2011; Gracia-Lor *et al.*, 2012) conclude that wastewater treatment installation owned by the hospital does not have the capability to reduce wastewater containing drugs.

The activities done in the hospital will produce the residue of drugs, one of them is antibiotics. Antibiotic is an important part of treatment because it helps to treat the humans' infection (Hirsch *et al.*, 1999). Wastewater from the health services, for example, from hospital containing chemicals and drugs which are carcinogenic and genotoxic which cause cancer and genetic abnormalities (Escher *et al.*, 2010; Orias and Perrodin, 2013; Bayer *et al.*, 2014; Sharma *et al.*, 2015).




Fig. 1. Pathways of Pollution in Water (Stuart *et al.*, 2012)

Yoon *et al.* (2010) found that 31 micropollutants (Atorvastatin, Octylphenol, Gemfibrozil, Triclosan, Fluoxetine, Musk ketone, Estradiol, Progesterone, Ibuprofen, Ethynylestradiol, BHA, Testosterone, Bisphenol A, Naproxen, Benzophenone, Estrone, the TCPP, Diazepam, Dilantin, Carbamazepine, Atrazine, DEET, Trimethoprim, Sulfa-methoxazole, TCEP, Primidone, Diclofenac, Meprobamate, Caffeine, Iopromide) which are compound of the endocrine system disorder and drugs in the surface water and in the river in Seoul, South Korea in the summer and winter condition. Hamjinda *et al.* (2015) also observe the character of wastewater hospital using antibiotic compound as the research subject in the seven hospitals in Bangkok city and found some of antibiotics compound (Levofloxacin, Ofloxacin, Quinolone, Ciprofloxacin, Norfloxacin, Sulfametoksazol, Norfloxacin) which have an effect on the toxicity on several types of microorganisms (*Chlorella vulgaris* and *Scenedesmus quadricauda*), and Microcrustacean (*Moina macrocopa*).

Prayitno *et al.* (2012) conduct a research about the quality of the hospital wastewater in Malang city and some of the result of the studies about wastewater treatment quality in several hospitals in Indonesia (RSUD Kelet Jepara, RSUD Dr. Muhammad Zein Painan Sumbar, RS Permadi, RSU Prov. NTB, RSUP Dr. Wahidin Sudirohusodo Sulsel, RS PKT Bontang, RSUP Persahabatan) conclude that facts and data are relatively similar (in aspect Physics, Biology, Chemistry, and Radioactivity) and they do not focus on researching the content of drugs in the wastewater because of the limitations of technology and the regulation done by Indonesia Government does not regulate specifically about the contaminant material, whereas in the world, zero tolerance of antibiotics effect is being socialized.

The detection of micropollutants content in wastewater with specific containing drugs can be done using High-Performance Liquid Chromatography (HPLC) (Ren *et al.*, 2008; Dolar *et al.*, 2013; Mendoza *et al.*, 2015; Sabry *et al.*, 2015). According to Ferrer and Thurman (2012), the use of High-Resolution Mass Spectrometry (LC/Q-TOF/MS) with the principle of mass spectrometer enables the identification and characterization of more than 100 drugs and metabolite in the water source at small concentrations in a part per trillion (ppt). The examination method can provide a good sensitivity and selectivity in identifying the types of antibiotics which are unknown its group (untargeted) in low

concentrations of the water sample.

In this research, screening and identification of the type of antibiotics in the hospital wastewater has been done in Indonesia. The identification of antibiotic which is detected can be used as the recommendations to the Government of Indonesia. The recommendation explains about the new element polluters in the hospital wastewater which needs special regulation to keep the environmental sustainability.

MATERIALS AND METHODS

This research was conducted during August – December 2016 by taking wastewater hospital in one of the hospital type B in Palembang City, South Sumatra Province, Indonesia. Wastewater sample used is a mixed of all activities done in the hospital which has liquid characteristic and hazardous and toxic material (B3). The assumption which can be used for the same activities having the similarities in terms of the characteristic properties of the wastewater produced. A sample of the waste used is the wastewater hospital in reception pond before entering to the installation process of wastewater treatment with size 2.4 m x 1.9 m x 4 m and volume $\pm 18,2 \text{ m}^3$. The selection of the sample research was done by using purposive sampling with the criteria the hospital which has technically wastewater treatment pond located in the underground so that the confounding variable, namely rainfall and extreme weather can be neglected.

The sampling instrumentation used vandorn sampler which can be used to a specific depth. Wastewater retrieval methods used grab method and composite. Wastewater sample mixed is taken from different points (basic, middle, and the surface of the pond) in different time span, with the same volume of 2 liters and homogenized. The sample is stored in the sample bottle of polyethylene as much as 2 liters and then sent it to the examination laboratory, Anglerbiochem Laboratory Surabaya, Indonesia.

The screening process will be done to predict qualitatively the types of antibiotics contained in the wastewater by using High-Resolution Mass Spectrometry (LC/Q-TOF/MS) 6500 Series for the Detection of Pharmaceuticals in Water.

Standard operating procedure using High-Resolution Mass Spectrophotometry (LC/Q-TOF/MS) 6500 Series to examine the hospital wastewater are as follows the analysis uses the Mobile Phase

and C18 column A (10% Methanol with 5 mM Ammonium acetate), Mobile Phase B (10% Methanol with 5 mM Ammonium acetate). The wastewater sample is prepared then done the evaporation at a temperature of 40°C until dry (residue), residue then is added to methanol solvent 50%. The process uses Vortec and Ultrasonic, Centrifuge 10,000 rpm for 5 minutes, Syringe with filter 0.22 µm, and injected to a High-Resolution Mass Spectrometry (LC/Q-TOF/MS) 6500 Series.

The use of the Agilent 6540 Q-TOF/LC/MS is an instrumentation system which is integrated with some supporting software, namely Molecular Feature Extractor, Molecular Formula Generator, and Accurate Mass Databases to detect, identify, and predict the concentration of antibiotics that exist in the water.

RESULTS AND DISCUSSION

The Qualitative Antibiotics Data Consumption in Hospital

The wastewater treatment system in Hospital X uses a combination of technology systems treatment processes in physics, chemistry and biology integratedly. The pool is made of cast concrete material and means residence time - approximately 30 minutes with a processing capacity of 1000 m³/day. The wastewater treatment process is not designed to reduce the level of medicines, especially antibiotics. Antibiotics is a pollutant compound that has characteristics and requires further processing (Aamand *et al.*, 2014).

Each hospital provides services including inpatient, emergency, operations, radiology, laboratory, pharmacy, kitchen, laundry, administrative, education and training, and other supporting services. The potential discovery of residual antibiotics use that comes with wastewater from the process of hospital services is by performing sampling on the primary pond.

The activities of the Hospital X, known as a type B-Hospital, which are mainly related to medical activities will be using the medicines - either antibiotics or other medicines. Primary data collection technique of antibiotics use applies to time series method, that is a data-retrieval of antibiotic use in a period of a month or a certain period (July – September 2016), so that this data can be regarded as historical data or time series. The results of qualitative data of antibiotics consumption in

Hospital X are presented in Table 1.

Based on Table 1, the use of antibiotics in Hospital X can be classified based on the mechanism of action of the usefulness of these antibiotics, such as groups of Beta-lactam, Cephalosporins, Lincosamide, Nitroimidazole aim to inhibit the synthesis or to damage the cell wall of bacteria or protozoa. Groups of Aminoglycosides, Thiamphenicol, Tetracycline, Macrolide, Clindamycin, aim to modify or to inhibit protein synthesis. Group of Sulfonamides aims to inhibit essential enzymes in the metabolism of folate. Group of Quinolones affects the synthesis or nucleic acid metabolism.

Molecular Feature Extractor (MFE)

Molecular Feature Extractor (MFE) produces the first output from the results of wastewater samples in the form of the chromatogram of the antibiotics compound (Fig. 2, Fig. 3, Fig. 4, Fig. 5 and Fig. 6).

Molecular Formula Generator (MFG)

Molecular Formula Generator (MFG) will identify the form of the *chromatogram* in order to be matched with a particular compound formula to produce the spectrum of antibiotics (Fig. 7 Cyprofloxacin, Fig. 8 Lincomycin, Fig. 9 Metronidazole, Fig. 10 Netilmicin, and Fig. 11 Ofloxacin/Levofloxacin).

Accurate Mass Database (AMD)

The final step of the instrumentation system is Accurate Mass Database. In this step, the spectrum from MFG will be matched with the mass of molecular compounds and retention time in the library tools, then detected antibiotics types will be obtained. Five Antibiotics are detected from Hospital Wastewater in the tropical country (Table 2).


Ciprofloxacin Issue

The identification result of antibiotics type detected in waste water samples of the hospital produces 5 types of antibiotics detected from 126 types of antibiotics in which screening are conducted. The use of five types of antibiotics to patients through oral (tablets, capsules, and syrups), injection and infuse will remain the rest of antibiotics due to the improper use and disposal, hence these remains are mixed with wastewater of the hospital.

The research conducted on wastewater effluent of the hospital found that even though the wastewater treatment process had been done, but the content of

Table 1. Qualitative Data of Antibiotics Consumption in Hospital X

Type of Antibiotics	Brand of Products	Type of Use	Group
Amikacin	Amikacin, Amiosin	Injection	Aminoglycosides
Amoxicillin	Amoxicillin, Amoxsan	Syrup, pill	Beta-lactam
Amoxicillin, Clavulanat	Co Amoxiclav, Capsinat	Syrup, pill	Beta-lactam
Azitromycin	Azomax, Infimycin, Zibramax, Zistic	Syrup, Pill, Injection	Macrolide
Ampicillin, Sulbactam	Bactesyn	Injection dan Tablet	Beta-lactam
Cefadroxy	Cefradoxyl, Cefat, Renasistin	Syrup, Pill	Cephalosporins
Cefixime	Cefixime, Fixacep, Nucef, Starcef	Syrup, Pill, drop	Cephalosporins
Cefuroxime	Anbacim	Injection	Cephalosporins
Cefotaxime	Cefotaxime, Biocef, Kalfoxim	Injection	Cephalosporins
Cefoperazone	Cefoperazone, Cefoject	Injection	Cephalosporins
Ceftriaxone	Broadced	Injection	Cephalosporins
Ceftazidime	Ceftazidime, Ceftum, Zibac	Injection	Cephalosporins
Ceftizoxime	Ceftizoxime	Injection	Cephalosporins
Ceftriaxone	Ceftriaxone, Elpicef	Injection	Cephalosporins
Clindamycin	Clindamycin	Pill	Clindamycin
Cotrimoxazol	Cotrimoxazol	Syrup, Tablet	Sulfonamides
Ciprofloxacin	Ciprofloxacin, Baquinor, Renator	Pill	Quinolones
Doxicyclin	Doxicyclin	Kapsul	Tetracycline
Erytromycin	Erytromycin, Erysanbe	Pill, Syrup	Macrolide
Kanamycin	Kanamycin	Injection	Aminoglycosides
Levofloxacin	Cravit, Volequin	Tablet, Infus	Quinolones
Lincomycin	Lincomycin	Pill, Syrup	Lincosamide
Metronidazole	Farnat	Infus	Nitroimidazole
Meropenem	Merem, Merofen, Meropenem	Injection	Beta-lactam
Netilmicin	Netromycin	Injection	Aminoglycosides
Ofloxacin	Ofloxacin, Pharflox	Pill	Quinolones
Primadex	Primadex	Pill, Syrup	Sulfonamides
Tetracycline	Tetracycline, Tetrasanbe	Pill	Tetracycline
Thiamphenicol	Thiamphenicol, Biothicol	Pill	Thiamphenicol

**Fig. 2.** Chromatogram of Ciprofloxacin

antibiotic Ciprofloxacin was still detected. Hamjinda *et al.* (2015) who conducted a research in Bangkok, Thailand found Kuinolon, Ofloxacin, Levofloxacin, Norfloksasin, Sulfametoksazol, Norfloksasin dan Ciprofloxacin. Mater *et al.* (2014) found Ciprofloxacin, Tamoxifen and Cyclo-phosphamide

in France. Furthermore, Varela *et al.* (2014) who investigated the sample from hospital in Portugal, detected Amoxicillin, Ciprofloxacin, Fluoroquinolones, Arsenic, Mercury, Metracyclines, Sulphonamides, and Penicillin G. Santos *et al.*, (2013) detected antibiotics such as Ciprofloxacin,




Fig. 3. Chromatogram of Lincomycin




Fig. 4. Chromatogram of Metronidazole




Fig. 5. Chromatogram of Netilmicin




Fig. 6. Chromatogram of Ofloxacin/Levofloxacin




Fig. 7. Spectrum of Ciprofloxacin




Fig. 8. Spectrum of Lincomycin




Fig. 9. Spectrum of Metronidazole

Table 2. The Results of Antibiotics Detection in the Wastewater Sample of Hospital X

Antibiotics	RT	Mass (m/z)	MW	Err/mDA	mSigma	Intenstiy	CF
Ciprofloxacin	4,38	C ₁₇ H ₁₉ FN ₃ O ₃ ¹⁺	332,140496	0,2	9,4	9,00E+04	100
Lincomycin	4,06	C ₁₈ H ₃₅ N ₂ O ₆ S ¹⁺	407,221034	-0,7	10,9	8,00E+05	100
Metronidazole	3,64	C ₆ H ₁₀ N ₃ O ₃ ¹⁺	172,071668	-0,2	12,1	3,00E+04	100
Netilmicin	4,74	C ₂₁ H ₄₂ N ₅ O ₇ ¹⁺	476,307875	1,4	14,5	3,00E+06	100
Ofloxacin/Levofloxacin	4,25	C ₁₈ H ₂₁ FN ₃ O ₄ ¹⁺	362,151061	-0,4	16,8	1,50E+05	100

Ofloxacin, Sulfamethoxazole, Azithromycin, Clarithro-mycin, Acetaminophen and Ibuprofen in Spain.

Identification of Ciprofloxacin detected in the hospital wastewater was due to the level of Ciprofloxacin use rate is very high. Ciprofloxacin has a broad spectrum, active against both negative and positive gram-bacteria. Ciprofloxacin is used for the treatment of respiratory tract infections, urinary tract, gastrointestinal and abdominal infections, including infections by negative gram-bacteria (*Escherichia coli*, *Haemophilus influenza*, *Klebsiella pneumoniae*, *Legionella pneumophila*, *Moraxella catarrhalis*, *Proteus mirabilis*, and *Pseudomonas aeruginosa*) and gram-positive (*Staphylococcus aureus*, *Streptococcus pneumonia*, *Staphylococcus epidermis*, *Enterococcus faecalis*, and *Streptococcus pyogenes*).

Results of laboratory examination of the content of some antibiotics in hospital wastewater showed that Ciprofloxacin has the highest intensity 9×10^4 . Intensity displays the estimation of the amount of Ciprofloxacin concentration as the highest antibiotic compounds in hospital wastewater. Considering the err-value (MDA) of the next data and Sigma, Ciprofloxacin has the smallest value that can be inferred that level of sensitivity and selectivity of detection of these compounds are better.

CONCLUSION

Hospital activities have the potential to generate waste water containing antibiotics. The rest of antibiotics use in health services of the hospital will be mixed with other remaining waste products which are then included into wastewater treatment plants and caused a negative impact on the environment. Early detection of identifying antibiotics contained in the hospital wastewater is required to provide input and suggestion to the regulations applicable in Indonesia. Agilent 6540 Q-TOF/LC /MS is an instrument that has a selectivity and excellent sensitivity to identify qualitatively untargeted antibiotics. There are 5 types of antibiotics detected in hospital wastewater where Ciprofloxacin has the highest concentration estimation due to the intensity value of these antibiotics.

ACKNOWLEDGEMENTS

The authors are very thankful to PT. Angler Biochem Laboratory in Surabaya, Indonesia for their supports

in terms of their laboratory that gives the final results of antibiotics screening process of the hospital wastewater. The authors are also thankful to their staffs: Mr. Bastian and Mr. Widodo for their contribution during the sampling process. We appreciate the Indonesian Directorate General of Higher Education (DIKTI) for the financial support granted based on research of Doctoral Dissertation.

REFERENCES

- Aamand, J., Gavrilescu, M., Agathos, S. and Fava, F. 2014. Emerging Pollutants in the Environment: Present and Future Challenges in Biomonitoring, Ecological Risks and Bioremediation. *New Biotechnology*. 1-10.
- Bayer, A., Asner, R., Schüssler, W., Kopf, W., Weib, K., Sengl, M. and Letzel, M. 2014. Behavior of sartans (antihypertensive drugs) in wastewater treatment plants , their occurrence and risk for the aquatic environment. *Environ Sci Pollut Res*. 21(18) : 10830-9.
- Behera, S. K., Kim, H. W., Oh, J. E. and Park, H. S. 2011. Occurrence and removal of antibiotics, hormones and several other pharmaceuticals in wastewater treatment plants of the largest industrial city of Korea. *Science of the Total Environment* (Volume 409). Elsevier B.V.
- Deegan, A. M., Shaik, B., Nolan, K., Urell, K., Tobin, J. and Morrissey, A. 2011. Treatment options for wastewater effluents from pharmaceutical companies. *International Journal Science Technology*. 8 (3) : 649-666.
- Dolar, D., Košutiæ, K., Periša, M. and Babiaæ, S. 2013. Photolysis of enrofloxacin and removal of its photodegradation products from water by reverse osmosis and nanofiltration membranes. *Separation and Purification Technology*. 115 : 1-8.
- Escher, B. I., Baumgartner, R., Koller, M., Treyer, K., Lienert, J. and Mcardell, C. S. 2010. Environmental toxicology and risk assessment of pharmaceuticals from hospital wastewater. *Water Research*. 45 (1) : 75-92.
- Ferrer, I. and Thurman, E.M. 2012. Analysis of 100 pharmaceuticals and their degradates in water samples by liquid chromatography/quadrupole time-of-flight mass spectrometry. *J. Chromatography A*. 1259 : 148-157.
- Gracia-Ior, E., Sancho, J. V., Serrano, R. and Hernández, F. 2012. Chemosphere Occurrence and removal of pharmaceuticals in wastewater treatment plants at the Spanish Mediterranean area of Valencia. *Chemosphere*. 87 (5) : 453-462.
- Hamjinda, N. S., Chiemchaisri, W., Watanabe, T. and Honda, R. 2015. Toxicological assessment of hospital wastewater in different treatment

- processes. *Environmental Science and Pollution Research*. Springer-Verlag Berlin Heidelberg.
- Hirsch, R., Ternes, T., Haberer, K. and Kratz, K. 1999. Occurrence of antibiotics in the aquatic environment. *The Science of the Total Environment*. 109-118.
- Jelic, A., Gros, M., Ginebreda, A., Cespedes-Sánchez, R., Ventura, F., Petrovic, M. and Barcelo, D. 2011. Occurrence, partition and removal of pharmaceuticals in sewage water and sludge during wastewater treatment. *Water Research*. 45 (3) : 1165-1176.
- Jiang, J., Zhou, Z. and Sharma, V. K. 2013. Occurrence, transportation, monitoring and treatment of emerging micro-pollutants in waste water- A review from global views. *Microchemical Journal*.
- Kosma, C. I., Lambropoulou, D.A. and Albanis, T. 2010. Occurrence and removal of PPCPs in municipal and hospital wastewaters in Greece. *Journal of Hazardous Materials*. 179 (3) : 804-17.
- Mater, N., Geret, F., Castillo, L., Faucet-marquis, V., Albasi, C. and Pfohl-leszkowicz, A. 2014. In vitro tests aiding ecological risk assessment of ciprofloxacin, tamoxifen and cyclophosphamide in range of concentrations released in hospital wastewater and surface water. *Environment International*. 63 : 191-200.
- Mendoza, A., Aceña, J., Pérez, S., Alda, M. L. De, Barceló, D., and Gil, A. 2015. Pharmaceuticals and iodinated contrast media in a hospital waste-water: A case study to analyse their presence and characterise their environmental risk and hazard. *Environmental Research*. 140 : 225-241.
- Orias, F., and Perrodin, Y. 2013. Science of the Total Environment Characterisation of the ecotoxicity of hospital effluents: A review. *Science of The Total Environment*. 455 : 250-276.
- Periasamy, D. and Sundaram, A. 2013. A novel approach for pathogen reduction in wastewater treatment. *Journal of Environmental Science and Engineering*, 1-9.
- Prayitno, Kusuma, Z., Yanuwadi, B., and Laksmono, R. W. 2012. Study of Hospital Wastewater Characteristic in Malang City. *International Journal of Engineering and Science*. 2 (2) : 13-16.
- Ren, S., Park, M., Sah, H. and Lee, B. 2008. Effect of pharmaceutical excipients on aqueous stability of rabeprazole sodium. *International Journal of Pharmaceutics*. 350: 197-204.
- Sabry, S. M., Belal, T. S. and Mahgoub, A. A. 2015. Development and validation of HPLC-DAD method for the simultaneous determination of amoxicillin, metronidazole and rabeprazole sodium. Application to spiked simulated intestinal fluid samples. *Annales Pharmaceutiques Francaises*.
- Santos, L. H. M. L. M., Gros, M., Rodriguez-Mozaz, S., Delerue-Matos, C., Pena, A., Barceló, D., and Montenegro, M. C. B. S. M. 2013. Contribution of hospital effluents to the load of pharmaceuticals in urban wastewaters: Identification of ecologically relevant pharmaceuticals. *Science of The Total Environment*. 461-462 : 302-316.
- Sharma, P., Mathur, N., Singh, A. and Sogani, M. 2015. Monitoring hospital wastewaters for their probable genotoxicity and mutagenicity. Environment Monitoring Assessment.
- Stuart, M., Lapworth, D., Crane, E. and Hart, A. 2012. Review of risk from potential emerging contaminants in UK groundwater. *Science of the Total Environment*. 416 : 1-21.
- Sui, Q., Cao, X., Lu, S., Zhao, W., Qiu, Z., and Yu, G. 2015. Occurrence, sources and fate of pharmaceuticals and personal care products (PPCPs) in the groundwater: A review. *Emerging Contaminants*. 1 (1) : 14-24.
- Verlicchi, P., Aukidy, M. Al, and Zambello, E. 2012. Occurrence of pharmaceutical compounds in urban wastewater: Removal, mass load and environmental risk after a secondary treatment -A review. *Science of the Total Environment*. 429 : 123-155.
- World Health Organization. 2015. Indonesia : WHO Statistical Profile.
- WWF, ToxicChemical. http://wwf.panda.org/about_our_earth/teacher_resources/webfieldtrips/toxics/.
- Xie, H. 2012. Occurrence, Ecotoxicology, and Treatment of Anticancer Agents as Water Contaminants. *Environmental and Analytical Toxicology*.
- Yamina, B., Tahar, B., Lila, M., Hocine, H. and Laure, F. M. 2014. Study on Cadmium Resistant-Bacteria Isolated from Hospital Wastewaters. *Advances in Bioscience and Biotechnology*, (July), 718-726.
- Yuan, Q., Guo, M. and Yang, J. 2015. Fate of Antibiotic Resistant Bacteria and Genes during Wastewater Chlorination: Implication for Antibiotic Resistance Control. *PLOS One*, 1-12.
- Yoon, Y., Ryu, J., Oh, J., Choi, B.G. and Snyder, S. A. 2010. Occurrence of endocrine disrupting compounds, pharmaceuticals, and personal care products in the Han River (Seoul, South Korea). *Science of The Total Environment*. 408 (3) : 636-643.
- Zorita, S., Mårtensson, L. and Mathiasson, L. 2009. Occurrence and removal of pharmaceuticals in a municipal sewage treatment system in the south of Sweden. *Science of The Total Environment*. 407(8): 2760-2770.
-

**LEMBAR
HASIL PENILAIAN SEJAWAT SEBIDANG ATAU PEER REVIEW
KARYA ILMIAH: JURNAL ILMIAH**

Judul Artikel Ilmiah : The Screening of Potential Antibiotics from Hospital Wastewater in Tropical Region (Case Study at Palembang, South Sumatra, Indonesia)

Penulis Artikel ilmiah : Ian Kurniawan, S Nasir, Hermansyah, **Mardiyanto**

Identitas Jurnal Ilmiah : a. Nama Jurnal : Jounal Pollution Research
b. Nomor/Volume/Hal : 36/2
c. Edisi (bulan/tahunan) : Agustus/2017
d. Penerbit : EM International
e. Jumlah halaman : 5

Kategori Publikasi Jurnal Ilmiah : Jurnal Ilmiah Internasional Bereputasi
(beri √ pada kategori yang tepat) Jurnal Ilmiah Internasional
 Jurnal Ilmiah Nasional Terakreditasi dan Terindex DOAJ
 Jurnal Ilmiah Nasional Tidak Terakreditasi
 Jurnal Ilmiah Nasional Terindeks DOAJ dll.

I. Hasil Penilaian Validasi:

No.	ASPEK	URAIAN/KOMENTAR PENILAI
1.	Indikasi Plagiasi	Tidak terindikasiplagiat
2.	Linearitas	Sesuai dengan bidang ilmu farmasi, khususnya bidang teknologifarmasi

II. Hasil Penilaian Peer Review:

Komponen yang dinilai	Nilai Maksimal Jurnal Ilmiah (isikan di kolom yang sesuai)					Nilai Akhir yang diperoleh
	Internasional Bereputasi	Internasional	Nasional Terakreditasi	Nasional tidak Terakreditasi	Nasional terindeks DOAJ	
Kelengkapan dan kesesuaian unsur isi jurnal (10%)	10	-	-	-	-	10
Ruang lingkup dan kedalaman pembahasan (30%)	29	-	-	-	-	29
Kecukupan dan kemutakhiran data/informasi dan metodologi (30%)	28	-	-	-	-	28
Kelengkapan unsur dan kualitas penerbit (30%)	29	-	-	-	-	29
Total = (100%)						96
Kontribusi pengusul (Penulis pertama/Anggota Utama)	Penulis sudah memberikan kontribusi dengan baik					
Komentar/Ulasan PEER REVIEW						
Kelengkapan dan kesesuaian unsur :	Lengkap dan sesuai dengan kaidah penulisan ilmiah internasional					
Ruang lingkup dan kedalaman Pembahasan :	Kedalaman pembahasan sangat baik					
Kecukupan dan kemutakhiran data dan metodologi	Data yang diperoleh dan metode yang digunakan sudah sangat baik					
Kelengkapan unsur dan kualitas penerbit	Sudah sangat baik					

Indralaya, 29 April 2020
Penilai II



Prof. Aldes Lesbani, Ph.D
NIP.197408121998021001
Unit Kerja: Kimia
Bidang Ilmu: Kimia Anorganik
Jabatan/Pangkat: Guru Besar

**LEMBAR
HASIL PENILAIAN SEJAWAT SEBIDANG ATAU PEER REVIEW
KARYA ILMIAH: JURNAL ILMIAH**

Judul Artikel Ilmiah : The Screening of Potential Antibiotics from Hospital Wastewater in Tropical Region (Case Study at Palembang, South Sumatra, Indonesia)

Penulis Artikel ilmiah : Ian Kurniawan, S Nasir, Hermansyah, **Mardiyanto**

Identitas Jurnal Ilmiah : a. Nama Jurnal : Jounal Pollution Research
b. Nomor/Volume/Hal : 36/2
c. Edisi (bulan/tahunan) : Agustus/2017
d. Penerbit : EM International
e. Jumlah halaman : 5

Kategori Publikasi Jurnal Ilmiah : Jurnal Ilmiah Internasional Bereputasi
(beri \surd pada kategori yang tepat) Jurnal Ilmiah Internasional
 Jurnal Ilmiah Nasional Terakreditasi dan Terindex DOAJ
 Jurnal Ilmiah Nasional Tidak Terakreditasi
 Jurnal Ilmiah Nasional Terindeks DOAJ dll.

I. Hasil Penilaian Validasi:

No.	ASPEK	URAIAN/KOMENTAR PENILAI
1.	Indikasi Plagiasi	Tidak terindikasi plagiat
2.	Linearitas	Sesuai dengan bidang ilmu farmasi, khususnya bidang teknologifarmasi

II. Hasil Penilaian Peer Review:

Komponen yang dinilai	Nilai Maksimal Jurnal Ilmiah (isikan di kolom yang sesuai)					Nilai Akhir yang diperoleh
	Internasional Bereputasi	Internasional	Nasional Terakreditasi	Nasional tidak Terakreditasi	Nasional terindeks DOAJ	
Kelengkapan dan kesesuaian unsur isi jurnal (10%)	10	-	-	-	-	10
Ruang lingkup dan kedalaman pembahasan (30%)	29	-	-	-	-	29
Kecukupan dan kemutakhiran data/informasi dan metodologi (30%)	28	-	-	-	-	28
Kelengkapan unsur dan kualitas penerbit (30%)	29	-	-	-	-	29
Total = (100%)						96
Kontribusi pengusul (Penulis pertama/Anggota Utama)	Penulis sudah memberikan kontribusi dengan baik					
Komentar/Ulasan PEER REVIEW						
Kelengkapan dan kesesuaian unsur :	Lengkap dan sesuai dengan kaidah penulisan ilmiah internasional					
Ruang lingkup dan kedalaman Pembahasan :	Kedalaman pembahasan sangat baik					
Kecukupan dan kemutakhiran data dan metodologi	Data yang diperoleh dan metode yang digunakan sudah sangat baik					
Kelengkapan unsur dan kualitas penerbit	Sudah sangat baik					

Indralaya, 29 April 2020
Penilai II



Dr.rer.nat Risfidian Mohadi, MSi
NIP.19771127200501003
Unit Kerja: Kimia
Bidang Ilmu: Polimeric Kimia
Jabatan/Pangkat: Lektor Kepala