

# PREVALENCE AND PREDICTORS OF EXCLUSIVE BREASTFEEDING PRACTICES IN SEBERANG ULU I, PALEMBANG: A CROSS- SECTIONAL STUDY

*by* Indah Purnama Sari

---

**Submission date:** 02-Jan-2021 04:46PM (UTC+0700)

**Submission ID:** 1482459793

**File name:** JIKM\_NOVEMBER\_2019.docx (38.19K)

**Word count:** 3387

**Character count:** 19131

# PREVALENCE AND PREDICTORS OF EXCLUSIVE BREASTFEEDING PRACTICES IN SEBERANG ULU I, PALEMBANG: A CROSS-SECTIONAL STUDY

Indah Purnama Sari,<sup>1</sup> Dewi Handayani,<sup>2</sup> Fatmalina Febry<sup>3</sup>

<sup>1,3</sup>Study Program of Nutrition, Public Health Faculty, Universitas Sriwijaya

<sup>2</sup>Master's Degree from Public Health Faculty, Universitas Sumatera Utara

## ABSTRACT

Exclusive breastfeeding is one of the successful indicators in improving the health of babies. From 2013 until 2014, the practice of exclusive breastfeeding has decreased in Indonesia and has not reached the national targets of around 80% of babies being exclusively breastfed. This study aimed to identify the prevalence and determinants of exclusive breastfeeding in Seberang Ulu I, Palembang. A community-based cross sectional study was executed from July 5, 2016 to August 5, 2016 among mothers who had infants aged less than six months. A purposive sampling technique was used to select a sample of 125 participants. Data were collected using a structured questionnaire by the face-to-face interview technique. Bivariate and multiple logistic regression analysis were used to determine factors associated with the practice of breastfeeding exclusively and to control confounding effects. The prevalence of exclusive breastfeeding was 26.4% (95% CI: 20.0%-35.1%). After adjusting for confounders (mothers' attitudes and health workers' support), mothers who had good knowledge were 11.66 times more likely to breastfeed exclusively than those who had poor knowledge (AOR: 11.66, 95% CI: 3.07-44.31). The prevalence of exclusive breastfeeding in Seberang Ulu I, Palembang, was still very low and has not reached the national target. The recommendation is to maximize the role of health workers in providing information about exclusive breastfeeding, so that they can continue providing motivation for mothers, husbands and families, so that the success and sustainability of exclusive breastfeeding practices could be achieved.

**Keywords:** Mother's knowledge, mother's attitude, health workers support, exclusive breastfeeding

## PREVALENSI DAN PREDIKTOR PEMBERIAN ASI EKSKLUSIF DI SEBERANG ULU I, PALEMBANG: STUDI CROSS-SECTIONAL

### ABSTRAK

Pemberian ASI eksklusif merupakan salah satu indikator keberhasilan upaya peningkatan kesehatan bayi. Dalam kurun waktu tahun 2013 sampai dengan tahun 2014, terjadi penurunan cakupan pemberian ASI eksklusif di Indonesia. Hasil cakupan tersebut masih dibawah target program secara nasional yaitu sebanyak 80% bayi diberikan ASI eksklusif. Penelitian ini bertujuan untuk mengidentifikasi prevalensi dan determinan praktik pemberian ASI eksklusif di Kecamatan Seberang Ulu I, Palembang. Studi potong lintang berbasis masyarakat telah dilaksanakan dari tanggal 5 Juli 2016 sampai 5 Agustus 2016 pada ibu yang memiliki bayi berusia 0-6 bulan. Teknik pengambilan sampel secara tidak acak digunakan untuk memilih 125 responden. Data dikumpulkan menggunakan kuisioner terstruktur melalui teknik tatap muka. Analisis bivariat dan regresi logistik ganda digunakan untuk mengetahui faktor yang berhubungan dengan praktik pemberian ASI eksklusif dan untuk mengendalikan efek perancu. Prevalensi pemberian ASI eksklusif sebesar 26.4% (95% CI: 20.0%-35.1%). Setelah mengontrol efek perancu (sikap ibu dan dukungan petugas kesehatan), ibu berpengetahuan baik berkemungkinan 11.66 kali lebih tinggi untuk dapat memberikan ASI eksklusif dibandingkan ibu berpengetahuan kurang baik (OR<sub>a</sub>: 11.66, 95% CI: 3.07-44.31). Prevalensi pemberian ASI eksklusif di Kecamatan Seberang Ulu I, Palembang masih rendah dan belum mencapai target nasional. Adapun saran yang dapat diberikan adalah memaksimalkan peran tenaga kesehatan dalam memberikan informasi mengenai ASI eksklusif sehingga tenaga kesehatan dapat terus memotivasi para ibu, suami dan keluarga, yang akhirnya keberhasilan dan keberlangsungan praktik pemberian ASI eksklusif dapat tercapai.

**Kata kunci:** Pengetahuan ibu, sikap ibu, dukungan tenaga kesehatan, ASI eksklusif

## INTRODUCTION

Exclusive breastfeeding is the practice<sup>24</sup> giving only breast milk to infants in their first six months of life (no other food or water).<sup>1</sup> Being breastfed exclusively is also a key indicator of childhood survival,<sup>4</sup> nutrition, early growth and development. It not only provides infants with the best start in their life but also has benefits for mothers, protects against non-communicable diseases and contributes to environmental sustainability.<sup>2</sup> Exclusive breastfeeding has a significant impact on the reduction of morbidity and mortality<sup>16</sup> from diarrhea and pneumonia.<sup>3</sup>

Improvements in breastfeeding rates are critical to the attainment of the Post-2015 Sustainable Development Goals due to the fact that global breastfeeding rates have seen only slow progress since 1995. Globally, fewer than 40% of infants aged 0 to six months are exclusively breastfed.<sup>2</sup> Indonesia, a developing country, has more infants who are non-exclusively breastfed than the Philippines and Vietnam.<sup>3</sup> In Indonesia, based on data from Ministry of Health, the coverage of exclusive breastfeeding<sup>21</sup> in 2014 (52.3%) decreased compared to the coverage of exclusive breastfeeding in 2013 (54.3%). This means that only 52.3% of infants are breastfed to six months of age. The coverage is still below the national target<sup>1</sup> of 80% of infants being given breast milk from birth to six months of age (exclusive breastfeeding).<sup>4</sup> Many factors can affect the practice of exclusive breastfeeding by mothers, such as socio-demographic, knowledge, attitude, and husbands', families' (mother's or mother-in-law's) and health workers' support.<sup>5-13</sup>

South Sumatera Province is one of the provinces that focus on exclusive breastfeeding due to the low coverage of exclusive breastfeeding in 2014, at 64.5%.<sup>4</sup> The coverage of exclusive breastfeeding for Palembang City in 2014 was 74.18%. Seberang Ulu I sub-district is one of the sub-districts under the auspices of the City Health

Office of Palembang. Based on the Health Profile of Palembang City in 2014, it was found that Seberang Ulu I sub-district was one of the sub-districts with exclusive breastfeeding coverage (72.87%) that was lower than Palembang City.<sup>14</sup> Therefore, strategies were needed to promote and support exclusive breastfeeding, not only to increase the coverage of infants less than six months being exclusively breastfed, but also to decrease the morbidity and mortality of infants. The aim of this study was to identify the prevalence of exclusive breastfeeding and to determine the predictors among mothers of infants less than six months of age in Seberang Ulu I sub-district, Palembang, Indonesia.

## METHOD

### Study Design and Sampling Procedure

This was a community-based cross-sectional study held from July 5 to August 5, 2016, in Seberang Ulu I sub-district, Palembang. The populations were all mothers of children who had been born alive. Samples were mothers of infants who had been aged 7 months until 11 months who fulfilled the inclusion criteria, such as infants with single birth or who were either the first or the last child in the family. While exclusion criteria were infants with infectious diseases (diarrhea and Acute Respiratory Duct Infection) and those who had a single parent. The sample size was calculated using hypothesis tests for two population proportions. In a previous study<sup>1</sup> conducted by Tewabe<sup>15</sup>, the prevalence of exclusive breastfeeding among mothers who had and had not received breastfeeding counseling during antenatal care was found to be about 64% and 38%, respectively. Given the 95% confidence intervals (CI), the 80% power of the test and a 10% non-response rate, the final sample size was about 125 eligible mothers, selected by purposive sampling.

## Data Collection Instruments and Data Collection Procedure

14

Data were collected during house visits by face-to-face interviews using a structured interviewer-administered questionnaire. A content and construct validity test was performed to ensure the validity and reliability of the questionnaire. To ensure that the questionnaire was systematic, it was divided into seven parts for the respondents to answer. Part A was about socio-demographic characteristics such as age of mother, occupation, monthly income, and education level, parity, birth attendants, place of birth, and types of birth. Part B was about exclusive breastfeeding practice, which included breastfeeding after birth, one hour after birth, until two months, two to four months, four to six months, and types of foods and drinks that were consumed by infants. Exclusive breastfeeding practice means that mothers give infants only breast milk until the age of six months. Infants who had received breast milk exclusively until six months of age were coded '1' while those that had not been exclusively breastfed were coded '0'.

Part C, D, E, F and G are categorized into 2 categories which previously tested using Kolmogorov-Smirnov test to investigate the normal distribution of the data. The mean was used if data normally distributed and the median if data didn't fitted normal distribution. Part C was about mothers' knowledge, which consisted of 21 questions; each correct answer was given a score of 1 and a score of 0 was given for each incorrect answer. After totaling the scores and median (18.00), it was dichotomized into good knowledge  $\geq$  median and poor knowledge  $<$  median. Part D was about mothers' attitudes, which consisted of 8 questions. Each item of the question was based on Likert's scale, ranging from 1 to 5, strongly disagree (1), disagree (2), neutral (3), agree (4), and strongly agree (5) for positive questions and

counterparts for negative questions. After totaling the scores and mean (56.37), it was dichotomized into positive  $\geq$  mean and less positive  $<$  mean. Parts E, F and G were about husbands' support (25 questions), families' support (20 questions), and health workers' support (17 questions) that used a scale ranging from 1 to 3 with never (1), sometimes (2) and often (3) for positive questions and counterparts for negative questions. After totaling the scores of husbands' support (mean (34.15)), families' support (median (31.00)), and health workers' support (mean (25.00)), it was dichotomized into  $\geq$  mean/median and no  $<$  mean/median. All respondents were briefed about the objectives of the study and written informed consent was obtained prior to data collection.

## Data Processing and Analysis

The collected data were checked and verified manually. Any inconsistencies and inaccuracies were corrected the following day. After that, completed data were coded and entered on the computer using statistical software. Before data analysis, the data were crosschecked for any unusual findings. Descriptive statistics were used to identify the socio-demographic characteristics, mothers' knowledge and attitudes, support of husbands, families and health workers, and the prevalence of exclusive breastfeeding. To determine factors associated with exclusive breastfeeding practice, binary logistic regression was used. Two models were run: the first model was performed to determine each independent variable with the outcome variables, which were included in the second, final model (using multiple logistic regression). The strength of association was measured using adjusted odds ratio (AOR) with 95% confidence intervals (CI). The p-value of less than 0.05 was declared as a significant association.

## RESULTS

### Socio-demographic Characteristics

A total of 125 mothers with infants of age 7-11 months were interviewed. Around seven-ninths (77.6%) of mothers were delivered at the age of 20-35 years. Regarding educational status, 74 (59.2%) of the mothers had higher than secondary education and a majority (86.4%) of the study participants were unemployed. Around three-quarters

(75.2%) of the mothers earned a monthly household income less than the City Minimum Wage of Palembang: about 169.30 USD. More than seven-tenths (70.4%) of mothers had more than two children and 105 (84%) mothers delivered by vaginal birth (normally). Most (96.8% and 98.4%) of the mothers were delivered in the health facility and had a health worker as a birth attendant (Table 1).

**Table 1.**  
**Socio-demographic characteristics among mothers of infants aged less than six months, Seberang Ulu I, Palembang, 2016 (n=125)**

Variables	Category	Total (n)	Percentage (%)
Maternal age (in years)	20-35	97	77.6
	<20 and or >35	28	22.4
Maternal education	Secondary education and above	74	59.2
	Primary education and lower	51	40.8
Maternal occupation	Unemployed	108	86.4
	Employed	17	13.6
Monthly household income (USD)	Less than 169.30	94	75.2
	More than 169.30	31	24.8
Number of children	2 and above	88	70.4
	Only 1	37	29.6
Mode of delivery	Normal/ vaginal	105	84
	Caesarian section	20	16
Place of birth	Health facility	121	96.8
	Home	4	3.2
Birth attendants	Health workers	123	98.4
	Non health workers	2	1.6

### Prevalence and Predictors of Exclusive Breastfeeding Practice

The prevalence of exclusive breastfeeding practice in Seberang Ulu I sub-district was 26.4% (95% CI: 20.0%-35.1%). More than half, 51.2% and 54.4%, of mothers reported they had good knowledge of and

attitudes about exclusive breastfeeding, respectively. While 66 (52.8%) mothers had their husbands' support, they also had the support of their families and health workers for feeding their infants only breast milk until the age of six months (Table 2).

**Table 2.**  
**Prevalence and predictors of exclusive breastfeeding practice among mothers of infants aged less than six months, Seberang Ulu I, Palembang, 2016 (n=125)**

Variables	Category	Total (n)	Percentage (%)
Exclusive breastfeeding practice	Yes	33	26.4
	No	92	73.6
Mothers' knowledge	Good knowledge	64	51.2
	Poor knowledge	61	48.8
Mothers' attitude	Positive	68	54.4
	Less positive	57	45.6
Husbands' support	Yes	66	52.8
	No	59	47.2
Family support	Yes	66	52.8
	No	59	47.2
Health workers' support	Yes	66	52.8
	No	59	47.2



## Factors Associated with Exclusive Breastfeeding Practice

The independent variables of the practice of exclusive breastfeeding were maternal age, education, occupation, monthly household income, number of children, mode of delivery, place of birth, birth attendants, mother's knowledge, mother's attitude, and husband's, family's and health workers' support. In the bivariate logistic regression, only mother's knowledge and attitude were significantly associated with exclusive breastfeeding practice ( $p$ -value<0.05). However, all of the independent variables

were included in the multiple logistic regression models. While each independent variable was adjusted for other variables, mother's knowledge was a dominant variable associated with exclusive breastfeeding practice after adjusted by mother's attitude and health workers' support (95% confidence level and a  $p$ -value < 0.05). Mothers who had good knowledge were 11.66 times more likely to breastfeed exclusively than those who had poor knowledge, after adjusted by mother's attitude and health worker's support (OR<sub>a</sub>: 11.66, 95% CI: 3.07-44.31) (Table 3).

**Table 3.**  
Factors associated with exclusive breastfeeding practice of mothers of infants aged less than six months, Seberang Ulu I, Palembang, 2016

Variables	Category	Exclusive Breastfeeding Practice		OR <sub>a</sub> (95% CI)	OR <sub>a</sub> (95% CI)
		Yes	No		
		(n & %)	(n & %)		
Maternal age (in years)	20-35	26 (26.8)	71 (73.2)	1.09 (0.42-2.89)	-
	<20 and or >35	7 (25.0)	21 (75.0)	1	-
	Secondary education and above	23 (31.1)	51 (68.9)	1.85 (0.79-4.32)	-
Maternal education	Primary education and lower	10 (19.6)	41 (80.4)	1	-
	Unemployed	28 (25.9)	80 (74.1)	0.84 (0.27-2.59)	-
Maternal occupation	Employed	5 (29.4)	12 (70.6)	1	-
	Less than 169.30	24 (25.5)	70 (74.5)	0.84 (0.34-2.07)	-
Monthly household income (USD)	More than 169.30	9 (29.0)	22 (71.0)	1	-
	2 and above	23 (26.1)	65 (73.9)	0.96 (0.40-2.28)	-
Number of children	Only 1	10 (27.0)	27 (73.0)	1	-
	Normal/ vaginal	29 (27.6)	76 (72.4)	1.53 (0.47-4.95)	-
Mode of delivery	Caesarian section	4 (20.0)	16 (80.0)	1	-
	Health facility	32 (26.4)	89 (73.6)	1.08 (0.11-10.75)	-
Place of birth	Home	1 (25.0)	3 (75.0)	1	-
	Health workers	33 (26.8)	90 (73.2)	-	-
Birth attendants	Non health workers	0 (0.0)	2 (100.0)	-	-
	Good knowledge	28 (43.8)	36 (56.3)	8.71 (3.08-24.64)*	11.66 (3.07-44.31)*
Mother's knowledge	Poor knowledge	5 (8.2)	56 (91.8)	1	1
	Good attitude	24 (35.3)	44 (64.7)	2.91 (1.22-6.93)*	0.82 (0.25-2.69)
Mother's attitude	Poor attitude	9 (15.8)	48 (84.2)	1	1
	Yes	21 (31.8)	45 (68.2)	1.83 (0.81-4.14)	-
Husband's support	No	12 (20.3)	47 (79.7)	1	-
	Yes	20 (30.3)	46 (69.7)	1.54 (0.69-3.46)	-
Family support	No	13 (22.0)	46 (78.0)	1	-
	Yes	19 (28.8)	47 (71.2)	1.29 (0.58-2.89)	0.64 (0.25-1.66)
Health worker's support	No	14 (23.7)	45 (76.3)	1	1

1 = reference, \* =  $p$ -value < 0.05, n = total, % = percent

## DISCUSSION

<sup>11</sup> Exclusive breastfeeding is the provision of breast milk given to infants from birth for six months, without adding any or replacing with other foods or beverages. The prevalence of exclusive breastfeeding in this study was less than one-third, 26.4%, which is lower than Indonesia's national target level of 80%.<sup>4</sup> It is also lower than declared in early international studies in Ethiopia (74.1%),<sup>16</sup> Hawassa Southern Ethiopia (60.9%),<sup>17</sup> Lebanon (27.4%),<sup>18</sup> Bahir Dar City Ethiopia (50.3%),<sup>19</sup> Halaba Ethiopia (70.5%),<sup>20</sup> Magelang City of Indonesia (27.4%),<sup>21</sup> Gianyar District of Indonesia (31.9%),<sup>22</sup> Ghana (27.7%).<sup>6</sup> But it is higher than the prevalence in Congo (2.8%),<sup>23</sup> Canada (15.3%),<sup>24</sup> and City of Palu, Central Sulawesi, Indonesia (26.2%).<sup>8</sup> This variation might be due to methodological differences, time, economic status, and social-cultural differences.

Mother's knowledge was one variable that became the focus in this research. Knowledge is the result of human sensing or one's experience of the object through the five senses (sight, smell, taste, touch and hearing) at the time of sensing and being strongly influenced by the intensity of attention and perception of the object.<sup>25</sup> In this study, mothers who had good knowledge practiced exclusive breastfeeding (51.2%), as opposed to those who had poor knowledge (48.8%). This result is similar to studies from Ethiopia,<sup>9,26</sup> Congo,<sup>23</sup> Myanmar,<sup>27</sup> Ghana,<sup>28</sup> and Indonesia.<sup>21</sup> The results of this study showed that about 16.0% of mothers said that colostrum should not be given to infants and 14.4% of them said that colostrum should be discarded. This indicated that the mothers' knowledge about breast milk, especially colostrum, was inadequate. It is very important that mothers, especially young mothers, receive adequate information on the nutrition content of colostrum and breast milk

as a whole, so that infants can be exclusively breastfed until the age of six months. In addition to the mother's knowledge, the husband's and family's support plays an important role in the continuity of exclusive breastfeeding practices. Husbands can provide support to improve mothers' knowledge by helping to find out information about the importance of exclusive breastfeeding and how to breastfeed the baby properly. The results showed that only 16.8% of mothers receive this support from their husbands.

After an adjustment was made using a multiple logistic regression model, mothers' attitudes associated with breastfed exclusively for infants in the first their six months. More than half (54.4%) of mothers in Seberang Ulu I sub-district, Palembang had a positive attitude to the practice of exclusive breastfeeding. Bivariate logistic regression showed that mothers with a positive attitude were 2.91 times more likely to breastfeed their infants exclusively compared to those who with a negative attitude (OR: 2.91, 95% CI: 1.22-6.93). While in multiple logistic regression, mothers' attitude was a confounder. This result was consistent with the previous studies done in Indonesia,<sup>29</sup> Ireland,<sup>30</sup> Lebanon,<sup>18</sup> and Ethiopia.<sup>26</sup> This study also found that almost four-fifths (77.6%) of mothers showed that they had a feeling of excitement when giving breastfed exclusively to their babies and 63.2% of mothers believed that infants who had been exclusively breastfed were healthier than those who had been fed formula milk. However, the results showed that among mothers with a positive attitude toward exclusive breastfeeding practices, 64.7% of mothers did not exclusively breastfed. This shows that not all positive attitudes will result in positive behavior. This is because that attitude is a reaction or response to a stimulus or object that is still undiscovered.<sup>25</sup>

Along with mothers' knowledge and attitude, health workers' support was one of

the supporting factors for the continuity of exclusive breastfeeding. The results showed that almost all mothers received information on exclusive breastfeeding from health workers at the time of pregnancy. About 92.0% of mothers said that health workers provided information about giving breastfeeding to infants, and 94.4% of mothers said that health workers recommended giving colostrum to babies. This study is also consistent with previous studies in Ethiopia<sup>15</sup> and Myanmar<sup>27</sup> that concluded that health workers' support is associated with the exclusive breastfeeding of infants. It means that health workers not only have an important role to play in improving the quality of health services to mothers, but also play a role in raising people's awareness, willingness and ability to live healthily. The optimum degree of health is an investment in the development of human resources, so that people are productive, both socially and economically.

## CONCLUSION

This study concludes that the prevalence of exclusive breastfeeding in Seberang Ulu I sub-district of Palembang is still very low and has not reached the national target. Mothers' knowledge is a dominant variable with exclusive breastfeeding, while mothers' attitude and health workers' support also affects exclusive breastfeeding practices. Mothers' knowledge and attitude are also related to husbands', families' and health workers' support, which can motivate mothers to breastfeed their babies exclusively. The advice from this study is to maximize the role of health workers in providing information about exclusive breastfeeding, aimed not only at the mother, but also at the nearest family members (husband, mother, mothers-in-law). In addition, health workers need to take a refresher course related to the practice of exclusive breastfeeding, so that they can continue to provide motivation for mothers, husbands and families, leading to the success and sustainability of exclusive breastfeeding practices.



# PREVALENCE AND PREDICTORS OF EXCLUSIVE BREASTFEEDING PRACTICES IN SEBERANG ULU I, PALEMBANG: A CROSS-SECTIONAL STUDY

## ORIGINALITY REPORT

17%

SIMILARITY INDEX

8%

INTERNET SOURCES

12%

PUBLICATIONS

6%

STUDENT PAPERS

## PRIMARY SOURCES

1

Gedion Asnake Azeze, Kelemu Abebe Gelaw, Natnael Atnafu Gebeyehu, Molalegn Mesele Gesese, Taklu Marama Mokonnen. "Exclusive Breastfeeding Practice and Associated Factors among Mothers in Boditi Town, Wolaita Zone, Southern Ethiopia, 2018: A Community-Based Cross-Sectional Study", International Journal of Pediatrics, 2019

Publication

2%

2

Submitted to Universitas Muslim Indonesia

Student Paper

2%

3

Tilahun Tewabe. "Timely initiation of breastfeeding and associated factors among mothers in Motta town, East Gojjam zone, Amhara regional state, Ethiopia, 2015: a cross-sectional study", BMC Pregnancy and Childbirth, 2016

Publication

1%

Submitted to Middle Tennessee State University

4

Student Paper

1%

5

[article.sciencepublishinggroup.com](https://article.sciencepublishinggroup.com)

Internet Source

1%

6

[lemlit.unsri.ac.id](https://lemlit.unsri.ac.id)

Internet Source

1%

7

[www.ncbi.nlm.nih.gov](https://www.ncbi.nlm.nih.gov)

Internet Source

1%

8

140.122.83.189:8080

Internet Source

1%

9

Asrat Sonko, Amare Worku. "Prevalence and predictors of exclusive breastfeeding for the first six months of life among women in Halaba special woreda, Southern Nations, Nationalities and Peoples' Region/SNNPR/, Ethiopia: a community based cross-sectional study", Archives of Public Health, 2015

Publication

1%

10

Erwin Halim, Gen Norman Thomas, Daphne Ratna Hidayat, Diurman Gea. "“Smart Healthcare” a Medical Record System for Effective Health Services", 2020 International Conference on Information Management and Technology (ICIMTech), 2020

Publication

1%

11	<a href="http://journal.umpo.ac.id">journal.umpo.ac.id</a> Internet Source	1 %
12	<a href="http://journal.unhas.ac.id">journal.unhas.ac.id</a> Internet Source	1 %
13	Submitted to Western Governors University Student Paper	<1 %
14	<a href="http://www.researchgate.net">www.researchgate.net</a> Internet Source	<1 %
15	"Handbook of dietary and nutritional aspects of bottle feeding", Wageningen Academic Publishers, 2014 Publication	<1 %
16	Submitted to The University of Manchester Student Paper	<1 %
17	Zemene Tigabu Kebede, Kassahun Alemu Gelaye, Mehari Woldemariam Merid, Temesgen Yihunie Akalu et al. "High common childhood illnesses while the treatment-seeking behavior of mothers/caregivers' was low in rural northwest Ethiopia in 2018. A community-based cross-sectional study", Research Square, 2019 Publication	<1 %
18	<a href="http://scidoc.org">scidoc.org</a> Internet Source	<1 %
19	<a href="http://nurfaizinyunus.blogspot.com">nurfaizinyunus.blogspot.com</a>	

<1 %

20

[www.researchsquare.com](http://www.researchsquare.com)

Internet Source

<1 %

21

Nia Dwi Yuliaty, Hadi Hadi, Sri Rahayu, Noor Pramono, Donny Kristanto Mulyantoro. "THE IMPACT OF COMBINATION OF ROLLING AND OKETANI MASSAGE ON PROLACTIN LEVEL AND BREAST MILK PRODUCTION IN POST-CESAREAN SECTION MOTHERS", Belitung Nursing Journal, 2017

Publication

<1 %

22

[www.panafrican-med-journal.com](http://www.panafrican-med-journal.com)

Internet Source

<1 %

23

Submitted to King's College

Student Paper

<1 %

24

Submitted to Laureate Higher Education Group

Student Paper

<1 %

25

[jnc.stikesmaharani.ac.id](http://jnc.stikesmaharani.ac.id)

Internet Source

<1 %

26

Tam, Sing-Fai, and David Watkins. "DEVELOPMENT AND VALIDATION OF A SELF-CONCEPT QUESTIONNAIRE FOR HONG KONG CHINESE ADULTS WHO HAVE PHYSICAL DISABILITY", Social Behavior and

<1 %

- 
- |           |  |                |
|-----------|--|----------------|
| <b>27</b> | Hanulan Ulan Septiani, Artha Budi, Karbito Karbito. "Faktor-Faktor yang Berhubungan dengan Pemberian ASI Eksklusif Oleh Ibu Menyusui yang Bekerja Sebagai Tenaga Kesehatan", Jurnal Aisyah : Jurnal Ilmu Kesehatan, 2017<br><small>Publication</small> | <b>&lt;1 %</b> |
|-----------|--|----------------|
- 
- |           |  |                |
|-----------|--|----------------|
| <b>28</b> | <a href="http://cheelious.blogspot.com">cheelious.blogspot.com</a><br><small>Internet Source</small> | <b>&lt;1 %</b> |
|-----------|--|----------------|
- 
- |           |  |                |
|-----------|--|----------------|
| <b>29</b> | <a href="http://repository.unhas.ac.id">repository.unhas.ac.id</a><br><small>Internet Source</small> | <b>&lt;1 %</b> |
|-----------|--|----------------|
- 
- |           |  |                |
|-----------|--|----------------|
| <b>30</b> | <a href="http://www.hindawi.com">www.hindawi.com</a><br><small>Internet Source</small> | <b>&lt;1 %</b> |
|-----------|--|----------------|
- 
- |           |  |                |
|-----------|--|----------------|
| <b>31</b> | <a href="http://es.scribd.com">es.scribd.com</a><br><small>Internet Source</small> | <b>&lt;1 %</b> |
|-----------|--|----------------|
- 
- |           |   |                |
|-----------|---|----------------|
| <b>32</b> | Herman Herman, Yulfiana Yulfiana, Nurdin Rahman, Ahmad Yani. "Perilaku Ibu Menyusui dalam Keberhasilan Pemberian ASI Eksklusif di Wilayah Kerja Puskesmas Tawaeli Kota Palu", MPPKI (Media Publikasi Promosi Kesehatan Indonesia): The Indonesian Journal of Health Promotion, 2018<br><small>Publication</small> | <b>&lt;1 %</b> |
|-----------|---|----------------|
- 
- |           |   |  |
|-----------|---|--|
| <b>33</b> | Mizanur Rahman, Sharmin Mizan, Razitasham |  |
|-----------|---|--|



Binti Safii, Akhtar Ahmad. "Are the urban poor satisfied with health care services for child delivery? Evidence from an Urban Primary Health Care (UPHC) Project in Bangladesh", Family Medicine & Primary Care Review, 2018

Publication

<1%

Exclude quotes

On

Exclude matches

< 5 words

Exclude bibliography

On