



First record of Hagen's batwing *Atrophaneura hageni* (Rogenhofer, 1889) (Lepidoptera: Papiolinidae) in southern Sumatra, Indonesia

DONI SETIAWAN¹, INA APRILIA², MUHAMMAD IQBAL^{3*}, GUNTUR PRAGUSTIANDI²,
ARUM SETIAWAN¹ & INDRA YUSTIAN¹

¹*Department of Biology, Faculty of Science, Sriwijaya University, Jalan Raya Palembang-Prabumulih km 32, Indralaya, Sumatera Selatan 30662, Indonesia*

²*Conservation Biology Program, Faculty of Science, Sriwijaya University, Jalan Padang Selasa 524, Palembang, Sumatera Selatan 30129, Indonesia*

³*Biology Program, Faculty of Science, Sriwijaya University, Jalan Padang Selasa 524, Palembang, Sumatera Selatan 30129, Indonesia*

*Corresponding author: kpbsos26@yahoo.com

Received 24 December 2019 | Accepted by V. Pešić: 7 February 2020 | Published online 13 February 2020.

Abstract

Hagen's batwing *Atrophaneura hageni* (Rogenhofer, 1889) (Lepidoptera: Papiolinidae) is endemic to Sumatra and it was previously known only from North Sumatra and West Sumatra provinces. Here we report the first record of this species from South Sumatra. A male individual of *A. hageni* was recorded on 19 April 2018 in the Pagar Alam District, South Sumatra Province, Indonesia. Our record represent the most southern locality in the range of this rare highland species.

Key words: *Atrophaneura hageni*, batwing butterflies, distribution, Rhopalocera.

Introduction

The swallowtail butterflies (Lepidoptera: Papilionidae) is a group with more than 500 species globally (Kirton 2014; Hoskins 2015) that contains many of the largest and most beautiful butterfly species (Hill & Abang 2010). This family is known as swallowtail butterflies because many species have a tail at the hindwing tornus (Holloway *et al.* 2011). There are 24 Papilionidae species listed as precious and protected Indonesian butterflies, and these are the butterflies of highest conservation concern in Indonesia (Peggie 2011).

The genus *Atrophaneura* Reakirt, [1865] contains tailless butterflies somewhat resembling the genus *Troides* Hübner, [1819] at all stages of the life cycle. The larvae have large spiny tubercles, and feed on Aristolochiaceae (known as Birthwort) species. The adults are smaller than those of *Troides* and have shorter wings (Ek-Amnuay 2012). There are up to five *Atrophaneura* species in Sumatra. *Atrophaneura hageni* is a rare endemic butterfly having a range restricted to mountain forest in northern Sumatra (North Sumatra to West Sumatra provinces) (de Niceville & Martin 1896; D'Abrera 1982; Tsukada & Nishiyama 1982).

This paper report the first record of *A. hageni* in the South Sumatra Province, providing an evidence that the range of this species may be broader than it was expected.

Materials and Methods

A field survey was conducted on 19 April 2018 in Kampung Empat Village ($4^{\circ}02'83"S$, $103^{\circ}08'99"E$), Dempo Selatan Subdistrict, Pagar Alam District, South Sumatra Province (Figs. 1-2). This site is situated within the Dempo Mountain Protected Forest Area, in plant cover of height canopy which various plant species are abundant, e.g. *Ficus* spp., *Clibadium surinamensis* L., *Pandanus* sp., *Alsophila junghuhniana* Kunze, *Litsea* sp., *Lithocarpus* sp., *Schima wallichii* (DC.) Korth., and *Vanda* sp. The area is very close to the Dempo Mountain, bordering with a tea plantation. The altitude is 1,600 m above sea level, an area of lower montane forest in Sumatra (1.200 - 2.100 m) as defined by Whitten *et al.* (2000). On one occasion, our team observed and photographed a large black butterfly perching at *Ficus* sp. and *Alsophila junghuhniana*. Unfortunately, we are unable to collect a specimen due to a large distance and rapid movements of the butterfly. However, it was identified as *A. hageni* based on the combination of specific morphological characters using the series of photographic images (Figs. 3-4).




Figure 1. Distribution of *A. hageni* (map modified after Tsukada & Nishiyama 1982). Gray filling and area contoured by a dashed line indicate the range of *A. hageni*, and red circle indicates our new record in the Pagar Alam District, South Sumatra Province, Indonesia.

Results and Discussion

The butterfly was identified as a member of the genus *Atrophaneura* by its large-medium size, white dorsal side of the head, entirely black body, black forewing with whitish shadow on venations, black hindwing with a white large spot in the post discal and submarginal area, and with four series of black spots (Fig. 3-4). These patterns are constant for the *Atrophaneura priapus* group. In Sumatra, there are three species of white headed *Atrophaneura*, that is: *A. hageni*, *A. priapus* (Boisduval, 1836), and *A. sycorax* (Grose-Smith, 1885) (D'Abrera 1982; Tsukada & Nishiyama 1982).



Figure 2. Habitat of *A. hageni* in the Pagar Alam District, South Sumatra Province, Indonesia (Photo: Doni Setiawan).

The white headed *Atrophaneura* individual found in the Pagar Alam District was identified as *A. hageni* by its specific characters compared to *A. priapus* and *A. sycorax*. The butterfly differs from *Atrophaneura sycorax* by an entirely black dorsal surface of the abdomen and from *A. sycorax* by having a more than apical half of the body yellowish-white. The butterfly is very similar to *Atrophaneura priapus* by sharing white head and specific wing patterns. However, the latter species has more elongate hindwing, a white spot marking in hindwing extending between radius and discocellulars zone, and a smaller series of black spots within the white area in the hindwing (D'Abrera 1982; Tsukada & Nishiyama 1982; Harada *et al.* 2012). The butterfly from Pagar Alam District, South Sumatra differs from a set of characters listed above by its distinct white color in the head area, a more rounded hindwing, a white spot marking in hindwing absent or not extends to the area between radius and discocellulars zone, and by a larger series of black spots within the white area in the hindwing. Based on the specific characters listed above and appropriate guides (D'Abrera 1982; Tsukada & Nishiyama 1982; Harada *et al.* 2012), the butterfly found in the Pagar Alam district was identified as a male of *A. hageni*. The male was considered rarely seen than female (de Niceville & Martin 1896).

Atrophaneura hageni is endemic to the high-altitude regions of Sumatra. It was found in mountain jungles above 1,000 m altitude in the Karo Hill (3°N), North Sumatra Province, while its range extends to the Padang District (0°S), West Sumatra Province, in which this species occurred above 1,500 m altitude (Tsukada & Nishiyama 1982). However, it has never been reported from South Sumatra, and it was assumed that in the southern part of the island this species is replaced by *Atrophaneura priapus* (Tsukada & Nishiyama 1982). An old record labelled as Teluk Betung (Lampung Province) in 1928, coastal zone area of southern Sumatra (Creuwels 2020), must be error, because *A. hageni* is high altitude species. The record of *A. hageni* in the Pagar Alam District (4°S) represents the most southern occurrence of this species; approximately 1,000 km south of the Padang District.



Figure 3. Male of *A. hageni* found on 19 April 2018 in the Pagar Alam District, South Sumatra Province, Indonesia (Photo: Doni Setiawan).




Figure 4. The same male of *A. hageni* in flight when approaching by team, Pagar Alam District, South Sumatra Province, Indonesia (Photo: Guntur Pragustiandi).

Additional field surveys and collecting efforts are needed to estimate the status of *A. hageni* at the Pagar Alam District, i.e. if it is a local resident population or a single vagrant individual of this rare taxon. In our opinion, *A. hageni* was not previously recorded from South Sumatra due to the lack of entomological surveys in mountain forests of this area. This species is likely threatened by forest fragmentation and other kinds of anthropogenic impacts. Finally, our record is of importance for future checklists of Papilionidae at regional and global levels (Scriber 1995), particularly as a tropic archipelagic country, Indonesia has high level of endemism in animal distributions, including butterflies (Peggie 2011).

Acknowledgements

Our field survey in the Pagar Alam District was funded by the PT. Pertamina (Persero) Integrated Terminal Palembang, and by a PNBP Faculty of Science of Sriwijaya University research grant 2020 to the first author. We also like to thank anonymous reviewers for their constructive comments on earlier versions of this manuscript. We thank Colin Trainor (Charles Darwin University, Darwin, Australia) for checking English grammar to make this paper improve.

References

- Creuwels, J. (2020) *Naturalis Biodiversity Center (NL) - Lepidoptera*. *Naturalis Biodiversity Center*. Occurrence dataset <https://doi.org/10.15468/n4q0sa> accessed via GBIF.org on 2020-02-07. <https://www.gbif.org/occurrence/2443598972>
- D'Abrera, B. (1982) *Butterflies of the Oriental Region, Part I*. Hill House, Victoria, 244 pp.
- Ek-Amnuay, P. (2012) *Butterflies of Thailand*. Baan Lae Suan Amarin Printing and Publishing, Bangkok, 943 pp.
- Harada, M., Teshirogi, M., Ozawa, H. & Yago, M. (2012) *Catalogue of the Suguru Igarashi Insect Collection, The University Museum, The University of Tokyo. Part I. Lepidoptera, Papilionidae*. The University of Tokyo, Tokyo. <http://umdb.um.u-tokyo.ac.jp/DDoubutu/igarashi/en/cabinet2.php?Number=1063&skip=0&max=200> (Accessed on 02.01.2020).
- Hill, D.S. & Abang, F. (2010) *The insects of Borneo (including South-east Asia)*. Universiti Malaysia Sarawak, Sarawak, 435 pp.
- Holloway, J.D., Kibby, G. & Peggie, D. (2001) *The families of Malesian moths and butterflies*. Brill, Leiden, 455 pp.
- Hoskins, A. (2015) *Butterflies of the world*. Reed New Holland Publishers Pty Ltd, London, 312 pp.
- Kirton, L.G. (2014) *A naturalist's guide to the butterflies of Peninsular Malaysia, Singapore and Thailand*. John Beaufoy Publishing, Oxford, 176 pp.
- de Niceville, L. & Martin, L. (1896). A list of the butterflies of Sumatra, with especial reference to the species occurring in the north-east of the Island. *Journal of the Asiatic Society of Bengal [II]*, 64, 357–555.
- Peggie, D. (2011) *Precious and protected Indonesian butterflies*. PT Binamitra Megawarna, Jakarta, 72 pp.
- Scriber, J.M. (1995) Overview of swallowtail butterflies: Taxonomic and distributional latitude. In: Scriber, J.M., Tsubaki, Y. & Lederhouse, R.C. (Eds.), *Swallowtail butterflies: Their ecology and evolutionary biology*. Scientific Publishers, Gainesville, pp. 3–15.
- Tsukada, E. & Nishiyama, Y. (1982) *Butterflies of the South East Asian islands: Volume I Family Papilionidae*. In: Tsukada, E (Ed.), *Butterflies of the South East Asian islands*. Plapac Co., Ltd, Tokyo, pp. 1–457.
- Whiten, T., Damanik, S.J., Anwar, J. & Hisyam, N. (2000) *The ecology of Sumatra*. Periplus, Singapore, 478 pp.

**SJR**

Scimago Journal & Country Rank

Enter Journal Title, ISSN or Publisher Name

[Home](#)[Journal Rankings](#)[Country Rankings](#)[Viz Tools](#)[Help](#)[About Us](#)

Ads by Google

[Stop seeing this ad](#)[Why this ad?](#)

Ecologica Montenegrina

7

H Index

Country	Montenegro - SIR Ranking of Montenegro
Subject Area and Category	Agricultural and Biological Sciences Animal Science and Zoology Ecology, Evolution, Behavior and Systematics Insect Science Plant Science
Publisher	
Publication type	Journals
ISSN	23370173, 23369744
Coverage	2014-2020
Scope	Ecologica Montenegrina (ISSN 2336-9744 (online) ISSN 2337-0173 (print)) is peer-reviewed journal in which scientific articles and reports are quickly published. The papers are in the fields of taxonomy, biogeography and ecology (for example: new taxa for science, taxonomic revision, and/or fundamental ecology and biogeography papers). Open access publishing option is strongly encouraged for authors with research grants and other funds. For those without grants/funds, all accepted manuscripts will be published but access is secured for subscribers only.
Homepage	
How to publish in this journal	
Contact	
Join the conversation about this journal	











Ads by Google

[Stop seeing this ad](#)[Why this ad?](#)


Quartiles







ECOLOGICA MONTENEGRINA



Volume 28
February 2020

Ecologica Montenegrina

HOME ABOUT LOGIN REGISTER SEARCH
 CURRENT ARCHIVES ANNOUNCEMENTS

Home > About the Journal > Editorial Team

Editorial Team

Editor in Chief

Vladimir Pešić, Department of Biology, University of Montenegro, Montenegro

Editorial Board

Badamdorj Bayartogtokh, National University of Mongolia, Mongolia
Holger Braun, Museo de La Plata, Argentina

Dr Patricio De los Rios Escalante, Universidad Católica de Temuco, Chile

Sergey G. Ermilov, Tyumen State University, Tyumen, Russia, Russian Federation

Igor Dovgal, A. Kovalevsky Institute of Marine Biological Research, Russian Federation

Thibault Datry, Irstea, UR MALY, Centre de Lyon-Villeurbanne, France

Zoltán Fehér, Natural History Museum Vienna 3rd Zoology Department, Austria

Dilian Georgiev, University of Plovdiv, Bulgaria

Reinhard Gerecke

Wolfram Graf, University of Natural Resources and Life Sciences, Vienna, Austria

Michał Grabowski, Department of Invertebrate Zoology and Hydrobiology, University of Łódź, Poland

Clemens Grosser, Germany

Borislav Gueorguiev, National Museum of Natural History, Sofia, Bulgaria

Sead Hadžiblahović, Environmental Protection Agency of Montenegro, Montenegro

Dr Laith A. Jawad, Flat Bush, Manukau, Auckland, New Zealand, New Zealand

Dr Roman Yakovlev, Altai State University, Russian Federation

Max Kasparek, Zoology in the Middle East, Germany

Gordan S. Karaman, Montenegrin Academy of Sciences, Montenegro

Jeno Kontschán, Plant Protection Institute, Centre for Agricultural Research, Hungarian Academy of Sciences, Hungary

Andrey Kostianoy, P.P.Shirshov Institute of Oceanology, Russian Academy of Sciences, Moscow, Russia, Russian Federation

Katarina Ljubisavljević, Montenegro

Dr Andrzej Zawal, Department of Invertebrate Zoology & Limnology, University of Szczecin, 71-415 Szczecin, Waska 13, Poland, Poland

Slobodan Evgenije Makarov, University of Belgrade Faculty of Biology, Serbia

Dávid Murányi, Hungarian Natural History Museum Department of Zoology

Baross u. 13, H-1088 Budapest, Hungary, Hungary

Momir M Paunović, Institute for Biological Research "Sinisa Stankovic" University of Belgrade Republic of Serbia Bulevar despota Stefana 142, 11000 Belgrade, Serbia, Serbia

Radmila Petanović, University of Belgrade - Faculty of Agriculture Department of Entomology and Agricultural Zoology Nemanjina 6, P.O. BOX 127 11080 Belgrade - Zemun SERBIA, Serbia

Prof. Alireza Saboori, University of Tehran, Iran, Islamic Republic of

Zoltán Sándor Varga, University of Debrecen, Hungary

Dr Anton V. Volynkin, Altai State University

Paul Selden, University of Kansas, United States

Nelli Grigoryevna Sergeeva, Institute of Marine Biological Researches RAS, Russian Federation

Harry Smit, Naturalis Biodiversity Center, Netherlands

Boris Sket, Univerza v Ljubljani, Slovenia

Vesna Vukasinovic Petic, Department of Technology, University of Montenegro, Podgorica, Montenegro

OPEN JOURNAL SYSTEMS

Journal Help

USER

Username

Password

Remember me

NOTIFICATIONS

- View
- Subscribe

LANGUAGE

Select Language

JOURNAL CONTENT

Search

Search Scope

Browse

- By Issue
- By Author
- By Title
- Other Journals

FONT SIZE

INFORMATION

- For Readers
- For Authors
- For Librarians

KEYWORDS

Amphipoda Black Sea.
Bythinella Carabidae.
China China. Croatia.
Gastropoda Greater Antilles.
Greece Halacaroidea.
Hirudinea Hungary.
Hydrobiidae.
Hydryphantidae.
Macedonia Mollusca.
Montenegro.
Montenegro. New species.
Republic of.
Adygaea Salicidae.
Serbia. Taxonomy.
Turkey. Ukraine.
biodiversity checklist.
distribution ecology.
faunistics female.
identification key.
jumping spiders.

ISSN: 2336-9744

Ecologica Montenegrina

HOME ABOUT LOGIN REGISTER SEARCH
 CURRENT ARCHIVES ANNOUNCEMENTS

Home > Archives > Vol 28 (2020)

Vol 28 (2020)

Table of Contents

Articles

A new genus and two new species (Gastropoda, Hydrobiidae) from Morocco	PDF 1-6
Peter Glöer, Youness Mabrouki, Abdelkhaleq Fouzi Taybi	
Communities of the Bidentetea class of small coastal river valleys of the Western Pomerania (Poland)	PDF 8-19
Edyta Stępień, Stanisław Rosadziński	
First reliable record of many-plumed moth <i>Pterotopteryx spilodesma</i> (Lepidoptera, Alucitidae) on Sakhalin Island	PDF 23-25
Olga Titova, Peter Ustjuzhanin, Vasiliy Kovtunovich, Alexander Streltsov	
First record of Hagen's batwing <i>Atrophaneura hageni</i> (Rogenhofer, 1889) (Lepidoptera: Papilionidae) in southern Sumatra, Indonesia	PDF 26-30
Doni Setiawan, Ina Aprilia, Muhammad Iqbal, Guntur Pragustiandi, Arum Setiawan, Indra Yustian	
Recent invasion of the Lime Swallowtail <i>Papilio demoleus</i> (Lepidoptera: Papilionidae) to Seychelles	PDF 31-39
Yulia S. Kolosova, Ivan N. Bolotov	
Contribution to the knowledge of the Carpenter Moths (Lepidoptera, Cossidae) of the Maputo Special Reserve in South Mozambique with description of two new species	PDF 40-52
Roman V. Yakovlev, Gyula M. László, Alvaro A. Vetina	
New data on <i>Ovalisia (Palmar) festiva</i> (Linnaeus) (Coleoptera: Buprestidae) and its natural enemies reported from Bulgaria	PDF 53-60
Stoyka Ruseva, Ivaylo Todorov, Anelya Pencheva	
New data on the family Himantariidae Bollman, 1893 (Chilopoda: Geophilomorpha) from Kazakhstan	PDF 61-66
Yuri V. Dyachkov	

Correspondence

<i>Bythinella rethymnonensis</i> n. sp., replacement name for <i>Bythinella magdalena</i> Glöer & Hirschfelder 2019	PDF 7
Peter Glöer, Hans-Jürgen Hirschfelder	
New site of <i>Cordulegaster heros</i> Theischinger, 1979 in western Slovakia – the confirmation of its northern border of range of occurrence in Europe (Odonata: Cordulegastridae)	PDF 20-22
Paweł Buczyński, Grzegorz Michoński, Andrzej Zawal	

ISSN: 2336-9744

[OPEN JOURNAL SYSTEMS](#)

[Journal Help](#)

USER

Username
 Password
 Remember me

NOTIFICATIONS

- [View](#)
- [Subscribe](#)

LANGUAGE

Select Language

JOURNAL CONTENT

Search
 Search Scope

Browse

- [By Issue](#)
- [By Author](#)
- [By Title](#)
- [Other Journals](#)

FONT SIZE

INFORMATION

- [For Readers](#)
- [For Authors](#)
- [For Librarians](#)

KEYWORDS

[Amphipoda](#) [Black Sea](#).
[Bythinella](#) [Carabidae](#).
[China](#) [China](#). [Croatia](#).
[Gastropoda](#) [Greater Antilles](#).
[Greater Antilles](#).
[Greece](#) [Halacaroidea](#).
[Hungary](#).
[Hydrobiidae](#).
[Hydryphantidae](#).
[Macedonia](#) [Mollusca](#).
[Montenegro](#).
[Montenegro](#). [New species](#) [Republic of Adygeya](#).
[Salticidae](#).
[Serbia](#). [Taxonomy](#).
[Turkey](#). [Ukraine](#).
[biodiversity](#). [checklist](#).
[distribution](#). [ecology](#).
[faunistics](#). [female identification key](#).
[jumping spiders](#).

First record of Hagen's batwing *Atrophaneura hageni* (Rogenhofer, 1889) (Lepidoptera Papiolinidae) in southern Sumatra, Indonesia

By Arum Setiawan



Ecologica Montenegrina
ISSN 2339-8744 (online); ISSN 2339-0173 (print)

Ecologica Montenegrina 28: 26-30 (2020)
This journal is available online at: www.biota.org/em

Article

First record of Hagen's batwing *Atrophaneura hageni* (Rogenhofer, 1889) (Lepidoptera: Papiolinidae) in southern Sumatra, Indonesia

DONI SETIAWAN¹, INA APRILIA², MUHAMMAD IQBAL^{3*}, GUNTUR PRAGUSTIANDI²,
ARUM SETIAWAN¹ & INDRA YUSTIAN¹

¹Department of Biology, Faculty of Science, Sriwijaya University, Jalan Raya Palembang-Prabumulih km 32, Indralaya, Sumatera Selatan 30662, Indonesia

²Conservation Biology Program, Faculty of Science, Sriwijaya University, Jalan Padang Selasa 524, Palembang, Sumatera Selatan 30129, Indonesia

³Biology Program, Faculty of Science, Sriwijaya University, Jalan Padang Selasa 524, Palembang, Sumatera Selatan 30129, Indonesia

*Corresponding author: kpbsos26@yahoo.com

Received 24 December 2019 | Accepted by V. Pešić: 7 February 2020 | Published online 13 February 2020.

Abstract

Hagen's batwing *Atrophaneura hageni* (Rogenhofer, 1889) (Lepidoptera: Papiolinidae) is endemic to Sumatra and it was previously known only from North Sumatra and West Sumatra provinces. Here we report the first record of this species from South Sumatra. A male individual of *A. hageni* was recorded on 19 April 2018 in the Pagar Alam District, South Sumatra Province, Indonesia. Our record represent the most southern locality in the range of this rare highland species.

Key words: *Atrophaneura hageni*, batwing butterflies, distribution, Rhopalocera.

Introduction

The swallowtail butterflies (Lepidoptera: Papilionidae) is a group with more than 500 species globally (Kirton 2014; Hoskins 2015) that contains many of the largest and most beautiful butterfly species (Hill & Abang 2010). This family is known as swallowtail butterflies because many species have a tail at the hindwing tornus (Holloway *et al.* 2011). There are 24 Papilionidae species listed as precious and protected Indonesian butterflies, and these are the butterflies of highest conservation concern in Indonesia (Peggie 2011).

The genus *Atrophaneura* Reakirt, [1865] contains tailless butterflies somewhat resembling the genus *Troides* Hübner, [1819] at all stages of the life cycle. The larvae have large spiny tubercles, and feed on Aristolochiaceae (known as Birthwort) species. The adults are smaller than those of *Troides* and have shorter wings (Ek-Amnuay 2012). There are up to five *Atrophaneura* species in Sumatra. *Atrophaneura hageni* is a rare endemic butterfly having a range restricted to mountain forest in northern Sumatra (North Sumatra to West Sumatra provinces) (de Niceville & Martin 1896; D'Abrera 1982; Tsukada & Nishiyama 1982).

This paper report the first record of *A. hageni* in the South Sumatra Province, providing an evidence that the range of this species may be broader than it was expected.

Materials and Methods

A field survey was conducted on 19 April 2018 in Kampung Empat Village ($4^{\circ}02'83"S$, $103^{\circ}08'99"E$), Dempo Selatan Subdistrict, Pagar Alam District, South Sumatra Province (Figs. 1-2). This site is situated within the Dempo Mountain Protected Forest Area, in plant cover of height canopy which various plant species are abundant, e.g. *Ficus* spp., *Clibadium surinamensis* L., *Pandanus* sp., *Alsophila junghuhniana* Kunze, *Litsea* sp., *Lithocarpus* sp., *Schima wallichii* (DC.) Korth., and *Vanda* sp. The area is very close to the Dempo Mountain, bordering with a tea plantation. The altitude is 1,600 m above sea level, an area of lower montane forest in Sumatra (1.200 - 2.100 m) as defined by Whitten *et al.* (2000). On one occasion, our team observed and photographed a large black butterfly perching at *Ficus* sp. and *Alsophila junghuhniana*. Unfortunately, we are unable to collect a specimen due to a large distance and rapid movements of the butterfly. However, it was identified as *A. hageni* based on the combination of specific morphological characters using the series of photographic images (Figs. 3-4).




Figure 1. Distribution of *A. hageni* (map modified after Tsukada & Nishiyama 1982). Gray filling and area contoured by a dashed line indicate the range of *A. hageni*, and red circle indicates our new record in the Pagar Alam District, South Sumatra Province, Indonesia.

Results and Discussion

The butterfly was identified as a member of the genus *Atrophaneura* by its large-medium size, white dorsal side of the head, entirely black body, black forewing with whitish shadow on venations, black hindwing with a white large spot in the post discal and submarginal area, and with four series of black spots (Fig. 3-4). These patterns are constant for the *Atrophaneura priapus* group. In Sumatra, there are three species of white headed *Atrophaneura*, that is: *A. hageni*, *A. priapus* (Boisduval, 1836), and *A. sycorax* (Grose-Smith, 1885) (D'Abrera 1982; Tsukada & Nishiyama 1982).



Figure 2. Habitat of *A. hageni* in the Pagar Alam District, South Sumatra Province, Indonesia (Photo: Doni Setiawan).

The white headed *Atrophaneura* individual found in the Pagar Alam District was identified as *A. hageni* by its specific characters compared to *A. priapus* and *A. sycorax*. The butterfly differs from *Atrophaneura sycorax* by an entirely black dorsal surface of the abdomen and from *A. sycorax* by having a more than apical half of the body yellowish-white. The butterfly is very similar to *Atrophaneura priapus* by sharing white head and specific wing patterns. However, the latter species has more elongate hindwing, a white spot marking in hindwing extending between radius and discocellulars zone, and a smaller series of black spots within the white area in the hindwing (D'Abrera 1982; Tsukada & Nishiyama 1982; Harada *et al.* 2012). The butterfly from Pagar Alam District, South Sumatra differs from a set of characters listed above by its distinct white color in the head area, a more rounded hindwing, a white spot marking in hindwing absent or not extends to the area between radius and discocellulars zone, and by a larger series of black spots within the white area in the hindwing. Based on the specific characters listed above and appropriate guides (D'Abrera 1982; Tsukada & Nishiyama 1982; Harada *et al.* 2012), the butterfly found in the Pagar Alam district was identified as a male of *A. hageni*. The male was considered rarely seen than female (de Nicewill & Martin 1896).

Atrophaneura hageni is endemic to the high-altitude regions of Sumatra. It was found in mountain jungles above 1,000 m altitude in the Karo Hill (3°N), North Sumatra Province, while its range extends to the Padang District (0°S), West Sumatra Province, in which this species occurred above 1,500 m altitude (Tsukada & Nishiyama 1982). However, it has never been reported from South Sumatra, and it was assumed that in the southern part of the island this species is replaced by *Atrophaneura priapus* (Tsukada & Nishiyama 1982). An old record labelled as Teluk Betung (Lampung Province) in 1928, coastal zone area of southern Sumatra (Crewwels 2020), must be error, because *A. hageni* is high altitude species. The record of *A. hageni* in the Pagar Alam District (4°S) represents the most southern occurrence of this species; approximately 1,000 km south of the Padang District.



Figure 3. Male of *A. hageni* found on 19 April 2018 in the Pagar Alam District, South Sumatra Province, Indonesia (Photo: Doni Setiawan).




Figure 4. The same male of *A. hageni* in flight when approaching by team, Pagar Alam District, South Sumatra Province, Indonesia (Photo: Guntur Pragustiandi).

Additional field surveys and collecting efforts are needed to estimate the status of *A. hageni* at the Pagar Alam District, i.e. if it is a local resident population or a single vagrant individual of this rare taxon. In our opinion, *A. hageni* was not previously recorded from South Sumatra due to the lack of entomological surveys in mountain forests of this area. This species is likely threatened by forest fragmentation and other kinds of anthropogenic impacts. Finally, our record is of importance for future checklists of Papilionidae at regional and global levels (Scriber 1995), particularly as a tropic archipelagic country, Indonesia has high level of endemism in animal distributions, including butterflies (Peggie 2011).

Acknowledgements

Our field survey in the Pagar Alam District was funded by the PT. Pertamina (Persero) Integrated Terminal Palembang, and by a PNBP Faculty of Science of Sriwijaya University research grant 2020 to the first author. We also like to thank anonymous reviewers for their constructive comments on earlier versions of this manuscript. We thank Colin Trainor (Charles Darwin University, Darwin, Australia) for checking English grammar to make this paper improve.

References

- Creuwels, J. (2020) *Naturalis Biodiversity Center (NL) - Lepidoptera*. Naturalis Biodiversity Center. Occurrence dataset <https://doi.org/10.15468/n4q0sa> accessed via GBIF.org on 2020-02-07. <https://www.gbif.org/occurrence/2443598972>
- D'Abrera, B. (1982) *Butterflies of the Oriental Region, Part I*. Hill House, Victoria, 244 pp.
- Ek-Amnuay, P. (2012) *Butterflies of Thailand*. Baan Lae Suan Amarin Printing and Publishing, Bangkok, 943 pp.
- Harada, M., Teshirogi, M., Ozawa, H. & Yago, M. (2012) *Catalogue of the Suguru Igarashi Insect Collection, The University Museum, The University of Tokyo. Part I. Lepidoptera, Papilionidae*. The University of Tokyo, Tokyo. <http://umdb.um.u-tokyo.ac.jp/DDoubutu/igarashi/en/cabinet2.php?Number=1063&skip=0&max=200> (Accessed on 02.01.2020).
- Hill, D.S. & Abang, F. (2010) *The insects of Borneo (including South-east Asia)*. Universiti Malaysia Sarawak, Sarawak, 435 pp.
- Holloway, J.D., Kibby, G. & Peggie, D. (2001) *The families of Malesian moths and butterflies*. Brill, Leiden, 455 pp.
- Hoskins, A. (2015) *Butterflies of the world*. Reed New Holland Publishers Pty Ltd, London, 312 pp.
- Kirton, L.G. (2014) *A naturalist's guide to the butterflies of Peninsular Malaysia, Singapore and Thailand*. John Beaufoy Publishing, Oxford, 176 pp.
- de Niceville, L. & Martin, L. (1896). A list of the butterflies of Sumatra, with especial reference to the species occurring in the north-east of the Island. *Journal of the Asiatic Society of Bengal [II]*, 64, 357–555.
- Peggie, D. (2011) *Precious and protected Indonesian butterflies*. PT Binamitra Megawarna, Jakarta, 72 pp.
- Scriber, J.M. (1995) Overview of swallowtail butterflies: Taxonomic and distributional latitude. In: Scriber, J.M., Tsubaki, Y. & Lederhouse, R.C. (Eds.), *Swallowtail butterflies: Their ecology and evolutionary biology*. Scientific Publishers, Gainesville, pp. 3–15.
- Tsukada, E. & Nishiyama, Y. (1982) *Butterflies of the South East Asian islands: Volume I Family Papilionidae*. In: Tsukada, E. (Ed.), *Butterflies of the South East Asian islands*. Plapac Co., Ltd, Tokyo, pp. 1–457.
- Whiten, T., Damanik, S.J., Anwar, J. & Hisyam, N. (2000) *The ecology of Sumatra*. Periplus, Singapore, 478 pp.

First record of Hagen's batwing *Atrophaneura hageni* (Rogenhofer, 1889) (Lepidoptera Papiolinidae) in southern Sumatra, Indonesia

ORIGINALITY REPORT

2%
SIMILARITY INDEX

MATCH ALL SOURCES (ONLY SELECTED SOURCE PRINTED)

★ "Carbon Sequestration Potential of Agroforestry Systems", Springer
Nature, 2011 1%

[Crossref](#)

EXCLUDE QUOTES ON

EXCLUDE MATCHES < 1%

EXCLUDE
BIBLIOGRAPHY ON

FORMAT PENILAIAN (VALIDASI & PEER REVIEW)
LEMBAR
HASIL PENILAIAN SEJAWAT SEBIDANG ATAU PEER REVIEW
KARYA ILMIAH : JURNAL ILMIAH

Jurnal Artikel Ilmiah	: First record of Hagen's batwing Atrophaneura hageni (Rogenhofer, 1889) (Lepidoptera: Papilionidae) in southern Sumatra, Indonesia
Penulis Artikel Ilmiah	: Arum Setiawan
Identitas Jurnal Artikel Ilmiah	: a. Nama Jurnal : Ecologica Montenegrina b. Nomor/Volume/Hal : 1/28/26-30 c. Edisi (bulan/tahun) : Februari 2020 d. Penerbit : Center for Biodiversity of Montenegro e. Jumlah Halaman 5
Kategori Publikasi Jurnal Ilmiah (beri ✓ pada kategori yang tepat)	: <input checked="" type="checkbox"/> Jurnal Ilmiah Internasional Bereputasi <input type="checkbox"/> Jurnal Ilmiah Internasional <input type="checkbox"/> Jurnal Ilmiah Nasional Terakreditasi S1, S2 <input type="checkbox"/> Jurnal Ilmiah Nasional Terakreditasi S3, S4 <input type="checkbox"/> Jurnal Ilmiah Nasional Tidak Terakreditasi.

I. Hasil Penilaian Validasi :

No.	ASPEK	URAIAN/KOMENTAR PENILAIAN
1.	Indikasi Plagiasi	2 %
2.	Linearitas	Sudah linier dengan bidang biologi konservasi

II. Hasil Penilaian Peer Review :

Komponen Yang Dinilai	Nilai Maksimal Jurnal Ilmiah (isikan di kolom yang sesuai)					Nilai Akhir Yang Diperoleh
	Internasional Bereputasi (Maks 40)	Internasional (Maks 20)	Nasional Terakreditasi S1, S2 Maks 25	Nasional Terakreditasi S3, S4 Maks 20	Nasional tidak Terakreditasi (maks 10)	
Kelengkapan dan Kesesuaian unsur isi jurnal (10%)	4					4
Ruang lingkup dan kedalaman pembahasan (30%)	12					12
Kecukupan dan Kemutahiran data/informasi dan metodologi (30%)	12					12
Kelengkapan unsur dan kualitas penerbit (30%)	12					12
Total = (100%)	40					40
Kontribusi Pengusul (Penulis Pertama /Anggota Utama)	Anggota Utama $(0,4 \times 40) / 5 = 3,2$					3,2

KOMENTAR/ULASAN PEER REVIEW

• Kelengkapan dan Kesesuaian Unsur:	Paper terkait deskripsi kupu-kupu Atrophaneura hageni di Sumatera Selatan. Isi paper sudah memenuhi kaidah-kaidah karya ilmiah dan sudah sesuai dengan bidang biologi konservasi
• Ruang Lingkup dan Kedalaman Pembahasan:	Hasil penelitian dibahas secara komprehensif dengan penyampaian pembanding dari temuan-temuan penelitian lainnya dan teori terkait. Referensi yang diacu dalam pembahasan sudah cukup update untuk bidang kajian ini.
• Kecukupan & Kemutahiran Data & Metodologi:	Data-data hasil penelitian sudah baik dan didukung peta lokasi sampling dan gambar yang ditampilkan menarik. Data didapatkan dengan menggunakan metode yang sudah standard.
• Kelengkapan Unsur & Kualitas Penerbit:	Penerbit Center for Biodiversity of Montenegro berkualitas baik, tidak termasuk predatory publisher, dan jurnal masuk di Q2.

Surabaya, 15 Mei 2020
Penilai 1



Prof. Hery Purnobasuki, M.Si., Ph.D.
NIP 196705071991021001
Unit Kerja : Jurusan Biologi FST Unair
Bidang Ilmu : Biologi
Jabatan/Pangkat : Guru Besar/ Pembina Utama Madya

FORMAT PENILAIAN (VALIDASI & PEER REVIEW)

LEMBAR

HASIL PENILAIAN SEJAWAT SEBIDANG ATAU PEER REVIEW

KARYA ILMIAH : JURNAL ILMIAH

Jurnal Artikel Ilmiah

: First record of Hagen's batwing Atrophaneura hageni (Rogenhofer, 1889)
 (Lepidoptera: Papiolinidae) in southern Sumatra, Indonesia

Penulis Artikel Ilmiah

: Arum Setiawan

Identitas Jurnal Artikel Ilmiah

: a. Nama Jurnal : Ecologica Montenegrina

b. Nomor/Volume/Hal : 1/28/26-30

c. Edisi (bulan/tahun) : Februari 2020

d. Penerbit : Center for Biodiversity of Montenegro

e. Jumlah Halaman : 5

Kategori Publikasi Jurnal Ilmiah
 (beri V pada kategori yang tepat)

: Jurnal Ilmiah Internasional Bereputasi

Jurnal Ilmiah Internasional

Jurnal Ilmiah Nasional Terakreditasi S1, S2

Jurnal Ilmiah Nasional Terakreditasi S3, S4

Jurnal Ilmiah Nasional Tidak Terakreditasi


I. Hasil Penilaian Validasi :

No.	ASPEK	URAIAN/KOMENTAR PENILAIAN
1.	Indikasi Plagiasi	2 %
2.	Linearitas	V

II. Hasil Penilaian Peer Review :

Komponen Yang Dinilai	Nilai Maksimal Jurnal Ilmiah (isikan di kolom yang sesuai)					Nilai Akhir Yang Diperoleh
	Internasional Bereputasi (Maks 40)	Internasional (Maks 20)	Nasional Terakreditasi S1, S2 Maks 25	Nasional Terakreditasi S3, S4 Maks 20	Nasional tidak Terakreditasi (maks 10)	
Kelengkapan dan Kesesuaian unsur isi jurnal (10%)	4					3
Ruang lingkup dan kedalaman pembahasan (30%)	12					11
Kecukupan dan Kemutahiran data /informasi dan metodologi (30%)	12					12
Kelengkapan unsur dan kualitas penerbit (30%)	12					12
Total = (100%)	40					36
Kontribusi Pengusul (Penulis Pertama /Anggota Utama)	Journal of Ecologica Montenegrina Vol. 28(1) 2020. SJR 0,37; IF 0,79. Penulis ke 5 dari 6 penulis. Nilai maksimal 90%. Nilai pengusul = $(0,4 \times 0,9 \times 40)/5 = 2,88$					2,88
KOMENTAR/ULASAN PEER REVIEW						
• Kelengkapan dan Kesesuaian Unsur:	Format kurang lengkap.					
• Ruang Lingkup dan Kedalaman Pembahasan	Ruang lingkup masih terkait. Dalam judul tertulis Sumatra (yang baku adalah Sumatera). Pembahasan cukup mendalam.					
• Kecukupan & Kemutakhiran Data & Metodologi:	Kesimpulan kurang jelas dan tidak sambung dengan latar belakang. Similarity hanya 2% artinya tidak ada plagiasi.					
• Kelengkapan Unsur & Kualitas Penerbit:	Penerbit sangat baik.					

Yogyakarta, 5 Juni 2020

Penilai


tanda tangan

Prof. Dr. Suwarno Hadisusanto

NIP 195411161983031002

Unit Kerja : Fakultas Biologi UGM

Bidang Ilmu : Biologi/Ekologi

Jabatan/Pangkat : Guru Besar/ Pembina Utama Madya