

Bioactive Compounds Evaluation of the Mudskippers in the Estuarine Area of Musi River, South Sumatera, Indonesia

Mohammad Rasyid Ridho^{1*}, Arum Setiawan¹, Sarno¹, Arwinskyah¹, Enggar Patriono¹, Sulistiono¹

¹ Department of Biology, Faculty of Mathematic and Natural Sciences, Sriwijaya University, Palembang 30139, Indonesia

* Corresponding author's e-mail: moh.rasyidridho@mipa.unsri.ac.id

ABSTRACT

This research was conducted to assess various bioactive compounds in the meat of mudskippers found in the Sungsang 2 Village, Banyuasin regency, South Sumatra. The research had been synthesized using n-hexane from February to May 2018 at Agricultural Technology Production Laboratory, Sriwijaya University. Subsequently, the mudskippers (*Periophthalmus chrysospilos*, *Boleophthalmus boddarti*, and *Periophthalmodon schlosseri*) extract was synthesized and further investigate by conducting the GC-MS analysis. The study of bioactive compounds in mudskippers was carried out by evaporating in vacuum evaporation at 30°C. The results showed that *Periophthalmus chrysospilos* has five classes of bioactive compounds such as steroids, terpenoids, carotenoids, cannabinoids, and alkaloids and *Boleophthalmus boddarti* has five classes of bioactive compounds including steroids, carotenoids, terpenoids, bufadienolides, and carotatoxins. Furthermore, *Boleophthalmus dussumieri* has four classes of bioactive compounds such as steroids, carotenoids, terpenoids, and carotatoxins and *Periophthalmodon schlosseri* has three classes of bioactive compounds including steroids, carotenoids and terpenoids. This research discussed the potential benefit of all bioactive compounds in mudskippers for alleviating and treating children with enuresis through the review of literature on the most cited bioactive compounds.

Keywords: mudskippers, Musi river, bioactive compounds

INTRODUCTION

Mudskipper is a genus from the family of Gobiidae and subfamily Oxudercinae that is habitually known as ikan gelodok at Musi river, South Sumatera (Shukla et al., 2014; Polgar et al., 2017; Lauriano et al., 2018; Ridho et al., 2019). Mudskippers are commonly found in brackish river water and beaches with certain salinity concentration, as well as mangrove areas of the Indo-Pacific region (Tytler and Vaughan, 1983). Mudskippers are amphibious fish that are able to tolerate both fresh and marine water on muddy and sandy bottom habitat, spawning in estuarine areas (Hu et al., 2016; Hong et al., 2017). Mudskippers are also capable of spending extended periods out of water (Martin et al., 2013; Konstantinidis et al., 2016; Martinez et al., 2018). They appear during low

tide, foraging on tropical mudflats, seeking mate and preserving territories for some portion of their daily cycle (Dabruzzi et al., 2019). According to the classification of amphibious gobies, 40 species of mudskippers have been currently identified in the world, which are divided into four genera of Oxudercinae, namely *Boleophthalmus*, *Periophthalmus*, *Periophthalmodon* and *Scartelaos* (Clayton 1993; Graham 1997; Graham and Wegner 2010; Ishimatsu and Gonzales 2011; Takeda et al., 2012; Murdy 2011a; You et al., 2018).

Juveniles of this Gobiidae are herbivores that eat more diatoms and filamentous algae, while the major food of adult Gobiidae are crustaceans (95.5%), polychaetas (86.4%) and land insects (85%) (Bob-Manuel, 2011). The burrows of these habitats are categorically suitable for the mudskippers for feeding grounds, spawning

areas and nursery grounds (Masuda et al., 1984; Bob-Manuel, 2011; Satapoomin and Poovachiranon, 1997; Redjeki, 2013). Mudskippers are commonly consumed and employed in traditional medicine in many countries such as China, Malaysia and Japan (Feulner 2013; Gadhavi et al., 2017; Ikram et al., 2010; looi et al., 2016 and Yang, X. X., 2017). The medicinal mudskippers were often prepared from the natural products which contain different bioactive compounds (Mahadevan et al., 2019; Wu et al., 2009). On the other hand, many fishermen and general public around the Musi river ignore the existence and benefits of mudskippers.

Corticosteroids, one of bioactive compounds in mudskippers, are steroid hormone class for overcoming inflammations (Panda and Mabalirajan, 2018; Heughten et al., 2018). These hormones are a contributing factor to alleviating enuresis (bedwetting) in child. Enuresis is most common among children or young persons and is denoted by repeated voiding of urine into the bed or clothing (Apos et al., 2017 and Shafi et al., 2019). The study area corresponded to bioactive compounds like corticosteroid compounds in mudskippers that can reduce the enuresis problem in children (Jain et al., 2016). The authors intended to evaluate the bioactive compounds in the four species mudskippers including *Periophthalmus chrysospilos*, *Boleophthalmus boddarti*, *Boleophthalmus dussumieri* and *Periophthalmodon schlosseri*.

MATERIALS AND METHODS

Fishes and experimental design

Mudskippers were obtained from an estuarine area around the Sungai 2 village, Banyuasin regency, South Sumatera. Mudskippers, apparently healthy, were randomly caught using fishing equipment and transported to the Agricultural Technology Production Laboratory, Sriwijaya University for cleaning and removing the bones from the fish. After the filleting process, the mudskippers were divided into four groups with *Periophthalmus chrysophilus* (8 grams), *Boleophthalmus boddarti* (100 grams), *Boleophthalmus dussumieri* (13.8 grams) and *Periophthalmodon schlosseri* (100 grams) which were homogenized.

The crushed fish flesh dissolved in n-hexane solvent with a sample ratio of 1: 4 and macerated for 24 hours. The next step involved filtering with calico cloth and filter paper and evaporating with vacuum evaporator at 30°C with a speed of 200 rpm. Each extract sample was reconstituted using n-hexane solvent, then each 1 µL was taken to be injected and analyzed in GC-MS alternately.

Analysis technique

The data in this research were expressed in the form of chromatograms that contain graphs and were equipped with a list of detected chemical compounds, along with the structure of compounds, retention times, areas, and probability of the compound types. The data analysis was performed by identifying the chemical compounds detected to find out the natural name (nature of compound name), synonym, and the class of compounds of these chemical compounds. Identification was done by comparing the compound name detected with a data base on the literature in the form of books and journals. The studies reported as full-text articles, those published as abstracts only and unpublished data were included as well.

Gas chromatography and mass spectroscopy (GC-MS) analysis

The qualitative analyses of the mudskipper extract were expressed with a GCMS-QP 2010 Plus (Shimadzu, Japan) system equipped under computer control at 70 eV (Mohammed and Imad, 2013). The methanol extract in the amount of 1 µl was entered into the GC-MS with a micro injector and followed by scanning for 45 min. The temperature was maintained in the oven at 100°C and Helium gas was utilized as a carrier as well as an eluent. The helium flow was arranged to 1 ml per min and electron energy was fired by a mass detector about 70 eV. The compound of elite 1 was used as separation column (100% dimethyl poly siloxane) (Hameed et al., 2016; Kareem et al., 2015). The identity of the compounds in the extracts was assigned by the comparison of their retention indices and mass spectra fragmentation patterns with those stored on the computer library and also with the data published in the literature. The compounds were identified by comparing their spectra to those of the Wiley and NIST/EPA/NIH mass spectral libraries.

RESULTS AND DISCUSSIONS

GC-MS results of the *periophthalmus chrysospilos* extract

The gas chromatography and mass spectroscopy analysis of bioactive compounds were carried out in the *Periophthalmus chrysospilos* extract that was presented in Figure 1. It can be seen that the vertical lines in the chromatogram pattern point out the short periods of time when the ion beam is deflected-off the plate (and thus off the beam monitor) which is necessary when changing the position of the photographic plate, a process that takes about 2–3 seconds with the present arrangement. This pen deflection creates a useful signal of the exact quantities of the chromatogram being noted with the mass spectrometer and facilitates the reconnection of the spectra on the plate with the chromatogram.

The GC-MS chromatogram showed that the presence of five major peaks of bioactive compounds was detected and the components corresponding to the peaks were determined as follows. The highest peak in chromatogram pattern could be seen that the extract of *Periophthalmus chrysospilos* has retention time 14.44 minutes. On the basis of the GC-MS signal, the sample has many bioactive compounds which were indicated by the signals detected. It shows that the GC-MS analysis successfully proves that the *Periophthalmus chrysospilos* extract is suitable for treating the people with inflammatory diseases, especially enurosis.

Bioactive compounds in the *Periophthalmus chrysospilos* extract

The results of *Periophthalmus chrysospilos* extract revealed the amounts of bioactive compounds such as terpenoids, alkaloids, steroids,

carotenoids and cannabinoids. As many as 4 types of terpenoid compounds were detected, , with a total of 4 compounds. One type of carotenoid compounds was detected with a total of 4 compounds. Additionally, one type of cannabinoid compounds was detected with a total of 2 compounds; and 1 alkaloid compound was detected with a total of 1 compound. The cannabinoid bioactive compounds were also detected in this sample with a total of 1 compound. Cannabinoid compounds include secondary metabolites which are usually found in cannabis plants. According to Maggyvin and Sinuraya (2017), cannabinoids are classified by source and origin. Phytocannabinoids are derivative compounds from cannabis and other plants, including delta-9-tetrahydrocannabinol (THC), cannabidiol (CBD), and cannabinol (CBN). The details of *Boleophthalmus boddarti* extract were elucidated in the Table 1.

The highest amount of bioactive compounds in *Periophthalmus chrysospilos* extract corresponded to the steroid compounds; as many as 10 types of compounds were detected with a total of 24 compounds. The total number of compounds detected are named Androst-7-ene-6,17-dione, 2,3,14-trihydroxy-, where the total amount of compounds is 10 compounds. According to Moss (1989), androstanone is anabolic steroid group that belongs to the basic carbon cycle, which is unsaturated and is an alkyl substitute in the 17th carbon chain.

GC-MS Results of the *Boleophthalmus boddarti* extract

The result of GC-MS analysis of *Boleophthalmus boddarti* extract was described as the chromatogram form for all chemical compounds that were read by the detector (Fig. 2). The vertical lines were found in this chromatogram




Figure 1. Chromatogram result of peak chemical compounds in *Periophthalmus chrysospilos*
(Source: LHP, 2018)




Figure 2. Chromatogram result of peak chemical compounds in *Boleophthalmus boddarti*
(Source: LHP, 2018)

Table 1. The amount of bioactive compounds in the *Periophthalmus chrysospilos* extract

No.	Compound names	Compound classes	Total numbers
1.	D-Limonene	Terpenoids	1
2.	Limonene	Terpenoids	1
3.	Aspidospermidin-17-ol, 1-acetyl-19,21-epoxy-15, 16-dimethoxy-	Alkaloids (Aspidospermidin)	1
4.	Choltesan-3-ol, 2-methylene-	Steroids	6
5.	Azulene, 1,4-dimethyl-7-(1-methylethyl)-	Terpenoids	1
6.	17.alfa.,21a-28, 30-Bisnorhopane	Steroids	1
7.	4-Androsten-6a-ol-3,17-dione	Steroids	3
8.	3,9-Epoxy pregn-16-ene-14,20-diol, 7,11,18-triacetoxy-3-methoxy-	Steroids	1
9.	Cholestan-26-oic acid, 3,7,12-trihydroxy-	Steroids	1
10.	Rhodopin	Carotenoids	1
11.	Cannabinol, trifluoroacetate	Cannabinoids	4
12.	Cholestan-3,5-diol, 5-acetate	Steroids	2
13.	Milbemycin B, 5-demethoxy-5-one-6,28-anhydro-25-ethyl-4-methyl-13-chloro-oxime	Steroids	1
14.	9,19-Cyclolanostan-3-ol, 24,24-epoxymethano-acetate	Terpenoids	1
15.	8,14-Seco-3,19-epoxyandrostane-8,14-dione, 17-acetoxy-3 α -methoxy-4,4-dimethyl-	Steroids	11
16.	Androst-7-ene-6,17-dione, 2,3,14-trihydroxy-	Steroids	10
17.	2-Phenanthrenecarboxylic acid, 1-(1,3-dithian-2-ylmethyl)-7-hydroxy-2,4b-dimethyl-1,2,3,4,4a,4b,5,6,7,8,10,10a-dodecahydro-,methyl ester	Steroids	1

pattern. Those indicate more than one bioactive compound in the *Boleophthalmus boddarti* extract. The presence of five major peaks of bioactive compounds was detected and the most compound are from steroids. The highest peak in chromatogram pattern could be seen that the *Boleophthalmus boddarti* extract has retention time of 14.44. On the basis of the GC-MS signal, the sample has many bioactive compounds, which was shown by some signals detected. It shows that the *Boleophthalmus boddarti* extract can be successfully used for treating people with inflammatory diseases, especially enuresis, as proven by the GC-MS analysis.

Bioactive compounds in the *Boleophthalmus boddarti* extract

The results of *Boleophthalmus boddarti* extract revealed the amounts of bioactive compounds such as steroids, carotenoids, terpenoids, bufadienolide, carotatoxin, cholesterol, and corticosteroid shown in Table 2. The compounds of steroid groups were detected as many as 24 types of compounds with a total of 44 compounds. Specific steroid compounds, namely corticosteroids, were detected with 2 types of compounds and a total of 3 compounds. The details of *Boleophthalmus boddarti* extract were elucidated in Table 2.

Corticosteroids are drugs containing steroid hormones that are useful in increasing the needed steroid hormones in the body, relieving inflammation and helping in alleviate asthma, allergy and multiple sclerosis (Cidlowski, 2013; Panda and Mabalirajan, 2018; Rozaliyani, 2011). According to the analysis of Prasad *et al.* (2018), corticosteroids also reduce the mortality risk from tuberculosis effect at least in the short term. A quite unique compound, called Gamabufotalin, was detected in the *Boleophthalmus boddarti* extract; it is a bioactive compound of the bufadienolide type. According to Dmitrieva *et al.* (2000), Bufadienolides are biosynthetic steroids that are independent of the side chain of cholesterol division. Bufadienolides (BDs) also include the corticic steroid groups,

which are usually obtained from amphibian poison extracts, these compounds are often used as traditional medicine in Asia.

GC-MS Results of *Boleophthalmus dussumieri* extract

The result of the GC-MS analysis of *Boleophthalmus dussumieri* extract was described as the chromatogram form for all chemical compounds that were read by the detector (Fig. 3). The vertical lines were found in this chromatogram pattern. Those indicate more than one bioactive compound in the *Boleophthalmus dussumieri* extract. The presence of four major peaks of bioactive compounds was detected by GC-MS like steroids, terpenoids,

Table 2. The amount of bioactive compounds in the *Boleophthalmus boddarti* extract

No.	Compound names	Compound classes	Total numbers
1.	<i>Curan-17-oic acid, 19,20-dihydroxy-, methyl ester</i>	Steroids	2
2.	<i>Cholestan-3-ol, 2-methylene-</i>	Steroids	12
3.	<i>Citronellol epoxide</i>	Terpenoids	1
4.	<i>Falcarinol</i>	Carotatoxin (secondary cancer anti metabolites)	1
5.	<i>Cholestan-3-one, cyclic 1,2-ethanediyl aetal</i>	Steroids	1
6.	<i>..psi.,psi.-Carotene, 1,1',2,2'-tetrahydro-1,1'-dimethoxy</i>	Carotenoids	8
7.	<i>Androsten-4-en-11-ol-3,17-dione, 9-thiocyanato-</i>	Steroids	2
8.	<i>Androsten-5-en-17-one-3,11-bis[(trimethylsilyl)oxy]-, O-(phenylmethyl)oxime</i>	Steroids	1
9.	<i>Prosta-5,13-dien-1-oic acid, 9,11,15-tris[(trimethylsilyl)oxy]-, trimethylsilyl ester</i>	Steroids	6
10.	<i>Rhodoxanthin</i>	Carotenoids	2
11.	<i>Rhodopin</i>	Carotenoids	2
12.	<i>5Á-Ergost-24-en-26-oic acid, 5,6Á-epoxy-4Á, 18,22-trihydroxy-3-methoxy-1-ox o-, é-lactone, diacetate,</i>	Steroids	1
13.	<i>Ethyl iso-allocholate</i>	Steroids	1
14.	<i>Gamabufotalin</i>	Bufadienolide (steroid aglycones in frogs)	1
15.	<i>Betamethasone acetate</i>	Corticosteroid	2
16.	<i>Cholestan-26-oic acid, 3,7,12-trihydroxy-</i>	Steroids	2
17.	<i>..psi.,psi.-Carotene, 3,4-didehydro-1,2,7',8'-tetrahydro-1-methoxy-2-ox o</i>	Carotenoids	1
18.	<i>..psi.,psi.-Carotene, 3,3',4,4'-tetrahydro-1',2'-dihydro-1-hydroxy-1'methoxy</i>	Carotenoids	1
19.	<i>Prosta-5,13-dien-1-oic acid, 9,11,15-tris[(trimethylsilyl)oxy]-, trimethylsilyl ester</i>	Steroids	2
20.	<i>17Á-Acetoxy-1',1'-dicarboethoxy-1Á,2Á-dihydrocy cloprop[1,2]-5Á-androst-1-en-3-one</i>	Steroids	2
21.	<i>Cholesterol</i>	Cholesterol	1
22.	<i>26-Nor-5-cholesten-3Á-ol-25-one</i>	Steroids	1
23.	<i>17-(1,5-Dimethylhexyl)-10,13-dimethyl-2,3,4,7,8,9,10,11,12,13,14,15,16,17-tetradecahydro-1Hcyclopenta[a]phenanthren-3-ol</i>	Steroids	1
24.	<i>Androst-4-en-9-thiocyanomethyl-11-ol-3,17-dione</i>	Steroids	1
25.	<i>Androst-5-en-3-one, 19-acetoxy-4,4-dimethyl-,oxime</i>	Steroids	1
26.	<i>3-Isopropyl-6a,7,10b-trimethyl-8-(2-oxo-2-phenylethyl)dodecahydrobenzo[f]chromene-7-carboxylic acid, methyl ester</i>	Corticosteroid	1
27.	<i>Androst-8-en-3-ol, 4,4,14Á-trimethyl-17-(2-bromo-1-methylethyl)</i>	Steroids	1
28.	<i>1',1'-Dicarboethoxy-1Á,2Á-dihydro-3'H-cycloprop[1,2]cholest-1,4,6-trien-3-one</i>	Steroids	1
29.	<i>Prosta-5,13-dien-1-oic acid, 9,11,15,19-tetrakis[(trimethylsilyl)oxy]-, methyl ester</i>	Steroids	1




Figure 3. Chromatogram result of peak chemical compounds in *Boleophthalmus dussumieri*
(Source: LHP, 2018)

carotenoids and carotatoxins. The highest peak in the chromatogram pattern could be seen that *Boleophthalmus dussumieri* extract has retention time of 22.61. On the basis of the GC-MS signal, the sample has significant bioactive compounds, which was shown in Table 3.

Bioactive compounds in the *Boleophthalmus dussumieri* extract

The bioactive compounds detected from the *Boleophthalmus dussumieri* extract include steroids, terpenoids, carotenoids and carotatoxins. The compounds of steroid classes include as many as 11 types of compounds, with a total of 26 compounds. The group of carotenoid compounds was detected in as many as 6 types of compounds, with a total of 13 compounds. The group of terpenoid compounds was detected by 2 types of compounds, with a total of 2 compounds. The carotatoxin compounds were also detected in the *Boleophthalmus dussumieri* extract, with a total of 1 compound. The details of *Boleophthalmus dussumieri* extract were elucidated in Table 3.

The name of steroid compounds with the highest total number of compounds is Spirost-8-en-11-one, 3-hydroxy. According to Rajendran et al. (2017), Spirost-8-en-11-one, 3-hydroxy included the steroid compounds which have an anti-cancer bioactivity. Spirost-8-en-11-one, 3-hydroxy has the chemical formula C₂₇H₄₀O₄. The group of carotenoid compounds detected with the highest amount is Rhodopin, where the total amount of these compounds is 5 compounds. According to Komori et al. (1998), Rhodopin is a carotenoid compound which is usually produced by *Rhodospirillum rubrum* bacteria with the chemical formula C₄₀H₅₈O.

Bioactive carotatoxin alkaloids were also detected in the *Boleophthalmus dussumieri* extract. The name of the carotatoxin group is falcarinol and the total of compound is only 1 compound. According to Zaini et al. (2012), falcarinol (FaOH) is one of the compounds found in carrots; this compound has shown bioactive action in several types of cell cultures. Falcarinol has been shown to be cytotoxic to the culture of acute lymphoblastic leukemia cells.

GC-MS Results of the *Periophthalmodon schlosseri* extract

The result of GC-MS analysis of *Periophthalmodon schlosseri* extract was described as the chromatogram form for all chemical compounds that was read by the detector (Fig. 4). The vertical lines were found in this chromatogram pattern. Those indicate more than one bioactive compound in *Boleophthalmus boddarti* extract. The presence of five major peaks of bioactive compounds was detected and the most compound are from steroids. The highest peak in the chromatogram pattern indicated that the *Boleophthalmus boddarti* extract has retention time of 14.44. On the basis of the GC-MS signal, the sample has many bioactive compounds which was shown by some signals detected. It shows that the GC-MS analysis proves that the *Boleophthalmus boddarti* extract can be successfully used for treating people with inflammatory diseases especially enurosis.

Bioactive compounds in the *Periophthalmodon Schlosseri* extract

The bioactive compounds detected from *Periophthalmodon schlosseri* extract including steroids, carotenoids and terpenoids. As many as

Table 3. The amount of bioactive compounds in the *Boleophthalmus boddarti* extract

No.	Compound names	Compound classes	Total numbers
1.	<i>Limonene</i>	Terpenoids	1
2.	<i>D-Limonene</i>	Terpenoids	1
3.	<i>Corynan-17-ol, 18,19-didehydro-10-methoxy-,acetate (ester)</i>	Steroids	2
4.	<i>Rhodopin</i>	Carotenoids	5
5.	<i>Ç-Sitosterol</i>	Steroids	1
6.	<i>Spirost-8-en-11-one, 3-hydroxy</i>	Steroids	13
7.	<i>Cholest-5-en-3-ol, 24-propylidene-</i>	Steroids	2
8.	<i>Stigmasta-5,24(28)-dien-3-ol</i>	Steroids	1
9.	<i>..psi.,.psi.-Carotene, 1,1',2,2'-tetrahydro-1,1'-dimethoxy</i>	Carotenoids	4
10.	<i>5á-Ergost-24-en-26-oic acid, 5,6á-epoxy-4á, 18,22-trihydroxy-3-methoxy-1-ox o-, é-lactone, diacetate,</i>	Steroids	1
11.	<i>..psi.,.psi.-Carotene, 3,4-didehydro-1,2,7',8'-tetrahydro-1-methoxy-2-ox o</i>	Carotenoids	1
12.	<i>5-Chloro-6beta-nitro-5alpha-cholestane-3-one</i>	Steroids	1
13.	<i>Prosta-5,13-dien-1-oic acid, 9,11,15-tris[(trimethylsilyl)oxy]-, trimethylsilyl ester</i>	Steroids	1
14.	<i>Cholest-4-ene, 3á-(methoxymethoxy)-</i>	Steroids	1
15.	<i>Rhodoxanthin</i>	Carotenoids	1
16.	<i>4'-Apo-á,.psi.-carotenoic acid</i>	Carotenoids	1
17.	<i>Astaxanthin</i>	Carotenoids	1
18.	<i>Androst-7-ene-6,17-dione, 2,3,14-trihydroxy</i>	Steroids	2
19.	<i>Prosta-5-en-1-oic acid, 9,11,15-tris[(trimethylsilyl)oxy]-, methylsilyl ester</i>	Steroids	1
20.	<i>Falcarinol</i>	Carotatoxin	1

11 bioactive steroid compounds were detected, with a total of 21 compounds; 3 types of the carotenoid group compounds, with a total of 6 compounds; while only 1 type of terpenoid group compounds were detected with a total of 1 compound. The details of the *Periophthalmodon schlosseri* extract were elucidated in Table 4.

The steroid compounds in the *Periophthalmodon schlosseri* extract are named Cholestan-3-ol, 2-methylene-, Cholest-5-en-3-ol (3á)-, tetradecanoate, Cholesta-3,5-diene, Cholesteryl

benzoate, 17.alfa., 21á-28, 30-Bisnorhopane, Cholesterol, 26-Nor-5-cholest-3á-ol-25-one, Cholestane-3,5-diol, 5-acetate, Campesterol, Ergost-5-en-3-ol, and Androst- 7-ene-6,17-dione, 2,3,14-trihydroxy-. According to Salempa and Muhamram (2016), steroids are natural compounds which consist of a carbon skeleton and comprise three circles of six perdidro phenanthrene fused into one circle of five. Saturated-click hydrocarbons have a circumference system consisting of 17 carbon atoms.




Figure 4. Chromatogram result of peak chemical compounds in *Periophthalmodon schlosseri*
(Source: LHP, 2018)

Table 4. The amount of bioactive compounds in the *Periophthalmodon schlosseri* extract

No.	Compound names	Compound classes	Total numbers
1.	D-Limonene	Terpenoids	1
2.	Cholestan-4-ol, 2-methylene-	Steroids	4
3.	Cholest-5-en-3-ol (3 α)-, tetradecanoate	Steroids	1
4.	Cholesta-3,5-diene	Steroids	1
5.	Cholesteryl benzoate	Steroids	1
6.	Rhodopin	Carotenoids	3
7.	Azafrin	Carotenoids	1
8.	17.alfa.,21 α -28,30-Bisnorhopane	Steroids	2
9.	Cholesterol	Steroids	2
10.	26-Nor-5-cholest-3-ol-25-one	Steroids	1
11.	Cholestane-3,5-diol, 5-acetate	Steroids	1
12.	Campesterol	Steroids	1
13.	Ergost-5-en-3--ol	Steroids	1
14.	Astaxanthin	Carotenoids	2
15.	Androst-7-ene-6,17-dione, 2,3,14-trihydroxy-	Steroids	6

D-Limonene is the only group of terpenoid compounds which was detected in the *Periophthalmodon schlosseri* extract. This compound was also detected in samples of *Periophthalmus chrysospilos* extract and *Boleophthalmus dussumieri* extract. According to Sun (2007), D-Limonene is a monocyclic terpenoid with a lemon-like odor and is a major constituent of several essential oils of Citrus sp. (orange, lemon, mandarin, lime and grapefruit). D-Limonene is widely used as an additive (additive) in perfumes, soap, food, gum and drinks. This compound is also listed in the Code of Federal Regulation as a material that is known to be safe or generally recognized as safe (GRAS) as a flavoring agent.

The group of carotenoid compounds detected includes rhodopin, azafrin and astaxanthin. The rhodopin compounds were detected in all samples with a sufficient number of total compounds in each sample. The astaxanthin compounds were also detected in the flesh extracts of *Boleophthalmus boddarti* and *Boleophthalmus dussumieri*, with a total of 1 compound each in each sample. In turn, the azafrin compounds were only detected in *Periophthalmodon schlosseri* extract. According to Gopalakrishnan and Kalaiarasi (2012), azafrin is a compound of the carotenoid ester group with the molecular formula C28H40O4. The azafrin compounds include the bioactive compounds that have the potential as antioxidants, anti-inflammatory, anticancer, anti-criticism, natural dyes, eye pain medications, and as compounds that can reduce hypertension.

Physical and chemical properties of the environment at sampling locations

The main point of mudskippers sampling based on the GPS is in the coordinates S 02° 21'57.5 "E 104° 54'00.1". The environment measurements of the physical and chemical properties were carried out at the main point of sampling and obtained as presented in Table 5.

The measurement of the physical and chemical properties of the environment was carried out at 3 different times. The first measurements were made in late March, precisely in the afternoon, and obtained soil moisture data of 100%, soil pH of 6.5, water pH of 6.82, and water salinity of 0.1‰. The second measurement was carried out in an April morning and 100% of soil moisture was obtained, soil pH was 6.5, water pH was 7.32, and water salinity was 0‰. The third measurement was carried out in the afternoon of April and obtained 100% soil moisture data, a soil pH of 6.5, a water pH of 6.7, and water salinity of 0‰.

Table 5. Physical and chemical properties of the environment at the sampling location

Parameter	Measurement Time		
	March	April Morning	April Afternoon
Soil moisture (%)	100	100	100
Soil pH	6.5	6.5	6.5
Water pH	6.82	7.32	6.7
Waters salinity (%)	0.1	0	0

On the basis of the data of the physical and chemical properties of the environment, it can be seen that the soil moisture at the main sampling point shows 100% at 3 times measurements with different times. This shows that the sampling location has a substrate in the form of mud which is always wet. According to Kurnia et al. (2006), the soil moisture is one of the key variables in the hydrological process that plays an important role in determining the availability of water as a very fundamental element in the life of living things. The 100% soil moisture indicates that the soil is saturated with water.

The water salinities at 3 measurements with different times showed the numbers 0.1‰, 0‰ and 0‰. This shows that the waters around the sampling location tend to be tasteless. According to Masuda et al. (1984) and Redjeki (2013), the gudder is an amphidromous demersal fish that tolerates freshwater and sea, lives on muddy and sandy bottom and spawns in estuarine areas.

CONCLUSIONS

The bioactive compound analysis of the *Periophthalmus chrysospilos* extract showed that it consists of steroids, terpenoids, carotenoids, cannabinoids, and alkaloids. As many as 10 types of hormones were found in the *Periophthalmus chrysospilos* extract, with a total of 24 compounds. The *Boleophthalmus boddarti* extract comprised steroids, carotenoids, terpenoids, bufadienoloid, and carotatoksin. There were 24 types of compounds detected with a total of 44 compounds in the *Boleophthalmus boddarti* extract. The *Boleophthalmus dussumieri* extract has bioactive compounds from the groups of steroids, carotenoids, terpenoids, and carotatoksin. The *Boleophthalmus dussumieri* extract has as many as 11 types of compounds with a total of 26 compounds. The *Periophthalmodon schlosseri* extract contained the bioactive compounds including steroids, carotenoids and terpenoids. As many as 11 types of compounds were detected with a total of 21 compounds. The higher concentrations of bioactive compounds were found in the mudskippers that are possibly consumed for treating inflammatory diseases, enuresis in children and allergy.

Acknowledgments

The authors would like to thank Dr. Arum Setiawan and Sulistiono, Dr. Sarno, Dr. Arwinskyah,

Enggar Patriono M.Si., College of Agricultural Technology Production Laboratory, Sriwijaya University for his help in specimen preparation. We also thank anonymous referees for constructive criticism on the manuscript. This work was supported by the projects of Kemenristekdikti.

REFERENCES

1. Bob-Manuel, F.G., 2011. Food and feeding ecology of the mudskipper *Periophthalmus koelreuteri* (Pallas) Gobiidae at Rumuolumeni Creek Niger Delta, Nigeria. Agriculture and Biology Journal of North America. 2, 897–901.
2. Clayton DA. 1993. Mudskippers. Oceanogr Mar Biol Annu. Rev 31:507–77.
3. Dabruzzii, T. F., Fangue, N. A., Kadir, N. N, and Bennett, W.A., 2019. Thermal Niche Adaptations of Common Mudskipper (*Periophthalmus kalolo*) and Barred Mudskipper (*Periophthalmus argentilineatus*) in Air and Water. Journal of Thermal Biology. 81, 170–177.
4. DKP Provinsi Sumatera Selatan, 2017. Sumberdaya: Pengelolaan Pesisir Provinsi Sumatera Selatan 1. [online] Tersedia: dislutan.sumselprov.go.id/page/38/sumberdaya#. [11 November 2017].
5. Dmitrieva, R. I. Bagrov, A. Y., Lalli, E., Sassone-Corsi, P., Stocco, D., dan Doris, P. A., 2000. Mammalian Bufadienolide is Synthesized from Cholesterol in the Adrenal Cortex by Pathway that is Independent of Cholesterol Side-chain Cleavage Hypertension. 36, 3, 442–8.
6. Feulner, G. R., and Binish R, 2013. Re-discovery of the mudskipper *Periophthalmus waltoni* Koumans, 1941 in the United Arab Emirates. Tribulus, vol. 21, 2013, p. 42+. Gale Academic Onefile, Accessed 13 Oct. 2019
7. Gadnavi, M.K., Kukadia, D., Dar, S., Gokulakannan, N., Talukdar, G., Sivakumar, K., and Gopi, G. V., 2017. Indigenous techniques of catching mudskipper in Bhavnagar and Bharuch districts, Gujarat. Indian Journal of Traditional Knowledge. 16(3), pp. 533–538.
8. Gopalakrishnan, S., and Kalaiarasi, T., 2012. Identification of Chemical Compound from the Fruits of *Cucumis trigonus* Roxb. By GC-MS Analysis. International Journal of Phytopharmacy. 2 (5) 122–128.
9. Graham, J. B., 1997. Air-breathing fishes: evolution, diversity, and adaptation. San Diego: Academic Press. p. 299.
10. Graham, J. B., and Wegner, N. C., 2010. Breathing air in water and in air: the air-breathing fishes. In: Nilsson GE, editor. Respiratory physiology of vertebrates: life with and without oxygen. Cambridge:

- Cambridge University Press. p. 174–221.
11. Hameed, I. H., Jebor, M. A., and Ommer, A. J., 2016. Haplotype data of mitochondrial DNA coding region encompassing nucleotide positions 11,719–12,184 and evaluate the importance of these positions for forensic genetic purposes in Iraq. *Mitochondrial DNA Part A*. 27(2), 1324–1327.
 12. Huang, M. C., and Lee C. L., 2019. Thermostability of tropomyosins from the fast skeletal muscles of tropical fish species. *Fish Physiology and Biochemistry*. 45, 1189–1202.
 13. Hu, W., Zhang, J., Kang, B., 2016. Structure and function of corneal surface of mudskipper fishes. *Fish Physiology and Biochemistry*. 42, 1481–1489.
 14. Ikram, M. M., Ismail, A., Yap, C. K., and Azwady, A. A. N., 2013. Levels of heavy metals (Zn, Cu, Cd, and Pb) in mudskippers (*Periophthalmodon schlosseri*) and sediments collected from intertidal areas at Morib and Remis, Peninsular Malaysia. *Toxicology & Environmental Chemistry*. 92(8), 1471–1486.
 15. Ishimatsu, A., and Graham, J. B., TT. 2011. Mudskippers: front runners in the modern invasion of land. In: Patzner R, Van Tassell JL, Kovac M, Kapoor BG, editors. *The biology of gobies*. Enfield: Science Publishers. p. 603–32.
 16. Kareem, M. A., Abdulzahra, A. I., Hameed, I. H., and Jebor, M. A., 2015. A New Polymorphic Positions Discovered in Mitochondrial DNA Hypervariable Region HVIII From Central and North-Central of Iraq. *Mitochondrial DNA Part A*. 27(5), 3250–3254.
 17. Komori, M. R., Ghosh, S., Takachi, S., Hu, Y., Mizoguchi, T., Koyama, Y., and Kuki, 1998. A Null Lesion in the Rhodopin 3,4-desaturase of *Rhodospirillum rubrum* Unmask a Cryptic Branch of the Carotenoid Biosynthetic Pathway. *Biochemistry*. 37: 8987–8994.
 18. Kurnia, U., Agus, F., Adimihardja, A., dan Dariah, A. 2006. *Sifat Fisik Tanah dan Metode Analisisnya*. BBSLDP: Banjarmasin.
 19. Lauriano, E. R., Faggio, C., Capillo, G., Spanò, N., Kuciel, M., Aragona, M., Pergolizzi, S., 2018. Immunohistochemical characterization of epidermal dendritic-like cells in giant mudskipper, *Periophthalmodon schlosseri*. *Fish & Shellfish Immunology*. 74, 380–385
 20. Looi, L. J., Aris, A. Z., Haris, H., Yusoff, F. Md., and Hasim, Z., 2016. The levels of mercury, methylmercury and selenium and the selenium health benefit value in grey-eel catfish (*Plotosus canius*) and giant mudskipper (*Periophthalmodon schlosseri*) from the Strait of Malacca. *Journal of Chemosphere*. 152, 265–273
 21. Maggyvin, E. dan Sinuraya, R. K. 2017. Marijuana dan Autisme: Sebuah Literatur Review. *Jurnal Farmaka*. 15 (1): 70–92.
 22. Mahadevan, G., Mohan, K., Vinoth, J., and Ravi, V., 2019. Biotic potential of mucus extracts of giant mudskipper *Periophthalmodon schlosseri* (Pallas, 1770) from Pichavaram, southeast coast of India. *The journal of basic and applied zoology*. 80(13).
 23. Martin, K. L., 2013. Theme and variations: amphibious air-breathing intertidal fishes
 24. Martínez, H. L., Agorreta, A., Sánchez, M. T., and Mauro, D. S., 2018. Evidence of positive selection suggests possible role of aquaporins in the water to land transition of mudskippers. *Organisms Diversity and Evolution*. 18(4), 499–414.
 25. Moss, S.M. 1989. Effect of Exogenous Androgens on Growth, Biochemical Composition and Reproduction of the Coot Calm *Mulinia literalis*. Pac SCI.
 26. Mohammed, A., and Imad, H., 2013. Autosomal STR: From locus information to next generation sequencing technology. *Res. J. Biotechnol.* 8(10):92–105.
 27. Murdy, E.O., 1989. A taxonomic revision and cladistic analysis of the Oxudercine gobies (Gobiidae: Oxudercinae). *Rec Aust Mus Suppl* 11: 1–93.
 28. Panda, L., and Mabalirajan, U., 2018. Recent updates on corticosteroid resistance in asthma. *European Medical Journal*. 3(3), 49–57.
 29. Polgar, G., Ghanbarifardi, M., Milli, S., Agorreta, A., Aliabadi, M., Esmaili, H. R., and Khang, T. F., 2017. Ecomorphological adaptation in three mudskippers (Teleostei: Gobioidei: Gobiidae) from the Persian Gulf and the Gulf of Oman. *Hydrobiologia*. 795(1), 91–111.
 30. Prasad, K., Singh, M. B., and Ryan, H., 2016. Corticosteroids for managing tuberculous meningitis (Review). *European Medical Journal*. 3(3), 49–57.
 31. Rajendran, P., Bharatidhasan, R., dan Sureshkumar, K. 2017. GC-MS Analysis of Phyto-Components in Raw and Treated Sugarcane Juice. *Int.J.Curr.Microbiol.App.Sci.* 6 (7): 51–61.
 32. Rozaliyani, A., Susanto, A. D., Swidarmoko, B., dan Yunus, F. 2011. Mekanisme Resistens Kartikosteroid pada Asma. *Jurnal Respir Indo*. 31 (4): 210–223.
 33. Ridho, M. R., Patrono, E., Sholikah, M., 2019. Food habits of three species of mudskippers in the Musi River Estuary, South Sumatra, Indonesia. *Biodiversitas*. 20(8), 2368–2374.
 34. Salempa, P. dan Muhamram, H. 2016. *Senyawa Steroid dalam Tumbuhan Bayur*. Badan penerbit UNM: Makasar.
 35. Sari, M., 2012. *Jenis Fauna Hutan Mangrove di Areal PT. Bina Ovipipari Semesta dan Sekitaranya*. Penerbit PT. Bina Ovipipari Semesta: Pontianak.
 36. Sun, J. 2007. D-Limonene: Safety and Clinical Application. *Alternative Medicine Review*. 12 (3): 259–264
 37. Takeda, T., Hayashi, M., Toba, A., Soyano, K., and Ishimatsu, A., 2012. *Ecology of the Australian*

- mudskipper *Periophthalmus minutus*, an amphibious fish inhabiting a mudflat in the highest intertidal zone. *Australian Journal of Zoology*. 57: 312–320.
38. Wu, R. X., Hong, W. S., Zhang, Q. Y., Chen, S. X., 2009. Comparative enzyme activities of the intestinal brush border membranes of the herbivorous mudskipper *Boleophthalmus pectinirostris* and the carnivorous Chinese black sleeper *Bostrichthys siensis*. *Journal of Applied Ichthyology*. 25, 571–575.
39. Yang, X. X., Gu, W., Liang, L., Yan, H. L., Wang, Y. F., Bi, Q., Zhang, T., Yu, J., and Rao, G. X., 2017. Screening for the bioactive constituents of traditional Chinese medicines-progress and challenges. Royal society of chemistry. 7, 3089–3100.
40. You, X. X., Sun, M., Li, J., Bian, C., Chen, J., Yi, Y., Yu, H., and Shi, Q., 2018. Mudskippers and Their Genetic Adaptations to an Amphibious Lifestyle.
41. Zaini, R. G., Brandt, K., Clench, M. R., dan Maitre, C. L. L. 2012. Effect of Bioactive Compounds from Carrots (*Daucus carota L.*), Polyacetylenes, Beta-carotene and Lutein on Human Lymphoid Leukemia Cells. *Medical Chemistry Journal*. 12 (0): 2–13.



(<http://www.jeeng.net/>)

All issues

< (<http://www.jeeng.net/Issue-2-2020,7955>)

> (<http://www.jeeng.net/Issue-4-2020,8477>)

Volume 21, Issue 3, 2020

Analysis of Partially Evacuated Solar Still System under Jordan Climate

(<http://www.jeeng.net/Analysis-of-Partially-Evacuated-Solar-Still-System-under-Jordan-Climate,116324,0,2.html>)

Nader Aljabarin (<http://www.jeeng.net/Author-Nader-Aljabarin/66044>)

J. Ecol. Eng. 2020; 21(3):1-9

DOI: <https://doi.org/10.12911/22998993/116324> (<https://doi.org/10.12911/22998993/116324>)

Abstract

(<http://www.jeeng.net/Analysis-of-Partially-Evacuated-Solar-Still-System-under-Jordan-Climate,116324,0,2.html>)

Article (PDF) (<http://www.jeeng.net/pdf-116324-48114?filename=Analysis%20of%20Partially.pdf>)

Stats

Macrozoobentos Distribution as a Bioindicator of Water Quality in the Upstream of the Citarum River

(<http://www.jeeng.net/Macrozoobentos-Distribution-as-a-Bioindicator-of-Water-Quality-in-the-Upstream-of,116335,0,2.html>)

Nurul Chazanah (<http://www.jeeng.net/Author-Nurul-Chazanah/133289>), Barti Setiani Muntalif (<http://www.jeeng.net/Author-Barti-Muntalif/133290>), Priana Sudjono (<http://www.jeeng.net/Author-Priana-Sudjono/133291>)

J. Ecol. Eng. 2020; 21(3):10-17

DOI: <https://doi.org/10.12911/22998993/116335> (<https://doi.org/10.12911/22998993/116335>)
 EN (<http://www.jeeng.net/>) PL (<http://www.jeeng.net/pl>)



- ❑ Abstract
(<http://www.jeeng.net/Macrozoobentos-Distribution-as-a-Bioindicator-of-Water-Quality-in-the-Upstream-of,116335,0,2.html>)
- ❑ Article (PDF) (<http://www.jeeng.net/pdf-116335-48033?filename=Macrozoobentos.pdf>)

Results of Surface Water Quality Monitoring of the Western Bug River Basin in Lviv Region

(<http://www.jeeng.net/Results-of-Surface-Water-Quality-Monitoring-of-the-Western-Bug-River-Basin-in-Lviv,118303,0,2.html>)

Zoriana Odnorih (<http://www.jeeng.net/Author-Zoriana-Odnorih/136211>), Roman Manko (<http://www.jeeng.net/Author-Roman-Manko/136212>), Myroslav Malovanyy (<http://www.jeeng.net/Author-Myroslav-Malovanyy/94769>), Khrystyna Soloviy (<http://www.jeeng.net/Author-Khrystyna-Soloviy/136213>)

J. Ecol. Eng. 2020; 21(3):18–26

DOI: <https://doi.org/10.12911/22998993/118303> (<https://doi.org/10.12911/22998993/118303>)

- ❑ Abstract
(<http://www.jeeng.net/Results-of-Surface-Water-Quality-Monitoring-of-the-Western-Bug-River-Basin-in-Lviv,118303,0,2.html>)
- ❑ Article (PDF) (<http://www.jeeng.net/pdf-118303-48026?filename=Results%20of%20Surface%20Water.pdf>)

The Influence of Anaerobic Digestion on Selected Heavy Metals Fractionation in Sewage Sludge

(<http://www.jeeng.net/The-Influence-of-Anaerobic-Digestion-on-Selected-Heavy-Metals-Fractionation-in-Sewage,118302,0,2.html>)

Magdalena Zdeb (<http://www.jeeng.net/Author-Magdalena-Zdeb/126869>), Małgorzata Pawłowska (<http://www.jeeng.net/Author-Małgorzata-Pawłowska/10369>), Justyna Pacan (<http://www.jeeng.net/Author-Justyna-Pacan/136210>)

J. Ecol. Eng. 2020; 21(3):27–35

DOI: <https://doi.org/10.12911/22998993/118302> (<https://doi.org/10.12911/22998993/118302>)

- ❑ Abstract
(<http://www.jeeng.net/The-Influence-of-Anaerobic-Digestion-on-Selected-Heavy-Metals-Fractionation-in-Sewage,118302,0,2.html>)

in-

[Sewage,118302,0,2.html](http://www.jeeng.net/Sewage,118302,0,2.html)[EN \(http://www.jeeng.net/\)](http://www.jeeng.net/)[PL \(http://www.jeeng.net/pl/\)](http://www.jeeng.net/pl/)[Article \(PDF\) \(http://www.jeeng.net/pdf-118302-48030?filename=The Influence of.pdf\)](http://www.jeeng.net/pdf-118302-48030?filename=The%20Influence%20of.pdf)

Sonification Energy in the Process of Ultrasonic Disintegration

[\(http://www.jeeng.net/Sonification-Energy-in-the-Process-of-Ultrasonic-Disintegration,118301,0,2.html\)](http://www.jeeng.net/Sonification-Energy-in-the-Process-of-Ultrasonic-Disintegration,118301,0,2.html)Paweł Wolski (<http://www.jeeng.net/Author-Pawel-Wolski/136209>)

J. Ecol. Eng. 2020; 21(3):36–40

DOI: <https://doi.org/10.12911/22998993/118301> (<https://doi.org/10.12911/22998993/118301>)[Abstract](#)

Stats

[\(http://www.jeeng.net/Sonification-Energy-in-the-Process-of-Ultrasonic-Disintegration,118301,0,2.html\)](http://www.jeeng.net/Sonification-Energy-in-the-Process-of-Ultrasonic-Disintegration,118301,0,2.html)[Article \(PDF\) \(http://www.jeeng.net/pdf-118301-48010?filename=Sonification Energy in.pdf\)](http://www.jeeng.net/pdf-118301-48010?filename=Sonification%20Energy%20in.pdf)

Changes in the Quality of Domestic Sewage Caused by the Reduction of Water Consumption in Households

[\(http://www.jeeng.net/Changes-in-the-Quality-of-Domestic-Sewage-Caused-by-the-Reduction-of-Water-Consumption,118300,0,2.html\)](http://www.jeeng.net/Changes-in-the-Quality-of-Domestic-Sewage-Caused-by-the-Reduction-of-Water-Consumption,118300,0,2.html)Jarosław Chudzicki (<http://www.jeeng.net/Author-Jarosław-Chudzicki/136207>), Katarzyna Umiejewska (<http://www.jeeng.net/Author-Katarzyna-Umiejewska/136208>)

J. Ecol. Eng. 2020; 21(3):41–50

DOI: <https://doi.org/10.12911/22998993/118300> (<https://doi.org/10.12911/22998993/118300>)[Abstract](#)

Stats

[\(http://www.jeeng.net/Changes-in-the-Quality-of-Domestic-Sewage-Caused-by-the-Reduction-of-Water-Consumption,118300,0,2.html\)](http://www.jeeng.net/Changes-in-the-Quality-of-Domestic-Sewage-Caused-by-the-Reduction-of-Water-Consumption,118300,0,2.html)[Article \(PDF\) \(http://www.jeeng.net/pdf-118300-48025?filename=Changes in the Quality of.pdf\)](http://www.jeeng.net/pdf-118300-48025?filename=Changes%20in%20the%20Quality%20of.pdf)

Features of Heavy Metals Accumulation in Bottom Sediments of the Southern Bug Hydroecosystem

[\(http://www.jeeng.net/Features-of-Heavy-Metals-Accumulation-in-Bottom-Sediments-of-the-Southern-Bug-Hydroecosystem,118299,0,2.html\)](http://www.jeeng.net/Features-of-Heavy-Metals-Accumulation-in-Bottom-Sediments-of-the-Southern-Bug-Hydroecosystem,118299,0,2.html)Olena Mitryasova (<http://www.jeeng.net/Author-Olena-Mitryasova/126567>), Piotr Koszelnik (<http://www.jeeng.net/Author-Piotr-Koszelnik/37570>), Renata Gruca-Rokosz (<http://www.jeeng.net/Author-Renata-Gruca-Rokosz/71631>), Victor Smirnov (<http://www.jeeng.net/Author-Victor-Smirnov/136203>), Svetlana Smirnova (<http://www.jeeng.net/Author-Svetlana-Smirnova/136204>), Yevgen Bezsonov (<http://www.jeeng.net/Author-Yevgen>)

Bezsonov/136205), Monika Zdeb (<http://www.jeeng.net/Author-Monika-Zdeb/37474>), Sabina Ziembowicz (<http://www.jeeng.net/Author-Sabina-Ziembowicz/136206>) EN (<http://www.jeeng.net/>) PL (<http://www.jeeng.net/pl>)



J. Ecol. Eng. 2020; 21(3):51–60

DOI: <https://doi.org/10.12911/22998993/118299> (<https://doi.org/10.12911/22998993/118299>)

✉ Abstract
(<http://www.jeeng.net/Features-of-Heavy-Metals-Accumulation-in-Bottom-Sediments-of-the-Southern-Bug-Hydroecosystem,118299,0,2.html>)

📄 Article (PDF) (<http://www.jeeng.net/pdf-118299-48039?filename=Features%20of%20Heavy%20Metals.pdf>)

Grazing of Native Livestock Breeds as a Method of Grassland Protection in Roztocze National Park, Eastern Poland

(<http://www.jeeng.net/Grazing-of-Native-Livestock-Breeds-as-a-Method-of-Grassland-Protection-in-Roztocze,118298,0,2.html>)

Mariusz Kulik (<http://www.jeeng.net/Author-Mariusz-Kulik/136199>), Marianna Warda (<http://www.jeeng.net/Author-Marianna-Warda/136200>), Adam Gawryluk (<http://www.jeeng.net/Author-Adam-Gawryluk/128291>), Andrzej Bochniak (<http://www.jeeng.net/Author-Andrzej-Bochniak/41523>), Krzysztof Patkowski (<http://www.jeeng.net/Author-Krzysztof-Patkowski/136201>), Antoni Lipiec (<http://www.jeeng.net/Author-Antoni-Lipiec/136202>), Tomasz M. Gruszecki (<http://www.jeeng.net/Author-Tomasz-Gruszecki/91345>), Michał Pluta (<http://www.jeeng.net/Author-Michał-Pluta/72666>), Elżbieta Bielińska (<http://www.jeeng.net/Author-Elżbieta-Bielinska/62516>), Barbara Futa (<http://www.jeeng.net/Author-Barbara-Futa/28104>)

J. Ecol. Eng. 2020; 21(3):61–69

DOI: <https://doi.org/10.12911/22998993/118298> (<https://doi.org/10.12911/22998993/118298>)

✉ Abstract
(<http://www.jeeng.net/Grazing-of-Native-Livestock-Breeds-as-a-Method-of-Grassland-Protection-in-Roztocze,118298,0,2.html>)

📄 Article (PDF) (<http://www.jeeng.net/pdf-118298-48115?filename=Grazing%20of%20Native.pdf>)

Bioactive Compounds Evaluation of the Mudskippers in the Estuarine Area of Musi River, South Sumatera, Indonesia

(<http://www.jeeng.net/Bioactive-Compounds-Evaluation-of-the-Mudskippers-in-the-Estuarine-Area-of-Musi-River,118296,0,2.html>)

Mohammad Rasyid Ridho (<http://www.jeeng.net/Author-Mohammad-Ridho/136195>), Arum Setiawan (<http://www.jeeng.net/Author-Arum-Setiawan/138108>), Sarno Sarno (<http://www.jeeng.net/Author-Sarno-Sarno/138112>), Arwinskyah Arwinskyah (<http://www.jeeng.net/Author-Arwinskyah-Arwinskyah/138113>), Enggar Patriono (<http://www.jeeng.net/Author-Enggar-Patriono/138114>), Sulistiono Sulistiono (<http://www.jeeng.net/Author-Sulistiono-Sulistiono/138115>)

J. Ecol. Eng. 2020; 21(3):70–80

- DOI:** <https://doi.org/10.12911/22998993/118296> (<https://doi.org/10.12911/22998993/118296>)
EN (<http://www.jeeng.net/>) PL (<http://www.jeeng.net/pl>)
- Abstract
(<http://www.jeeng.net/Bioactive-Compounds-Evaluation-of-the-Mudskippers-in-the-Estuarine-Area-of-Musi-River,118296,0,2.html>)
 Article (PDF) (<http://www.jeeng.net/pdf-118296-48116?filename=Bioactive%20Compounds.pdf>)
-

First Report on Fusarium Brachygibbosum Isolate FIR 16_ITS Isolated from Iraqi Wheat Plant

(<http://www.jeeng.net/First-Report-on-Fusarium-Brachygibbosum-Isolate-FIR-16-ITS-Isolated-from-Iraqi-Wheat,118295,0,2.html>)
Hamdia Z. Ali (<http://www.jeeng.net/Author-Hamdia-Ali/136194>), Mahdi S. Hameed (<http://www.jeeng.net/Author-Mahdi-Hameed/138116>), Abdulrahman A. Abdulrahman (<http://www.jeeng.net/Author-Abdulrahman-Abdulrahman/138119>), Hutham M. Saood (<http://www.jeeng.net/Author-Hutham-Saood/138118>)

J. Ecol. Eng. 2020; 21(3):81–86

DOI: <https://doi.org/10.12911/22998993/118295> (<https://doi.org/10.12911/22998993/118295>)

- Abstract
(<http://www.jeeng.net/First-Report-on-Fusarium-Brachygibbosum-Isolate-FIR-16-ITS-Isolated-from-Iraqi-Wheat,118295,0,2.html>)
 Article (PDF) (<http://www.jeeng.net/pdf-118295-48459?filename=First%20Report%20on%20Fusarium.pdf>)
-

Soybean Yield and Vegetative Water Consumption in Various Irrigation Methods

(<http://www.jeeng.net/Soybean-Yield-and-Vegetative-Water-Consumption-in-Various-Irrigation-Methods,118294,0,2.html>)

Rakhiya Yelnazarkyzy (<http://www.jeeng.net/Author-Rakhiya-Yelnazarkyzy/121881>), Serik Kenenbayev (<http://www.jeeng.net/Author-Serik-Kenenbayev/136191>), Z.O. Ospanbayev (<http://www.jeeng.net/Author-Z.O.-Ospanbayev/136192>), P.A. Kalashnikov (<http://www.jeeng.net/Author-P.A.-Kalashnikov/136193>)

J. Ecol. Eng. 2020; 21(3):87–93

DOI: <https://doi.org/10.12911/22998993/118294> (<https://doi.org/10.12911/22998993/118294>)

- Abstract
(<http://www.jeeng.net/Soybean-Yield-and-Vegetative-Water-Consumption-in-Various-Irrigation-Methods,118294,0,2.html>)
 Article (PDF) (<http://www.jeeng.net/pdf-118294-48119?filename=Soybean%20Yield%20and.pdf>)
-

[EN \(http://www.jeeng.net/\)](http://www.jeeng.net/)[PL \(http://www.jeeng.net/pl\)](http://www.jeeng.net/pl)

The Performance of Hybrid Constructed Wetland System for Treating Batik Wastewater

(<http://www.jeeng.net/The-Performance-of-Hybrid-Constructed-Wetland-System-for-Treating-Batik-Wastewater,118292,0,2.html>)

Erina Rahmadyanti (<http://www.jeeng.net/Author-Erina-Rahmadyanti/136189>), Oktavia Audina (<http://www.jeeng.net/Author-Oktavia-Audina/136600>)

J. Ecol. Eng. 2020; 21(3):94–103

DOI: <https://doi.org/10.12911/22998993/118292> (<https://doi.org/10.12911/22998993/118292>)

Abstract

Stats

(<http://www.jeeng.net/The-Performance-of-Hybrid-Constructed-Wetland-System-for-Treating-Batik-Wastewater,118292,0,2.html>)

Article (PDF) (<http://www.jeeng.net/pdf-118292-48011?filename=The%20Performance%20of%20Hybrid.pdf>)

The Characteristics of Organic Compounds in Landfill Leachate Biologically Treated in Different Technological Conditions

(<http://www.jeeng.net/The-Characteristics-of-Organic-Compounds-in-Landfill-Leachate-Biologically-Treated,118291,0,2.html>)

Justyna Koc-Jurczyk (<http://www.jeeng.net/Author-Justyna-Koc-Jurczyk/596>), Łukasz Jurczyk (<http://www.jeeng.net/Author-Łukasz-Jurczyk/589>)

J. Ecol. Eng. 2020; 21(3):104–111

DOI: <https://doi.org/10.12911/22998993/118291> (<https://doi.org/10.12911/22998993/118291>)

Abstract

Stats

(<http://www.jeeng.net/The-Characteristics-of-Organic-Compounds-in-Landfill-Leachate-Biologically-Treated,118291,0,2.html>)

Article (PDF) (<http://www.jeeng.net/pdf-118291-48120?filename=The%20Characteristics%20of.pdf>)

Fractionation of Cadmium, Lead and Copper in Municipal Solid Waste Incineration Bottom Ash

(<http://www.jeeng.net/Fractionation-of-Cadmium-Lead-and-Copper-in-Municipal-Solid-Waste-Incineration-Bottom,118290,0,2.html>)

Adam Łukowski (<http://www.jeeng.net/Author-Adam-Łukowski/44131>), Joanna Irena Olejniczak (<http://www.jeeng.net/Author-Joanna-Olejniczak/137983>)

J. Ecol. Eng. 2020; 21(3):112–116

DOI: <https://doi.org/10.12911/22998993/118290> (<https://doi.org/10.12911/22998993/118290>)
 EN (<http://www.jeeng.net/>) PL (<http://www.jeeng.net/pl>)



✉ Abstract
 (<http://www.jeeng.net/Fractionation-of-Cadmium-Lead-and-Copper-in-Municipal-Solid-Waste-Incineration-Bottom,118290,0,2.html>)

✉ Article (PDF) (http://www.jeeng.net/pdf-118290-48009?filename=Fractionation%20of%20Cadmium_.pdf)

Experimental Equipment for Investigation of Soybean Storage with Active Ventilation

(<http://www.jeeng.net/Experimental-Equipment-for-Investigation-of-Soybean-Storage-with-Active-Ventilation,118289,0,2.html>)

Sholpan Duisenova (<http://www.jeeng.net/Author-Sholpan-Duisenova/136183>), Aibek Atykhanov (<http://www.jeeng.net/Author-Aibek-Atykhanov/136184>), Dimitar Karaivanov (<http://www.jeeng.net/Author-Dimitar-Karaivanov/136185>), Bekbossyn Kassymbayev (<http://www.jeeng.net/Author-Bekbossyn-Kassymbayev/136186>), Kabdryakhim Kalym (<http://www.jeeng.net/Author-Kabdryakhim-Kalym/136187>), Aigul Sagyndikova (<http://www.jeeng.net/Author-Aigul-Sagyndikova/136188>)

J. Ecol. Eng. 2020; 21(3):117–127

DOI: <https://doi.org/10.12911/22998993/118289> (<https://doi.org/10.12911/22998993/118289>)

✉ Abstract
 (<http://www.jeeng.net/Experimental-Equipment-for-Investigation-of-Soybean-Storage-with-Active-Ventilation,118289,0,2.html>)

✉ Article (PDF) (<http://www.jeeng.net/pdf-118289-48028?filename=Experimental%20Equipment.pdf>)

Study of Liquid Waste Quality and Potential Pollution Load of Motor Vehicle Wash Business in Bekasi City (Indonesia)

(<http://www.jeeng.net/Study-of-Liquid-Waste-Quality-and-Potential-Pollution-Load-of-Motor-Vehicle-Wash,118288,0,2.html>)

Elanda Fikri (<http://www.jeeng.net/Author-Elanda-Fikri/64655>), Arifah Nur Shabrina Putri (<http://www.jeeng.net/Author-Arifah-Shabrina-Putri/136182>), Teguh Budi Prijanto (<http://www.jeeng.net/Author-Teguh-Prijanto/138106>), Osman Syarieff (<http://www.jeeng.net/Author-Osman-Syarieff/138107>)

J. Ecol. Eng. 2020; 21(3):128–134

DOI: <https://doi.org/10.12911/22998993/118288> (<https://doi.org/10.12911/22998993/118288>)

✉ Abstract
 (<http://www.jeeng.net/Study-of-Liquid-Waste-Quality-and-Potential-Pollution-Load-of-Motor-Vehicle-Wash,118288,0,2.html>)





Vehicle-

Wash,118288,0,2.html)

EN (<http://www.jeeng.net/>)PL (<http://www.jeeng.net/pl>)[Article \(PDF\) \(<http://www.jeeng.net/pdf-118288-48110?filename=Study%20of%20Liquid%20Waste.pdf>\)](http://www.jeeng.net/pdf-118288-48110?filename=Study%20of%20Liquid%20Waste.pdf)

The Relationship among Dissolved Inorganic Phosphate, Particulate Inorganic Phosphate, and Chlorophyll-a in Different Seasons in the Coastal Seas of Semarang and Jepara

(<http://www.jeeng.net/The-Relationship-among-Dissolved-Inorganic-Phosphate-Particulate-Inorganic-Phosphate,118287,0,2.html>)

Lilik Maslukah (<http://www.jeeng.net/Author-Lilik-Maslukah/97391>), Muhammad Zainuri (<http://www.jeeng.net/Author-Muhammad-Zainuri/97394>), Anindya Wirasatriya (<http://www.jeeng.net/Author-Anindya-Wirasatriya/119152>), Siti Maisyarah (<http://www.jeeng.net/Author-Siti-Maisyarah/136181>)

J. Ecol. Eng. 2020; 21(3):135–142

DOI: <https://doi.org/10.12911/22998993/118287> (<https://doi.org/10.12911/22998993/118287>)

[Abstract](#)

Stats

(<http://www.jeeng.net/The-Relationship-among-Dissolved-Inorganic-Phosphate-Particulate-Inorganic-Phosphate,118287,0,2.html>)

[Article \(PDF\) \(<http://www.jeeng.net/pdf-118287-48111?filename=The%20Relationship%20among.pdf>\)](#)

Manual Treatment of Urban Wastewater by Chemical Precipitation for Production of Hydroponic Nutrient Solutions

(<http://www.jeeng.net/Manual-Treatment-of-Urban-Wastewater-by-Chemical-Precipitation-for-Production-of,118286,0,2.html>)

Tânia Correia (<http://www.jeeng.net/Author-Tânia-Correia/136175>), Mariana Regato (<http://www.jeeng.net/Author-Mariana-Regato/136178>), Adelaide Almeida (<http://www.jeeng.net/Author-Adelaide-Almeida/136180>), Teresa Santos (<http://www.jeeng.net/Author-Teresa-Santos/136179>), Leonor Amaral (<http://www.jeeng.net/Author-Leonor-Amaral/136177>), Maria de Fátima Nunes Carvalho (<http://www.jeeng.net/Author-Maria%20de%20F%C3%A1tima%20Carvalho/136176>)

J. Ecol. Eng. 2020; 21(3):143–152

DOI: <https://doi.org/10.12911/22998993/118286> (<https://doi.org/10.12911/22998993/118286>)

[Abstract](#)

Stats

(<http://www.jeeng.net/Manual-Treatment-of-Urban-Wastewater-by-Chemical-Precipitation-for-Production-of,118286,0,2.html>)

[Article \(PDF\) \(<http://www.jeeng.net/pdf-118286-48982?filename=Manual%20Treatment%20of%20Urban.pdf>\)](#)

Study on the Application of Floating Beds of Macrophytes (*Vetiveria zizanioides* and *Phragmites australis*) in Pilot Scale for the Removal of Heavy Metals from Águas Fortes Stream (Alentejo-Portugal)

(<http://www.jeeng.net/Study-on-the-Application-of-Floating-Beds-of-Macrophytes-Vetiveria-zizanioides-and,118285,0,2.html>)

Teresa Borralho (<http://www.jeeng.net/Author-Teresa-Borralho/114505>), David Gago (<http://www.jeeng.net/Author-David-Gago/137159>), Adelaide Almeida (<http://www.jeeng.net/Author-Adelaide-Almeida/136180>)

J. Ecol. Eng. 2020; 21(3):153–163

DOI: <https://doi.org/10.12911/22998993/118285> (<https://doi.org/10.12911/22998993/118285>)

📄 Abstract

(<http://www.jeeng.net/Study-on-the-Application-of-Floating-Beds-of-Macrophytes-Vetiveria-zizanioides-and,118285,0,2.html>)

📄 Article (PDF) (<http://www.jeeng.net/pdf-118285-48122?filename=Study%20on%20the%20Application.pdf>)

📊 Stats

Effect of Diesel Fuel Temperature on the Nitrogen Oxides Emission from a Compression-Ignition Engine

(<http://www.jeeng.net/Effect-of-Diesel-Fuel-Temperature-on-the-Nitrogen-Oxides-Emission-from-a-Compression,118283,0,2.html>)

Mirosław Czechlowski (<http://www.jeeng.net/Author-Mirosław-Czechlowski/136174>)

J. Ecol. Eng. 2020; 21(3):164–170

DOI: <https://doi.org/10.12911/22998993/118283> (<https://doi.org/10.12911/22998993/118283>)

📄 Abstract

(<http://www.jeeng.net/Effect-of-Diesel-Fuel-Temperature-on-the-Nitrogen-Oxides-Emission-from-a-Compression,118283,0,2.html>)

📄 Article (PDF) (<http://www.jeeng.net/pdf-118283-48109?filename=Effect%20of%20Diesel%20Fuel.pdf>)

📊 Stats

The State and Needs of the Development of Water Supply and Sewerage Infrastructure in the Radzyń District

(<http://www.jeeng.net/The-State-and-Needs-of-the-Development-of-Water-Supply-and-Sewerage-Infrastructure,118282,0,2.html>)

Monika Bogusz (<http://www.jeeng.net/Author-Monika-Bogusz/136173>), Michał Marzec (<http://www.jeeng.net/Author-Michał-Marzec/465>), Arkadiusz Malik (<http://www.jeeng.net/Author-Arkadiusz-Malik/66955>), Krzysztof Jóźwiakowski (<http://www.jeeng.net/Author-Krzysztof-Jóźwiakowski/39779>)

J. Ecol. Eng. 2020; 21(3):171–179

DOI: <https://doi.org/10.12911/22998993/118282> (<https://doi.org/10.12911/22998993/118282>)

✉ Abstract
[\(http://www.jeeng.net/The-State-and-Needs-of-the-Development-of-Water-Supply-and-Sewerage-Infrastructure,118282,0,2.html\)](http://www.jeeng.net/The-State-and-Needs-of-the-Development-of-Water-Supply-and-Sewerage-Infrastructure,118282,0,2.html)

[EN \(http://www.jeeng.net/\)](http://www.jeeng.net/)

[PL \(http://www.jeeng.net/pl/\)](http://www.jeeng.net/pl/)

✉ Article (PDF) ([http://www.jeeng.net/pdf-118282-48123?filename=The State and Needs of.pdf](http://www.jeeng.net/pdf-118282-48123?filename=The%20State%20and%20Needs%20of.pdf))

Effect of Unitary Soil Tillage Energy on Soil Aggregate Structure and Erosion Vulnerability

(<http://www.jeeng.net/Effect-of-Unitary-Soil-Tillage-Energy-on-Soil-Aggregate-Structure-and-Erosion-Vulnerability,118281,0,2.html>)

Tomasz Wojciechowski (<http://www.jeeng.net/Author-Tomasz-Wojciechowski/136171>), Andrzej Mazur (<http://www.jeeng.net/Author-Andrzej-Mazur/520>), Andrzej Przybylak (<http://www.jeeng.net/Author-Andrzej-Przybylak/40201>), Jarosław Piechowiak (<http://www.jeeng.net/Author-Jarosław-Piechowiak/136172>)

J. Ecol. Eng. 2020; 21(3):180–185

DOI: <https://doi.org/10.12911/22998993/118281> (<https://doi.org/10.12911/22998993/118281>)

✉ Abstract
[\(http://www.jeeng.net/Effect-of-Unitary-Soil-Tillage-Energy-on-Soil-Aggregate-Structure-and-Erosion-Vulnerability,118281,0,2.html\)](http://www.jeeng.net/Effect-of-Unitary-Soil-Tillage-Energy-on-Soil-Aggregate-Structure-and-Erosion-Vulnerability,118281,0,2.html)

✉ Article (PDF) ([http://www.jeeng.net/pdf-118281-48124?filename=Effect of Unitary Soil.pdf](http://www.jeeng.net/pdf-118281-48124?filename=Effect%20of%20Unitary%20Soil.pdf))

Assessment of the “Oława” Smelter (Oława, Southwest Poland) on the Environment with Ecotoxicological Tests

(<http://www.jeeng.net/Assessment-of-the-Oława-Smelter-Oława-Southwest-Poland-on-the-Environment-with-Ecotoxicological,118280,0,2.html>)

Klaudia Radlińska (<http://www.jeeng.net/Author-Klaudia-Radlińska/136170>), Magdalena Wróbel (<http://www.jeeng.net/Author-Magdalena-Wróbel/26424>), Agnieszka Stojanowska (<http://www.jeeng.net/Author-Agnieszka-Stojanowska/93766>), Justyna Rybak (<http://www.jeeng.net/Author-Justyna-Rybak/28>)

J. Ecol. Eng. 2020; 21(3):186–191

DOI: <https://doi.org/10.12911/22998993/118280> (<https://doi.org/10.12911/22998993/118280>)

✉ Abstract
[\(<http://www.jeeng.net/Assessment-of-the-Oława-Smelter-Oława-Southwest-Poland-on-the-Environment>\)](http://www.jeeng.net/Assessment-of-the-Oława-Smelter-Oława-Southwest-Poland-on-the-Environment)

with-

[N \(http://www.jeeng.net/\)](http://www.jeeng.net/)[PL \(http://www.jeeng.net/pl/\)](http://www.jeeng.net/pl/)[Article \(PDF\) \(http://www.jeeng.net/pdf-118280-48112?filename=Assessment of the _Olawa_.pdf\)](http://www.jeeng.net/pdf-118280-48112?filename=Assessment%20of%20the%20_Olawa_.pdf)

The Sprouting Activity of Oil Flax Seeds after Treatment with a Preparation Made of Bottom Ash and Technical Carbon

(<http://www.jeeng.net/The-Sprouting-Activity-of-Oil-Flax-Seeds-after-Treatment-with-a-Preparation-Made,118279,0,2.html>)

Abilzhan Khusainov (<http://www.jeeng.net/Author-Abilzhan-Khusainov/136165>), Leonid Skipin (<http://www.jeeng.net/Author-Leonid-Skipin/136166>), Gulmira Kyzdarbekova (<http://www.jeeng.net/Author-Gulmira-Kyzdarbekova/136167>), Razia Khusainova (<http://www.jeeng.net/Author-Razia-Khusainova/136168>), Anuar Kakabaev (<http://www.jeeng.net/Author-Anuar-Kakabaev/136169>)

J. Ecol. Eng. 2020; 21(3):192–201

DOI: <https://doi.org/10.12911/22998993/118279> (<https://doi.org/10.12911/22998993/118279>)

[Abstract](#)

Stats

(<http://www.jeeng.net/The-Sprouting-Activity-of-Oil-Flax-Seeds-after-Treatment-with-a-Preparation-Made,118279,0,2.html>)

[Article \(PDF\) \(http://www.jeeng.net/pdf-118279-48125?filename=The%20Sprouting%20Activity%20of.pdf">Article \(PDF\) \(http://www.jeeng.net/pdf-118279-48125?filename=The Sprouting Activity of.pdf\)](#)

The Issues of Catchment Boundaries Delimitation in a Suburban Area – the Case Study of Poznań Metropolitan Area, Poland

(<http://www.jeeng.net/The-Issues-of-Catchment-Boundaries-Delimitation-in-a-Suburban-Area-the-Case-Study,118278,0,2.html>)

Karol Dawid Mrozik (<http://www.jeeng.net/Author-Karol-Mrozik/454>), Michał Napierała (<http://www.jeeng.net/Author-Michał-Napierała/453>), Anna Adamska (<http://www.jeeng.net/Author-Anna-Adamska/15603>)

J. Ecol. Eng. 2020; 21(3):202–208

DOI: <https://doi.org/10.12911/22998993/118278> (<https://doi.org/10.12911/22998993/118278>)

[Abstract](#)

Stats

(<http://www.jeeng.net/The-Issues-of-Catchment-Boundaries-Delimitation-in-a-Suburban-Area-the-Case-Study,118278,0,2.html>)

[Article \(PDF\) \(http://www.jeeng.net/pdf-118278-48113?filename=The%20Issues%20of%20Catchment.pdf">Article \(PDF\) \(http://www.jeeng.net/pdf-118278-48113?filename=The Issues of Catchment.pdf\)](#)

Rainwater Management Solutions and their Impact on Shaping Inner City Areas Undergoing Transformation (Case Study of the ZAC Clichy-Batignolles Area in Paris)

(<http://www.jeeng.net/Rainwater-Management-Solutions-and-their-Impact-on-Shaping-Inner-City-Areas-Undergoing,118277,0,2.html>) EN (<http://www.jeeng.net/>) PL (<http://www.jeeng.net/pl>)  

Renata Józwik (<http://www.jeeng.net/Author-Renata-Józwik/136164>)

J. Ecol. Eng. 2020; 21(3):209–219

DOI: <https://doi.org/10.12911/22998993/118277> (<https://doi.org/10.12911/22998993/118277>)

 Abstract

 Stats

(<http://www.jeeng.net/Rainwater-Management-Solutions-and-their-Impact-on-Shaping-Inner-City-Areas-Undergoing,118277,0,2.html>)

 Article (PDF) ([http://www.jeeng.net/pdf-118277-48126?filename=Rainwater Management.pdf](http://www.jeeng.net/pdf-118277-48126?filename=Rainwater%20Management.pdf))

The Effects of Water Extracts from Tansy on Pea Leaf Weevil and Black Bean Aphid

(<http://www.jeeng.net/The-Effects-of-Water-Extracts-from-Tansy-on-Pea-Leaf-Weevil-and-Black-Bean-Aphid,118275,0,2.html>)

Natalia Kwiecień (<http://www.jeeng.net/Author-Natalia-Kwiecień/137427>), Janina Gospodarek

(<http://www.jeeng.net/Author-Janina-Gospodarek/8189>), Elżbieta Boligłowa (<http://www.jeeng.net/Author-Elżbieta-Boligłowa/136264>)

J. Ecol. Eng. 2020; 21(3):220–227

DOI: <https://doi.org/10.12911/22998993/118275> (<https://doi.org/10.12911/22998993/118275>)

 Abstract

 Stats

(<http://www.jeeng.net/The-Effects-of-Water-Extracts-from-Tansy-on-Pea-Leaf-Weevil-and-Black-Bean-Aphid,118275,0,2.html>)

 Article (PDF) ([http://www.jeeng.net/pdf-118275-48035?filename=The Effects of Water.pdf](http://www.jeeng.net/pdf-118275-48035?filename=The%20Effects%20of%20Water.pdf))

Sweet Corn Yield Simulation Using Normalized Difference Vegetation Index and Leaf Area Index

(<http://www.jeeng.net/Sweet-Corn-Yield-Simulation-Using-Normalized-Difference-Vegetation-Index-and-Leaf,118274,0,2.html>)

Pavlo Volodymyrovych Lykhovyd (<http://www.jeeng.net/Author-Pavlo-Lykhovyd/89354>)

J. Ecol. Eng. 2020; 21(3):228–236

DOI: <https://doi.org/10.12911/22998993/118274> (<https://doi.org/10.12911/22998993/118274>)

 Abstract

 Stats

(<http://www.jeeng.net/Sweet-Corn-Yield-Simulation-Using-Normalized-Difference-Vegetation-Index-and-Leaf,118274,0,2.html>)



Difference-
Vegetation-Index-
and-
Leaf,118274,0,2.html)

EN (<http://www.jeeng.net/>)

PL (<http://www.jeeng.net/pl>)

Article (PDF) (<http://www.jeeng.net/pdf-118274-48127?filename=Sweet%20Corn%20Yield.pdf>)

The Use of Zeta Potential Measurement as a Control Tool of Surface Water Coagulation

(<http://www.jeeng.net/The-Use-of-Zeta-Potential-Measurement-as-a-Control-Tool-of-Surface-Water-Coagulation,118273,0,2.html>)

Izabela Teresa Zimoch (<http://www.jeeng.net/Author-Izabela-Zimoch/34570>), Dominik Mroczko (<http://www.jeeng.net/Author-Dominik-Mroczko/84275>)

J. Ecol. Eng. 2020; 21(3):237–242

DOI: <https://doi.org/10.12911/22998993/118273> (<https://doi.org/10.12911/22998993/118273>)

Abstract
(<http://www.jeeng.net/The-Use-of-Zeta-Potential-Measurement-as-a-Control-Tool-of-Surface-Water-Coagulation,118273,0,2.html>)

Article (PDF) (<http://www.jeeng.net/pdf-118273-48128?filename=The%20Use%20of%20Zeta%20Potential.pdf>)

Stats

Submit your paper (<https://www.editorialsystem.com/jeeng>)

Instructions for Authors (<http://www.jeeng.net/Instructions-for-Authors,25.html>)

All issues (<http://www.jeeng.net/Archive>)

Articles in press (<http://www.jeeng.net/Online-first>)

Most read



Month Year

Impact of Municipal Solid Waste Landfill on Environment – a Case Study (<http://www.jeeng.net/Impact-of-Municipal-Solid-Waste-Landfill-on-Environment-a-Case-Study,89664,0,2.html>)

Wastewater Treatment Methods for Effluents from the Confectionery Industry – an Overview (<http://www.jeeng.net/Wastewater-Treatment-Methods-for-Effluents-from-the-Confectionery-Industry-an-Overview,112557,0,2.html>)

Design of Biofilter Odor Removal System for Conventional Wastewater Treatment Plant
EN (<http://www.jeeng.net/>) PL (<http://www.jeeng.net/pl>)
(<http://www.jeeng.net/Design-of-Biofilter-Odor-Removal-System-for-Conventional-Wastewater-Treatment-Plant,89791,0,2.html>)



Indexes



Keywords index (<http://www.jeeng.net/Keywords>)

Authors index (<http://www.jeeng.net/Authors>)



Copyright © 2006-2020 Bentus (<http://www.bentus.com>) All rights reserved.

**SJR**

Scimago Journal & Country Rank

Enter Journal Title, ISSN or Publisher Name

[Home](#)[Journal Rankings](#)[Country Rankings](#)[Viz Tools](#)[Help](#)[About Us](#)

Ads by Google

[Stop seeing this ad](#) [Why this ad? ⓘ](#)

Journal of Ecological Engineering

Country	Poland - SIR Ranking of Poland	10
Subject Area and Category	Agricultural and Biological Sciences Ecology, Evolution, Behavior and Systematics	
	Environmental Science Environmental Science (miscellaneous)	H Index
Publisher	Polskie Towarzystwo Inżynierii Ekologicznej	
Publication type	Journals	
ISSN	22998993	
Coverage	2013-ongoing	
Scope	- Industrial and municipal waste management - Pro-ecological technologies and products - Energy-saving technologies - Environmental landscaping - Environmental monitoring - Climate change in the environment - Sustainable development - Processing and usage of mineral resources - Recovery of valuable materials and fuels - Surface water and groundwater management - Water and wastewater treatment - Smog and air pollution prevention - Protection and reclamation of soils - Reclamation and revitalization of degraded areas - Heavy metals in the environment - Renewable energy technologies - Environmental protection of rural areas - Restoration and protection of urban environment - Prevention of noise in the environment - Environmental life-cycle assessment (LCA) - Simulations and computer modeling for the environment	
	Homepage How to publish in this journal Contact	
	Join the conversation about this journal	


IJENS Journals

IJENS publisher SCOPUS - indexed; Call For Papers


ijens.org

OPEN


Quartiles




SJR




Citations per document



Total Cites




% International Collaboration



External Cites per Doc

Cites per Doc






Citable documents

Non-citable documents

Cited documents

Uncited documents



← Show this widget in your own website

Just copy the code below and paste within your html code:

```
<a href="https://www.scimagojr.com/j...>
```



Dr. MOHAMMAD AL ZUBI 1 year ago

please what is the impact factor of this journal

reply



Elena Corera 1 year ago

SCImago Team

Dear Dr Mohammad,

thank you very much for your request. You can check that information in SJR website
<https://www.scimagojr.com>

Best Regards,

SCImago Team

Leave a comment

Name

Email

(will not be published)

I'm not a robot

reCAPTCHA
[Privacy](#) - [Terms](#)

Submit

The users of Scimago Journal & Country Rank have the possibility to dialogue through comments linked to a specific journal. The purpose is to have a forum in which general doubts about the processes of publication in the journal, experiences and other issues derived from the publication of papers are resolved. For topics on particular articles, maintain the dialogue through the usual channels with your editor.

IJENS Journals

IJENS publisher SCOPUS - indexed; Call For Papers


ijens.org

OPEN

Developed by:



Powered by:



Follow us on @ScimagoJR

Scimago Lab, Copyright 2007-2020. Data Source: Scopus®





(<http://www.jeeng.net/>)

About the Journal

Journal of Ecological Engineering (JEE) is a peer-reviewed international journal that publishes original research and review articles in the areas of the protection and restoration of the natural environment.

All articles are published full in English.

The articles are published in an electronic version, and the printed version does not appear.

JEE is a member of CrossRef – the citation-linking backbone for online publications.

iThenticate – a plagiarism detection service

iThenticate service was developed in partnership with CrossRef. It is the most prominent similarity check service of plagiarism detection include intellectual property protection and document-versus-documents analysis. The editorial board undertake the comprehensive manuscripts screening, aiming to eliminate plagiarism and provide a high standard and quality peer-review process.

Topics of the Journal

- Industrial and municipal waste management
- Pro-ecological technologies and products
- Energy-saving technologies
- Environmental landscaping
- Environmental monitoring
- Climate change in the environment
- Sustainable development
- Processing and usage of mineral resources
- Recovery of valuable materials and fuels
- Surface water and groundwater management
- Water and wastewater treatment
- Smog and air pollution prevention
- Protection and reclamation of soils
- Reclamation and revitalization of degraded areas
- Heavy metals in the environment



- Renewable energy technologies
- Environmental protection of rural areas [EN](http://www.jeeng.net/)
- Restoration and protection of urban environment
- Prevention of noise in the environment
- Environmental life-cycle assessment (LCA)
- Simulations and computer modeling for the environment

Abstracting and Indexing

- Emerging Sources Citation Index (ESCI) – Web of Science Core Collection
- SCOPUS
- EBSCOhost
- J-Gate
- BazTech
- IndexCopernicus
- Google Scholar
- DOAJ – Directory of Open Access Journals
- CNKI Scholar

The frequency of issues

Quarterly, from 2013 to 2016

Bimonthly, from 2017, continuing

Journal Metrics (<http://www.jeeng.net/Journal-Metrics,1909.html>)

Submit your paper (<https://www.editorialsystem.com/jeeng>)

Instructions for Authors (<http://www.jeeng.net/Instructions-for-Authors,25.html>)

All issues (<http://www.jeeng.net/Archive>)

Articles in press (<http://www.jeeng.net/Online-first>)

Most read



Month Year

Impact of Municipal Solid Waste Landfill on Environment – a Case Study (<http://www.jeeng.net/Impact-of-Municipal-Solid-Waste-Landfill-on-Environment-a-Case-Study,89664,0,2.html>)

Wastewater Treatment Methods for Effluents from the Confectionery Industry – an Overview (<http://www.jeeng.net/Wastewater-Treatment-Methods-for-Effluents-from-the-Confectionery-Industry-an-Overview,112557,0,2.html>)

The Impact of Biopreparations and Sowing Dates on the Productivity of Fennel (*Foeniculum vulgare* Mill.)

(<http://www.jeeng.net/The-Impact-of-Biopreparations-and-Sowing-Dates-on-the-Productivity-of-Fennel-Foeniculum,119802,0,2.html>)



Indexes



Keywords index (<http://www.jeeng.net/Keywords>)

Authors index (<http://www.jeeng.net/Authors>)



Copyright © 2006-2020 Bentus (<http://www.bentus.com>) All rights reserved.



Editorial Board

EDITOR-IN-CHIEF:

Gabriel Borowski – Lublin University of Technology, Faculty of Environmental Engineering, Poland
e-mail: g.borowski@pollub.pl (<mailto:g.borowski@pollub.pl>)

SCIENTIFIC BOARD MEMBERS:

- Maria de Fátima Nunes de Carvalho** – Polytechnic Institute of Beja, Portugal
- Myroslav S. Malovany** – Lviv Polytechnic National University, Ukraine
- Maria Cristina Lavagnolo** – University of Padova, Italy
- Antonio Joao Carvalho de Albuquerque** – University of Beira Interior, Covilhã, Portugal
- Sameh Alsaqoor** – Tafila Technical University, Jordan
- Süer Anaç** – Ege University, Izmir, Turkey
- Rune Bakke** – University College of Southeast Norway, Norway
- Nelson Barros** – University of Fernando Pessoa, Porto, Portugal
- Zhihong Cao** – Institute of Soil Sciences, Chinese Academy of Sciences, Nanjing, China
- Aneta Czechowska-Kosacka** – Lublin University of Technology, Poland
- Magdalena Gajewska** – Gdańsk University of Technology, Poland
- Joan Garcia** – Polytechnic University of Catalonia, Barcelona, Spain
- Hassimi Abu Hasan** – National University of Malaysia
- Faruque Hossain** – New York University, New York, USA
- Katarzyna Ignatowicz** – Białystok University of Technology, Poland
- Krzysztof Jóźwiakowski** – University of Life Sciences in Lublin, Poland
- Aleksander Kiryluk** – Białystok University of Technology, Poland
- Michał Kopeć** – University of Agriculture in Kraków, Poland
- Joanna Kostecka** – University of Rzeszów, Poland
- Jana Kotovicová** – Mendel University in Brno, Czech Republic
- Peter Kováčik** – Slovak University of Agriculture (SUA) in Nitra, Slovak Republic
- Grzegorz Kusza** – Opole University, Poland
- Fabio Masi** – IRIDRA S.r.l., Florence, Italy
- Yuriy A. Mazhaysky** – Ryazan State Agricultural Academy, Ryazan, Russia
- Álvaro Monteiro** – University of Fernando Pessoa, Porto, Portugal
- Adam M. Paruch** – Norwegian Institute for Agricultural and Environmental Research – Bioforsk, Norway

Lucjan Pawłowski – Lublin University of Technology, Poland

Ryszard Pokładek – Wrocław University of Environmental and Life Sciences, Poland
EN (<http://www.jeeng.net/>) PL (<http://www.jeeng.net/pl>)

Katerina Pozachen'yuk – Taurida National V.I. Vernadsky University, Ukraine

Harsha Ratnaweera – Norwegian Institute for Water Research – NIVA, Oslo, Norway

Czesława Rosik-Dulewska – Opole University, Poland

Hynek Roubík – Czech University of Life Sciences Prague, Czech Republic

Pavel Ryant – Mendel University in Brno, Czech Republic

Heralt Schöne – Neubrandenburg University of Applied Sciences, Germany

László Simon – University College of Nyíregyháza, Hungary

Elżbieta Skorbiłowicz – Białystok University of Technology, Poland

Vladimir Soldatov – National Academy of Sciences of Belarus, Minsk, Belarus

Jung-Jeng Su – National Taiwan University, Taipei, Taiwan

Agata Szymańska-Pulikowska – Wrocław University of Environmental and Life Sciences, Poland

Alexander Tsyganov – Belarusian State Agricultural Academy, Gorki, Belarus

Tomasz Tymiński – Wrocław University of Environmental and Life Sciences, Poland

Sylvia Waara – Halmstad University, Sweden

Andrzej Wac-Włodarczyk – Lublin University of Technology, Poland

Raoul Weiler – University of Leuven, Belgium

Józefa Wiater – Białystok University of Technology, Poland

Barbara Wiśniowska-Kielian – University of Agriculture in Kraków, Poland

Xiaoping Zhu – Hunter College of The City University of New York, USA

CONSULTING EDITORS:

Izolda Gorgol – Department of Applied Mathematics, Lublin University of Technology, Poland

Szymon Obrusiewicz – Faculty of Environmental Engineering, Lublin University of Technology, Poland

Submit your paper (<https://www.editorialsystem.com/jeeng>)

Instructions for Authors (<http://www.jeeng.net/Instructions-for-Authors,25.html>)

All issues (<http://www.jeeng.net/Archive>)

Articles in press (<http://www.jeeng.net/Online-first>)

Most read



Month Year

Impact of Municipal Solid Waste Landfill on Environment – a Case Study (<http://www.jeeng.net/Impact-of-Municipal-Solid-Waste-Landfill-on-Environment-a-Case-Study,89664,0,2.html>)

Wastewater Treatment Methods for Effluents from the Confectionery Industry – an Overview

(<http://www.jeeng.net/Wastewater-treatment-methods-for-Effluents-from-the-Confectionery-Industry-an-Overview,112557,0,2.html>)

The Impact of Biopreparations and Sowing Dates on the Productivity of Fennel (*Foeniculum vulgare* Mill.)

(<http://www.jeeng.net/The-Impact-of-Biopreparations-and-Sowing-Dates-on-the-Productivity-of-Fennel-Foeniculum,119802,0,2.html>)

Indexes

Keywords index (<http://www.jeeng.net/Keywords>)

Authors index (<http://www.jeeng.net/Authors>)



Copyright © 2006-2020 Bentus (<http://www.bentus.com>) All rights reserved.



(<http://www.jeeng.net/>)

Editorial Office and Publisher

Editorial Office:

Lublin University of Technology
Environmental Engineering Faculty
Nadbystrzycka 40B St.
20-618 Lublin
POLAND
e-mail: office@jeeng.net (<mailto:office@jeeng.net>)

Publisher:

Polish Society of Ecological Engineering (PTIE)
Branch of PTIE in Lublin
Nadbystrzycka 40B St.
20-618 Lublin
POLAND

Co-publisher:

Scientific Publishing House
Lazurowa 26/1 St.
20-781 Lublin
POLAND

Submit your paper (<https://www.editorialsystem.com/jeeng>)

Instructions for Authors (<http://www.jeeng.net/Instructions-for-Authors,25.html>)

All issues (<http://www.jeeng.net/Archive>)

Articles in press (<http://www.jeeng.net/Online-first>)

[Most read](#)[EN \(<http://www.jeeng.net/>\)](#)[PL \(<http://www.jeeng.net/pl>\)](#)[Month](#) [Year](#)

Impact of Municipal Solid Waste Landfill on Environment – a Case Study (<http://www.jeeng.net/Impact-of-Municipal-Solid-Waste-Landfill-on-Environment-a-Case-Study,89664,0,2.html>)

Wastewater Treatment Methods for Effluents from the Confectionery Industry – an Overview (<http://www.jeeng.net/Wastewater-Treatment-Methods-for-Effluents-from-the-Confectionery-Industry-an-Overview,112557,0,2.html>)

The Impact of Biopreparations and Sowing Dates on the Productivity of Fennel (*Foeniculum vulgare* Mill.) (<http://www.jeeng.net/The-Impact-of-Biopreparations-and-Sowing-Dates-on-the-Productivity-of-Fennel-Foeniculum,119802,0,2.html>)

[Indexes](#)

Keywords index (<http://www.jeeng.net/Keywords>)

Authors index (<http://www.jeeng.net/Authors>)



Copyright © 2006-2020 Bentus (<http://www.bentus.com>) All rights reserved.

Journal of Ecological Engineering


JEE



CROSSMARKED



Scopus



Google

Scopus
CrossMark

Scopus
CrossMark

BibTeX

Scopus
CrossMark



CrossMark

Bioactive Compounds Evaluation of the Mudskippers in the Estuarine Area of Musi River, South Sumatera, Indonesia

By Arum Setiawan

Bioactive Compounds Evaluation of the Mudskippers in the Estuarine Area of Musi River, South Sumatera, Indonesia

Mohammad Rasyid Ridho^{1*}, Arum Setiawan¹, Sarno¹, Arwinskyah¹, Enggar Patriono¹, Sulistiono¹

¹ Department of Biology, Faculty of Mathematic and Natural Sciences, Sriwijaya University, Palembang 30139, Indonesia

* Corresponding author's e-mail: moh.rasyidridho@mipa.unsri.ac.id

ABSTRACT

This research was conducted to assess various bioactive compounds in the meat of mudskippers found in the Sungsang 2 Village, Banyuasin regency, South Sumatra. The research had been synthesized using n-hexane from February to May 2018 at Agricultural Technology Production Laboratory, Sriwijaya University. Subsequently, the mudskippers (*Periophthalmus chrysospilos*, *Boleophthalmus boddarti*, and *Periophthalmodon schlosseri*) extract was synthesized and further investigate by conducting the GC-MS analysis. The study of bioactive compounds in mudskippers was carried out by evaporating in vacuum evaporation at 30°C. The results showed that *Periophthalmus chrysospilos* has five classes of bioactive compounds such as steroids, terpenoids, carotenoids, cannabinoids, and alkaloids and *Boleophthalmus boddarti* has five classes of bioactive compounds including steroids, carotenoids, terpenoids, bufadienolides, and carotatoxins. Furthermore, *Boleophthalmus dussumieri* has four classes of bioactive compounds such as steroids, carotenoids, terpenoids, and carotatoxins and *Periophthalmodon schlosseri* has three classes of bioactive compounds including steroids, carotenoids and terpenoids. This research discussed the potential benefit of all bioactive compounds in mudskippers for alleviating and treating children with enuresis through the review of literature on the most cited bioactive compounds.

Keywords: mudskippers, Musi river, bioactive compounds

INTRODUCTION

Mudskipper is a genus from the family of Gobiidae and subfamily Oxudercinae that is habitually known as ikan gelodok at Musi river, South Sumatera (Shukla et al., 2014; Polgar et al., 2017; Lauriano et al., 2018; Ridho et al., 2019). Mudskippers are commonly found in brackish river water and beaches with certain salinity concentration, as well as mangrove areas of the Indo-Pacific region (Tytler and Vaughan, 1983). Mudskippers are amphibious fish that are able to tolerate both fresh and marine water on muddy and sandy bottom habitat, spawning in estuarine areas (Hu et al., 2016; Hong et al., 2017). Mudskippers are also capable of spending extended periods out of water (Martin et al., 2013; Konstantinidis et al., 2016; Martinez et al., 2018). They appear during low

tide, foraging on tropical mudflats, seeking mate and preserving territories for some portion of their daily cycle (Dabruzzi et al., 2019). According to the classification of amphibious gobies, 40 species of mudskippers have been currently identified in the world, which are divided into four genera of Oxudercinae, namely *Boleophthalmus*, *Periophthalmus*, *Periophthalmodon* and *Scartelaos* (Clayton 1993; Graham 1997; Graham and Wegener 2010; Ishimatsu and Gonzales 2011; Takeda et al., 2012; Murdy 2011a; You et al., 2018).

Juveniles of this Gobiidae are herbivores that eat more diatoms and filamentous algae, while the major food of adult Gobiidae are crustaceans (95.5%), polychaetas (86.4%) and land insects (85%) (Bob-Manuel, 2011). The burrows of these habitats are categorically suitable for the mudskippers for feeding grounds, spawning

areas and nursery grounds (Masuda et al., 1984; Bob-Manuel, 2011; Satapoomin and Poovachiranon, 1997; Redjeki, 2013). Mudskippers are commonly consumed and employed in traditional medicine in many countries such as China, Malaysia and Japan (Feulner 2013; Gadhavi et al., 2017; Ikram et al., 2010; looi et al., 2016 and Yang, X. X., 2017). The medicinal mudskippers were often prepared from the natural products which contain different bioactive compounds (Mahadevan et al., 2019; Wu et al., 2009). On the other hand, many fishermen and general public around the Musi river ignore the existence and benefits of mudskippers.

Corticosteroids, one of bioactive compounds in mudskippers, are steroid hormone class for overcoming inflammations (Panda and Mabalirajan, 2018; Heughten et al., 2018). These hormones are a contributing factor to alleviating enuresis (bedwetting) in child. Enuresis is most common among children or young persons and is denoted by repeated voiding of urine into the bed or clothing (Apos et al., 2017 and Shafi et al., 2019). The study area corresponded to bioactive compounds like corticosteroid compounds in mudskippers that can reduce the enuresis problem in children (Jain et al., 2016). The authors intended to evaluate the bioactive compounds in the four species mudskippers including *Periophthalmus chrysophilus*, *Boleophthalmus boddarti*, *Boleophthalmus dussumieri* and *Periophthalmodon schlosseri*.

MATERIALS AND METHODS

Fishes and experimental design

Mudskippers were obtained from an estuarine area around the Sungsang 2 village, Banyuasin regency, South Sumatera. Mudskippers, apparently healthy, were randomly caught using fishing equipment and transported to the Agricultural Technology Production Laboratory, Sriwijaya University for cleaning and removing the bones from the fish. After the filleting process, the mudskippers were divided into four groups with *Periophthalmus chrysophilus* (8 grams), *Boleophthalmus boddarti* (100 grams), *Boleophthalmus dussumieri* (13.8 grams) and *Periophthalmodon schlosseri* (100 grams) which were homogenized.

The crushed fish flesh dissolved in n-hexane solvent with a sample ratio of 1: 4 and macerated for 24 hours. The next step involved filtering with calico cloth and filter paper and evaporating with vacuum evaporator at 30°C with a speed of 200 rpm. Each extract sample was reconstituted using n-hexane solvent, then each 1 µL was taken to be injected and analyzed in GC-MS alternately.

Analysis technique

The data in this research were expressed in the form of chromatograms that contain graphs and were equipped with a list of detected chemical compounds, along with the structure of compounds, retention times, areas, and probability of the compound types. The data analysis was performed by identifying the chemical compounds detected to find out the natural name (nature of compound name), synonym, and the class of compounds of these chemical compounds. Identification was done by comparing the compound name detected with a data base on the literature in the form of books and journals. The studies reported as full-text articles, those published as abstracts only and unpublished data were included as well.

Gas chromatography and mass spectroscopy (GC-MS) analysis

The qualitative analyses of the mudskipper extract were expressed with a GCMS-QP 2010 Plus (Shimadzu, Japan) system equipped under computer control at 70 eV (Mohammed and Imad, 2013). The methanol extract in the amount of 1 µl was entered into the GC-MS with a micro injector and followed by scanning for 45 min. The temperature was maintained in the oven at 100°C and Helium gas was utilized as a carrier as well as an eluent. The helium flow was arranged to 1 ml per min and electron energy was fired by a mass detector about 70 eV. The compound of elite 1 was used as separation column (100% dimethyl poly siloxane) (Hameed et al., 2016; Kareem et al., 2015). The identity of the compounds in the extracts was assigned by the comparison of their retention indices and mass spectra fragmentation patterns with those stored on the computer library and also with the data published in the literature. The compounds were identified by comparing their spectra to those of the Wiley and NIST/EPA/NIH mass spectral libraries.

RESULTS AND DISCUSSIONS

GC-MS results of the *periophthalmus chrysospilos* extract

The gas chromatography and mass spectroscopy analysis of bioactive compounds were carried out in the *Periophthalmus chrysospilos* extract that was presented in Figure 1. It can be seen that the vertical lines in the chromatogram pattern point out the short periods of time when the ion beam is deflected-off the plate (and thus off the beam monitor) which is necessary when changing the position of the photographic plate, a process that takes about 2–3 seconds with the present arrangement. This pen deflection creates a useful signal of the exact quantities of the chromatogram being noted with the mass spectrometer and facilitates the reconnection of the spectra on the plate with the chromatogram.

The GC-MS chromatogram showed that the presence of five major peaks of bioactive compounds was detected and the components corresponding to the peaks were determined as follows. The highest peak in chromatogram pattern could be seen that the extract of *Periophthalmus chrysospilos* has retention time 14.44 minutes. On the basis of the GC-MS signal, the sample has many bioactive compounds which were indicated by the signals detected. It shows that the GC-MS analysis successfully proves that the *Periophthalmus chrysospilos* extract is suitable for treating the people with inflammatory diseases, especially enuresis.

Bioactive compounds in the *Periophthalmus chrysospilos* extract

The results of *Periophthalmus chrysospilos* extract revealed the amounts of bioactive compounds such as terpenoids, alkaloids, steroids,

carotenoids and cannabinoids. As many as 4 types of terpenoid compounds were detected, with a total of 4 compounds. One type of carotenoid compounds was detected with a total of 4 compounds. Additionally, one type of cannabinoid compounds was detected with a total of 2 compounds; and 1 alkaloid compound was detected with a total of 1 compound. The cannabinoid bioactive compounds were also detected in this sample with a total of 1 compound. Cannabinoid compounds include secondary metabolites which are usually found in cannabis plants. According to Maggyvin and Sinuraya (2017), cannabinoids are classified by source and origin. Phytocannabinoids are derivative compounds from cannabis and other plants, including delta-9-tetrahydrocannabinol (THC), cannabidiol (CBD), and cannabinol (CBN). The details of *Boleophthalmus boddarti* extract were elucidated in the Table 1.

The highest amount of bioactive compounds in *Periophthalmus chrysospilos* extract corresponded to the steroid compounds; as many as 10 types of compounds were detected with a total of 24 compounds. The total number of compounds detected are named Androst-7-ene-6,17-dione, 2,3,14-trihydroxy-, where the total amount of compounds is 10 compounds. According to Moss (1989), androstane is an anabolic steroid group that belongs to the basic carbon cycle, which is unsaturated and is an alkyl substitute in the 17th carbon chain.

GC-MS Results of the *Boleophthalmus boddarti* extract

The result of GC-MS analysis of *Boleophthalmus boddarti* extract was described as the chromatogram form for all chemical compounds that were read by the detector (Fig. 2). The vertical lines were found in this chromatogram




Figure 1. Chromatogram result of peak chemical compounds in *Periophthalmus chrysospilos*
(Source: LHP, 2018)




Figure 2. Chromatogram result of peak chemical compounds in *Boleophthalmus boddarti*
(Source: LHP, 2018)

Table 1. The amount of bioactive compounds in the *Periophthalmus chrysospilos* extract

No.	Compound names	Compound classes	Total numbers
1.	D-Limonene	Terpenoids	1
2.	Limonene	Terpenoids	1
3.	Aspidospermidin-17-ol, 1-acetyl-19,21-epoxy-15, 16-dimethoxy-	Alkaloids (Aspidospermidin)	1
4.	Choltesan-3-ol, 2-methylene-	Steroids	6
5.	Azulene, 1,4-dimethyl-7-(1-methylethyl)-	Terpenoids	1
6.	17.alfa.,21a-28, 30-Bisnorhopane	Steroids	1
7.	4-Androsten-6-ol-3,17-dione	Steroids	3
8.	3,9-Epoxy pregn-16-ene-14,20-diol, 7,11,18-triacetoxy-3-methoxy-	Steroids	1
9.	Cholestan-26-oic acid, 3,7,12-trihydroxy-	Steroids	1
10.	Rhodopin	Carotenoids	1
11.	Cannabinol, trifluoroacetate	Cannabinoids	4
12.	Cholestan-3,5-diol, 5-acetate	Steroids	2
13.	Milbemycin B, 5-demethoxy-5-one-6,28-anhydro-25-ethyl-4-methyl-13-chloro-oxime	Steroids	1
14.	9,19-Cyclolanostan-3-ol, 24,24-epoxymethano-acetate	Terpenoids	1
15.	8,14-Seco-3,19-epoxyandrostan-8,14-dione, 17-acetoxy-3 α -methoxy-4,4-dimethyl-	Steroids	11
16.	Androst-7-ene-6,17-dione, 2,3,14-trihydroxy-	Steroids	10
17.	2-Phenanthrenecarboxylic acid, 1-(1,3-dithian-2-ylmethyl)-7-hydroxy-2,4b-dimethyl-1,2,3,4,4a,4b,5,6,7,8,10,10a-dodecahydro-,methyl ester	Steroids	1

pattern. Those indicate more than one bioactive compound in the *Boleophthalmus boddarti* extract. The presence of five major peaks of bioactive compounds was detected and the most compound are from steroids. The highest peak in chromatogram pattern could be seen that the *Boleophthalmus boddarti* extract has retention time of 14.44. On the basis of the GC-MS signal, the sample has many bioactive compounds, which was shown by some signals detected. It shows that the *Boleophthalmus boddarti* extract can be successfully used for treating people with inflammatory diseases, especially enuresis, as proven by the GC-MS analysis.

Bioactive compounds in the *Boleophthalmus boddarti* extract

The results of *Boleophthalmus boddarti* extract revealed the amounts of bioactive compounds such as steroids, carotenoids, terpenoids, bufadienolide, carotaxin, cholesterol, and corticosteroid shown in Table 2. The compounds of steroid groups were detected as many as 24 types of compounds with a total of 44 compounds. Specific steroid compounds, namely corticosteroids, were detected with 2 types of compounds and a total of 3 compounds. The details of *Boleophthalmus boddarti* extract were elucidated in Table 2.

Corticosteroids are drugs containing steroid hormones that are useful in increasing the needed steroid hormones in the body, relieving inflammation and helping in alleviate asthma, allergy and multiple sclerosis (Cidlowski, 2013; Panda and Mabalirajan, 2018; Rozaliyani, 2011). According to the analysis of Prasad *et al.* (2018), corticosteroids also reduce the mortality risk from tuberculosis effect at least in the short term. A quite unique compound, called Gamabufotalin, was detected in the *Boleophthalmus boddarti* extract; it is a bioactive compound of the bufadienolide type. According to Dmitrieva *et al.* (2000), Bufadienolides are biosynthetic steroids that are independent of the side chain of cholesterol division. Bufadienolides (BDs) also include the corticosteroid groups,

which are usually obtained from amphibian poison extracts, these compounds are often used as traditional medicine in Asia.

GC-MS Results of *Boleophthalmus dussumieri* extract

The result of the GC-MS analysis of *Boleophthalmus dussumieri* extract was described as the chromatogram form for all chemical compounds that were read by the detector (Fig. 3). The vertical lines were found in this chromatogram pattern. Those indicate more than one bioactive compound in the *Boleophthalmus dussumieri* extract. The presence of four major peaks of bioactive compounds was detected by GC-MS like steroids, terpenoids,

Table 2. The amount of bioactive compounds in the *Boleophthalmus boddarti* extract

No.	Compound names	Compound classes	Total numbers
1.	<i>Curan-17-oic acid, 19,20-dihydroxy-, methyl ester</i>	Steroids	2
2.	<i>Cholestan-3-ol, 2-methylo-</i>	Steroids	12
3.	<i>Citronellol epoxide</i>	Terpenoids	1
4.	<i>Falcarinol</i>	Carotatoxin (secondary cancer anti metabolites)	1
5.	<i>Cholestan-3-one, cyclic 1,2-ethanediyl aetal</i>	Steroids	1
6.	<i>..psi.,psi.-Carotene, 1,1',2,2'-tetrahydro-1,1'-dimethoxy</i>	Carotenoids	8
7.	<i>Androsten-4-en-11-ol-3,17-dione, 9-thiocyanato-</i>	Steroids	2
8.	<i>Androsten-5-en-17-one-3,11-bis[(trimethylsilyl)oxy]-, O-(phenylmethyl)oxime</i>	Steroids	1
9.	<i>Prosta-5,13-dien-1-oic acid, 9,11,15-tris[(trimethylsilyl)oxy]-, trimethylsilyl ester</i>	Steroids	6
10.	<i>Rhodoxanthin</i>	Carotenoids	2
11.	<i>Rhodopin</i>	Carotenoids	2
12.	<i>5α-Ergost-24-en-26-oic acid, 5,6α-epoxy-4α, 18,22-trihydroxy-3-methoxy-1-oxo-, β-lactone, diacetate,</i>	Steroids	1
13.	<i>Ethyl iso-allocholate</i>	Steroids	1
14.	<i>Gamabufotalin</i>	Buadienolide (steroid aglycones in frogs)	1
15.	<i>Betamethasone acetate</i>	Corticosteroid	2
16.	<i>Cholestan-26-oic acid, 3,7,12-trihydroxy-</i>	Steroids	2
17.	<i>..psi.,psi.-Carotene, 3,4-didehydro-1,2,7,8'-tetrahydro-1-methoxy-2-oxo</i>	Carotenoids	1
18.	<i>..psi.,psi.-Carotene, 3,3',4,4'-tetrahydro-1',2'-dihydro-1-hydroxy-1'methoxy</i>	Carotenoids	1
19.	<i>Prosta-5,13-dien-1-oic acid, 9,11,15-tris[(trimethylsilyl)oxy]-, trimethylsilyl ester</i>	Steroids	2
20.	<i>17α-Acetoxy-1',1'-dicarboethoxy-1α,2α-dihydroxy cycloprop[1,2]-5α-androst-1-en-3-one</i>	Steroids	2
21.	<i>Cholesterol</i>	Cholesterol	1
22.	<i>26-Nor-5-cholest-3-ol-25-one</i>	Steroids	1
23.	<i>17-(1,5-Dimethylhexyl)-10,13-dimethyl-2,3,4,7,8,9,10,11,12,13,14,15,16,17-tetradecahydro-1Hcyclopenta[a]phenanthren-3-ol</i>	Steroids	1
24.	<i>Androst-4-en-9-thiocyanomethyl-11-ol-3,17-dione</i>	Steroids	1
25.	<i>Androst-5-en-3-one, 19-acetoxy-4,4-dimethyl-, oxime</i>	Steroids	1
26.	<i>3-Isopropyl-6α,7,10b-trimethyl-8-(2-oxo-2-phenylethyl)dodecahydrobenzo[f]chromene-7-carboxylic acid, methyl ester</i>	Corticosteroid	1
27.	<i>Androst-8-en-3-ol, 4,4,14α-trimethyl-17-(2-bromo-1-methylethyl)</i>	Steroids	1
28.	<i>1',1'-Dicarboethoxy-1α,2α-dihydro-3'H-cycloprop[1,2]cholest-1,4,6-trien-3-one</i>	Steroids	1
29.	<i>Prosta-5,13-dien-1-oic acid, 9,11,15,19-tetrakis[(trimethylsilyl)oxy]-, methyl ester</i>	Steroids	1




Figure 3. Chromatogram result of peak chemical compounds in *Boleophthalmus dussumieri*

(Source: LHP, 2018)

carotenoids and carotatoxins. The highest peak in the chromatogram pattern could be seen that *Boleophthalmus dussumieri* extract has retention time of 22.61. On the basis of the GC-MS signal, the sample has significant bioactive compounds, which was shown in Table 3.

Bioactive compounds in the *Boleophthalmus dussumieri* extract

The bioactive compounds detected from the *Boleophthalmus dussumieri* extract include steroids, terpenoids, carotenoids and carotatoxins. The compounds of steroid classes include as many as 11 types of compounds, with a total of 26 compounds. The group of carotenoid compounds was detected in as many as 6 types of compounds, with a total of 13 compounds. The group of terpenoid compounds was detected by 2 types of compounds, with a total of 2 compounds. The carotatoxin compounds were also detected in the *Boleophthalmus dussumieri* extract, with a total of 1 compound. The details of *Boleophthalmus dussumieri* extract were elucidated in Table 3.

The name of steroid compounds with the highest total number of compounds is Spirost-8-en-11-one, 3-hydroxy. According to Rajendran et al. (2017), Spirost-8-en-11-one, 3-hydroxy included the steroid compounds which have an anti-cancer bioactivity. Spirost-8-en-11-one, 3-hydroxy has the chemical formula C₂₇H₄₀O₄. The group of carotenoid compounds detected with the highest amount is Rhodopin, where the total amount of these compounds is 5 compounds. According to Komori et al. (1998), Rhodopin is a carotenoid compound which is usually produced by Rhodospirillum rubrum bacteria with the chemical formula C₄₀H₅₈O.

Bioactive carotatoxin alkaloids were also detected in the *Boleophthalmus dussumieri* extract. The name of the carotatoxin group is falcarinol and the total of compound is only 1 compound. According to Zaini et al. (2012), falcarinol (FaOH) is one of the compounds found in carrots; this compound has shown bioactive action in several types of cell cultures. Falcarinol has been shown to be cytotoxic to the culture of acute lymphoblastic leukemia cells.

GC-MS Results of the *Periophthalmodon schlosseri* extract

The result of GC-MS analysis of *Periophthalmodon schlosseri* extract was described as the chromatogram form for all chemical compounds that was read by the detector (Fig. 4). The vertical lines were found in this chromatogram pattern. Those indicate more than one bioactive compound in *Boleophthalmus boddarti* extract. The presence of five major peaks of bioactive compounds was detected and the most compound are from steroids. The highest peak in the chromatogram pattern indicated that the *Boleophthalmus boddarti* extract has retention time of 14.44. On the basis of the GC-MS signal, the sample has many bioactive compounds which was shown by some signals detected. It shows that the GC-MS analysis proves that the *Boleophthalmus boddarti* extract can be successfully used for treating people with inflammatory diseases especially enurosis.

Bioactive compounds in the *Periophthalmodon Schlosseri* extract

The bioactive compounds detected from *Periophthalmodon schlosseri* extract including steroids, carotenoids and terpenoids. As many as

Table 3. The amount of bioactive compounds in the *Boleophthalmus boddarti* extract

No.	Compound names	Compound classes	Total numbers
1.	<i>Limonene</i>	Terpenoids	1
2.	<i>D-Limonene</i>	Terpenoids	1
3.	<i>Corynan-17-ol, 18,19-didehydro-10-methoxy-, acetate (ester)</i>	Steroids	2
4.	<i>Rhodopin</i>	Carotenoids	5
5.	<i>β-Sitosterol</i>	Steroids	1
6.	<i>Spirost-8-en-11-one, 3-hydroxy</i>	Steroids	13
7.	<i>Cholest-5-en-3-ol, 24-propylidene-</i>	Steroids	2
8.	<i>Stigmasta-5,24(28)-dien-3-ol</i>	Steroids	1
9.	<i>..psi.,psi.-Carotene, 1,1',2,2'-tetrahydro-1,1'-dimethoxy</i>	Carotenoids	4
10.	<i>5α-Ergost-24-en-26-oic acid, 5,6α-epoxy-4α, 18,22-trihydroxy-3-methoxy-1-ox o-β-lactone, diacetate,</i>	Steroids	1
11.	<i>..psi.,psi.-Carotene, 3,4-didehydro-1,2,7',8'-tetrahydro-1-methoxy-2-ox o</i>	Carotenoids	1
12.	<i>Chloro-6beta-nitro-5alpha-cholest-3-one</i>	Steroids	1
13.	<i>Prosta-5,13-dien-1-oic acid, 9,11,15-tris[(trimethylsilyl)oxy]-, trimethylsilyl ester</i>	Steroids	1
14.	<i>Cholest-4-ene, 3α-(methoxymethoxy)-</i>	Steroids	1
15.	<i>Rhodoxanthin</i>	Carotenoids	1
16.	<i>4'-Apo-α.,psi.-carotenoic acid</i>	Carotenoids	1
17.	<i>Astaxanthin</i>	Carotenoids	1
18.	<i>Androst-7-ene-6,17-dione, 2,3,14-trihydroxy</i>	Steroids	2
19.	<i>Prosta-5-en-1-oic acid, 9,11,15-tris[(trimethylsilyl)oxy]-, methylsilyl ester</i>	Steroids	1
20.	<i>Falcarinol</i>	Carotatoxin	1

11 bioactive steroid compounds were detected, with a total of 21 compounds; 3 types of the carotenoid group compounds, with a total of 6 compounds; while only 1 type of terpenoid group compounds were detected with a total of 1 compound. The details of the *Periophthalmodon schlosseri* extract were elucidated in Table 4.

The steroid compounds in the *Periophthalmodon schlosseri* extract are named Cholestan-3-ol, 2-methylene-, Cholest-5-en-3-ol (3α)-, tetradecanoate, Cholesta-3,5-diene, Cholesteryl

benzoate, 17.alfa., 21α-28, 30-Bisnorhopane, Cholesterol, 26-Nor-5-cholest-3α-ol-25-one, Cholestane-3,5-diol, 5-acetate, Campesterol, Ergost-5-en-3-ol, and Androst-7-ene-6,17-dione, 2,3,14-trihydroxy-. According to Salempa and Muhamarram (2016), steroids are natural compounds which consist of a carbon skeleton and comprise three circles of six peridro phenanthrene fused into one circle of five. Saturated-click hydrocarbons have a circumference system consisting of 17 carbon atoms.




Figure 4. Chromatogram result of peak chemical compounds in *Periophthalmodon schlosseri*
(Source: LHP, 2018)

Table 4. The amount of bioactive compounds in the *Periophthalmodon schlosseri* extract

No.	Compound names	Compound classes	Total numbers
1.	D-Limonene	Terpenoids	1
2.	Cholestan-4-ol, 2-methylene-	Steroids	4
3.	Cholest-5-en-3-ol (3 α)-, tetradecanoate	Steroids	1
4.	Cholesta-3,5-diene	Steroids	1
5.	Cholesteryl benzoate	Steroids	1
6.	Rhodopin	Carotenoids	3
7.	Azafrin	Carotenoids	1
8.	17.alfa.,21 α -28,30-Bisnorhopane	Steroids	2
9.	Cholesterol	Steroids	2
10.	26-Nor-5-cholest-3 α -ol-25-one	Steroids	1
11.	Cholestane-3,5-diol, 5-acetate	Steroids	1
12.	Campesterol	Steroids	1
13.	Ergost-5-en-3-ol	Steroids	1
14.	Astaxanthin	Carotenoids	2
15.	Androst-7-ene-6,17-dione, 2,3,14-trihydroxy-	Steroids	6

D-Limonene is the only group of terpenoid compounds which was detected in the *Periophthalmodon schlosseri* extract. This compound was also detected in samples of *Periophthalmus chrysospilos* extract and *Boleophthalmus dussumieri* extract. According to Sun (2007), D-Limonene is a monocyclic terpenoid with a lemon-like odor and is a major constituent of several essential oils of Citrus sp. (orange, lemon, mandarin, lime and grapefruit). D-Limonene is widely used as an additive (additive) in perfumes, soap, food, gum and drinks. This compound is also listed in the Code of Federal Regulation as a material that is known to be safe or generally recognized as safe (GRAS) as a flavoring agent.

The group of carotenoid compounds detected includes rhodopin, azafrin and astaxanthin. The rhodopin compounds were detected in all samples with a sufficient number of total compounds in each sample. The astaxanthin compounds were also detected in the flesh extracts of *Boleophthalmus boddarti* and *Boleophthalmus dussumieri*, with a total of 1 compound each in each sample. In turn, the azafrin compounds were only detected in *Periophthalmodon schlosseri* extract. According to Gopalakrishnan and Kalaiarasi (2012), azafrin is a compound of the carotenoid ester group with the molecular formula C₂₈H₄₀O₄. The azafrin compounds include the bioactive compounds that have the potential as antioxidants, anti-inflammatory, anticancer, anti-criticism, natural dyes, eye pain medications, and as compounds that can reduce hypertension.

Physical and chemical properties of the environment at sampling locations

The main point of mudskippers sampling based on the GPS is in the coordinates S 02° 21'57.5 "E 104° 54'00.1". The environment measurements of the physical and chemical properties were carried out at the main point of sampling and obtained as presented in Table 5.

The measurement of the physical and chemical properties of the environment was carried out at 3 different times. The first measurements were made in late March, precisely in the afternoon, and obtained soil moisture data of 100%, soil pH of 6.5, water pH of 6.82, and water salinity of 0‰. The second measurement was carried out in an April morning and 100% of soil moisture was obtained, soil pH was 6.5, water pH was 7.32, and water salinity was 0‰. The third measurement was carried out in the afternoon of April and obtained 100% soil moisture data, a soil pH of 6.5, a water pH of 6.7, and water salinity of 0‰.

Table 5. Physical and chemical properties of the environment at the sampling location

Parameter	Measurement Time		
	March	April Morning	April Afternoon
Soil moisture (%)	100	100	100
Soil pH	6.5	6.5	6.5
Water pH	6.82	7.32	6.7
Waters salinity (%)	0.1	0	0

On the basis of the data of the physical and chemical properties of the environment, it can be seen that the soil moisture at the main sampling point shows 100% at 3 times measurements with different times. This shows that the sampling location has a substrate in the form of mud which is always wet. According to Kurnia et al. (2006), the soil moisture is one of the key variables in the hydrological process that plays an important role in determining the availability of water as a very fundamental element in the life of living things. The 100% soil moisture indicates that the soil is saturated with water.

The water salinities at 3 measurements with different times showed the numbers 0.1‰, 0‰ and 0‰. This shows that the waters around the sampling location tend to be tasteless. According to Masuda et al. (1984) and Redjeki (2013), the gudder is an amphidromous demersal fish that tolerates freshwater and sea, lives on muddy and sandy bottom and spawns in estuarine areas.

CONCLUSIONS

The bioactive compound analysis of the *Periophthalmus chrysospilos* extract showed that it consists of steroids, terpenoids, carotenoids, cannabinoids, and alkaloids. As many as 10 types of hormones were found in the *Periophthalmus chrysospilos* extract, with a total of 24 compounds. The *Boleophthalmus boddarti* extract comprised steroids, carotenoids, terpenoids, bufadienoloid, and carotatoksin. There were 24 types of compounds detected with a total of 44 compounds in the *Boleophthalmus boddarti* extract. The *Boleophthalmus dussumieri* extract has bioactive compounds from the groups of steroids, carotenoids, terpenoids, and carotatoksin. The *Boleophthalmus dussumieri* extract has as many as 11 types of compounds with a total of 26 compounds. The *Periophthalmodon schlosseri* extract contained the bioactive compounds including steroids, carotenoids and terpenoids. As many as 11 types of compounds were detected with a total of 21 compounds. The higher concentrations of bioactive compounds were found in the mudskippers that are possibly consumed for treating inflammatory diseases, enuresis in children and allergy.

Acknowledgments

The authors would like to thank Dr. Arum Setiawan and Sulistiono, Dr. Sarno, Dr. Arwinsky,

Enggar Patriono M.Si., College of Agricultural Technology Production Laboratory, Sriwijaya University for his help in specimen preparation. We also thank anonymous referees for constructive criticism on the manuscript. This work was supported by the projects of Kemenristekdikti.

REFERENCES

1. Bob-Manuel, F.G., 2011. Food and feeding ecology of the mudskipper *Periophthalmus koelreuteri* (Pallas) Gobiidae at Rumuolumeni Creek Niger Delta, Nigeria. Agriculture and Biology Journal of North America. 2, 897–901.
2. Clayton DA. 1993. Mudskippers. Oceanogr Mar Biol Annu. Rev 31:507–77.
3. Dabruzzi, T. F., Fangue, N. A., Kadir, N. N, and Bennett, W. A., 2019. Thermal Niche Adaptations of Common Mudskipper (*Periophthalmus kalolo*) and Barred Mudskipper (*Periophthalmus argentilineatus*) in Air and Water. Journal of Thermal Biology. 81, 170–177.
4. DKP Provinsi Sumatera Selatan, 2017. Sumberdaya: Pengelolaan Pesisir Provinsi Sumatera Selatan 1. [online] Tersedia: dislutan.sumselprov.go.id/page/38/sumberdaya#. [11 November 2017].
5. Dmitrieva, R. I. Bagrov, A. Y., Lalli, E., Sassone-Corsi, P., Stocco, D., dan Doris, P. A., 2000. Mammalian Bufadienolide is Synthesized from Cholesterol in the Adrenal Cortex by Pathway that is Independent of Cholesterol Side-chain Cleavage Hypertension. 36, 3, 442–8.
6. Feulner, G. R., and Binish R, 2013. Re-discovery of the mudskipper *Periophthalmus waltoni* Koumans, 1941 in the United Arab Emirates. Tribulus, vol. 21, 2013, p. 42+. Gale Academic Onefile, Accessed 13 Oct. 2019
7. Gadnavi, M.K., Kukadia, D., Dar, S., Gokulakannan, N., Talukdar, G., Sivakumar, K., and Gopi, G. V., 2017. Indigenous techniques of catching mudskipper in Bhavnagar and Bharuch districts, Gujarat. Indian Journal of Traditional Knowledge. 16(3), pp. 533–538.
8. Gopalakrishnan, S., and Kalaiarasi, T., 2012. Identification of Chemical Compound from the Fruits of *Cucumis trigonus* Roxb. By GC-MS Analysis. International Journal of Phytopharmacy. 2 (5) 122–128.
9. Graham, J. B., 1997. Air-breathing fishes: evolution, diversity, and adaptation. San Diego: Academic Press, p. 299.
10. Graham, J. B., and Wegner, N. C., 2010. Breathing air in water and in air: the air-breathing fishes. In: Nilsson GE, editor. Respiratory physiology of vertebrates: life with and without oxygen. Cambridge:

- Cambridge University Press. p. 174–221.
11. Hameed, I. H., Jebor, M. A., and Ommer, A. J., 2016. Haplotype data of mitochondrial DNA coding region encompassing nucleotide positions 11,719–12,184 and evaluate the importance of these positions for forensic genetic purposes in Iraq. *Mitochondrial DNA Part A*. 27(2), 1324–1327.
 12. Huang, M. C., and Lee C. L., 2019. Thermostability of tropomyosins from the fast skeletal muscles of tropical fish species. *Fish Physiology and Biochemistry*. 45, 1189–1202.
 13. Hu, W., Zhang, J., Kang, B., 2016. Structure and function of corneal surface of mudskipper fishes. *Fish Physiology and Biochemistry*. 42, 1481–1489.
 14. Ikram, M. M., Ismail, A., Yap, C. K., and Azwady, A. A. N., 2013. Levels of heavy metals (Zn, Cu, Cd, and Pb) in mudskippers (*Periophthalmus schlosseri*) and sediments collected from intertidal areas at Morib and Remis, Peninsular Malaysia. *Toxicology & Environmental Chemistry*. 92(8), 1471–1486.
 15. Ishimatsu, A., and Graham, J. B., TT. 2011. Mudskippers: front runners in the modern invasion of land. In: Patzner R, Van Tassell JL, Kovac M, Kapoor BG, editors. *The biology of gobies*. Enfield: Science Publishers. p. 603–32.
 16. Kareem, M. A., Abdulzahra, A. I., Hameed, I. H., and Jebor, M. A., 2015. A New Polymorphic Positions Discovered in Mitochondrial DNA Hypervariable Region HVIII From Central and North-Central of Iraq. *Mitochondrial DNA Part A*. 27(5), 3250–3254.
 17. Komori, M. R., Ghosh, S., Takachi, S., Hu, Y., Mizoguchi, T., Koyama, Y., and Kuki, 1998. A Null Lesion in the Rhodopin 3,4-desaturase of Rhodospirillum rubrum Unmask a Cryptic Branch of the Carotenoid Biosynthetic Pathway. *Biochemistry*. 37: 8987–8994.
 18. Kurnia, U., Agus, F., Adimihardja, A., dan Dariah, A. 2006. *Sifat Fisik Tanah dan Metode Analisisnya*. BBSIDP: Banjarmasin.
 19. Lauriano, E. R., Faggio, C., Capillo, G., Spanò, N., Kuciel, M., Aragona, M., Pergolizzi, S., 2018. Immunohistochemical characterization of epidermal dendritic-like cells in giant mudskipper, *Periophthalmus schlosseri*. *Fish & Shellfish Immunology*. 74, 380–385.
 20. Looi, L. J., Aris, A. Z., Haris, H., Yusoff, F. Md., and Hasim, Z., 2016. The levels of mercury, methylmercury and selenium and the selenium health benefit value in grey-eel catfish (*Plotosus canius*) and giant mudskipper (*Periophthalmus schlosseri*) from the Strait of Malacca. *Journal of Chemosphere*. 152, 265–273.
 21. Maggyvin, E. dan Sinuraya, R. K. 2017. Marijuana dan Autisme: Sebuah Literatur Review. *Jurnal Farmaka*. 15 (1): 70–92.
 22. Mahadevan, G., Mohan, K., Vinoth, J., and Ravi, V., 2019. Biotic potential of mucus extracts of giant mudskipper *Periophthalmus schlosseri* (Pallas, 1770) from Pichavaram, southeast coast of India. *The journal of basic and applied zoology*. 80(13).
 23. Martin, K. L., 2013. Theme and variations: amphibious air-breathing intertidal fishes
 24. Martínez, H. L., Agorreta, A., Sánchez, M. T., and Mauro, D. S., 2018. Evidence of positive selection suggests possible role of aquaporins in the water to land transition of mudskippers. *Organisms Diversity and Evolution*. 18(4), 499–414.
 25. Moss, S.M. 1989. Effect of Exogenous Androgens on Growth, Biochemical Composition and Reproduction of the Coot Calm *Mulinia literalis*. Pac SCI.
 26. Mohammed, A., and Imad, H., 2013. Autosomal STR: From locus information to next generation sequencing technology. *Res. J. Biotechnol.* 8(10):92–105.
 27. Murdy, E.O., 1989. A taxonomic revision and cladistic analysis of the Oxudercine gobies (Gobiidae: Oxudercinae). *Rec Aust Mus Suppl* 11: 1–93.
 28. Panda, L., and Mabalirajan, U., 2018. Recent updates on corticosteroid resistance in asthma. *European Medical Journal*. 3(3), 49–57.
 29. Polgar, G., Ghanbarifardi, M., Milli, S., Agorreta, A., Aliabadi, M., Esmaeili, H. R., and Khang, T. F., 2017. Ecomorphological adaptation in three mudskippers (Teleostei: Gobioidei: Gobiidae) from the Persian Gulf and the Gulf of Oman. *Hydrobiologia*. 795(1), 91–111.
 30. Prasad, K., Singh, M. B., and Ryan, H., 2016. Corticosteroids for managing tuberculous meningitis (Review). *European Medical Journal*. 3(3), 49–57.
 31. Rajendran, P., Bharatidhasan, R., dan Sureshkumar, K. 2017. GC-MS Analysis of Phyto-Components in Raw and Treated Sugarcane Juice. *Int.J.Curr.Microbiol.App.Sci.* 6 (7): 51–61.
 32. Rozaliyani, A., Susanto, A. D., Swidarmoko, B., dan Yunus, F. 2011. Mekanisme Resistens Kartikosteroid pada Asma. *Jurnal Respir Indo*. 31 (4): 210–223.
 33. Ridho, M. R., Patrono, E., Sholikah, M., 2019. Food habits of three species of mudskippers in the Musi River Estuary, South Sumatra, Indonesia. *Biodiversitas*. 20(8), 2368–2374.
 34. Salempa, P. dan Muhamarram, H. 2016. *Senyawa Steroid dalam Tumbuhan Bayur*. Badan penerbit UNM: Makasar.
 35. Sari, M., 2012. *Jenis Fauna Hutan Mangrove di Areal PT. Bina Ovipipari Semesta dan Sekitarinya*. Penerbit PT. Bina Ovipipari Semesta: Pontianak.
 36. Sun, J. 2007. D-Limonene: Safety and Clinical Application. *Alternative Medicine Review*. 12 (3): 259–264
 37. Takeda, T., Hayashi, M., Toba, A., Soyano, K., and Ishimatsu, A., 2012. *Ecology of the Australian*

- mudskipper *Periophthalmus minutus*, an amphibious fish inhabiting a mudflat in the highest intertidal zone. *Australian Journal of Zoology*. 57: 312–320.
38. Wu, R. X., Hong, W. S., Zhang, Q. Y., Chen, S. X., 2009. Comparative enzyme activities of the intestinal brush border membranes of the herbivorous mudskipper *Bolcophthalmus pectinirostris* and the carnivorous Chinese black sleeper *Bostrichthys siensis*. *Journal of Applied Ichthyology*. 25, 571–575.
39. Yang, X. X., Gu, W., Liang, L., Yan, H. L., Wang, Y. F., Bi, Q., Zhang, T., Yu, J., and Rao, G. X., 2017.
- Screening for the bioactive constituents of traditional Chinese medicines—progress and challenges. Royal society of chemistry. 7, 3089–3100.
40. You, X. X., Sun, M., Li, J., Bian, C., Chen, J., Yi, Y., Yu, H., and Shi, Q., 2018. Mudskippers and Their Genetic Adaptations to an Amphibious Lifestyle.
41. Zaini, R. G., Brandt, K., Clench, M. R., dan Maitre, C. L. L. 2012. Effect of Bioactive Compounds from Carrots (*Daucus carota L.*), Polyacetylenes, Beta-carotene and Lutein on Human Lymphoid Leukemia Cells. *Medical Chemistry Journal*. 12 (0): 2–13.

Bioactive Compounds Evaluation of the Mudskippers in the Estuarine Area of Musi River, South Sumatera, Indonesia

ORIGINALITY REPORT

11 %

SIMILARITY INDEX

MATCH ALL SOURCES (ONLY SELECTED SOURCE PRINTED)

★Yalkowsky, . "Solubility Data", Handbook of Aqueous Solubility Data, 1%
2003.

[Crossref](#)

EXCLUDE QUOTES

ON

EXCLUDE MATCHES

< 1%

EXCLUDE

ON

BIBLIOGRAPHY

FORMAT PENILAIAN (VALIDASI & PEER REVIEW)
LEMBAR
HASIL PENILAIAN SEJAWAT SEBIDANG ATAU PEER REVIEW
KARYA ILMIAH : JURNAL ILMIAH

Jurnal Artikel Ilmiah	: Bioactive Compounds Evaluation of the Mudskippers in the Estuarine Area of Musi River, South Sumatera, Indonesia
Penulis Artikel Ilmiah	: Arum Setiawan
Identitas Jurnal Artikel Ilmiah	: a. Nama Jurnal : Journal of Ecological Engineering b. Nomor/Volume/Hal : 3/21/70-80 c. Edisi (bulan/tahun) : April 2020 d. Penerbit : Polskie Towarzystwo Inżynierii Ekologicznej e. Jumlah Halaman 11
Kategori Publikasi Jurnal Ilmiah (beri ✓ pada kategori yang tepat)	: <input checked="" type="checkbox"/> Jurnal Ilmiah Internasional Bereputasi <input type="checkbox"/> Jurnal Ilmiah Internasional <input type="checkbox"/> Jurnal Ilmiah Nasional Terakreditasi S1, S2 <input type="checkbox"/> Jurnal Ilmiah Nasional Terakreditasi S3, S4 <input type="checkbox"/> Jurnal Ilmiah Nasional Tidak Terakreditasi

I. Hasil Penilaian Validasi :

No.	ASPEK	URAIAN/KOMENTAR PENILAIAN
1.	Indikasi Plagiasi	11 %
2.	Linearitas	Paper kurang linier dengan bidang biologi konservasi

II. Hasil Penilaian Peer Review :

Komponen Yang Dinilai	Nilai Maksimal Jurnal Ilmiah (isikan di kolom yang sesuai)					Nilai Akhir Yang Diperoleh
	Internasional Bereputasi (Maks 40)	Internasional (Maks 20)	Nasional Terakreditasi S1, S2 Maks 25	Nasional Terakreditasi S3, S4 Maks 20	Nasional tidak Terakreditasi (maks 10)	
Kelengkapan dan Kesesuaian unsur isi jurnal (10%)	4					3
Ruang lingkup dan kedalaman pembahasan (30%)	12					10
Kecukupan dan Kemutahiran data/informasi dan metodologi (30%)	12					12
Kelengkapan unsur dan kualitas penerbit (30%)	12					12
Total = (100%)	40					37
Kontribusi Pengusul (Penulis Pertama /Anggota Utama)	Anggota Utama (0,4x37)/5= 2,96					2,96

KOMENTAR/ULASAN PEER REVIEW

• Kelengkapan dan Kesesuaian Unsur:	Paper terkait senyawa bioactive dr ikan glodok di sungai Musi. Isi paper sudah memenuhi kaidah-kaidah karya ilmiah namun kurang sesuai dengan bidang biologi konservasi
• Ruang Lingkup dan Kedalaman Pembahasan:	Hasil penelitian dibahas cukup komprehensif dengan penyampaian pembanding dari temuan-temuan penelitian lainnya dan teori terkait. Referensi yang diacu dalam pembahasan sudah cukup update untuk bidang kajian ini.
• Kecukupan & Kemutahiran Data & Metodologi:	Data-data hasil penelitian sudah baik dan didukung peta lokasi sampling dan gambar yang ditampilkan menarik. Data didapatkan dengan menggunakan metode yang sudah standard.
• Kelengkapan Unsur & Kualitas Penerbit:	Penerbit Polskie Towarzystwo Inżynierii Ekologicznej berkualitas baik, tidak termasuk predatory publisher, dan jurnal masuk di Q3.

Surabaya, 15 Mei 2020
Penilai 1



Prof. Hery Purnobasuki, M.Si., Ph.D.
NIP 196705071991021001
Unit Kerja : Jurusan Biologi FST Unair
Bidang Ilmu : Biologi
Jabatan/Pangkat : Guru Besar/ Pembina Utama Madya

FORMAT PENILAIAN (VALIDASI & PEER REVIEW)
LEMBAR HASIL PENILAIAN SEJAWAT SEBIDANG ATAU PEER REVIEW
KARYA ILMIAH : JURNAL ILMIAH

Jurnal Artikel Ilmiah	: Bioactive Compounds Evaluation of the Mudskippers in the Estuarine Area of Musi River, South Sumatera, Indonesia
Penulis Artikel Ilmiah	: Arum Setiawan
Identitas Jurnal Artikel Ilmiah	: <ul style="list-style-type: none"> a. Nama Jurnal : Journal of Ecological Engineering b. Nomor/Volume/Hal : 3/21/70-80 c. Edisi (bulan/tahun) : April 2020 d. Penerbit : Polskie Towarzystwo Inżynierii Ekologicznej e. Jumlah Halaman : 11
Kategori Publikasi Jurnal Ilmiah (beri V pada kategori yang tepat)	: <ul style="list-style-type: none"> <input checked="" type="checkbox"/> Jurnal Ilmiah Internasional Bereputasi <input type="checkbox"/> Jurnal Ilmiah Internasional <input type="checkbox"/> Jurnal Ilmiah Nasional Terakreditasi S1, S2 <input type="checkbox"/> Jurnal Ilmiah Nasional Terakreditasi S3, S4 <input type="checkbox"/> Jurnal Ilmiah Nasional Tidak Terakreditasi

I. Hasil Penilaian Validasi :

No.	ASPEK	URAIAN/KOMENTAR PENILAIAN
1.	Indikasi Plagiasi	11 %
2.	Linearitas	

II. Hasil Penilaian Peer Review :

Komponen Yang Dinilai	Nilai Maksimal Jurnal Ilmiah (isikan di kolom yang sesuai)					Nilai Akhir Yang Diperoleh
	Internasional Bereputasi (Maks 40)	Internasional (Maks 20)	Nasional Terakreditasi S1, S2 Maks 25	Nasional Terakreditasi S3, S4 Maks 20	Nasional tidak Terakreditasi (maks 10)	
Kelengkapan dan Kesesuaian unsur isi jurnal (10%)	4					3
Ruang lingkup dan kedalaman pembahasan (30%)	12					10
Kecukupan dan Kemutakhiran data/informasi dan metodologi (30%)	12					10
Kelengkapan unsur dan kualitas penerbit (30%)	12					12
Total = (100%)	40					35
Kontribusi Pengusul (Penulis Utama/ Anggota Utama)	Journal of Ecological Engineering, SJR 0,31, IF 3,032. Pengusul (Penulis ke 2 dari 6). Nilai maksimal 87,5%. Nilai pengusul : $(0,4 \times 0,875 \times 40)/5 = 2,8$					2,8
KOMENTAR/ULASAN PEER REVIEW						
• Kelengkapan dan Kesesuaian Unsur:	Format tidak semuanya dipenuhi. Sistem penulisan author berbeda antara dalam Narasi dengan Referensi. Ada author dalam Narasi tetapi tidak ada dalam Referensi dan sebaliknya.					
• Ruang Lingkup dan Kedalaman Pembahasan:	Ruang lingkup masih terkait dengan tugas di Departemen Biologi. Tujuan tidak jelas, kesimpulan kurang tegas. Pembahasan sudah cukup walau tidak mendalam.					
• Kecukupan & Kemutakhiran Data & Metodologi:	Metode sudah biasa dilakukan peneliti. Data untuk penelitian ini termasuk baru. Cara penulisan Rumus Molekul tidak baku.					
• Kelengkapan Unsur & Kualitas Penerbit:	Jurnal sudah memenuhi kriteria. Penerbit cukup bagus.					

Yogyakarta, 1 Juni 2020

Penilai 2

tanda tangan

Prof. Dr. Suwarno Hadisusanto

NIP 195411161983031002

Unit Kerja : Fakultas Biologi UGM

Bidang Ilmu : Biologi/Ekologi

Jabatan/Pangkat : Guru Besar/ Pembina Utama Madya