

The distribution and composition of Odonata (Dragonfly and Damselfly) in Sriwijaya University, Inderalaya Campus South Sumatera

by Dr.ir.yulia Pujiastuti, Ms.

Submission date: 08-May-2020 03:06PM (UTC+0700)

Submission ID: 1319278948

File name: 13._Biol_research_Yp.pdf (901.7K)

Word count: 2552

Character count: 13819

The distribution and composition of Odonata (Dragonfly and Damselfly) in Sriwijaya University, Inderalaya Campus South Sumatera

Yu. Pujiastuti^{1*}, Yuanita Windusari², Muhammad Agus³

¹Department of Plant Protection, Faculty of Agriculture, Sriwijaya University

²Department of Biology, Faculty of Mathematics and Natural Sciences, Sriwijaya University

³Department of Environmental Management, Graduate School of Sriwijaya University

Abstract

The information about distribution and composition of Odonata in Sriwijaya University campus area is still not much. The decrease in areas supporting the growth of Odonata influence distribution and composition of these species. The objective of research was to analyze the distribution and composition of Odonata in Sriwijaya University, Inderalaya Campus. This research had been conducted from December 2016 until February 2017 and continued in April 2017 by using visual observation, direct capture, and sticky traps. The sampling locations were determined by five points of observation station by using purposive sampling method. The result revealed Odonata living in Sriwijaya University, Inderalaya Campus is aggregated and consisted of 22 species belonged to five families. The highest composition of species was found in Libellulidae (77.65%) followed by Coenagrionidae (17.86%).

Keywords: composition, distribution, Odonata, Sriwijaya University.

Received: 16 June 2017 Revised: 21 August 2017 Accepted: 02 September 2017

Introduction

Sriwijaya University campus area consists of buildings, swamp areas, arboretum, agriculture land, oil palm plantation, green spaces and retention ponds. This campus area has high diversity of flora and fauna due to the variety of land conditions. Odonata is one of important components of this ecosystem. The environmental quality affects existence of Odonata (Suhonen *et al.*, 2010).

Odonata is one of insects which play a role as environmental contamination indicator. Fresh environment can be seen from the presence of a living Odonata species in the area because larvae phase of Odonata species live only in undisturbed water conditions (Berquier *et al.*, 2016). The loss of Odonata indicates the environmental disturbance, especially the waters of the area.

Odonata is one of the insects with high mobility. Its ability to fly and move from one place to another place in short time bring through a suitable place that has appropriate environmental conditions. The superiority of the dragonfly lies on the vision and ability of flying. Odonata is able to see in 360-degree angles and they are very sensitive to movement, leading these insects as a reliable predator (Sigit *et al.*, 2013).

Diversity of Odonata will be high in areas where water sources and habitat conditions are still in the natural or uncontaminated category (Siregar and Bakti, 2016; Suhonen *et al.*, 2010; Khan, 2015). The existence of pool, flooded area, and heterogeneous vegetated location will support high diversity and the abundance of Odonata.

Good environmental conditions are indicated by the pattern of Odonata distribution. Data about odonata in Sriwijaya University Inderalaya Campus is very rare, therefore observation of distribution of these insects is important to be conducted.

Method

This study was conducted from December 2016 until February 2017 and continued in April 2017. Data collection was performed at Sriwijaya University Inderalaya Campus area, Ogan Ilir District, South Sumatera Province. The sampling location was determined by five points of observation station. Observation of Odonata was done by purposive sampling method (Figure 1).

Visual observations were performed at each location with a frequency of once a week. There was 5 locations of observation considered as replications. Observation was performed purposely to find perch Odonata. They were well documented using camera photo. These photographs were useful to identify Odonata. The observation time was done at 7-10 AM and 2-6 PM since they were active during these time.

The capture of the odonata was directly carried out by using an insect net, a cone-shaped net with a height of 60 cm, a diameter of 30 cm and a net length of 1 meter. The sticky trap to capture Odonata was made of bamboo with a length of 1.5 m and each bamboo was smeared with glue. Installation of sticky trap was conducted once a week and placed one tap at each location (Figure 2).

Odonata captured was identified by Theischinger and Hawking (2006); Paulson, (2009); Sigit *et al.* (2013), and Setiyono *et al.* (2015). Distribution pattern was analyzed by Morisita index (Id) (Krebs, 1999) (Equations 1)

$$Id = \frac{n \sum x_i (x_i - 1)}{N (N - 1)} \quad (1)$$

* Corresponding Author:

Yu. Pujiastuti

Department of Plant Protection, Faculty of Agriculture,
Sriwijaya University

Phone: 0711-580663 Fax: 0711-580276

e-mail: ypujiastuti@unsri.ac.id

The magnitude of the distribution Morisita index as follows:
 value $I_d < 1$ indicate distribution of species on area is uniform,
 value $I_d = 1$ indicate distribution of species on area is random and
 value $I_d > 1$ indicate distribution of species on area is aggregated (Krebs, 1999).




Figure 1. Map of research location on Sriwijaya University Campus, Indralaya




Figure 2. Sticky Trap to capture Odonata




Figure 3. Member of Odonata order at Sriwijaya University campus area (a) *Acisoma panorhoides* (b) *Anax guttatus* (c) *Brachythemis contaminata* (d) *Brachydiplax chalybea* (e) *Crocothemis servilia* (f) *Diplacodes nebulosa* (g) *Diplacodes trivalis* (h) *Ictagomphus decatus* (i) *Lathrecista asiatica* (j) *Nannophya pygmaena* (k) *Neurothemis ramburii* (l) *Orthetrum pruinosum* (m) *Orthetrum sabina* (n) *Rhodothemis rufa* (o) *Rhyothemis phyllis* (p) *Trithemis pallidinervis* (q) *Agriocnemis femina* (r) *Agriocnemis pygmaena* (s) *Ceriagrion auranticum* (t) *Ischnura senegalensis* (u) *Lestes praemorsus* (v) *Pseudagrion microcephala*

Results

Based on observation on Odonata by using visual observation method, direct capture and sticky trap from December 2016 until February 2017 and continued at April 2017 was found 22 species (Table 1 & Figure 3).

These species were belonged to three families of Sub-Order Anisoptera namely Aeshidae, Gomphidae, and Libellulidae, and two families of Sub-Order Zygoptera namely such as Coenagrionidae and Lestidae.

Table 1. Distribution index of Odonata at Sriwijaya University campus area.

No	Species	December '16			January 2017			February 2017			April 2017		
		Xi	Id	D	Xi	Id	D	Xi	Id	D	Xi	Id	D
Anisoptera													
Aeshnidae													
1	<i>Anax guttatus</i>	-	-	-	2	0	-	-	-	-	-	-	-
Gomphidae													
2	<i>Ictinogomphus decatus</i>	-	-	-	24	1.54	A	13	2.95	A	8	1.96	A
Libellulidae													
3	<i>Acisoma panorphoides</i>	3	5.00	A	18	4.44	A	4	5.00	A	9	3.06	A
4	<i>Brachythemis contaminata</i>	104	2.11	A	175	1.75	A	231	1.79	A	160	4.07	A
5	<i>Brachydiplax chalybea</i>	-	-	-	-	-	-	-	-	-	2	0	-
6	<i>Diplacodes nebulosa</i>	-	-	-	-	-	-	-	-	-	1	0	-
7	<i>Diplacodes trivalis</i>	5	1.00	R	8	5.00	A	10	2.12	A	6	5.00	A
8	<i>Lathrecista asiatica</i>	-	-	-	2	5.00	-	-	-	-	-	-	-
9	<i>Neurothemis ramburii</i>	46	1.00	R	94	1.78	A	47	1.44	A	21	0.95	U
10	<i>Nannophya pygmaena</i>	-	-	-	-	-	-	-	-	-	2	5.00	A
11	<i>Trithemis pallidinervis</i>	14	5.00	A	21	5.00	A	8	5.00	A	-	-	-
12	<i>Orthetrum pruinosum</i>	-	-	-	11	2.82	A	-	-	-	2	5.00	A
13	<i>Orthetrum sabina</i>	70	1.92	A	81	1.43	A	72	1.45	A	36	2.02	A
14	<i>Rhyothemis phyllis</i>	50	3.72	A	60	3.52	A	85	3.49	A	21	3.10	A
15	<i>Crocothemis servilia</i>	15	1.90	A	23	2.51	A	23	2.31	A	9	1.39	A
16	<i>Rhodothemis rufa</i>	-	-	-	1	0	-	-	-	-	-	-	-
Zygoptera													
Coenagrionidae													
17	<i>Agriocnemis femina</i>	8	2.32	A	12	2.05	A	3	1.67	A	-	-	-
18	<i>Agriocnemis pygmaena</i>	7	3.57	A	36	1.88	A	5	1.00	A	6	0.67	U
19	<i>Ceriagrion auranticum</i>	-	-	-	1	0	-	1	-	-	3	5.00	A
20	<i>Ischnura senegalensis</i>	12	1.59	A	36	1.40	A	24	1.85	A	8	3.75	A
21	<i>Pseudagrion microcephalum</i>	25	2.45	A	65	1.57	A	58	1.37	A	13	2.50	A
Lestidae													
22	<i>Lestes praemorsus</i>	-	-	-	15	3.76	A	18	5.00	A	10	5.00	A

Species found in sub-Order of Anisoptera were *Acisoma panorphoides*, *Anax guttatus*, *Brachythemis contaminata*, *Brachydiplax chalybea*, *Crocothemis servilia*, *Diplacodes nebulosa*, *D. trivalis*, *Ictinogomphus decorates*, *Lathrecista asiatica*, *Nanophya pygmaea*,

Neurothemis ramburii, *Orthetrum pruinosum*, *O. Sabina*, *Rhodothemis rufa*, *Rhyothemis Phyllis*, *Trithemis pallidinervis*. Species that were found in sub-Order of Zygoptera were *Agriocnemis femina*, *A. pygmaena*, *Ceriagrion latericum*, *Ischnura senegalensis*, *Lestes*

praemorsus, *Pseudagrion microcephalum* (Table 1 & Figure 3)

The distribution patterns of the three most commonly encountered species with the largest number of individuals from December 2016 to April 2017 were shown in Figure 4. Based on Figure 4, it was shown that

the distribution patterns was aggregated. Species of *B. contaminata* was always found in every location and time of observations, except in April 2017. *B. contaminata* was highly aggregated in the fifth location and not visible in the second and third locations. It was predicted that the fifth location was lack of food sources.




Figure 4. Distribution of *B. contaminata*, *O. sabina* dan *N. ramburii* pada bulan (a) Desember 2016 (b) Januari 2017 (c) Februari 2017 dan (d) April 2017

Discussions

The population of a species, when it reached the high level of density, and limitation of environmental carrying capacity, will experience spreading. Behavior to spread and form a particular pattern could be caused by the drive for food, avoidance of predators, climatic influences, marriage habits and other physical factors (Odum, 1998). The dispersion pattern of a species will follow good environmental conditions (Suhonen *et al.*, 2010 and Khan, 2015). Species spread to search typical food sources and conditions for appropriate activities (Alikodra, 2010; Indrawan *et al.*, 2007)

Based on measurement of abiotic conditions, campus area of Sriwijaya University was categorized as having clean water condition. It might be related to the condition of the existence of living things, especially Odonata order in the region. The areas with **unclean waters** would remove species from Odonata (Suhonen *et al.*, 2010). The olfactory sensory organs present in the antenna of Odonata will detect the condition of a watery area capable of supporting the life of the eggs to be placed. Therefore, Odonata will move from a dirty habitat to a clean habitat (Setiyono *et al.*, 2015).

Species of *B. contaminata* was aggregated at the fifth location in April 2017. It was predicted that the fifth location was a place for *B. contaminata* to find partners.

Animal aggregation was behavior in searching of spouses and other social behaviors (Campbell *et al.* 2003). Furthermore, Harrington and Stork (1995) explained that distribution of an animal can be caused by many factors such as changes in temperature, humidity, food sources, or the presence of interference.

Species of *B. contaminata*, *O. sabina* and *N. ramburii* which were part of sub-order Anisoptera found in Sriwijaya University campus area possessed a larger number of species when compared with sub-order Zygoptera. This might be happened due to the limited ability to spread of this sub-order (Harisah, 2016). Zygoptera was very sensitive to environmental conditions (Pamungkas and Ridwan, 2015) and preferred vegetated and shaded areas (Khan, 2015).

The fourth location was predicted as the most stable condition. It was showed number of *B. contaminata*, *O. sabina* and *N. ramburii* which were approximately similar and no dominating species (Figure 4). A stable condition of an ecosystem can create harmony condition among predators, preys and the environment. The preservation of population numbers would be able to maintain the diversity and evenness of species in an area. Balanced ecosystem conditions will reflect good environmental conditions.

The highest population growth in April 2017 was *B. contaminata*. Species of *B. contaminata* was feared to be uncontrollable and will lead to overpopulation. This condition will caused a negative effect, due to the increasingly tight competition. Excessive population can cause weak species will move to another place and the worse impact was extinction of species.

The existence of the Odonata could be influenced by natural enemy factors (Kalkman *et al.*, 2010). The natural enemies of the Odonata Order were birds, frogs, lizards,

larger carnivorous insects, even fellow Odonata (Paulson, 2009). The amount of *O. sabina* and *N. ramburii* was not as much as *B. contaminata*, but its presence can be found at each study site. The ability to fly of *O. sabina* and *N. ramburii* will lead to their survival. Species of *O. sabina* and *N. ramburii* were common species. They can be found every year, with high tolerance, and wide distribution (Kosterin, 2012; Pamungkas and Ridwan, 2015)

References

- Acharjee, B., and Karzee, L. 2016. A Checklist of Dragonfly (Odonata: Anisoptera) Diversity in The Campus of University of Science and Technology, Meghalaya (USTM), RI Bhoi District, Meghalaya, India. *Journal of Entomology and Zoology Studies*. 4:124-127.
- 1 Alikodra, H. 2010. *Teknik Pengelolaan Satwa liar, dalam Rangka Mempertahankan Keanekaragaman Hayati Indonesia*. IPB Press. Bogor: 68. (In Indonesian)
- Berquier, C., Orsini, A., and Andrei-Ruiz, M. 2016. Odonata Community Index – Corsica (OCIC): a New Biological Index Based on Adult Odonata Populations for Assessment of The Ecological Status of Watercourses in Corsica. *Journal Ecological Indicator*. 66:163-172.
- 1 Bismak, M. 2011. *Prosedur Operasi Standar (SOP) Untuk Survei Keragaman Jenis Pada Kawasan Konservasi*. Badan Penelitian dan Pengembangan Kehutanan. Bogor. (In Indonesian)
- Campbell, N., Reece., and Mitchell. 2003. *Biologi Edisi Kelima, Jilid II*. Erlangga. Jakarta. (In Indonesian)
- 1 Harington, N. dan Stork, N. 1995. *Insect in a Changing Environment*. Academic Press. New York.
- Indrawan, M., Primack, R., dan Supriatna, J. *Biologi Konservasi*. 2007. Yayasan Obor Indonesia. Jakarta: 626. (In Indonesian)
- Khan, M. 2015. Dragonflies and Damselflies (Insect: Odonata) of The Northeastern Region of Bangladesh With Five New Additions to The Odonata Fauna of Bangladesh. *Journal of Threatened Taxa*. 7: 7795-7804.
- Krebs, C. 1999. *Ecology Methodology, Second Edition*. An Imprint of Addison Wesley Longman, Inc. California.
- 5 Kosterin, O. 2012. A rapid Survey of Odonata on Bokor Olateau, Preah Monivong National Park, Cambodia. *Cambodian Journal of Natural History*. 1:75-86.
- 1 Odum, E. 1998. *Dasar-dasar Ekologi, Edisi Ketiga*. Gadjah Mada University Press. Yogyakarta. (In Indonesian)
- Pamungkas, D., and Ridwan, M. 2015. Keragaman Jenis Capung dan Capung Jarum (Odonata) di Beberapa Sumber Air di Magetan, Jawa Timur. *Prosiding Seminar Nasional Masyarakat Biodiveristas Indonesia*. 1:1295-1301. (In Indonesian)
- 1 Paulson, D. 2009. *Dragonflies and Damselflies of The West*. Princeton University Press. United Kingdom.
- 1 Setiyono, J., Siti, D., Husaini., Evi, N., Wahyu, S., and Nanang, K. 2015. *Keanekaragaman Capung, Kupu-kupu dan Burung Pegunungan Karst Kendeng Pati Jawa Tengah*. Sheep Indonesia Foundation. Yogyakarta. (In Indonesian)
- 1 Sigit, W., Bambang, F., Magdalena, P., Bernadeta, P., and Tabita, M. 2013. Naga Terbang Wendit, *Keanekaragaman Capung Perairan Wendit, Malang, Jawa Timur*. Indonesia Dragonfly Society. Malang. (In Indonesian)
- Siregar, A. and Bakti, D. 2016. Diversity and Distribution Of Odonata In University Sumatera Utara, Medan, Indonesia. *International Journal Of Scientific and Technology Research*. 24: 229-234.
- 1 Suhonen, J., Milla, H., Esa, K., Markku, K., Johanna, K., Jouni, P. and Jukka, S. 2010. Local Extinction of Dragonfly and Damselfly Populations in Low and High Quality Habitat Patches. *Journal of Conservation Biology*, 24: 1148-1153.
- 8 Theischinger, G. and Hawking, 2006. *The Complete Field Guide to Dragonflies of Australia*. CSIRO Publishing. Australia.

The distribution and composition of Odonata (Dragonfly and Damselfly) in Sriwijaya University, Inderalaya Campus South Sumatera

ORIGINALITY REPORT

19%

SIMILARITY INDEX

18%

INTERNET SOURCES

5%

PUBLICATIONS

9%

STUDENT PAPERS

PRIMARY SOURCES

1	sciencetechindonesia.com Internet Source	12%
2	eprints.umm.ac.id Internet Source	2%
3	Submitted to iGroup Student Paper	1%
4	Yulia Pujiastuti, Siti Masyitah, Sarah Dirgahayu, Suparman Surya Hadikusuma, Effendy .. "THE USE OF GOLDEN SNAIL MEAL TO ENRICH BACILLUS THURINGIENSIS CULTURE MEDIA AND ITS EFFECT ON THE BACTERIAL TOXICITY AGAINST SPODOPTERA LITURA", JURNAL HAMA DAN PENYAKIT TUMBUHAN TROPIKA, 2018 Publication	1%
5	O. E. Kosterin. "Occasional photographic records of butterflies (Lepidoptera, Papilionoidea) in Cambodia. 1. The coastal	1%

Cardamom foothills (SW Cambodia), 2010-2018", Acta Biologica Sibirica, 2019

Publication

6	mafiadoc.com Internet Source	1%
7	repository.unair.ac.id Internet Source	1%
8	www.journals.uchicago.edu Internet Source	1%
9	Fauziyah, Nurhayati, S M Bernas, A Putera, Y Suteja, F Agustiani. "Biodiversity of fish resources in Sungsang Estuaries of South Sumatra", IOP Conference Series: Earth and Environmental Science, 2019 Publication	1%

Exclude quotes On
Exclude bibliography Off

Exclude matches < 1%