

PAPER • OPEN ACCESS

Serum C-Reactive Protein in chronic kidney disease patients undergoing hemodialysis and correlation with dialytic age

To cite this article: Zulkhair Ali *et al* 2019 *J. Phys.: Conf. Ser.* **1246** 012002

View the [article online](#) for updates and enhancements.



240th ECS Meeting ORLANDO, FL

Orange County Convention Center Oct 10-14, 2021



Abstract submission due: April 9

SUBMIT NOW

Serum C-Reactive Protein in chronic kidney disease patients undergoing hemodialysis and correlation with dialytic age

Zulkhair Ali¹, Muhammad R Ridha², Erial Bahar³

¹ Division of Nephrology and Hypertension, Department of Internal Medicine, Faculty of Medicine, Universitas Sriwijaya, Mohammad Hoesin Hospital, Palembang, Indonesia

² Department of Internal Medicine, Faculty of Medicine Universitas Sriwijaya – Mohammad Hoesin Hospital, Palembang, Indonesia

³ Department of Anatomy, Faculty of Medicine, Universitas Sriwijaya, Palembang, Indonesia

E-mail: riset.drzulkhair@gmail.com

Abstract. Mortality in CKD patients undergoing hemodialysis considerably high. Previous study showed that mortality in patients undergoing hemodialysis was strongly correlated with chronic inflammation. The current study examined whether the serum c-reactive protein concentration (hsCRP) changed from pre- to post-dialysis. In addition, the current study also investigated the correlation between dialysis age and hsCRP in chronic kidney disease patients undergoing hemodialysis. The study was a case-series study of 30 patients with hemodialysis. The patients were divided into three groups based on dialysis age. Hemodialysis had been undergone for 4 hours, using *hemoflow F8HPS*, polysulfone synthetic dialysis membrane, and bicarbonate dialysis fluid. The primary outcome was serum concentration of high-sensitivity C-reactive protein (hsCRP), which was assessed in the pre-and-post dialysis, measured using *chemiluminescent method*. Dialysis does increase the hsCRP post-dialysis. It resulted in significant ($p < 0.05$) increment of serum hsCRP concentration, pre-dialysis 1.04 ± 0.90 mg/L compared to post-dialysis mean value 1.84 ± 2.01 mg/dL. There was no significant difference among serum hsCRP concentration mean value among groups of dialysis age. The correlations between various measurements of serum hsCRP concentration and dialysis age were statistically insignificant.

1. Introduction

Chronic kidney disease (CKD) is a current global health issue with estimated global prevalence between 11 to 13% [1]. CKD leads to a gradual decline of kidney function, with renal replacement therapy eventually demanded to substitute the function of the kidneys. The rate of decline in kidney function differs among patients, and the initiation of renal replacement therapy relies on the presence of symptoms related to kidney failure [2]. A decline in the glomerular filtration rate to less than 30 mL/min/1.73 m² is associated with severely decreased kidney function, described as stage four CKD, and preparation for renal replacement therapy is recommended [3-4]. The modalities of renal replacement therapy comprise transplantation, peritoneal dialysis, and hemodialysis, with 85% of patients with terminal stage of CKD in 53 countries undergoing hemodialysis [5].

Mortality among patients with hemodialysis is considerably high, in part due to the cardiovascular complications [6]. Several studies suggest that cardiovascular-related mortality among patients with hemodialysis was associated with chronic inflammation [7-9]. Inflammation during hemodialysis may



occur in some manners. Bio-incompatibility between dialyzer and blood, the endotoxin in dialysis fluid, access-related infections, and the glucose degradation products have been contributed to the inflammation responses during hemodialysis [10]. A study reports that about 35–65% of CKD patients undergoing haemodialysis exhibit the signs of inflammation, considering the prevalence in patients before dialysis is lower [11].

High-sensitivity C-reactive protein (hsCRP) is a well-known marker of systemic inflammation. Some epidemiologic studies have revealed that hsCRP may independently predicts vascular risk and may provide prognostic information [12-13]. In contrast to several other parameters that also constitute biological aspects of inflammation, hsCRP measurement is standardized and commonly available [14]. A study reports that postdialysis increment of hsCRP is associated with increased mortality among patients with hemodialysis [15]. The aim of current study was to determine the effect of hemodialysis on pre-to-post dialysis changes of serum c-reactive protein concentration (hsCRP), a well-known marker of systemic inflammation. In addition, the current study also identified the correlation between dialysis age and serum hsCRP concentration among chronic kidney disease patients undergoing hemodialysis.

2. Methods

The subjects were consecutively recruited from chronic kidney disease (CKD) patients who regularly attended the session of hemodialysis in Hemodialysis Unit of Mohammad Hoesin General Hospital Palembang. The prospective subjects were 25-60 years old and having hemodialysis session for at least 2 times. Patients with liver function disorders, lupus nephritis, and acute infection manifestations were excluded from the study. Patients were selected from a pool of 108 patients.

Dialysis age was determined by clinical records. Dialysis age was group into less than three months, between three and twelve months, and more than one year. Prior to hemodialysis, blood was taken via venipuncture to measure complete blood count and hsCRP. hsCRP was also measured four hours after hemodialysis. Hemodialysis had been undergone for 4 hours, using *hemoflow F8HPS*, polysulfone synthetic dialysis membrane, and bicarbonate dialysis fluid (all products were Nipro's, Akita, Japan). hsCRP was measured based on a blood specimens and was determined by *chemiluminescent method* (hsCRP, Cat.No,LKCRP1, Siemens Diagnostics).

Normal distribution of data was deduced with Shapiro-Wilk test. Kruskal Wallis non-parametric test and one-way ANOVA were used to compare data among more than two groups. Pearson correlation test was used to determine the correlation between variable. P-values of less than 0.05 were considered to be significant. Statistical analysis was performed using *SPSS 22.0 for Windows*.

3. Results

There were 30 patients recruited in total for the analysis of the study. General characteristics of patients were shown in Table 1. Mean of the patients' body mass index was normoweight range. Most of the patients were between 51-years-old and 60-years-old. There were 17 females and 13 males. The commonest etiology was hypertension. Hemoglobin, serum ureum, serum creatinine, random blood glucose, serum natrium, serum potassium, serum albumin, SGOT, and SGPT were normally distributed.

The measurements of hsCRP were shown in table 2. Hemodialysis resulted in significant ($p < 0.05$) increment of serum hsCRP concentration, pre-dialysis 1.04 ± 0.90 mg/L compared to post-dialysis mean value 1.84 ± 2.01 mg/dL. There was no significant difference among serum hsCRP concentration mean value among group of dialysis age.

The correlations between various measurement of serum hsCRP concentration and dialysis age were shown in table 3. Prehemodialytic, posthemodialytic, and pre-to-post hemodialytic hsCRP value were having insignificant negative correlation toward dialytic age. Pre-to-post hemodialytic hsCRP changes percentage was having insignificant weak positive correlation toward dialytic age.

4. Discussions

CRP has been popular as one of acute phase proteins and used as a biomarker of inflammation in clinical settings. CRP is synthesized in liver, mostly induced by interleukin-6. It is produced during acute phase response to infection or other inflammatory episodes [16]. An in vitro study showed that CRP activates the expression of adhesion molecules in human monocytes that involved in

inflammation [17]. CRP also contributes in scavenge processes of cell debris, conducting classical pathway of complement system [18]. The measurement of hsCRP has been suggested to be important biomarker in the initiation and progression of atherosclerosis [19]. Some studies have revealed that chronic hsCRP elevation of the vascular endothelium has a pivotal role in the development of cerebral and cardiac vascular disease, mainly in patients with chronic hemodialysis [20-22]. Since cerebral and cardiac vascular disease were main cause of death in patients with hemodialysis, hsCRP have been established as a useful tool in predicting mortality in hemodialysis.

Table 1. General characteristics of the subjects.

Characteristics	Total (n=30)	Median Value (min-max)	Mean±SD
Age (years)			
≤30	3(10%)		
31 - 40	6(20%)		
41 - 50	9(30%)		
51 - 60	12(40%)		
Sex			
Male	13(43.3%)		
Female	17(56.7%)		
Body Mass Index (kg/m²)			21.79±3.51
Duration on HD (months)		10(1-58)	
<3	10(33.3%)		
3-12	10(33.3%)		
>12	10(33.3%)		
Cause of HD			
Hypertension	19(63.3%)		
Diabetes Mellitus	5(16.7%)		
Renal Infection	3(10%)		
Gout Nephropathy	1(3.33%)		
Renal Cyst	1(3.33%)		
Obstructive Nephropathy	1(3.33%)		
Hemoglobin (g/dL)			8.27±1.14
Leucocytes (cell/mm ³) ^a		7150(4200-10500)	
Trombocytes (/mm ³) ^a		185000(111000-431000)	
Serum Ureum (mg/dL)			140.83±52.36
Serum Creatinine (mg/dL)			11.81±4.15
Random Blood Glucose (mg/dL)			130.23±64.29
Serum Sodium (mEq/L) ^a		143(133-149)	
Serum Potassium (mEq/L)			4.69±0.73
Serum Albumin (mg/dL)			3.85±0.42
SGOT (μg/L)			22±11.01
SGPT (μg/L)			23.47±15.21

^a Shapiro-Wilk test for normality <0.05

Table 2. Serum hsCRP concentration profile in subjects.

Duration on HD	n (%)	Prehemodialysis hsCRP		Posthemodialysis hsCRP	
		Mean(mg/dL)	p value	Mean(mg/dL)	p value
<3 months	10(33.%)	1.09±0.49		1.75±1.07	
3-12 months	10(33.%)	1.16±1.25	0.224 ^a	1.60±1.30	0.254 ^a
>12 months	10(33.%)	0.85±0.89		2.17±3.15	
All Subjects	30(100%)	1.04±0.90		1.84±2.01	0.008 ^b

^aKruskal-Wallis Test among three group, ^bDependent t-test for pre-to-post hemodialysis mean value

In current study, there was significant increment in four hours postdialysis serum hsCRP concentration compared to predialysis value. The similar finding was also found in other study conducted by Nagane et al [23], in spite of the increment of hsCRP was relatively small compared to the study conducted in European [24]. Previous study showed that the prevalence of chronic inflammation in American and European patients were considerably higher than Asians [24]. The genetic factors was hypothesized to be taken the role.

The current study showed that there was no significant correlation between dialysis age and serum hsCRP concentration among chronic kidney disease patients undergoing hemodialysis. This finding may relate to the theory that CRP concentrated optimally after 24 hours [25]. Serial monitoring of CRP may be used to in determining the magnitude of inflammation in patients undergoing hemodialysis.

Table 3. The correlation between dialysis age and various measurements values of hsCRP.

Measurement Values of hsCRP	Duration on Hemodialysis	
	r	p
Prehemodialytic hsCRP	-0.227 ^a	0.228
Posthemodialytic hsCRP	-0.134 ^a	0.481
Pre-to-post hemodialytic hsCRP changes	-0.059 ^a	0.769
Pre-to-post hemodialytic hsCRP changes percentage	0.029 ^a	0.945

^aPearson Correlation Test

5. Conclusion

In conclusion, dialysis increased the hsCRP post-dialysis significantly. The current study shows that there was no significant correlation between dialysis age and hsCRP.

6. References

- [1] Hill NR, Fatoba ST, Oke JL, Hirst JA, O'Callaghan CA, Lasserson DS, Hobbs FR 2016 Global prevalence of chronic kidney disease—a systematic review and meta-analysis *PloS one* **11** e0158765.
- [2] Daugirdas JT, Depner TA, Inrig J, Mehrotra R, Rocco MV, Suri RS, Weiner DE, Greer N, Ishani A, MacDonald R and Olson C 2015 KDOQI clinical practice guideline for hemodialysis adequacy: 2015 update *Am. J. Kidney Dis.* **66** 884-930.
- [3] Stevens PE and Levin A 2013 Evaluation and management of chronic kidney disease: synopsis of the kidney disease: improving Global Outcomes 2012 clinical practice guideline *Ann. Intern. Med.* **158** 825–30
- [4] Inker LA, Astor BC, Fox CH, Isakova T, Lash JP, Peralta CA, Tamura MK and Feldman HI 2014 KDOQI US commentary on the 2012 KDIGO clinical practice guideline for the evaluation and management of CKD *Am. J. Kidney Dis.* **63** 713–35.
- [5] Abubakar II, Tillmann T and Banerjee A 2015 Global, regional, and national age-sex specific all-cause and cause-specific mortality for 240 causes of death, 1990-2013: a systematic analysis for the Global Burden of Disease Study 2013 *Lancet* **385** 117-71.
- [6] Collins AJ, Foley RN, Herzog C, Chavers B, Gilbertson D, Herzog C, Ishani A, Johansen K, Kasiske B, Kutner N and Liu J 2013 US Renal Data System 2012 annual data report *Am. J. Kidney Dis.* **61** E1-459.
- [7] Nusair MB, Rajpurohit N and Alpert MA 2012 Chronic inflammation and coronary atherosclerosis in patients with end-stage renal disease. *Cardiorenal Med.* **2** 117-24.
- [8] Razeghi E, Omati H, Maziar S, Khashayar P and Mahdavi-Mazdeh M 2008 Chronic inflammation increases risk in hemodialysis patients *Saudi J. Kidney Dis. Transpl.* **19** 785.
- [9] Krane V and Wanner C 2011 Statins, inflammation and kidney disease 2011 *Nat. Rev. Nephrol.* **7** 385.
- [10] Kato S, Chmielewski M, Honda H, Pecoits-Filho R, Matsuo S, Yuzawa Y, Tranaeus A, Stenvinkel P and Lindholm B 2008 Aspects of immune dysfunction in end-stage renal disease. *Clin. J. Am. Soc. Nephrol.* **3** 1526-33.
- [11] Stenvinkel P 2001 Inflammatory and atherosclerotic interactions in the depleted uremic patient. *Blood Purif.* **19** 53-61.
- [12] Libby P and Ridker PM 2004 Inflammation and atherosclerosis: role of C-reactive protein in risk assessment. *Am. J. Med.* **116** 9-16.
- [13] Ridker PM 2004 High-sensitivity C-reactive protein, inflammation, and cardiovascular risk: from concept to clinical practice to clinical benefit. *Am Heart J.* **148** S19-26.

- [14] Hoefler IE, Steffens S, Ala-Korpela M, Bäck M, Badimon L, Bochaton-Piallat ML, Boulanger CM, Caligiuri G, Dimmeler S, Egido J and Evans PC 2015 Novel methodologies for biomarker discovery in atherosclerosis *Eur. Heart J.* **36** 2635-42.
- [15] Malaponte G, Bevelacqua V, Fatuzzo P, Rapisarda F, Emmanuele G, Travali S and Mazzarino MC 2002 IL-1 β , TNF- α and IL-6 release from monocytes in haemodialysis patients in relation to dialytic age *Nephrol. Dial. Transplant.* **17** 1964-70
- [16] Pepys MB and Hirschfield GM 2003 C-reactive protein: a critical update *J. Clin. Invest.* **111** 1805–12.
- [17] Gosset P, Wallaert B, Tonnel AB and Fourneau C 1999 Thiol regulation of the production of TNF α , IL-6 and IL-8 by human alveolar macrophages. *Eur. Respir. J.* **14** 98-105.
- [18] Volanakis JE 2001 Human C-reactive protein: expression, structure, and function. *Mol. Immunol.* **38** 189– 97.
- [19] Penttinen MO, Oorni K, Ala-Korpela M and Kovanen PT 2000 Modified LDL-trigger of atherosclerosis and inflammation in the arterial intima *J. Intern. Med.* **247** 359–70.
- [20] Chen T, Hassan HC, Qian P, Vu M and Makris A 2018 High-Sensitivity Troponin T and C-Reactive Protein Have Different Prognostic Values in Hemo-and Peritoneal Dialysis Populations: A Cohort Study. *J. Am. Heart Assoc.* **7** e007876.
- [21] Koenig W, Sund M, Frohlich M, Fischer HG, Lowel H, Doring A, Hutchinson WL and Pepys MB 1999 C-reactive protein, a sensitive marker of inflammation, predicts future risk of coronary heart disease in initially healthy middle-aged men *Circulation* **99** 237– 42.
- [22] Biasucci LM, Liuzzo G, Grillo RL, Caligiuri G, Rebuzzi AG, Buffon A, Summaria F, Ginnetti F, Fadda G and Maseri A 1999 Elevated levels of C-reactive protein at discharge in patients with unstable angina predict recurrent instability *Circulation* 1999 **99** 855– 60.
- [23] Nagane NS, Ganu JV and Jagtap PE 2013 Study of oxidative stress in pre-and post-hemodialysis in chronic renal failure patients *Biomed. Res.* **20** 13-24
- [24] Rysz J, Banach M, Cialkowska-Rysz A, Stolarek R, Barylski M, Drozd J and Okonski P 2006 Blood serum levels of IL-2, IL-6, IL-8, TNF- α and IL-1 β in patients on maintenance hemodialysis *Cell. Mol. Immunol.* **3** 151-4.
- [25] Hisamuddin E, Hisam A, Wahid S and Raza G 2015 Validity of C-reactive protein (CRP) for diagnosis of neonatal sepsis *Pakistan J. Med. Sci.* **31** 527.