

# risk assement of air pollution exposure (no2,so2,total suspended particulate,and particulet matter 10 micron)and smoking habits on the lung function of bus drivers in palembang city

*by* Elvi 4 Sunarsih

---

**Submission date:** 22-Jun-2020 11:33AM (UTC+0700)

**Submission ID:** 1347829549

**File name:** Jurnal\_UI\_2019.pdf (598.04K)

**Word count:** 3999

**Character count:** 20528

# Risk Assessment of Air Pollution Exposure (NO<sub>2</sub>, SO<sub>2</sub>, Total Suspended Particulate, and Particulate Matter 10 micron) and Smoking Habits on the Lung Function of Bus Drivers in Palembang City

## Analisis Risiko Paparan Polusi Udara (NO<sub>2</sub>, SO<sub>2</sub>, Total Suspended Particulate dan Particulate Matter 10 mikron) dan Kebiasaan Merokok terhadap Fungsi Paru pada Sopir Bus di Kota Palembang

Elvi Sunarsih\*, Suheryanto\*\*, Rini Mutahar\*\*\*, Rahmi Garmini\*\*\*\*

\*Department of Occupational Health and Environmental Safety, Faculty of Public Health, Universitas Sriwijaya, Palembang, Indonesia \*\*Faculty of Mathematics and Natural Sciences, Universitas Sriwijaya, Palembang, Indonesia \*\*\*Department of Epidemiology and Biostatistics, Faculty of Public Health, Universitas Sriwijaya, Palembang, Indonesia \*\*\*\*Department of Environmental Health, STIKes Muhammadiyah Palembang, Indonesia

### Abstract

Nitrogen dioxide (NO<sub>2</sub>), sulfur dioxide (SO<sub>2</sub>), particulate matter (PM<sub>10</sub>), and total suspended particulate (TSP) are the most common and harmful air pollutants to humans. In a short period, air pollution exposure at 5 ppm for 10 minutes in humans causes dyspnea, and when the level is increased to 800 ppm could cause 100% mortality in animals. This study was an analytical study with implementing a cross-sectional design and risk analysis. One hundred subjects were involved in this study. The results showed that the mean value of the non-cancer Hazard Index (HI) for real-time exposure was NO<sub>2</sub>: 1.85; SO<sub>2</sub>: 2.92; TSP: 7.09; and PM<sub>10</sub>: 11.7 (HI value ≥1). Test for forced vital capacity lung capacity to non-cancer risk estimation of NO<sub>2</sub>, SO<sub>2</sub>, TSP, and PM<sub>10</sub> indicated that there was no significant relationship (p-value >0.05). The variable of smoking habit is the most dominant variable (odds ratio [OR] = 12.542) which affects respiratory disorders. The exposure to NO<sub>2</sub>, SO<sub>2</sub>, TSP, and PM<sub>10</sub> in Palembang City bus drivers is considered hazardous to the health of subjects without cancer.

**Keywords:** Health risk analysis, nitrogen dioxide (NO<sub>2</sub>), particulate matter (PM<sub>10</sub>), sulfur dioxide (SO<sub>2</sub>), total suspended particulate

### Abstrak

Nitrogen dioksida (NO<sub>2</sub>), sulfur dioksida (SO<sub>2</sub>), Particulate Matter (PM<sub>10</sub>) dan Total Suspended Particulate (TSP) merupakan pencemar udara yang paling umum dan berbahaya bagi manusia. Dalam jangka waktu pendek, pemajanan polusi udara dengan kadar 5 ppm selama 10 menit terhadap manusia mengakibatkan kesulitan dalam bernapas, dan kadar sebesar 800 ppm akan mengakibatkan 100% kematian pada binatang-binatang. Penelitian analitik, dengan desain potong lintang dan analisis risiko. Besar sampel yang diambil sebanyak 100 sampel. Hasil penelitian menunjukkan bahwa non-cancer Hazard Index (HI) untuk pajanan *realtime* dengan nilai rata-rata adalah NO<sub>2</sub> sebesar 1,85; SO<sub>2</sub> sebesar 2,92; TSP sebesar 7,09; dan PM<sub>10</sub> sebesar 11,7 (nilai HI ≥1). Hasil uji analisis hubungan nilai kapasitas paru dengan estimasi risiko non-kanker NO<sub>2</sub>, SO<sub>2</sub>, TSP dan PM<sub>10</sub> menunjukkan bahwa tidak terdapat hubungan yang signifikan (nilai p > 0,05). Variabel kebiasaan merokok merupakan variabel yang paling dominan (OR = 12,542) yang memengaruhi gangguan pemapasan. Pemajanan NO<sub>2</sub>, SO<sub>2</sub>, TSP dan PM<sub>10</sub> pada sopir bus di Kota Palembang sudah dianggap berisiko terhadap kesehatan non-kanker.

**Kata kunci:** Analisis risiko kesehatan, nitrogen dioksida, particulate matter, sulfur dioksida, total suspended particulate

**How to Cite:** Sunarsih E, Suheryanto, Mutahar R, Garmini R. Risk Analysis of Air Pollution Exposure (NO<sub>2</sub>, SO<sub>2</sub>, TSP, and PM<sub>10</sub>) and Smoking Habits on the Lung Function of Bus Drivers in Palembang City. *Kesmas: Public Health Journal*. 2019; 13 (4): 202-206. (doi:10.21109/kesmas.v13i4.1923)

**Correspondence:** Elvi Sunarsih, Department of Occupational Health and Environmental Safety, Faculty of Public Health, University of Sriwijaya, Phone: (0711) 580645 e-mail: elvisunarsih@fkm.unsri.ac.id

Received: December 11<sup>th</sup> 2017

Revised: April 17<sup>th</sup> 2018

Accepted: March 28<sup>th</sup> 2019

## Introduction

Air pollution has been a matter of great concern globally because of the associated health risks to individuals. The situation is getting worse in developing countries with more urbanization, industrialization, and more importantly, the rapidly growing population, posing a threat to human life in the form of pulmonary, cardiovascular, carcinogenic, or asthmatic diseases by accumulating toxic pollutants, harmful gases, metals, hydrocarbons, and other dangerous substances.<sup>1</sup>

The rise in the number of vehicles also affects the increase in vehicle exhaust emissions level. The United States Environmental Protection Agency released six types of pollutants that dominate the air and are most commonly encountered, which are ozone (O<sub>3</sub>), particulate matter (PM), carbon monoxide (CO), nitrogen oxides (NO<sub>x</sub>), sulfur dioxide (SO<sub>2</sub>), and lead (Pb). Approximately 50% of NO<sub>x</sub> emissions are generated by vehicles. The effect of exposure to air pollution can occur in chronic or acute conditions. In the short term, air pollutant gas (Pb, NO<sub>2</sub>, SO<sub>2</sub>, TSP, and dust) may cause respiratory disorders such as weakness, coughing, shortness of breath, bronchopneumonia, pulmonary edema, cyanosis, and methemoglobinemia.

The increasing amount of gases and solid pollutants emitted into the atmosphere has a considerable impact on both human health and environmental ecosystems. Specifically, long exposure to high concentrations of pollutants may lead to a shortening of life by several years because of cardiovascular and respiratory diseases and contributes to an increased risk of death. Recent studies also indicate that cancer occurrences are related to some pollutants, namely PM. These pollutants are hazardous because of not only their physical parameters, but mainly to their content of many dangerous organic compounds, such as hydrocarbons, polyaromatic hydrocarbons and their derivatives, and both inorganic elements such as Pb, Pt, Cd, As, Cu, Zn, and compounds such as nitrates, ammonia ions, sulfates, and various others.<sup>2</sup>

According to Ministry of Transportation (2009), the current growth rate of vehicles in Indonesia has increased significantly. This number increased in 2009, 2010, and 2011 by the amount of 67,290,816 units, 77,170,306 units and 85,601,351 units, respectively. Until 2012, the total number of motor vehicles in Indonesia has increased almost half the times the number of motor vehicles has in 2008.

Smoking habits have a negative impact on health, especially on the respiratory system, but the prevalence of the number of smokers continues to increase. According to the data from the World Health Organization (WHO) in 2013, the highest prevalence of smoking was in Europe with 28% and Asia was fifth at

19%. Tobacco-related illnesses will be the leading major health problems and cause 8.4 million deaths every year. This study aimed to create a model of air pollution exposure (NO<sub>2</sub>, SO<sub>2</sub>, TSP, and dust) and smoking habits on the lung function of bus drivers in Palembang City.

## Methods

This study was a risk assessment with cross-sectional study to estimate health risks of the population who were at risk because of exposure to NO<sub>2</sub>, SO<sub>2</sub>, TSP, and dust. The non-cancer risk characteristic was expressed as the non-cancer Hazard Index (HI). This index required the values of NO<sub>2</sub>, SO<sub>2</sub>, TSP, and dust as the Reference of Concentration (RFC), anthropometric data (inhalation rate and body weight), and exposure pattern (time, frequency, and duration of NO<sub>2</sub>, SO<sub>2</sub>, TSP, and dust exposure). The non-cancer value is the ratio between intake and RFC where the intake results from the calculation of exposure assessment. The hypothesis of the correlation between the risk value of the health status of NO<sub>2</sub>, SO<sub>2</sub>, TSP, dust, and lung capacity and respiratory disorders.<sup>3</sup> The study was conducted in four phases, including the initial survey and field observation of the study population for initial data collection of risk assessment, spirometry measurement and collection of selected air sample data (NO<sub>2</sub>, SO<sub>2</sub>, TSP, dust), laboratory analysis and spirogram readings, and modeling of risk factors of air pollution exposure and smoking habits of bus drivers.

The study site was in Palembang City, South Sumatra Province, Indonesia. The study subjects were all bus drivers in Palembang City who were a high-risk population. Subjects were taken using the purposive sampling technique. The minimum sample size was calculated using the sample size 2.0 application which obtained 78.88 (rounded to 80) subjects needed. In this study, the Z score at the 1 - /2 confidence interval of two-tailed hypothesis testing was 1.96 with a minimal error of 0.1 for a health study. The proportion of respiratory tract disorders (lung capacity function) was taken from the previous study on the risk analysis of respiratory tract disorders among bus drivers in Palembang City, which was 0.389.<sup>6,7</sup>

In consideration of conditions in the field to avoid any loss to follow up or sample drop out, the sample size was increased by at least 20% or 16 subjects. Therefore, the minimum sample size required in this study was 96 subjects, hence the sample size we accepted was 100 subjects. Proportional random sampling was applied as the sample collection method, which divided the number of samples proportionally in each terminal. Of nine bus stations in Palembang City, four bus stations were selected as study sites, which were Jakabaring Bus Station, 7 Ulu Bus Station, Ampera Bus Station, and

Plaju Bus Station. The number of respondents that have been calculated proportionally was taken from each bus station. Measurements of TSP and PM<sub>10</sub> concentrations were performed at 4-point measurements. The determination of the measurement point was based on SNI 19-7119.9-2005 using Haz-Dust EPAM-5000. OdaLog 7000 was used for the measurement of NO<sub>2</sub> and SO<sub>2</sub> concentrations was used. This study used univariate, bivariate, and multivariate data analysis. Univariate analysis was used to describe variables, bivariate analysis was used to see the relationship and significance between two variables, and multivariate analysis knew the dominant risk factors of lung capacity disorders and made a risk model of a non-cancer control.

## Results

The mean levels of NO<sub>2</sub>, SO<sub>2</sub>, TSP, and PM<sub>10</sub> at Palembang City Bus Station were 0.416 mg/m<sup>3</sup> for NO<sub>2</sub>; 0.848 mg/m<sup>3</sup> for SO<sub>2</sub>; 5.559 mg/m<sup>3</sup> for TSP, and 2.543 mg/m<sup>3</sup> for PM<sub>10</sub>.

The most critical very high-risk air pollutant is TSP (5.559 mg/m<sup>3</sup>) since every year the TSP HI increases. The air pollutant with a low risk to pulmonary function is NO<sub>2</sub> (0.416 mg/m<sup>3</sup>). Exposure assessment was used to calculate the dose or the number of risk agents (pollutants or toxic substances) received individually and expressed as "intake". Further exposure pattern assessment was determined by the frequency of exposure, length of exposure, and route or path of exposure. The assessment of exposure frequency, duration/time of exposure, and duration of exposure in this study were

obtained from the calculation of data directly. The exposure route of NO<sub>2</sub>, SO<sub>2</sub>, TSP, and PM<sub>10</sub> in this study was through inhalation. The mean exposure time was 8 hours/day, the minimum value of exposure frequency was 142 days/year, the duration of exposure was at least two years, and the mean weight of subjects was 58.19 kg. The inhalation rate in this study was based on the standard from the U.S. Environmental Protection Agency (EPA) by taking the value of the inhalation rate on a long-term exposure level 31–41 years. The recommended inhalation rate obtained was 16 m<sup>3</sup>/day. After unit conversion, the rate of inhalation per hour was 0.66 m<sup>3</sup>/hour. The exposure analysis to determine the value of the inhalation intake was calculated using the following formula below.

Distribution of Inhalation intake of NO<sub>2</sub>, SO<sub>2</sub>, TSP, and PM<sub>10</sub> for each subject can be seen in Table 2. The lowest inhalation intake of NO<sub>2</sub> was 0.022 mg/kg/day, and the highest intake was 0.056 mg/kg/day. The lowest SO<sub>2</sub> inhalation intake was 0.066 mg/kg/day, and the highest intake was 0.089 mg/kg/day. The lowest TSP

$$I = \frac{CRtefeDt}{Wbtavg}$$

Notes: I = intake [mg/(kg x hari)], C = concentration (mg/m<sup>3</sup>), R = rate of inhalation (m<sup>3</sup>/hour), te = time of exposure (hour/day), fe = frequency of exposure (day/year), Dt = duration of time (year), Wb = weight (kg), tavg = average time period (Dt x 365 day/year).

Table 1. Levels of NO<sub>2</sub>, SO<sub>2</sub>, TSP, and PM<sub>10</sub> at Bus Stations in Palembang City

Bus Station	Level of NO <sub>2</sub> (mg/m <sup>3</sup> )	Level of SO <sub>2</sub> (mg/m <sup>3</sup> )	Level of TSP (mg/m <sup>3</sup> )	Level of PM <sub>10</sub> (mg/m <sup>3</sup> )
Jakabaring Bus Station	0.42	0.84	4.81	2.38
7 Ulu Bus Station	0.41	0.83	6.8	2.65
Ampera Bus Station	0.40	0.91	8.09	2.84
Plaju Bus Station	0.37	0.86	4.65	2.31
Mean	0.41	0.84	5.55	2.54

Notes: NO<sub>2</sub>= Nitrogen Dioxide, CO= Carbon Monoxide, SO<sub>2</sub>= Sulfur Dioxide, TSP= Total Suspended Particulate, PM<sub>10</sub>= Particulate Matter 10 micron

Table 2. Distribution of Inhalation Intake of NO<sub>2</sub>, SO<sub>2</sub>, TSP, and PM<sub>10</sub> at the Bus Station in Palembang City

Description	NO <sub>2</sub> Intake (mg/kg/day)	SO <sub>2</sub> Intake (mg/kg/day)	TSP Intake (mg/kg/day)	PM <sub>10</sub> Intake (mg/kg/day)
Mean	0.037	0.077	0.504	0.230
Median	0.037	0.075	0.476	0.256
Minimum	0.022	0.064	0.284	0.091
Maximum	0.055	0.087	0.828	0.373

Notes: NO<sub>2</sub>= Nitrogen Dioxide, CO= Carbon Monoxide, SO<sub>2</sub>= Sulfur Dioxide, TSP= Total Suspended Particulate, PM<sub>10</sub>= Particulate Matter 10 micron



Table 3. Model of Multivariate Analysis

Variable	$\beta$	p-Value	OR	95% CI
Non-cancer risk PM <sub>10</sub>	0.156	0.026	1.168	1.018–1.341
Smoking habit	2.529	0.001	12.542	2.940–53.500
Exposure duration	1.453	0.006	4.275	1.529–11.951
Records of the disease	1.393	0.028	4.026	1.160–13.967
Constant	-9.326			

Notes:  $\beta$  = Beta, p value = probability value, OR = Odds Ratio, 95%CI = 95% confidence level

inhalation intake was 0.288 mg/kg/day, and the highest intake was 0.842 mg/kg/day. The lowest inhalation intake of PM<sub>10</sub> was 0.092 mg/kg/day, and the highest intake was 0.379 mg/kg/day.

Risk characteristics were derived from two parts of the assessment, exposure assessment and dose-response assessment. The value of HI for real-time exposure using the mean values as described previously was 1.85 for NO<sub>2</sub>; 2.92 for SO<sub>2</sub>; 7.09 for TSP; and 11.7 for PM<sub>10</sub> (HI-value = 1).<sup>8</sup>

Therefore, it can be concluded that exposure of bus drivers to NO<sub>2</sub>, SO<sub>2</sub>, TSP, and PM<sub>10</sub> in Palembang City is already considered hazardous to the health of drivers without cancer, so control is needed.

The result of the multivariate analysis shows that the smoking habit variable is the most dominant variable related to respiratory disorders in bus drivers, with an odds ratio (OR) = 12.542 means that a bus driver who has a smoking habit is 12.542 times more likely to suffer from a respiratory disorder compared with a bus driver who has no smoking habit (Table 3).

## Discussion

The risk analysis showed that the demand for risk management was greater if HI > 1. The results of this study indicated that risk management was needed to control the health effects caused by NO<sub>2</sub>, SO<sub>2</sub>, TSP, and PM<sub>10</sub> exposure. Control was done in various ways, which could be formulated quantitatively by modifying the standard or quality standard. HI > 1 could be controlled by controlling the intake value of NO<sub>2</sub> inhalation because the RFC value was derived from experimental data that could not be modified (because it was a reference). Therefore, the value that had to be lowered was I to make it harmless. The mean value of I was 0.038 with 0.056 being the highest. The same equation from the non-cancer risk analysis equation to decrease the value of I to the acceptable RFC level of 0.02. However, the RFC values of SO<sub>2</sub>, TSP, and PM<sub>10</sub> were taken from previous studies because they were not yet available in the EPA's Integrated Risk Information System list or the Minimum Risk Level ATSDR table.

When subjects were enrolled at age 16 years old,

then they would have about two years more of exposure to NO<sub>2</sub>, SO<sub>2</sub>, TSP, and PM<sub>10</sub> assuming that they did not quit or move their jobs. Then, there would be an increase in risk according to the scenario with a non-cancer risk control simulation framework of NO<sub>2</sub>, SO<sub>2</sub>, TSP, and PM<sub>10</sub>. At Palembang City bus stations, it was assumed that there was no interaction between existing risks when considering the mean concentrations of NO<sub>2</sub> (0.038 mg/m<sup>3</sup>), SO<sub>2</sub> (0.078 mg/m<sup>3</sup>), TSP (0.512 mg/m<sup>3</sup>), and PM<sub>10</sub> (0.234 mg/m<sup>3</sup>), respectively. These concentrations increase every year as the rate of vehicles increases, which has been 5% per year (considering the trend of the growth rate of vehicle data from the Central Statistics Agency 1987–2013). The total exposure was considered constant based on a mean value of 8 hours/day. The inhalation rate, using the default value of 0.66 m<sup>3</sup>/hr, and the frequency of working days was 142 days/year, and the mean body weight at the time of the study was 58.19 kg. The simulation model 1 for exposure started from the second year to the sixth year.

The results of the risk analysis showed that HI > 1. Based on the EPA, this indicated that risk management was needed to control the health effects due to exposure to NO<sub>2</sub>, SO<sub>2</sub>, TSP, and PM<sub>10</sub>. The risk control simulation framework also showed that there was another contributing source of exposure besides NO<sub>2</sub>, SO<sub>2</sub>, TSP, and PM<sub>10</sub> in Palembang City bus station, which was cigarette smoke that exposed subjects. For the primary source of exposure (NO<sub>2</sub>, SO<sub>2</sub>, TSP, and PM<sub>10</sub>) in bus stations, the control was done in various ways, but the aim was to reduce the exposure risk by formulating the recommended standard value quantitatively to modify the standard or quality standard.

The study from Zhengzhou City, China showed that the data from six monitoring stations in Zhengzhou City analyzed the changing trend in concentrations of SO<sub>2</sub>, NO<sub>x</sub>/NO<sub>2</sub>, and TSP/PM<sub>10</sub> in 1996–2008, based on the non-parametric Mann-Kendall test and Sen's slope estimator. The overall air pollution level was evaluated by the Multi-Pollutant Index. It was found that the concentration of each pollutant exceeded the WHO guideline value by an apparent amount, but the changing trend varied: SO<sub>2</sub> and NO<sub>2</sub> significantly increased mainly due

to an increase in coal consumption and vehicle number, while NO<sub>2</sub>, TSP, and PM<sub>10</sub> decreased. The air pollution was serious, and differed markedly among the three functional regions: it is the most severe in the Industrial and Residential Area, followed by the Transportation Hub and Business District, and then the High-tech, Cultural and Educational Area.<sup>4</sup> The same study on lung function showed that work period, the use of Personal Protective Equipment (PPE), and the total dust level were factors which affect pulmonary function impairment in workers.<sup>5</sup>

The decrease in lung function over time was associated with both smoking habits and duration of occupational dust exposure. Specifically, occupational quartz exposure negatively influenced annual lung function parameters. Implementation of stricter occupational limit values for dust exposure resulted in a highly significant deceleration of the annual decrease in respiratory function.<sup>6</sup> Cough-related symptoms were associated with incident obstructive lung physiology and emphysema, whereas shortness of breath was associated with restrictive lung physiology. When placed in a clinical context, these are not surprising. Cough or phlegm and episodes of bronchitis may be more "exposure-driven" (including exposure beyond merely cigarette smoking, such as dusty work environments or air pollution) and indicate susceptibility.<sup>7</sup>

The study of smokers in the Netherlands showed that a total of 255 participants were followed up (49%). The point prevalence rate of nonsmoking at follow up was 18% (N Z 47), and 9%, assuming that all non-respondents were smokers. This rate was not lower than the expected rate of quitting in the Dutch population (8%–9%), and primary "care as usual" in smokers screened with abnormal lung function (10%; p-value > 0.05 for all comparisons). The average decline in post-bronchodilator Forced Expiratory Volume (FEV1) was 26 mL/year, which was unrelated to smoking status at follow up. In contrast, non-smokers showed a clinically meaningful and statistically significant reduction in post-bronchodilator FEV1 (p-value < 0.001).<sup>8</sup>

Factory workers had lower spirometric functions, especially those with high dust exposure. There was a dose-response relation between the duration of dust exposure and FEV1 and Forced Vital Capacity (FVC), the adjusted effect of 21 years of exposure on FEV1 being - 240 mL (-100 to -380 mL) and on FVC -300 mL (-140 to -460 mL). The adverse effect of dust on lung function was more substantial in current smokers and suggested a synergism between smoking and tile dust exposure.<sup>9</sup>

The lung capacity disorders most bus drivers in Palembang City had were caused by smoking habits with exposure to NO<sub>2</sub>, SO<sub>2</sub>, TSP, and PM<sub>10</sub> with an average

length of exposure of 13 hours/day. Dust is among the strong risk factors for pulmonary dysfunction depending on both the level and duration of exposure. In public transportation drivers, workers are in high levels of pollutants for a long time. This prolonged exposure will cause a build up of pollutants with small particles in the alveoli, resulting in changes in lung function. The longer the bus driver works in the terminal area, the longer the exposure to the pollutants is, then the occurrence of lung function disorders will also be greater. However, it also depends on the concentration of pollutants and the mechanism of clearance of each individual.

## Conclusion

Most bus drivers who have lung capacity disorders are caused by smoking habits with exposure to NO<sub>2</sub>, SO<sub>2</sub>, TSP, and PM<sub>10</sub> with an average length of exposure of 13 hours/day. The highest risk air pollutant is TSP since every year the TSP HI increases. Smoking habit variable was the most dominant variable, which influences respiratory disorders.

## References

1. Rumana HS, Sharma RC, Beniwal V, Sharma AK. A retrospective approach to assess human health risks associated with growing air pollution in urbanized area of Thar Desert, Western Rajasthan, India. *Journal of Environmental Health Science & Engineering*. 2014; 12 : 23.
2. Adamec V, R. Ličbinský, Cholava R. Transport and health risks of transport. *Transactions on Transport Sciences*. 2011; 4(3): 115-54.
3. Putri, A Retno dan Driejana. Analisis Konsentrasi NOx di dalam ruangan pada rumah tinggal di tepi jalan raya (studi kasus: Wilayah Karees, Bandung). Bandung: ITB. 2012.
4. Xile, DU, Changhe LU, Hairong W, Jianhua MA. Trends of urban air pollution in Zhengzhou City in 1996–2008. *Chinese Geographical Science*. 2012; 22(4): 402-15.
5. Nafisa SF, Joko T, Setiani O. Hubungan paparan debu kayu di lingkungan kerja terhadap gangguan fungsi paru pada pekerja di PT. Arumbai Kasembadan, Banyumas. *Jurnal Kesehatan Masyarakat*. 2016; 4(5):178-86.
6. Haluza D, Moshhammer H, Hochgatterer K. Dust is in the air: effects of occupational exposure to mineral dust on lung function in a 9-year study. *Lung*. 2014; 192: 111-17.
7. Kalhan R, Dransfield MT, Colangelo LA, Cuttica MJ, Jacobs JDR, Thyagarajan B, et. al. Respiratory symptoms in young adults and future lung disease. *American Journal of Respiratory and Critical Care Medicine*. 2018: p.197.
8. Kotz D, Wesseling G, Aveyard P, Schayck OCP van. Smoking cessation and development of respiratory health in smokers screened with normal spirometry. *Respiratory Medicine*. 2011; 105: 243-49.
9. Jaakkola MS, Sripaiboonkij P, Jaakkola JJK. Effects of occupational exposures and smoking on lung function in tile factory workers. *International Archives of Occupational and Environmental Health*. 2011; 84: 151-8.

risk assement of air pollution exposure (no2,so2,total suspended particulate,and particulet matter 10 micron)and smoking habits on the lung function of bus drivers in palembang city

ORIGINALITY REPORT

4%

SIMILARITY INDEX

%

INTERNET SOURCES

%

PUBLICATIONS

4%

STUDENT PAPERS

MATCH ALL SOURCES (ONLY SELECTED SOURCE PRINTED)

1%

★ Submitted to Politeknik Negeri Bandung

Student Paper

Exclude quotes On

Exclude bibliography On

Exclude matches < 1%