

The morphometric variability of the mangrove horseshoe crab (*Carcinoscorpius rotundicauda*) from Banyuasin estuarine of South Sumatra, Indonesia

by Wike Ayu Eka Putri

Submission date: 04-Jun-2020 12:08PM (UTC+0700)

Submission ID: 1337492047

File name: EM_Mba_OJi_2.pdf (875.11K)

Word count: 4086

Character count: 21777

The morphometric variability of the mangrove horseshoe crab (*Carcinoscorpius rotundicauda*) from Banyuasin estuarine of South Sumatra, Indonesia

FAUZIYAH^{1*}, WIKE A. E. PUTRI¹, ANNA I. S. PURWIYANTO¹, FITRI AGUSTRIANI¹,
APON Z. MUSTOPA² & FATIMAH³

¹Marine Science Study Program, Faculty of Mathematics and Natural Sciences, University of Sriwijaya, Indonesia;

²Research Center for Biotechnology, Indonesian Institute of Sciences (LIPI)

³Indonesian Center for Agricultural Biotechnology and Genetic Resources Research and Development, Ministry of Agriculture

*Corresponding author. E-mail: siti_fauziyah@yahoo.com

Received 21 October 2019 | Accepted by V. Pešić: 10 November 2019 | Published online 27 November 2019.

6

Abstract

The allometric analysis was applied to record the morphometric variability in different sex of the mangrove horseshoe crab (*Carcinoscorpius rotundicauda*) collected from Banyuasin estuarine of South Sumatra, Indonesia during July 2019. The body parameter measurement of body weight (BW), total length (TL), prosoma width (PW), carapace length (CL), and telson length (TL) were recorded then pooled refer to the sex. The differences in each body parameter between the males and females were observed through the Student's t-test. The length/width-weight relationship for both sexes was analyzed using the power equations while the width/length-length relationship was analyzed using the linear equations. The Student's t-test was used to determine significant differences from the isometric value. The females of *C. rotundicauda* were significantly heavier and larger in size than males except for telson length ($p < 0.05$). Both sexes revealed a negative allometric growth for the TL/CL-BW relationships while the males indicated a negative allometric growth but the females indicated an isometric growth for the PW-BW relationships. Both sexes also indicated the isometric growth for the CL-PW relationship. The TL-PW/TEL relationship also revealed a negative allometric growth for both sexes. The study results were expected to be used as baseline data for the horseshoe crab management plan and action.

Key words: Banyuasin estuarine, Body parameters, *Carcinoscorpius rotundicauda*, Morphometric, South Sumatra.

Introduction

6

Morphometric can be used as an effective tool to study the variation in shape and size of organisms (Webster, 2007) as well as the morphometric studies also useful for comparing various living organisms through the quantitative studies of various body parts (Srijaya et al., 2010). While the relationship between the size and shape of an organism (physical of morphological changes) is described by allometric studies and

its results useful for assessing the body shape differences of species inhabiting in various ecosystems (Srijaya et al., 2010; Syuhaida et al., 2019). At the first, the allometric application was used for many organisms especially calculating the population growth characteristics, and furthermore applied to study the population diversity by many biologists extensively (Srijaya et al., 2010).

The mangrove horseshoe crab (*Carcinoscorpius rotundicauda*) is one of the Asian horseshoe crabs (Manca et al., 2017), and was classified in the IUCN red list as data deficient since 1996 (World Conservation Monitoring Centre, 1996). While in Indonesia, the Asian horseshoe crabs are the protected genetic resources based on Minister of Forestry Decree No. 12/Kpts-II/1987 and Government Regulation No. 7/1999. Their distribution includes India, Philippines, Japan, Korea, China, Thailand, Malaysia, Singapore, and Indonesia (Cartwright-Taylor et al., 2011; Chen et al., 2015). In Banyuasin Coastal Waters of South Sumatra (Indonesia), the horseshoe crab was found as a discard catch for the trammel net fishing (Fauziyah et al., 2018) and first investigation record on these horseshoe crabs (*Tachypleus gigas* and *C. rotundicauda*) were successful identified (Fauziyah et al., 2019). However there was no information in detail for the species of *C. rotundicauda* both in the terms of their morphology and population structure.

Generally, there was very limited data found on the horseshoe crab morphology in coastal water of Sumatra Island and this study was a first record on the horseshoe crab morphometric in the coastal waters of South Sumatra, Indonesia. This study's aim was to describe the allometric relationship and the morphometric variability for *C. rotundicauda* from Banyuasin estuarine of South Sumatra, Indonesia. The result could provide baseline information for the growth aspect as well as provide valuable information for future research in detail to create stakeholders awareness in order to save *C. rotundicauda* population.

Material and methods

The live sample was obtained from the Banyuasin estuarine of South Sumatra, Indonesia (Figure 1) during July 2019. The samples were collected using a trammel net fishing and the sampling sites referred to the local fishermen information.




Figure 1. The sampling sites map in Banyuasin Estuarine of South Sumatra, Indonesia (Fauziyah et al. 2019).

The cross-section morphology of the telson was used to distinguish between *C. rotundicauda*, *Tachypleus gigas*, and *Tachypleus tridentatus* (Yang and K²²2015). Their telson is triangular for both *Tachypleus* species whereas circular for *C. rotundicauda* (Cartwright-Taylor et al., 2009; Dolejš and Vaňousová, 2015; Tanacredi et al., 2009). Female has a chelate clasper like scissors while the male has a hemichelate clasper like hooks on the first and second walking legs. For this study, female and male were identified as well as morphometric measurements were recorded (Figure 2). The carapace length (CL), prosoma width (PW), telson length (TEL), and total length (TL) were recorded to 1 mm accuracy while body weight (BW) was recorded to 1 gram accuracy. All measurements data obtained were pooled refer to the species and sex. The difference in morphometric parameters between the males and females were observed through Student's t-test (Two-Sample Assuming Unequal Variances) using Excel program.




Figure 2. The morphometric measurement of the horseshoe crab; total length (TL), telson length (TEL), prosoma width (PW), and carapace length (CL). The measurement was recorded to 1 mm accuracy.

The length-weight relationship for males and females were analyzed using the power equations $W = aL^b$ (Le Cren, 1951; Froese, 2006; Graham et al., 2009; Jawahir et al., 2017) while the carapace width-length relationship for males and females were analyzed using the linear equations $L_y = a + bL_x$ (Amaral et al., 2014). Regarding the PW-CL, PW-TL and TEL-TL relationships (Amaral et al., 2014), the growth pattern could be isometric ($b = 1$), negative allometric ($b < 1$) or positive allometric ($b > 1$) whereas for the PW-BW, TL-BW, and CL-BW relationships, the growth pattern could be isometric ($b = 3$), negative allometric ($b < 3$), or positive allometric ($b > 3$). The b value is describing growth pattern (Syuhaida et al., 2019) and the Student's t-test (Hegele-drywa et al., 2014) was needed to determine significant differences from the isometric value ($b = 3$ or $b = 1$) with the significant level at 5% ($P < 0.05$). The Student's t-test or known as Bailey's t-test was expressed as follows (Nair et al., 2015; Thomas, 2013):

$$t_s = \left| \frac{a-b}{sb} \right| \quad \text{or} \quad t_s = \left| \frac{1-b}{sb} \right|$$

where: t_s = the Student's t-test or Bailey's t-test

sb = standard error of the b coefficients

b = slope or coefficients regression

3 = isometric value for the PW-BW, TL-BW, and CL-BW relationship

1 = isometric value for the PW-CL, PW-TL and TEL-TL relationship

Results

A total of 45 adults *C. rotundicauda* were found during the survey ($\sigma = 18$, $\phi = 27$) and no juvenile individuals were obtained. The specimen sizes were shown as the normal distribution histograms (Figure 3).

In the measurement results of this study (Table 1), the BW means of males and females were 119.61 ± 22.88 gram (ranged from 85 gram to 163 gram) and 202.26 ± 53.80 grams (ranged from 72 gram to 285 gram). The PW means of males and females were 114.17 ± 8.62 mm (ranged between 100-130 mm) and 130 ± 12.48 mm (ranged between 100-150 mm). The CL means of males and females were 110.83 ± 10.04 mm (ranged between 100-130 mm) and 131.48 ± 11.59 mm (ranged between 100-150 mm). The TEL means of males and females were 138.53 ± 18.53 mm (ranged between 80-160 mm) and 145.56 ± 19.87 mm (ranged between 105-180 mm). The TL means of males and females were 250 ± 22.01 mm (ranged between 190-280 mm) and 277.04 ± 26.94 mm (ranged between 215-325 mm).




Figure 3. The size distribution of *Carcinoscropsius rotundicauda*. Females were greater in size than males for each measurement of prosoma width (A), carapace length (B), telson length (C) and total length (D).

Table 1. Morphometric measurement records of *Carcinoscropsius rotundicauda* male and female collected from Banyuasin estuarine of South Sumatra, Indonesia. There was a significant difference between sex for the parameters of BW, CL, TL, and PW unless the TEL parameter.

Parameters	Males (N = 18)		Females (N = 27)	
	Mean	Std. Dev.	Mean	Std. Dev.
BW (gram)	119.61 ^a	22.88	202.26 ^b	53.80
CL (mm)	110.83 ^a	10.04	131.48 ^b	11.59
TEL (mm)	138.53 ^a	18.53	145.56 ^a	19.87
TL (mm)	250 ^a	22.01	277.04 ^b	26.94
PW (mm)	114.17 ^a	8.62	130 ^b	12.48

^a BW = body weight, CL = carapace length, TEL = telson length, TL = total length, PW = prosoma width, Std. Dev. = standard deviations, and the same superscripts of each parameters indicated not significant difference between sex at $\alpha = 0.05$.

The most males were found in the PW class 111-120 mm whereas most females were found in the PW class 121-130 mm (Figure 3). The most males were found in the CL-class 101-110 mm while most females were found in the two CL-class namely 121-130 mm and 131-140 mm. The most males were found in the TEL class 141-150 mm while most females were found in the TEL class 151-160 mm. The most males were found in the TL class 241-250 mm while most females were found in the TL class 281-290 mm. The body parameters measurement indicated that the females were greater in size than males for each measurement of prosoma width, carapace length, telson length, and total length. However statistically (Table 1), the females of *C. rotundicauda* were significantly heavier (BW parameter) and larger (CL, PW, and TL parameters) than males while the females were not significantly larger than males for the TEL size.

Table 2. Different relationships between body part parameters for *Carcinoscorpius rotundicauda* male and female. Samples were collected from Banyuasin estuarine of South Sumatra, Indonesia using trammel net fishing. All regressions are significant ($\alpha = 0.05$).

Body Parameter	Parameters of the relationship				Bailey's t-test				Growth pattern
	b	Sb	Sig. F	R ²	t _b	β	t _{tab}	t _s	
TL-BW (Males)	1.847	0.308	0.000	0.800	5.995 ^S	3	2.262	3.741 ^S	allometric (-)
TL-BW (Females)	2.191	0.174	0.000	0.909	12.621 ^S	3	2.120	4.662 ^S	allometric (-)
PW-BW (Males)	2.244	0.261	0.000	0.822	8.605 ^S	3	2.120	2.901 ^S	allometric (-)
PW-BW (Females)	3.012	0.217	0.000	0.885	13.892 ^S	3	2.060	0.054 ^{NS}	Isometric
CL-BW (Males)	1.347	0.275	0.000	0.649	4.907 ^S	3	2.160	6.021 ^S	allometric (-)
CL-BW (Females)	2.077	0.295	0.000	0.712	7.033 ^S	3	2.086	3.128 ^S	allometric (-)
CL-PW (Males)	0.745	0.123	0.000	0.722	6.034 ^S	1	2.145	2.064 ^{NS}	Isometric
CL-PW (Females)	0.884	0.099	0.000	0.815	8.918 ^S	1	2.101	1.171 ^{NS}	Isometric
TL-PW (Males)	0.508	0.062	0.000	0.828	8.203 ^S	1	2.145	7.946 ^S	allometric (-)
TL-PW (Females)	0.336	0.037	0.000	0.806	9.128 ^S	1	2.086	18.061 ^S	allometric (-)
TL-TEL (Males)	0.739	0.095	0.000	0.800	7.739 ^S	1	2.131	2.737 ^S	allometric (-)
TL-TEL (Females)	0.680	0.057	0.000	0.849	11.837 ^S	1	2.060	5.580 ^S	allometric (-)

² BW = body weight, CL = carapace length, TEL = telson length, TL = total length, PW = prosoma width, Sb = standard error of the b coefficients, Sig. F = significance F-test, R² = the coefficient of determination, t_b = the t-test statistic for H₀ of b = 0, β = allometric value, t_{tab} = critical values of the t distribution, t_s = Bailey's t-test for allometric values, S = Significant, and NS = Not significant.

The TL-BW, PW-BW, CL-²², CL-PW, TL-PW and TL-TEL relationships for both sexes were shown in Figures 4-6 and Table 2. All relationships between body parameters of *C. rotundicauda* showed a significant difference ($p < 0.05$). The TL/CL-BW relationship for both sexes presented a negative allometric growth ($b < 3$). It's indicated that the BW grew slowly with respect to the TL/CL after molting. The different growth patterns were shown by the PW-BW relationship where the males indicated a negative allometric growth whereas the females indicated an isometric growth (the PW and BW grew at the same rate). The isometric growth was occurrence on the CL-PW relationship ($b = 1$) for both sex (the PW and CL grew at the same rate). The TL-PW/TEL relationship for both sex ($b < 1$) indicated a negative allometric growth (the PW/TEL grew slowly with respect to the TL). The highest R² value in the regression model was shown in females for the TL-BW relationship ($R^2 = 0.909$) whereas the lowest was male for the CL-BW relationship ($R^2 = 0.649$). In terms of the R² value ($R^2 > 0.5$), all relations between the major body parameters of *C. rotundicauda* indicated a strong correlation with each other.




Figure 4. Total length-body weight (left) and prosoma width-body weight (right) relationship of males and females *Carcinoscopus rotundicauda* from Banyuasin estuarine of South Sumatra, Indonesia. Both sex indicated the body weight grown slowly with respect to the total length (negative allometric, $b < 3$, $p < 0.05$) while the body weight of females also grown slowly with respect to the prosoma width (negative allometric, $b < 3$, $p < 0.05$) but the prosoma width and body weight for males grown at the same rate (isometric, $b = 3$, $p < 0.05$).




Figure 5. Carapace length-body weight (left) and prosoma width-carapace length (right) relationship of males and females *Carcinoscopus rotundicauda* from Banyuasin estuarine of South Sumatra, Indonesia. The body weight grew slowly with respect to the carapace length (negative allometric, $b < 3$, $p < 0.05$) for both sexes while the carapace length and prosoma width grown at the same rate (isometric, $b = 1$, $p < 0.05$).

Discussion

This study results provided important data that related to the difference of various body parameters for *C. rotundicauda* males and females from Banyuasin Estuarine since no information and publication before thus became the first record on morphometrical and allometric of horseshoe crabs in South Sumatra, Indonesia. An understanding of the relationship between one body parameter and the other of horseshoe crab was essential to know their growth (Panda and Naik, 2017). The morphometric variation in weight, width, and length for *C. rotundicauda* indicated the populations structure from immature to mature (Syuhaida et al., 2019) and the *C. rotundicauda* populations in this study indicated a mature specimen for both sexes due to their prosomal width were 8 cm and above (Cartwright-Taylor et al., 2009).

The measurement of major body parameters (W, CL, TL) indicated that the females statistically significant larger in size than male, but statistically there was no a significant difference in size of the TEL

between females and males. Similarly, the females larger in size than males were found in another study such as Gelang Patah (Johor) and Setiu of Peninsular Malaysia (Srijaya et al., 2010), Bhitarkanika National Park of Odisha (Panda and Naik, 2017), and Singapore (Cartwright-Taylor et al., 2009). The molting activities and numerous egg in a prosomal cavity for females affecting the females larger in size than males (Graham et al., 2009; Tan et al., 2012) and the reproductive activities of the *C. rotundicauda* also affecting variation in maturity level among their populations (Syuhaida et al., 2019).




Figure 6. Prosoma width-total length (left) and telson length-total length (right) relationship of males and females *Carcinoscropsus rotundicauda* from Banyuasin estuarine of South Sumatra, Indonesia. The telson length or prosoma width grew slowly with respect to the total length (negative allometric, $b < 1$, $p < 0.05$) for both sexes.

The TL/CL-BW relationships indicated the same growth patterns (negative allometric) for both sexes whereas the PW-BW relationships presented the differences in growth patterns (isometric for females and negative allometric for males). The PW and BW parameters of females grew at the same rate (isometric growth) and the same pattern also occurs in the PW and CL parameters for both sexes. Similar to *C. rotundicauda* from Merlimau Merlimau of Melaka (West Coast of Peninsular Malaysia), the isometric growth in the PW-CL relationship for both sexes was also recorded (Syuhaida et al., 2019). The TL parameter grew rapidly than the PW/TL for both sexes. Nonetheless, females grew slightly greater than males that exhibiting by the value of b females were greater than males.

The negative allometric growth in the TL-BW relationships also recorded in the West Coast of Peninsular Malaysia (Merlimau and Pendas) except female *C. rotundicauda* in Pendas expressed an isometric growth (Syuhaida et al., 2019). The negative allometric growth in the PW-BW relationships also recorded in the West Coast of Peninsular Malaysia except for females in Merlimau expressed an isometric growth (Syuhaida et al., 2019). The female *C. rotundicauda* from Bintan Bay of Riau Islands Province indicated negative allometric growth in the CL-BW relationships but positive allometric growth for males (Anggraini et al., 2017). The differences in the relationship between body parameters for *C. rotundicauda* population could indirectly be affected by food availability, local environmental conditions, feeding efficiency, population density, and genetic effect (Le 6ren, 1951; Jawahir et al., 2017; Panda and Naik, 2017; Srijaya et al., 2010; Vijayakumar et al., 2000). Further detailed studies using appropriate molecular tools are needed to find out the growth in the body parameters from the different populations of *C. rotundicauda* (Srijaya et al., 2010). In order to conserve these horseshoe c 22s, monitoring the balance of sex ratio and juveniles is essential as an indicator of this population's health (Cartwright-Taylor et al., 2009). In addition, these study results can be used as the baseline data for the conservation management plan to prevent declining the horseshoe crabs population in the future.

Acknowledgment

The authors would like to thank Mr. Ardani, Mr. Langgeng, Mr. Saderun and the crews of the trammel net (Roni Hastra, Brenda Sellyndia, Nita Puspita Sari, Sri Wulandari, Ari Muzari, Noviantrio Gulo) for their help and collaboration. We are very grateful to the reviewers who provided invaluable suggestions for this paper. And special thanks to the Indonesian Ministry of Research, Technology and Higher Education (Kemenristek Dikti) for their National Competitive Fund so that this research can be accomplished.

References

- 7 Amaral, K. D. S., Vieira, I. M., Osório, F. M., Rocha, J. D. M., and Lima, J. de F. (2014) Bioecology of the crab *Ucides cordatus* (Crustacea, Decapoda) in mangroves influenced by the Amazon River, Brazil. *Acta Amazonica*, 44(2), 213–222. doi: 10.1590/S0044-59672014000200007
- 9 Anggraini, R., Bengen, D. G., and Natih, N. M. N. (2017) Population structure and morphometry of horseshoe crab *Carcinoscorpius rotundicauda*, Latreille 1802 in Kampung Gisi Coastal Area of Bintan Bay of Riau Islands Province. *Jurnal Ilmu dan Teknologi Kelaut Tropis*, 9(1), 211–220.
- Cartwright-Taylor, L., von Bing, Y., Chi, H. C., and Tee, L. S. (2011) Distribution and abundance of horseshoe crabs *Tachypleus gigas* and *Carcinoscorpius rotundicauda* around the main island of Singapore. *Aquatic Biology*, 13(2), 127–136. doi: 10.3354/ab00346
- Cartwright-Taylor, L., Lee, J., and Hsu, C. C. (2009) Population structure and breeding pattern of the mangrove horseshoe crab *Carcinoscorpius rotundicauda* in Singapore. *Aquatic Biology*, 8, 61–69. doi: 10.3354/ab00206
- 3 Chen, C. P., Yang, M. C., Fan, L. F., Qiu, G., Liao, Y. Y., and Hsieh, H. L. (2015) Co-occurrence of juvenile horseshoe crabs *Tachypleus tridentatus* and *Carcinoscorpius rotundicauda* in an estuarine bay, Southwestern China. *Aquatic Biology*, 24(2), 117–126. doi: 10.3354/ab00641
- 19 Le Cren, F. D. (1951). The length-weight relationship and seasonal cycle in gonad weight I and condition in the perch (*Perca fluviatilis*). *Journal of Animal Ecology*, 20, 201–219.
- 11 Dolejš, P., and Vaňousová, K. (2015) A collection of horseshoe crabs (Chelicerata: Xiphosura) in the National Museum, Prague (Czech Republic) and a review of their immunological importance. *Arachnologische Mitteilungen*, 49, 1–9. doi: 10.5431/aramit4901
- 10 Fauziyah, Agustriani, F., Putri, W. A. E., Purwiyanto, A. I. S., and Suteja, Y. (2018) Composition and biodiversity of shrimp catch with trammel net in Banyuasin coastal waters of South Sumatera, Indonesia. *AACL Bioflux*, 11(5), 1515–1524.
- 12 Fauziyah, Purwiyanto, A. I. S., Putri, W. A. E., Agustriani, F., Mustopa, A. Z., and Fatimah. (2019) The first investigation record of threatened horseshoe crabs in the Banyuasin estuarine, South Sumatra, Indonesia. *Ecologica Montenegrina*, 24, 17–22.
- 13 Froese, R. (2006) Cube law, condition factor and weight-length relationships: History, meta-analysis and recommendations. *Journal of Applied Ichthyology*, 22(4), 241–253. doi: 10.1111/j.1439-0426.2006.00805.x
- Graham, L. J., Botton, M. L., Hata, D., Loveland, R. E., and Murphy, B. R. (2009) Prosomal-width-to-weight relationships in american Horseshoe crabs (*Limulus polyphemus*): Examining conversion factors used to estimate landings. *Fishery Bulletin*, 107(2), 235–243.
- 4 Hegele-drywa, J., Normant, M., Szwarc, B., and Pudlusk, A. (2014) Population structure, morphometry and individual condition of the non-native crab *Rhithropanopeus harrisii* (Gould, 1841), a recent coloniser of the Gulf of Gdańsk (southern Baltic Sea). *Oceanologia*, 56(4), 805–824. doi: 10.5697/oc.56-4.805
- 14 Jawahir, A. R. N., Samsur, M., Shabdin, M. L., and Rahim, K. A. (2017) Morphometric allometry of horseshoe crab, *Tachypleus gigas* at west part of Sarawak waters, Borneo, East Malaysia. *AACL Bioflux*, 10(1), 18–24.
- 5 Manca, A., Mohamad, F., Ahmad, A., Afham Mohd Sofa, M. F., and Ismail, N. (2017) Tri-spine horseshoe crab, *Tachypleus tridentatus* (L.) in Sabah, Malaysia: the adult body sizes and population estimate. *Journal of Asia-Pacific Biodiversity*, 10(3), 355–361. doi: 10.1016/j.japb.2017.04.011
- 18 Nair, P., Joseph, S., and Pillai, V. (2015) Length-weight relationship and relative condition factor of *Stolephorus commersonii* (Lacepede, 1803) exploited along Kerala coast. *Journal of the Marine Biological Association of India*, 57(2), 27–31. doi: 10.24124/jmbai.2015.57.2.01856-04
- 24 Panda, S., and Naik, P. K. (2017) Morphometric Study of Horseshoe Crab (*Carcinoscorpius rotundicauda*)

- in Odisha. *IJSS Journal of Surgery*, 3(2), 48–53. doi: 10.17354/SUR/2017/81
- Srijaya, T. C., Pradeep, P. J., Shaharom, F. M., and Chatterji, A. (2010) A New Record on the Morphometric Variations in the Populations of Horseshoe Crab (*Carcinoscorpius rotundicauda* Latreille) Obtained from Two Different Ecological Habitats of Peninsular Malaysia. *Our Nature*, 8, 204–211. doi: 10.3126/on.v8i1.4329
- Syuhaida, N., Rozihan, M., Akbar, J., Akmal, M., and Joni, H. (2019) Allometry relationship of mangrove horseshoe crab, *Carcinoscorpius rotundicauda* from the West Coast of Peninsular Malaysia. *International Journal of Fisheries and Aquatic Studies*, 7(2), 223–228.
- Tan, A. N., Christianus, A., Shakibazadeh, S., and Hajeb, P. (2012) Horseshoe Crab, *Tachypleus gigas* (Müller, 1785) Spawning Population at Balok Beach, Kuantan, Pahang, Malaysia. *Pakistan Journal of Biological Sciences*, 15(13), 610–620. doi: 10.3923/pjbs.2012.610.620
- Tanacredi, J. T., Botton, M. L., and Smith, D. R. (2009) *Biology and conservation of horseshoe crabs. Biology and Conservation of Horseshoe Crabs*. Heidelberg London: Springer. doi: 10.1007/978-0-387-89959-6
- Thomas, S. (2013) Allometric relationships of short neck clam *Paphia malabarica* from Dharmadam estuary, Kerala. *Journal of the Marine Biological Association of India*, 55(1), 50–54. doi: 10.6024/jmbai.2013.55.1.01755-08
- Vijayakumar, R., Das, S., Chatterji, A., and Parulekar, A. H. (2000) Morphometric characteristics in the horseshoe crab *Tachypleus gigas* (Arthropoda: Merostomata). *Indian Journal of Marine Sciences*, 29, 333–335.
- Webster, M. (2007) A Cambrian Peak in Morphological Variation Within Trilobite Species. *Science*, 317, 449–502. doi: 10.1126/science.1142964
- World Conservation Monitoring Centre. (1996) *Carcinoscorpius rotundicauda*. The IUCN Red List of Threatened Species 1996: e.T3856A10123044. doi: http://dx.doi.org/10.2305/IUCN.UK.1996.RLTS.T3856A10123044.en
- Yang, K. C., and Ko, H. S. (2015) First Record of Tri-spine Horseshoe Crab, *Tachypleus tridentatus* (Merostomata: Xiphosurida: Limulidae) from Korean Waters. *Animal Systematics, Evolution and Diversity*, 31(1), 42–45. doi: 10.5635/ased.2015.31.1.042

The morphometric variability of the mangrove horseshoe crab (*Carcinoscorpius rotundicauda*) from Banyuasin estuarine of South Sumatra, Indonesia

ORIGINALITY REPORT

20%

SIMILARITY INDEX

18%

INTERNET SOURCES

19%

PUBLICATIONS

18%

STUDENT PAPERS

PRIMARY SOURCES

1

www.vliz.be

Internet Source

2%

2

www.bioflux.com.ro

Internet Source

1%

3

www.frontiersin.org

Internet Source

1%

4

Submitted to Turun yliopisto

Student Paper

1%

5

Submitted to City University of Hong Kong

Student Paper

1%

6

Srijaya, T.C., P.J. Pradeep, S. Mithun, A. Hassan, F. Shaharom, and A. Chatterji. "A New Record on the Morphometric Variations in the Populations of Horseshoe Crab (*Carcinoscorpius rotundicauda* Latreille) Obtained from Two Different Ecological Habitats of Peninsular Malaysia", Our Nature, 2011.

Publication

1%

7	www.pereiraphc.com Internet Source	1%
8	Submitted to Wageningen University Student Paper	1%
9	Xiaoyong Xie, Zhou Wu, Chun-Chieh Wang, Yijian Fu et al. "Nursery habitat for Asian horseshoe crabs along the northern Beibu Gulf, China: Implications for conservation management under baseline gaps", Aquatic Conservation: Marine and Freshwater Ecosystems, 2019 Publication	1%
10	Fauziyah, Nurhayati, S M Bernas, A Putera, Y Suteja, F Agustiani. "Biodiversity of fish resources in Sungsang Estuaries of South Sumatra", IOP Conference Series: Earth and Environmental Science, 2019 Publication	1%
11	arages.de Internet Source	1%
12	Lusita Meilana, Qinhua Fang. "Local knowledge-based study on the status of horseshoe crabs along the Indonesian coast", Regional Studies in Marine Science, 2020 Publication	1%

Submitted to Southern Cross University

13

Student Paper

1 %

14

Submitted to International Islamic University
Malaysia

Student Paper

1 %

15

Submitted to University of Kent at Canterbury

Student Paper

1 %

16

www.iucnredlist.org

Internet Source

1 %

17

Submitted to University of Malta

Student Paper

1 %

18

eprints.cmfri.org.in

Internet Source

1 %

19

aquaticcommons.org

Internet Source

1 %

20

irep.iium.edu.my

Internet Source

1 %

21

"Changing Global Perspectives on Horseshoe
Crab Biology, Conservation and Management",
Springer Science and Business Media LLC,
2015

Publication

1 %

22

Submitted to Universiti Malaysia Sarawak

Student Paper

1 %

Exclude quotes On

Exclude matches < 1%

Exclude bibliography Off