# THE MAIN CAUSE OF HIV/AIDS INFECTION DISEASE

by Hamzah Hasyim

Submission date: 18-Feb-2022 09:56AM (UTC+0700)

**Submission ID:** 1765110141

File name: The\_Main\_Cause\_of\_HIVAIDS\_Infection\_Disease\_compressed.pdf (560.72K)

Word count: 8162

**Character count: 49763** 



Available online at: http://ejurnal.stikesprimanusantara.ac.id/

## Jurnal Kesebatan

| ISSN (Print) 2085-7098 | ISSN (Online) 2657-1366 |



Literature Review

### THE MAIN CAUSE OF HIV/AIDS INFECTION DISEASE



Aannisah Fauzaania <sup>1</sup>, Hamzah Hasyim<sup>2</sup>, Atik Wulandari<sup>3</sup>, Deasy Sumarni<sup>4</sup>, Aidil Fitrianto<sup>5</sup>, Bayu Anggriani<sup>6</sup>, Ani Meliyani<sup>7</sup>, Ahmad Taher<sup>8</sup>

1.2.3.4.5.67.8 Faculty of Public Health, Universitas Sriwijaya, Sumatera Selatan, Indonesia

#### INFORMASI ARTIKEL

Received: November 26, 2021 Revised: November 28, 2021 Accepted: Desember 02, 2021 Available online: Desember 07, 2021

#### KATA KUNCI

Morbidity; Neglected Tropical Disease

#### Korespondensi

#### Hamzah Hasyim

E-mail: hamzah@fkm.unsri.ac.id

#### ABSTRAK

Background: The emergence of Human Immunodeficiency Virus (HIV) has delivered large fitness demanding situations to youngsters and young people round the sector manifesting as biological, psychological, and social problems. This then have become a worldwide project which isn't always simplest depending on the fitness quarter, however additionally the complete authorities quarter in widespread due to the effect of HIV sickness which has the capacity to assault all groups.

Purpose: behavior a literature overview on HIV/AIDS. This literature overview will describe the primary reasons of HIV primarily based totally on a few dependable references.

Method: the use of a literature overview with the database used is Pubmed, a literature seek method with the keywords "HIV", "AIDs", and "HIV/AIDs". Result: Our meta-evaluation discovered that the pooled occurrence estimate of AUD amongst PLWHA changed into determined to be 29.80% (95% CI; 24.10–35.76). Conclusion: Results discovered that the primary motive of HIV contamination among men changed into the injection of narcotics, and withinside the girls it changed into sexual sex with an inflamed individual.

#### INTRODUCTION

In Southeast Asia, the range of humans dwelling with HIV is 1.6 million and 110,000 humans have died. For Indonesia, Nadia said, the anticipated HIV instances in Indonesia till 2020 are 543,a hundred humans.[1]. Although it has a tendency to fluctuate, statistics on HIV AIDS instances in Indonesia keeps to growth from yr to yr. As In the determine below, it is able to be visible that over the last 11 years the range of HIV instances in Indonesia reached its height in 2019, which turned into 50,282 instances.[2] Based on WHO statistics in 2019, there are 78% of latest HIV infections withinside the Asia Pacific region. For the very best AIDS case for 11 years the final in 2013, which turned into 12,214 instances.

Since 2008 HIV occurrence fee started out to upward thrust with the aid of using 10% and variety of latest contamination identified every yr multiplied with the aid of using 36% amongst every age and doubled amongst adults. Nationally, 67% of humans residing with HIV realize their status, 88% of them are on remedy and 86% of humans on remedy have viral suppression.[3] From a

DOI: http://dx.doi.org/10.35730/jk.v12i0.625

realistic standpoint, ac- complishing this purpose could be a great challenge. As men and women residing with HIV or susceptible to obtaining HIV who've get entry to to remedy and prevention maintain to enjoy the culmination of scien-tific advances, we need to now no longer take our points of interest off or waver in our pursuit of the closing purpose of finishing the epidemic as a worldwide fitness catastrophe.

Despite the sizeable demanding situations that intellectual fitness affords to HIV prevention and treatment, there are numerous critical and unmet possibilities to combine intellectual healthcare with HIV care. [4] Results found out that the primary motive of HIV contamination among adult males became the injection of narcotics, and withinside the ladies it became sexual sex with an inflamed individual. This literature evaluate will describe the primary reasons of HIV primarily based totally on a few dependable references.

#### **METHODS**

<u>Jurnal Kesehatan</u> is licensed under <u>CC BY-SA 4.0</u> © <u>Jurnal Kesehatan</u> The approach used is literature evaluation, literature evaluation is a artificial evaluation and precis of what is thought and unknown approximately the subject of a group of medical papers, inclusive of the present day place of work withinside the present knowledge [3]. The literature evaluation makes use of the Pubmed database (https://pubmed.ncbi.nlm.nih.gov/) as a supply of

article searches. The articles used are articles from the closing 10 years with a complete of 30 articles reviewed. The article seek method on Pubmed makes use of three keywords, namely: "HIV", "AIDs", "HIV/AIDs". The operating process of this literature evaluation may be visible in Figure 1 below:




Figure 1: Literature Review Work Procedure

#### RESULT AND DISCUSSION

After obtaining 30 articles, then extraction of the articles was carried out, the results of the extraction from these articles were as follows:

No	Title/Years	Objective of Research	Type of Research	Method of Collecting Data	Population and sample	Result
1	Trend of HIV/AIDS for the ultimate 26 years and predicting success of the ninety-ninety-ninety HIV prevention goals with the aid of using 2020 in Ethiopia: A time collection analysis/ Girum, Tadele. Wasie, Abebaw. Worku, Abdulsemed/2018	determine the fashion of HIV contamination for the final 26 years and to expect the achievements	-			Since 1995, new contamination has declined with the aid of using 81% and on the grounds that 2002; quantity of HIV instances has declined with the aid of using 35.5%. However, after exceptional decline for decades, on the grounds that 2008 HIV prevalence price commenced to upward push with the aid of using 10% and quantity of recent contamination recognized every yr multiplied with the aid of using 36% amongst every age and doubled amongst adults. ART insurance has multiplied with the aid of using 90% amongst all age and tripled amongst pregnant ladies inside 6 years. Nationally, 67% of humans dwelling with HIV recognize their status, 88% of them are on remedy and 86% of humans on remedy have viral suppression. As a

result, AIDS-associated loss of life declined with the aid of using seventy seven and 79% amongst all age and kids respectively. By 2020, 79% of humans dwelling with HIV will recognize their HIV status, of which 96-99% of HIV inflamed humans might be on ART and extra than 86% can have viral suppression. Conclusion: After exceptional decline, HIV contamination began out to growth withinside the previous couple of years amongst adults. With the cutting-edge trend, Ethiopia will gain the second one and 0.33 90% HIV targets, whilst the primary goal isn't attainable and with out reaching this overarching intention manipulate of the epidemic will now no longer be achieved. Therefore due interest is wanted to avoid the cutting-edge epidemics and prognosis of instances.[1] Ending the HIV/AIDS From a theoretical standpoint, Pandemic/ Robert the aim of finishing the HIV/ AIDS pandemic is Walter Eisinger Fauci, Anthony achievable; however, it'll S./2018 require extra and sustained assets to make to be had the already current medical advances on local, regional, na- tional, and worldwide levels. From a sensible standpoint, ac- complishing this aim might be a tremendous challenge. In this regard, the crucial additives withinside the attempt in the direction of attaining this maximum hard aim could likely be improvement of a reasonably powerful HIV vaccine to- gether with surest implementation of current remedy and prevention modalities (32). As humans residing with HIV or vulnerable to obtaining HIV who've get admission to to remedy and prevention maintain to enjoy the culmination of scien- tific advances, we ought to now no longer take our attractions off or waver in our pursuit of the remaining aim of finishing the epidemic as a worldwide fitness catastrophe. [2] Spending on health advanced stepped amassed Between 1995 and 2015, and HIV/AIDS: forward posted facts global health spending in line domestic health estimates of on home with capita grew at an annualised fee of 3·1% (95% uncertainty interval [UI] 3·1 spending and fitness spending fitness development spending, from via way of means of 1995 to 2015, assistance in 188 to 3.2), with increase being countries, 1995largest in upper-middle-profits supply, along from a 2015/Dieleman, Joseph numerous global places (5.4% in line with capita [UI 5·3-5·5]) and L., et al/2018 improvement of worldwide help for fitness, and, for agencies lower-middle-profits global places  $(4\cdot2\%)$  in line with capita  $[4\cdot2-4\cdot3]$ ). In 2015, the primary \$9.7 trillion (9.7 trillion to time, anticipated DOI: http://dx.doi.org/10.35730/jk.v12i0.625

HIV/AIDS

spending on spent on health worldwide. prevention and High-profits global places remedy and via spent \$6.5 trillion (6.4 way of means trillion to 6.5 trillion) or of supply of funding, for  $66 \cdot 3\%$  (66 · 0 to  $66 \cdot 5$ ) of the entire in 2015, even as lowprofits global places spent \$70.3 billion (69.3 billion to 188 countries 71·3 billion) or 0·7% (0·7 to 0.7). Between 1990 and 2017, development assist for health prolonged with the useful resource of the use of 394.7% (\$29.9 billion), with an estimated \$37.4 billion of development assist being disbursed for health in 2017, of which \$9.1 billion (24·2%) targeted HIV/AIDS. Between 2000 and 2015, \$562.6 billion (531.1 billion to 621.9 billion) have become spent on HIV/AIDS worldwide. Governments financed 57.6% (52.0 to 60·8) of that popular. Global HIV/AIDS spending peaked at 49.7 billion (46.2-54.7) in 2013, decreasing to \$48.9 billion (45.2 billion to 54.2 billion) in 2015. That year, low-profits and lower-middleprofits global places represented 74.6% of all HIV/AIDS disability-adjusted life-years, but without a doubt 36.6% (34.4 to 38.7) of popular HIV/AIDS spending. In 2015, \$9.3 billion (8.5 billion to 10·4 billion) or 19·0% (17·6 to 20·6) of HIV/AIDS financing have become spent on prevention, and \$27.3 billion (24·5 billion to 31·1 billion) or 55·8% (53·3 to 57.9) have become devoted to care and treatment.[3]
The RWHAP records display Reducing HIVthat HIV-associated final associated fitness disparities withinside results disparities amongst the Health Resources PLWH throughout unique and Services sociodemographic strata are Administration's Ryan White HIV/AIDS de- creasing. HRSA's efforts and projects defined formerly program/Mandsager. and the continued sports of Paul.Marier, RWHAP recipients and sub-Allison.Cohen, recipients were essential for Stacy.Fanning, figuring out disparities in Miranda. Hauck, care and results amongst Heather.Cheever. RWHAP cli- ents and Laura W./2018. figuring out techniques for lowering those disparities a number of the low-profits customers receiving RWHAPfunded services. Al- aleven though excellent enhancements were made, awesome variations remain. Continued consciousness and attempt may be vital for elimi- nating final results disparities amongst HIV-impacted pop- ulations served via way of means of the RWHAP[4] A general of 25 research with 25,154 contributors This evaluate PubMed. The occurrence of systematically EMBASE. alcohol use problems aimed to review Psych INFO systematically throughout evolved and amongst human

9.8 trillion) have become

	beings dwelling with HIV/AIDS: A systematic overview and meta- analysis/Duko, Bereket	evaluate the to be had research on the superiority of AUD amongst PLWHA and		and SCOPUS databases were		growing nations have been covered withinside the very last evaluation. Our meta- evaluation found out that the pooled incidence estimate of AUD amongst PLWHA
	Ayano, Getinet/2019	ahead feasible suggestions for destiny scientific exercise and research.				changed into located to be 29.80% (95% CI; 24.10–35.76). The incidence of AUD changed into better in males (26.90%) than female (13.37% HIV patients. In this study, the pooled incidence of AUD changed into extensively better (31.52%) whilst measured via way of means of Alcohol Use Disorders Identification Test (AUDIT) compared to Composite International Diagnostic Interview (CIDI) (13.51%). In addition, the pooled incidence of AUD changed into better withinside the evolved nations (42.09%) at the same time as decrease for growing nations
6	The Interplay of HIV and Autophagy in Early Infection/Cabrera- Rodríguez, Romina, et al/2021	The identity and evaluation of HIV-1 effective people that evidently manage viral replication withinside the absence of antiretroviral remedy has supplied clues approximately mobile tactics that might engage with viral proteins and RNA and outline next viral replication and medical progression	-	-	-	(24.52%)[5] Confirming this interaction, techniques primarily based totally on autophagy modulation are capable of inhibit early steps of HIV-1 contamination. Therefore, the intention of this evaluation is to summarize the information of the interaction among autophagy and the early activities of HIV-1 contamination, and the way autophagy modulation should impair or gain HIV-1 contamination and persistence, impacting viral pathogenesis, immune manage of viral replication, and medical development of HIV-1 inflamed patients.[6]
,	Mental fitness and HIV/AIDS: The want for an incorporated response/Remien, Robert H/2019	•	Editorial review	-	-	Mental fitness issues (starting from misery to SMI) are multiplied amongst humans atdanger for HIV and people dwelling with HIV. We have the vital assessment (screening equipment and efficacious remedies to deal with intellectual fitness issues amongst humans dwelling with and at danger for HIV. Despite the massive demanding situations that intellectual fitness affords to HIV prevention and treatment, there are numerous essential and unmet possibilities to combine intellectual healthcare with
8	Global Health Burden and Needs of Transgender Populations: A Review/Sari L. Reisner, ScD, et	-	Review	-	-	HIV care.[7] Transgender human beings are a various populace laid low with plenty of terrible fitness signs throughout high, middle, and coffee profits settings.[8]
)	al?2016  Investigating the contributing factors to HIV/AIDS	-	The gift take a look at	-	150 patient	Results observed that the precept purpose of HIV infection amongst person men

	infection from the perspective of HIV- infected patients/Mehraeen, Morteza Heydari, Mohammadreza Lankarani, Kamran B. Joulaei, Hassan Faghih, Marjan/2021		investigated the reasons of contracting HIV/AIDS from the angle of HIV-inflamed patients		modified into the injection of narcotics, and withinside the girls it modified into sexual intercourse with an infected individual. Meanwhile, 57% of the girls and 66% of the person men blamed themselves for contracting and transmitting the disease. The patients stated that if they could pass lower back to pre-infection period, they could use one of the following strategies to prevent the disease: (a) they is probably privy to hygienic/sanitary principles; (b) they could now not get married; and (c) they could prevent drug addiction. Also simplest 44% of the human beings had a achievement siblings (people who had been neither addicts nor HIV/AIDS-infected human beings), which modified into an remark that emphasizes on the epidemic of high-danger behaviors withinside the
10	Vitamin D status impacts genital mucosal immunity and markers of HIV-1 susceptibility in women/Anderson, Sharon M/2020	•	RCT		In order to serum food regimen D degrees, genital mucosal endpoints, collectively with soluble mediators, immune mobileular populations, gene expression, and ex vivo HIV-1 infection, were assessed. While systemic food regimen D degrees showed a large growth following supplementation, the ones changes translated into modest outcomes on the cervicovaginal factors studied. Paradoxically, post-supplementation food regimen D degrees were decreased in cervicovaginal fluids. Given the sturdy correlation amongst food regimen D reputation and HIV-1 infection and the large nature of food regimen D deficiency, further knowledge of the placement of food regimen D immunoregulation withinside the woman reproductive tract is important.[10]
11	Magnitude and predictors of not unusualplace intellectual disease amongst humans with HIV/AIDS in Ethiopia: A systematic assessment and meta-evaluation	accomplishing a scientific assessment and meta-evaluation of present literatures will have a paramount significance to reveal its summarized figure.	Meta Analysis	(PubMed/Medline, - Science Direct and PsycINFO	A wellknown of 13 primary studies examine the inclusion requirements were included in this systematic review. The pooled prevalence of now no longer unusualplace highbrow ailment modified into positioned to be 28.83% (95% CI: 17.93, 39.73) among people with HIV/AIDS in Ethiopia. The most prevalence of now no longer unusualplace highbrow ailment (35.20%) modified into determined among studies in which Kessler-10 modified into used as a screening tool. Single marital status (OR = 1.83; 95 percentCI: 1.03, 3.27), HIV/ AIDS-related stigma (OR = 2.21; 95 percentCI: 1.68,

						2.00) 1 1 1
						2.90) and modem interest unavailability (OR = 1.38; 95percentCI: 1.01, 1.88) had statistically big association with now no longer unusualplace highbrow ailment.[11]
12	Factors affecting discrimination in the direction of humans with HIV/AIDS in Sarawak, Malaysia/	The take a look at aimed to decide the perceived stage of discrimination closer to humans with HIV/AIDS a few of the grownup humans in Sarawak, Malaysia and additionally to decide the elements related to it	cross-sectional study	face-to-face interview using interviewer- guided questionnaires.	900 adults	The advise (SD) age for male and woman respondents have become 41.57 (13.45) and 38.99 (13.09) years respectively. The composite advise score for standard discrimination have become higher among woman (advise = 2.66, SD = 0.6) than male (advise = 2.47, SD = 0.6) respondents. Domain-clever discrimination of HIV/AIDS found that isolation, verbal discrimination, loss of resources, loss of services and standard discrimination had been drastically immoderate maximum of the females (p < 0.05) in assessment to males. Stepwise multiple linear regression assessment found that ethnicity, religion, understanding on HIV, an acquaintance of HIV and own circle of relatives profits appeared to be capacity predictors for discrimination withinside the course of peoples residing with HIV/AIDS (p < 0.05).
13	Prevalence of Depression and Anxiety Disorders in People Living with HIV/AIDS in a Tertiary Hospital in South Western Nigeria/	The goal of this take a look at became to decide the superiority of despair and tension ailment amongst HIV sufferers attending a tertiary organization in south western Nigeria	cross-sectional study	A semi- structured questionnaire	424	A widespread of 424 HIV exceptional have a examine people were recruited. A corresponding age and sex matched 429 manage subjects were moreover enrolled. The mean age of HIV exceptional patients and controls are 42.2 ± 9.5 years and 43.4 ± 12.4 years respectively. There have become a female prepondence among every have a examine populations. The incidence of melancholy among PLHIV have become 39.6% whilst melancholy have become lower in the (22.0%) manage business enterprise. Likewise anxiety have become reported in PLHIV and manage business enterprise as 32.6% and 28.7% respectively. Female gender, illiteracy, being divorced/widowed, unemployed and espresso CD4 remember were associated with melancholy while factors associated with anxiety illness protected lower age, female gender, low earnings, and espresso CD4 remember.[13]
14	Fertility preference in Iranian girls with HIV: A qualitative study/	The goal of this take a look at turned into to discover fertility intentions and reviews of	Quantitative	in-depth interviews were	-	Analysis of the which means gadgets of interviews confirmed issues in describing of fertility dreams of HIV ladies as follows: 1) Motherhood as a manner for balance of life; 2)

 $DOI: \underline{http://dx.doi.org/10.35730/jk.v12i0.625}$ 

		HIV-inflamed ladies in Iran				Uncertainties approximately the future; and 3) Unpleasant revel in of being pregnant and delivery. This subject matter has 3 sub-issues as stigma, discrimination and weak point in fitness care system.[14]
15	Personality modifications in HIV-inflamed sufferers who've been receiving notably energetic antiretroviral therapy/Dobrakowski, Pawel P. Skalski, Sebastian/2019		-			The examine proved there are statistically massive variations among the Actual self and the Retrospective self profiles regardig the want for change, submission, and the Adapted Child ego. The degree of shallowness in HIV-inflamed sufferers emerged po be sinificantly decrease while in comparison to different scientific manipulate groups. Patients with HIV suggested psychosocial advantages concerning their lifestyles philosophy, advanced man or woman strength, and advanced interpersonal relationships. The cognitive appraisal of the HIV contamination correlated extensively with positive persona tendencies in addition to the extent of psychosocial advantages, the period of ARV therapy, and the CD4
16	Stigma and gender measurement primarily based totally on rural and concrete groups in stopping HIV and AIDS in childbearing age couples in Banyumas Regency/Sistiarani, Colti  Hariyadi, Bambang  Munasib, Munasib  Sari, Septi M/2020	The intention of this have a look at turned into to map the stigma and examine the gen- der dimensions consistent with access, roles, benefit, and control. Material	cross-sectional study	-	194	mobileular count.[15]  Based at the research, maximum of the respondents had been among 36 and forty nine years old (66.33%). Men and ladies ratio had been nearly equal, ensuing in forty nine.74% and 50.26%, respectively. The majority of respondents had a excessive degree of education (39.37%), now no longer operating ended in 53.88%, whilst the ones having youngsters accounted for 67.87%. The effects of the take a look at found out that there has been no stigma discount in position and gender equality development consistent with intercourse category. There had been variations in position discount stigma among city and rural communities, and additionally variations withinside the position of gender equality development in city and rural areas.[16]
17	Social capital and HIV/AIDS withinside the United States: Knowledge, gaps, and destiny directions//Ransome, Yusuf Thurber, Katherine A. Swen, Melody Crawford, Natalie D. German, Danielle Dean, Lorraine T./2018	Social capital is a well-installed predictor of numerous behavioral fitness effects. However, we recognise much less approximately the connection with prevention, transmission, and remedy of HIV/AIDS	scoping review of	PubMed, Embase, PsycINFO, Web of Science, and Sociological Abstracts with out a restrict on ebook date, for articles in English language	-	We identified 1581 particular manuscripts and reviewed 13 based totally mostly on eligibility criteria. The earliest eli-gible test turn out to be published in 2003. More than half (n=7/13) focused on HIV or AIDS diagnosis, then prescribing ART and/or adherence (n=5/13), then linkage and or engagement in HIV care (n=4/13). Fifty eight percent (58%) documented a protective association amongst at least

		effects withinside the United States (US).				one social capital diploma and an HIV/AIDS out- come. Seven studies used demonstrated social capital scales, however there was big model in conceptual/ operational definitions and measures used. Most studies were based totally mostly on samples from the Northeast. Three studies right now focused on or stratified analyses among subgroups or key populations. Studies were cross-sectional, so causal inference is unknown.[17]
18	nutrients Coffee Intake and Neurocognitive Performance/	This examine aimed to examine the connection among ordinary espresso consumption and neurocognitive performance (NCP) in HIV-HCV coinfected people	cross-sectional survey.		-	Our effects confirmed significant, high quality institutions among improved espresso intake (ECI) (3 or extra cups of espresso in keeping with day) and NCP in verbal fluency, psychomotor speed (coding) and govt functioning. ECI may consequently hold neurocognitive functioning in humans dwelling with HIV and HCV [18]
19	Application of CRISPR/Cas9-primarily based totally gene enhancing in HIV- 1/AIDS therapy	-	-	-	-	Nevertheless, the difficulty of goal web website online and excessive off-goal consequences make it tough to be implemented withinside the HIV-1/AIDS gene remedy field. It is the primary time that the CCR5[19]
20	dyslipidemia Nutritional intervention reduces , fasting glucose and blood strain in humans residing with hiv/aids in antiretroviral therapy: A randomized medical trial evaluating dietary interventions/Silveira, Erika Aparecida/2020	observe aimed to assess the effectiveness of dietary interventions on numerous cardiometabolic parameters in PLWHA handled with ART	RCT	-	88	the two investigated nutritional interventions had been effective in lowering some cardiometabolic danger factors in PLWHA. However, after modifications for covariates, the individualized dietary prescription showed significant reductions withinside the primary very last consequences and, also, in more cardiometabolic danger factors than the nutritional counseling, [20]
21	Future technology for tracking HIV drug resistance and cure/Parikh, Urvi M. McCormick, Kevin Van Zyl, Gert Mellors, John W./2017	This evaluate modern and rising technology which are relevant for each surveillance of HIV drug resistance (HIVDR) and characterization of HIV reservoirs that persist no matter antiretroviral remedy and are barriers to curing HIV infection	-	-	-	Current populace genotyping techniques for resistance tracking are excessive fee and coffee throughput. NGS, mixed with less complicated pattern series and garage matrices (e.g. dried blood spots), has huge capacity to expand international surveillance and affected person tracking for HIVDR. Recent adaptions of NGS to become aware of integration webweb sites of HIV withinside the human genome and to represent the incorporated HIV provinuses are probable to facilitate investigations of the effect of experimental 'curative' interventions on HIV reservoirs [21]
22	HIV and aging: Role of the microbiome/Desai, Seema N. Landay, Alan L./2018	The motive of this newsletter is to check age-related changes in microbiota	-	-	-	Increasing evidence shows a large link in modifications withinside the composition, range and sensible additives of intestinal microbiome with regular developing vintage and

 $DOI: \underline{http://dx.doi.org/10.35730/jk.v12i0.625}$ 

		composition, variety and purposeful				HIV infection. Data on association of metabolites produced thru the microbiome
		capabilities in context of immune senescence, continual infection and comorbidities related to HIV infection.				with HIV-associated non-AIDS comorbidities is mounting. The impact of the microbiome adjustments on inflammation, immune and organ senescence and mechanisms thru which bio- behavioral pathways will exacerbate the ones outcomes desires to be in addition evaluated.[22]
23	HIV treatment and prevention 2019: Current standards of care/ Phanuphak, Nittaya Gulick, Roy M./2020	The reason of this assessment is to summarize the modern-day requirements of take care of each HIV remedy and HIV prevention in 2019	-	-		HIV remedy and HIV prevention have progressed markedly in current years because of the improvement of oral antiretrovirals which might be potent, convenient, and usually nicely tolerated, and cause virologic suppression and reduced HIV transmission.  Keywords [23]
24	Ethical Issues in HIV Phylogenetics/ Mehta, Sanjay Schairer, Cynthia Little, Susan/2019	HIV phylogenetic and molecular epidemiology (ME) analyses are more and more more being accomplished with a aim of enhancing HIV prevention efforts. However, ethical, criminal and social problems are related to those analyses, and ought to be taken into consideration whilst accomplished	-	-	-	HIV ME for research and public health pose huge ethical issues that preserve to comply with improving technology, prolonged sampling, and a changing jail and social climate. Guidance surrounding the ones issues wants to be advanced for researchers and public health officials in an iterative and area particular manner that cash owed for the functionality blessings and risks of this technology.[24]
25	HIV replication: A game of hide and sense/ Sauter, Daniel Kirchhoff, Frank/2016	-	Current opinion	-	-	HIV has advanced present day but imperfect mechanisms to live farfar from and counteract innate sensing. Whether virus-triggered immune activation represents honestly a suboptimal version of HIV to its human host or perhaps allows HIV replication, for example with the resource of the usage of developing the amount of viral purpose cells, remains to be clarified.[25]
226	Phamacovigilance and international HIV/AIDS/	This evaluate specializes in contemporary status, progress, demanding situations and possibilities in worldwide pharmacovigilance for HIV/AIDS treatment.	Review	-	-	Over 6 million HIV-infected human beings worldwide are on treatment with more than 100 fifty innovator and universal antiretroviral drug products. This achievement is made possible through sponsorship of HIV/AIDS applications the deliver of universal fixed-dose mixture products. The WHO prequalifies universal drug products used by Global Fund applications; america Food and Drug Administration tentatively approves universal products used by PEPFAR

						applications through a fast-tune evaluation process. Adequate countrywide or close by awesome and safety monitoring systems after drug distribution are lacking. The pharmaceutical sector, a big player in pharmacovigilance withinside the superior world, has now now not been engaged. Innovative techniques based mostly on collaboration and partnerships may be needed. Clinic-based completely or program-based completely cohort studies, randomized scientific trials and virtual clinical information also can moreover make a contribution pharmacovigilance-relevant information [26]
27	Molecular epidemiology withinside the HIV and SARS-CoV-2 pandemics/	The intention of this evaluate changed into to evaluate and assessment the utility of molecular epidemiology techniques for the progressed control and information of the HIV as opposed to SARS-CoV-2 epidemics	Review			Molecular biology techniques, which incorporates PCR and complete genome sequencing (WGS), have turn out to be powerful tools for epidemiological investigation. PCR techniques form the concept for lots high-sensitivity diagnostic tests and may supplement traditional contact tracing and surveillance strategies to define chance networks and transmission patterns. WGS techniques can in addition define the causative shops of illness, trace the origins of the pathogen, and make clean routes of transmission. When coupled with scientific datasets, consisting of virtual medical record data, the ones techniques can check out co-correlates of illness and pathogenesis. [27]
28	How HIV-1 access mechanism and extensively neutralizing antibodies manual structure- primarily based totally vaccine design	An HIV-1 vaccine that elicits widely neutralizing antibodies (bNAbs) stays to be developed. Here, we overview how understanding of bNAbs and HIV-1 access mechanism is guiding the structure-primarily based totally layout of vaccine immunogens and immunization regimens.	-			Isolation of bNAbs from HIV-  1-infected donors has induced an great knowledge of the webweb webweb sites of vulnerability that the ones antibodies intention on the HIV-1 envelope (Env) further to of the immunological pathways, which antibody lineages observe to boom massive and robust neutralization. Sites of vulnerability, however, stay withinside the context of severa envelope states required for HIV-1 entry, along with a prefusion-closed state, a single-CD4-sure intermediate, a three-CD4-sure intermediate, a and postfusion states, and it is not always easy which structural state optimally gives a particular internet site on-line of vulnerability
29	HIV subtype diversity worldwide/ Bbosa, Nicholas Kaleebu, Pontiano Ssemwanga, Deografius/2019	To offer a precis of the cutting-edge facts on the worldwide HIV subtype range	-	-	-	for vaccine elicitation.[28] The dynamic trade in HIV subtype distribution offers future stressful conditions for diagnosis, treatment and vaccine format and development. An increase in

 $DOI: \underline{http://dx.doi.org/10.35730/jk.v12i0.625}$ 

and distribution with the aid of using region. HIV is one of the maximum genetically numerous pathogens because of its high-mutation and recombination huge rates, populace length and speedy replication rate. This speedy evolutionary method has led numerous HIV subtypes which are heterogeneously globally distributed.

recombinant viruses suggests that coinfection and superinfection via divergent HIV strains has come to be extra now no longer unusualplace necessitating nonprevent surveillance to maintain music of the viral diversity. Cheaper near full-period genome sequencing strategies are essential in improving HIV subtype estimations. However, missing subtype statistics and espresso collection sampling tiers are nevertheless a undertaking in some geographical regions [29]

30 The courting among monetary reviews and HIV and AIDS remedy policies/ Economics, and in particular financial evaluations, are increasingly more being carried out provide treatment insurance guidance to desire makers. evaluations art work that has contributed to understanding of the relationship.

Consideration of the processes in which economic critiques could have greater effect over HIV and AIDS suggestions is needed. The inclined courting a few of the 2 shows the complicated and multifaceted decision-making technique that is often stimulated thru socioeconomic and political factors. If an economic evaluation is to steer insurance, then attention this is important. Extending the economic toolkit to include broader- based absolutely models that encompass political monetary machine variables, but do now now not compromise on comprehension, validity and robustness, will offer better informed insurance recommendations.[30]

However, after brilliant decline for decades, due to the fact 2008 HIV prevalence rate began to rise via 10% and huge range of recent infection diagnosed each twelve months elevated via 36% among each age and doubled among adults. Nationally, 67% of people residing with HIV apprehend their status, 88% of them are on treatment and 86% of people on treatment have viral suppression. From a sensible standpoint, ac- complishing this purpose can be a big challenge. As humans residing with HIV or prone to acquiring HIV who have get proper of access to to treatment and prevention hold to experience the end result of scien- tific advances, we should now now not take our sights off or waver in our pursuit of the very last purpose of completing the epidemic as a global health catastrophe. In 2015, \$9.7 trillion (9.7 trillion to 9.8 trillion) become spent on health global. Between 2000 and 2015,

\$562.6 billion (531.1 billion to 621.9 billion) become spent on HIV/AIDS global.

The RWHAP information show that HIV-related very last effects disparities among PLWH in the course of unique sociodemographic strata are de- creasing. Our meta-assessment observed out that the pooled prevalence estimate of AUD among PLWHA have become determined to be 29.80% (95% CI; 24.10–35.76). In addition, the pooled prevalence of AUD have become higher withinside the advanced countries (42.09%) whilst lower for developing countries (24.52%)[5]. Confirming this interplay, strategies based totally mostly on autophagy modulation are able to inhibit early steps of HIV-1 infection. Therefore, the aim of this assessment is to summarize the information of the interplay amongst autophagy and the early sports of HIV-1 infection, and the manner autophagy modulation may also need to impair or advantage HIV-1

infection and persistence, impacting viral pathogenesis, immune manipulate of viral replication, and medical improvement of HIV-1 infected patients.[6]. We have the essential assessment (screening) tools and efficacious treatments to cope with highbrow health problems among humans living with and at threat for HIV.

Despite the significant annoying conditions that highbrow health presents to HIV prevention and treatment, there are various vital and unmet opportunities to mix highbrow healthcare with HIV care.[7]. Results located that the number one cause of HIV infection amongst guys modified into the injection of narcotics, and withinside the ladies it modified into sexual intercourse with an infected individual. In addition to serum vitamins D levels, genital mucosal endpoints, which incorporates soluble mediators, immune mobileular populations, gene expression, and ex vivo HIV-1 infection, were assessed. A universal of 13 primary studies look at the inclusion requirements were covered in this systematic review. The pooled occurrence of now no longer unusualplace highbrow disease modified into positioned to be 28.83% (95% CI: 17.93, 39.73) among people with HIV/AIDS in Ethiopia.

The mean (SD) age for male and female respondents turn out to be 41.57 (13.45) and 38.99 (13.09) years respectively. The composite mean score for ordinary discrimination turn out to be higher among female. Domainrealistic discrimination of HIV/AIDS positioned that isolation, verbal discrimination, loss of resources, loss of services and ordinary discrimination had been substantially immoderate maximum of the females (p < 0.05) in contrast to males. Stepwise a couple of linear regression assessment determined out that ethnicity, religion, know-how on HIV, an acquaintance of HIV and own circle of relatives profits appeared to be functionality predictors for discrimination toward peoples living with HIV/AIDS (p < 0.05). A ordinary of 424 HIV great examine people had been recruited. The mean age of HIV great patients and controls are  $42.2 \pm 9.5$  years and  $43.4 \pm 12.4$ years respectively.

The prevalence of despair among PLHIV become 39.6% whilst despair become lower in the (22.0%) control group. Analysis of the meaning devices of interviews showed problems in describing of fertility desires of HIV women as follows: 1) Motherhood as a way for stability of life; 2) Uncertainties about the future; and 3) Unpleasant revel in of pregnancy and delivery. The diploma of vanity in HIV-infected patients emerged po be sinificantly lower even as in evaluation to one of a kind clinical control groups. The cognitive appraisal of the HIV infection correlated significantly with 668 Fauzaania, Aannisah, Et Al

wonderful individual dispositions similarly to the volume of psychosocial benefits, the period of ARV therapy, and the CD4 mobileular count.[15]. Based on the research, most of the respondents were amongst 36 and 40 9 years old (66.33%). The outcomes of the study found that there was no stigma bargain in function and gender equality improvement constant with sex category.

We identified 1581 precise manuscripts and reviewed 13 based mostly on eligibility criteria. The earliest eli- gible check emerge as published in 2003. Our effects showed significant, excessive first-class establishments amongst accelerated coffee intake (ECI) (three or more cups of coffee in keeping with day) and NCP in verbal fluency, psychomotor speed (coding) and authorities functioning. ECI may also therefore preserve neurocognitive functioning in people living with HIV and HCV [18]. Nevertheless, the quandary of purpose net internet site online and immoderate off-purpose effects make it hard to be carried out withinside the HIV-1/AIDS gene treatment field. the two investigated nutritional interventions have been effective in lowering some cardiometabolic chance factors in PLWHA. NGS, combined with simpler sample collection and storage matrices (e.g. Recent adaptions of NGS to find out integration webweb webweb sites of HIV withinside the human genome and to suggest the protected HIV proviruses are probable to facilitate investigations of the impact of experimental 'curative' interventions on HIV reservoirs [21].

Data on association of metabolites produced via the microbiome with HIV-associated non-AIDS comorbidities is mounting. The impact of the microbiome adjustments on inflammation, immune and organ senescence and mechanisms via which bio- behavioral pathways will exacerbate the ones outcomes needs to be in addition evaluated.[22]. HIV ME for research and public health pose considerable ethical issues that hold to comply with improving technology, increased sampling, and a changing crook and social climate. HIV has advanced trendy but imperfect mechanisms to keep away from and counteract innate sensing.

This achievement is made feasible thru sponsorship of HIV/AIDS programs [The Global Fund to Fight AIDS, Tuberculosis and Malaria; the President's Emergency Program for AIDS Relief (PEPFAR)] and the availability of not unusual fixed-dose mixture products. The pharmaceutical sector, a big player in pharmacovigilance withinside the superior world, has now not been engaged. WGS procedures can further define the causative stores of disease, trace the origins of the pathogen, and make clean routes of transmission. In the persevering with HIV epidemic, the ones procedures have been effectively

DOI: http://dx.doi.org/10.35730/jk.v12i0.625

deployed to choose out treatment gaps, transmission clusters and risk factors, aleven aleven though big boundaries to rapid or real-time implementation stay critical to overcome. Isolation of bNAbs from HIV-1-infected donors has triggered an unheard of expertise of the webweb webweb sites of vulnerability that the ones antibodies purpose on the HIV-1 envelope (Env) further to of the immunological pathways, which antibody lineages observe to enlarge huge and sturdy neutralization.

Furthermore, designated information of immunological pathways has led to talk about among vaccine developers as to how masses of the natural antibody-developmental pathway immunogens want to mimic, beginning from best the recognized epitope to the complete antibody-virus co-evolution process.[28]. The dynamic extrade in HIV subtype distribution gives future worrying conditions for diagnosis, treatment and vaccine format and development. Consideration of the processes in which economic evaluations ought to have extra have an impact on over HIV and AIDS suggestions is needed.

#### CONCLUSION

Our meta-evaluation found out that the pooled occurrence estimate of AUD amongst PLWHA become determined to be 29.80% (95% CI; 24.10–35.76). Results found out that the principle motive of HIV contamination among men become the injection of narcotics, and withinside the girls it become sexual sex with an inflamed individual.

#### REFERENCE

- [1] T. Girum, A. Wasie, and A. Worku, 'Trend of HIV/AIDS for the last 26 years and predicting achievement of the 90-90-90 HIV prevention targets by 2020 in Ethiopia: A time series analysis', *BMC Infect. Dis.*, vol. 18, no. 1, pp. 1–10, 2018, doi: 10.1186/s12879-018-3214-6.
- [2] R. W. Eisinger and A. S. Fauci, 'Ending the V/AIDS Pandemic', *Emerg. Microbes Infect.*, vol. 7, no. 1, pp. 413–416, 2018, doi: 10.1038/s41426-018-0062-9.
- [3] J. L. Dieleman et al., 'Spending on health and HIV/AIDS: domestic health spending and development assistance in 188 countries, 1995–2015', Lancet, vol. 391, no. 10132, pp. 1799–1829, 2018, doi: 10.1016/S0140-6736(18)30698-6.
- [4] P. Mandsager, A. Marier, S. Cohen, M. Fanning, H. Hauck, and L. W. Cheever, 'Reducing HIV-related health disparities in the Health Resources and Services Administration's Ryan White HIV/AIDS program', Am. J. Public Health, vol. 108, pp. S246–S250, 2018, doi: 10.2105/AJPH.2018.304689.
- B. Duko, M. Ayalew, and G. Ayano, 'The prevalence of alcohol use disorders among people living with HIV/AIDS: A systematic review and meta-analysis',

- Subst. Abus. Treat. Prev. Policy, vol. 14, no. 1, pp. 1–9, 2019, doi: 10.1186/s13011-019-0240-3.
- [6] R. Cabrera-Rodríguez et al., 'The In play of HIV and Autophagy in Early Infection', Front. Microbiol., vol. 12, no. April, pp. 1–27, 2021, doi: 10.3389/fmicb.2021.661446.
- [7] R. H. Remien, M. J. Stirratt, N. Nguyen, R. N. Robbins, A. N. Pala, and C. A. Mellins, 'Mental health and HIV/AIDS: The need for an integrated response', *Aids*, vol. 33, no. 9, pp. 1411–1420, 2019, doi: 10.1097/OAD.0000000000002227.
- [8] S. Sari L. Reisner, 'Global Health Burden and Needs of Transgender Populations: A Review', *Physiol. Behav.*, vol. 176, no. 1, pp. 100–106, 2016, doi: 10.1016/S0140-6736(16)00684-X.Global.
- [9] M. Mehraeen, M. Heydari, K. B. Lankarani, H. Joulaei, and M. Faghih, 'Investigating the contributing factors to HIV/AIDS infection from the perspective of HIV-infected patients', *BMC Psychol.*, vol. 9, no. 1, pp. 1–8, 2021, doi: 10.1186/s40359-021-00513-w.
- [10] S. M. Anderson et al., 'Vitamin D status impacts genital mucosal immunity and markers of HIV-1 susceptibility in women', *Nutrients*, vol. 12, no. 10, pp. 1–15, 2020, doi: 10.3390/nu12103176.
- [11] Z. Belayneh, B. Mekuriaw, T. Mehare, S. Shumy, and M. Tsehay, 'Magnitude and predictors of common mental disorder among people with HIV/AIDS in Ethiopia: A systematic review and meta-analysis', *BMC Public Health*, vol. 20, no. 1, pp. 1–11, 2020, doi: 10.1186/s12889-020-08800-8.
- [12] M. M. Rahman, A. S. Lemin, and C. A. Pangarah, 'Factors affecting discrimination toward people with HIV/AIDS in Sarawak, Malaysia', HIV AIDS Rev., vol. 9, no. 1, pp. 49–55, 2020, doi: 10.5114/hivar.2019.85916.
- [13] A. Olatayo Adeoti, M. U. Dada, and J. O. Fadare, 'Prevalence of Depression and Anxiety Disorders in People Living with HIV/AIDS in a Tertiary Hospital in South Western Nigeria', *Med. Reports Case Stud.*, vol. 03, no. 01, pp. 3–7, 2018, doi: 10.4172/2572-5130.1000150.
- [14] Z. Behboodi-Moghadam, A. Nikbakht-Nasrabadi, A. Ebadi, S. Esmaelzadeh–Saeieh, and M. Mohraz, 'Fertility desire in Iranian women with HIV: A qualitative study', Iran. J. Public Health, vol. 44, no. 8, pp. 1126–1134, 2015.
- [15] P. P. Dobrakowski and S. Skalski, 'Personality changes in HIV-infected patients who have been receiving highly active antiretroviral therapy', HIV AIDS Rev., vol. 18, no. 3, pp. 165–175, 2019, doi: 10.5114/hivar.2019.88445.
- [16] C. Sistiarani, B. Hariyadi, M. Munasib, and S. M. Sari, 'Stigma and gender dimension based on rural and urban communities in preventing HIV and AIDS in childbearing age couples in Banyumas Regency', *HIV AIDS Rev.*, vol. 9, no. 1, pp. 61–66, 2020, doi: 10.5114/hivar.2020.93178.
- [17] Y. Ransome, K. A. Thurber, M. Swen, N. D. Crawford, D. German, and L. T. Dean, 'Social capital and HIV/AIDS in the United States: Knowledge, gaps, and future directions', SSM Popul. Heal., vol. 5, no. May, pp. 73–85, 2018, doi: 10.1016/j.ssmph.2018.05.007.
- [18] S. Antwerpes et al., 'nutrients Coffee Intake and Neurocognitive Performance', pp. 1–13, 2020.

- [19] Q. Xiao, D. Guo, and S. Chen, 'Application of CRISPR/2 s9-based gene editing in HIV-1/AIDS therapy', Front. Cell. Infect. Microbiol., vol. 9, no. MAR, pp. 1–15, 2019, doi: 10.3389/fcimb.2019.00069.
- [20] E. A. Silveira, M. O. Falco, A. S. e. A. de C. Santos, M. Noll, and C. de Oliveira, 'Nutritional intervention reduces dyslipidemia, fasting glucose and blood pressure in people living with hiv/aids in antiretroviral therapy: A randomized clinical trial comparing two nutritional interventions', *Nutrients*, vol. 12, no. 10, pp. 1–18, 2020, doi: 10.3390/nu12102970.
- [21] U. M. Parikh, K. McCormick, G. Van Zyl, and J. W. Mellors, 'Future technologies for monitoring HIV drug resistance and cure', Curr. Opin. HIV AIDS, vol. 12, no. 2, pp. 182–189, 2017, doi: 10.1097/COH.000000000000344.
- [22] S. N. Desai and A. L. Landay, 'HIV and aging: Role of the microbiome', Curr. Opin. HIV AIDS, vol. 13, no. 1, pp. 22–27, 2018, doi: 10.1097/COH.000000000000433.
- [23] N. Phanuphak and R. M. Gulick, 'HIV treatment and prevention 2019: Current standards of care', *Curr. Opin. HIV AIDS*, vol. 15, no. 1, pp. 4–12, 2020, doi: 10.1097/COH.0000000000000588.
- [24] S. Mehta, C. Schairer, and S. Little, 'Ethical Issues in HIV Phylogenetics', *Physiol. Behav.*, vol. 176, no. 1, pp. 139–148, 2019, doi: 10.1097/COH.0000000000000538.Ethical.

- [25] D. Sauter and F. Kirchhoff, 'HIV replication: A game of hide and sense', *Curr. Opin. HIV AIDS*, vol. 11, no. 2, pp. 173–181, 2016, doi: 10.1097/COH.00000000000000233.
- [26] V. Miller, J. Nwokike, and A. Stergachis, 'Pharmacovigilance and global HIV/AIDS', Curr. Opin. HIV AIDS, vol. 7, no. 4, pp. 299–304, 2012, doi: 10.1097/COH.0b013e328354d8e7.
- [27] R. Lorenzo-Redondo, E. A. Ozer, C. J. Achenbach, R. T. D'Aquila, and J. F. Hultquist, 'Molecular epidemiology in the HIV and SARS-CoV-2 pandemics', *Curr. Opin. HIV AIDS*, vol. 16, no. 1, pp. 11–24, 2021, doi: 10.1097/COH.0000000000000660.
- [28] M. Pancera, A. Changela, and P. D. Kwong, 'How HIV-1 entry mechanism and broadly neutralizing antibodies guide structure-based vaccine design', vol. 12, no. 3, pp. 229–240, 2018, doi: 10.1097/COH.0000000000000360.How.
- [29] N. Bbosa, P. Kaleebu, and D. Ssemwanga, 'HIV subtype diversity worldwide', Curr. Opin. HIV AIDS, vol. 14, no. 3, pp. 153–160, 2019, doi: 10.1097/COH.000000000000534.
- [30] S. J. Taleski, K. Ahmed, and A. Whiteside, 'The relationship between economic evaluations and HIV and AIDS treatment policies', *Curr. Opin. HIV AIDS*, vol. 5, no. 3, pp. 204–209, 2010, doi: 10.1097/COH.0b013e3283384b58.

DOI: http://dx.doi.org/10.35730/jk.v12i0.625

## THE MAIN CAUSE OF HIV/AIDS INFECTION DISEASE

**ORIGINALITY REPORT** 

%
SIMILARITY INDEX

3%
INTERNET SOURCES

0% PUBLICATIONS

**PRIMARY SOURCES** 

1

bmcinfectdis.biomedcentral.com

Internet Source

1 %

2

etheses.whiterose.ac.uk

Internet Source

**1** %

Exclude quotes

On

Exclude matches

< 1%

Exclude bibliography