




International Symposium on

“Marine Ecosystems, Natural Products And Their Bioactive Metabolites

25-27 October 2011
Bogor, INDONESIA



PROGRAMME & ABSTRACTS



2011-IPB-INOC



**Inter-Islamic Science & Technology
Network on Oceanography
TURKEY**



**Faculty of Fisheries & Marine Science
Bogor Agricultural University
INDONESIA**

International Symposium

on

“Marine Ecosystems, Natural Products and their Bioactive Metabolites”

25-27 October 2011, Bogor, INDONESIA

PROCEEDINGS

Edited par: A. CHOUIKHI



INOC-DECEMBRE-2011

**PROCEEDINGS of the INTERNATIONAL Symposium on
“Marine Ecosystems, Natural Products and their Bioactive Metabolites”
25-27 October 2011, Bogor-INDONESIA.**

Actes du Symposium INTERNATIONAL sur
" Les écosystèmes marins, Produits naturels et leurs métabolites bioactifs ”
25-27 Octobre 2011, Bogor-INDONESIE

International Symposium

on

“Marine Ecosystems, Natural Products and their Bioactive Metabolites”

25-27 October 2011, Bogor- INDONESIA

TABLE OF CONTENTS	Page
Biological and biomedical properties, Biotechnology of Bioactive Marine Natural Products, Bioactive marine toxins	
<u>Ahsana D Farooq, Talat Roome, Sabira Naqvi & Muhammad I Choudhary</u> <u>Mangroves: rich source of pharmacological activities and needs attention.</u> <u>Aegiceras corniculatum a rich source of anti-inflammatory properties</u>	01
<u>Janjarus Watanachote, Supanee Leethochavalit, Nareerat Rittirut</u> <u>Antibacterial proteins and agglutinin of the innate immune system from skin mucus of marine fishes in Thailand</u>	10
<u>Jalal, K.C.A., Akbar John, B., Suffi, N.H., Md. Hazman, M.S., Deny, S., Kamaruzzaman, B.Y.</u> <u>Chitin and Chitosan from potential sea foods of Malaysia- A comparative study</u>	18
<u>Samia K. Mikhail & Wagdy Labib</u> <u>High protein and β-carotene yield of <i>Dunaliella</i> spp. exposed to high light intensities and nutrient limitation</u>	23
<u>Melki A., Zaenal Mustopa, Hefni Effendi</u> <u>Bio-Potentials activity of <i>Sonneratia caseolaris</i> (mangrove) extract as antibacterial collected from South SUMATERA.</u>	34
<u>Hina Saeed Baig & Ahsana Dar Farooq</u> <u>Seasonal variation in anti-inflammatory activity of two species of <i>Sargassum</i> from Northern Arabian Sea bordering PAKISTAN.</u>	39

<u>Janjarus Watanachote , Nareerat Rittirut Supanee Leethochavalit ,</u> <u>Characterization of <i>Cryptocaryon</i> sp. isolated from marine fish in THAILAND and in vitro treatment.</u>	49
<u>Tahere Bagheri, Mohammad Reza Imanpoor, Aliakbar Hedayati</u> <u>Hematological responses of Persian sturgeon, <i>Acipenser persicus</i>, to anesthetic effects of clove essence.</u>	60
<u>Samia K. Mikhail & Wagdy Labib</u> <u><i>Achnanthes longipes</i> Agardh a biofouling organism: An experimental study.</u>	69
THEME 2: Marine Biodiversity	
<u>Fadila Tazerouti, Lassad Neifar, & Louis Euzet</u> <u>The enigma of <i>Monocotyle Myliobatis</i> Taschenberg, 1878 (Monogenea, Monocotylidae) is finally resolved: <i>Monocotyle myliobatis</i> rediscovered from off the type host <i>Myliobatis aquila</i> (Elasmobranchii, Myliobatidae) from Algerian coasts.</u>	78
<u>BELAIFA B. , BOUAMRA A. , ARAB A.</u> <u>Variation du régime alimentaire du Sandre (<i>Sander lucioperca</i>.)</u>	86
<u>Benmalek Yamina</u> <u>Study on halophilic bacteria isolated from the Mediterranean Sea.</u>	98
<u>Isdy Sulisty</u> <u>Performance reproductive du Tambour <i>Otolithes ruber</i> (Perciforme, Sciaenidae) des eaux de Tegal (Mer de Java) selon les critères morpho-anatomiques et histologiques.</u>	106
<u>Tazerouti Fadila</u> <u>La parasitofaune des sélaciens torpedinidae des côtes algériennes : biodiversité et spécificité parasitaire.</u>	112
<u>Sahar Fahmy Mehanna</u> <u>Population dynamics and management of Snubnose emperor <i>Lethrinus bungus</i> (<i>L. borbonicus</i>) from the Foul Bay, Red Sea.</u>	121

<u>Mahsa Haghi, Ahmad Savari, Preeta Kochanian, Seyed Mohammad Bagher Nabavi, Mohammad Hasan Nami, Mohammad Zakeri</u> <u>Shallow- water macrobenthos assemblages in relation to sediment characteristic in the south Qeshm Island, Persian Gulf.</u>	129
<u>Aymen HADJ TAIEB , Mohamed GHORBEL, Nader BEN HADJ HAMIDA & Othman JARBOUI</u> <u>Etude de la biométrie de <i>DIPLODUS VULGARIS</i> (E.Geoffry ST.-Hillaire, 1817) des côtes Sud de la TUNISIE.</u>	137
<u>Mahnaz Rabbaniha, Gholamreza Izadpanahi, Fereidoon Owfi, Fatemeh Mohsenizadeh</u> <u>Plankton community assemblage in surface layers of Northern part of the Persian Gulf (IRAN - Bushehr area), using by PCA.</u>	143
<u>EL HAOUATI H., ARAB A</u> <u>Etude de la flore diatomique du lac de Réghaia (Algérie).</u>	150
<u>Mohammad A. Salari-Aliabadi, Ehsan Abedi</u> <u>Population genetic structure of <i>Scomberomorus commerson</i> using microsatellite markers in the Persian Gulf.</u>	159
<u>Asma Daryanavard, Asghar Abdoli, Bahram Kiabi</u> <u>Spatial and temporal variation in the Ichthyofauna of rocky tidal pools from Qeshm Island.</u>	167
<u>Zeinalipour Mohammad, Javanshir Arash & Seyfabadi Sayed Jaafar</u> <u>The Comparison of bivalve, <i>Mytilaster lineatus</i>, growth and recruitment in three coastal regions (Amirabad, Khazarabad and Noor) of Mazandaran.</u>	173
<u>Mahsa Haghi; Ahmad Savari; Preeta Kochanian; Mohammad Bagher Nabavi; Mohammad Hasan Nami; Mohammad Zakeri</u> <u>Benthic habitat mapping on the south coast of the Qeshm Island Geopark, Persian Gulf, based on single beam acoustics.</u>	183
Theme 3: Ecology, Biological & Chemical investigations	
<u>Suijat Al Azad, EjriaSaleh and Justin Hii Yung Siang</u> <u>Effects of human induced activities on the sediment characteristics of Sulaman Bay lagoon, Kota Kinabalu, SABAH, MALAYSIA.</u>	191
<u>Hosny Ibrahim Emar</u>	199

<u>Seasonal variations of Aliphatic hydrocarbons in Mediterranean seawater West of Alexandria.</u>	
<u>Fereidoon Owfi, Mahnaz Rabbaniha, Zeinab Ansari, Morvarid Rahimi, Maisam Toosi</u> <u>Ecological classification of intertidal habitats (Iranian coasts of the Persian Gulf & Oman Sea), using by CMECS Model.</u>	206
<u>S. Lakkis, V. Novel Lakkis</u> <u>Global Warming inducing “Tropicalisation” of the East Mediterranean, Case study: Invasion of Marine Tropical species in the Levantine Basin.</u>	217
<u>Aliakbar Hedayati, Abdolali Movahedinia and Tahere Bagheri</u> <u>Enzymatic responses of Yellowfin Seabream (<i>Acanthopagrus latus</i>) during exposure to chronic mercury in Mahshahr creeks.</u>	226
<u>Kambiz Jahanbin and Aliakbar Hedayati</u> <u>The first application of a new prebiotic for the health improvement of Atlantic salmon exposed to Mercury chloride.</u>	233
<u>Nemati H. , Shokri MR., Pazoopki J.</u> <u>Are beach seine fishing activities altering benthic communities in Caspian Sea?</u>	241
<u>Amir Jaferian, Hossein Zolgharnein, Mehdi Mohammadi, Mohamad-Ali Salari-Aliabadi, Seyed-Javad Hossini</u> <u>Fourfinger threadfin fish (<i>Eleutheronema tetradactylum</i>) in Persian Gulf: weight length model, morphological specification & optimized method of genomic DNA isolation.</u>	245
<u>Aliakbar Hedayati, Alireza Safahieh, Tahere Bagheri , Abdolali Movahedinia</u> <u>Effect of sub-lethal Mercury toxicity on hormonal indices of Yellowfin Seabream.</u>	255
<u>Mostafa M. El-Sheekh , Hanan M. Khairy , Rania El-Shenody</u> <u>Influence of the production of extra and intra-cellular polysaccharides from some algal species under toxic <i>Microcystis aeruginosa</i> (Kützing) crude extracts.</u>	263
<u>Fatemeh Pourjomeh, Mohammad Reza Shokri and Bahram Kiabi</u> <u>Running head: Surrogacy value of artificial reefs for natural reefs. Are cement boulders mimicking natural boulders for macrobenthos in southern Caspian</u>	274

Sea?	
Mohammad Zakeri, Jasem G. Marammazi , Preeta Kochanian Dynamics of total lipids and fatty acids composition during endogenous period in yellowfin seabream (<i>Acanthopagrus latus</i>) eggs and larvae.	283
Theme 4. Marine Aquaculture, Fisheries, Oceanography and Climate Change	
Tahere Bagheri, Mohammad Reza Imanpoor , Aliakbar Hedayati Replacement of soybean meal as alternative protein sources for improvement of marine aquaculture.	291
Mohammad Al-Zibdah, Mohammad Amir and Sayeda Mier Quality of rearing water and the growth profile of Gilthead Bream utilizing mega flow re-circulating aquaculture system (RAS).	298
BOUAMRA A., BELAIFA B., ARAB A. Dynamique de la population du Sandre (<i>Sander lucioperca</i>).	305
Mohamed El Houssein Ould Mohamed La pêche en Mauritanie: état des lieux et perspectives.	317
Lakkis S. & Sabour W. Marine Fish Resources in Lebanese coastal waters.	329
CLOSING SESSION : Recommendations	340
Opening statement By Dr. H.H. Kouyoumjian	341
Closure address by Prof.Dr. A. Chouikhi	342
List of Participants	343
Photos: participants to the symposium	358

BIO-POTENTIALS ACTIVITY OF *Sonneratia caseolaris* (MANGROVE) EXTRACT AS ANTIBACTERIAL COLLECTED FROM SOUTH SUMATERA

Melki ⁽¹⁾, A Zaenal Mustopa ⁽²⁾, Hefni Effendi ⁽³⁾

¹ (M.Si) (Department of Marine Science, Sriwijaya University) – (Jalan Raya Palembang Prabumulih Km. 32 Inderalaya, 30662) – (South Sumatera-Indonesia) – (Telp/fax. +62711581118) – (www.unsri.ac.id) – (melki@mipa.unsri.ac.id)

² (M.Si) (Research Center for Biotechnology, Indonesian Institut of Sciences) – (Jalan Raya Bogor Km 46, Cobinong, 16911) – (Cibonong-Bogor, Indonesia) – (Telp. +6221-8754587, Fax. +6221-8754588) – (www.biotek.lipi.go.id) – (azmustopa@yahoo.com)

³ (Dr.rer.nat) (Department of Aquatic Resources Management, Faculty of Fisheries and Marine Science, Bogor Agricultural University) – (Kampus IPB Darmaga, Bogor 16680) – (Bogor, Indonesia) – (Telp. +622518621262, Fax. +62251 8622134) – (www.ipb.ac.id) – (hefni_effendi@yahoo.com)

Summary

Crude extracts of four mangrove species (leaf, fruit, bark and root), i.e. *Avicennia alba*, *A. marina*, *Rhizophora mucronata*, and *Sonneratia caseolaris* collected from Teluk Payo, Banyuasin, South Sumatera was extracted in methanol, ethyl acetate, hexane and tested for antibacterial (*Escherichia coli* and *Staphylococcus aureus* pathogen), and brine shrimp cytotoxic assay. The highest activity recorded was methanol extract of *S. caseolaris* in *E. coli* isolates (18 mm inhibition) and in *S. aureus* isolates (19 mm inhibition), exhibiting relatively high biopotency. Brine Shrimp Lethality Test showed that leaf of *S. caseolaris* methanol extract was not toxic to *Artemia salina*. The high bioactive mangrove extract evaluated further by HPLC showed that mangrove extracts likely contains flavonoid.

Keywords: Extract of *Sonneratia caseolaris*, antibacterial, BSLT, column chromatography, TLC, and HPLC

Introduction

Microorganisms have potential to cause human diseases. Most of the time viruses, bacteria and fungi act as major pathogenic organisms. The discovery of antibiotics in the early twentieth century provided an increasingly important tool to combat bacterial diseases. As antibiotics are increasingly used and misused, the bacterial strains become resistant to antibiotics rapidly. Therefore, antibacterial activity of medicinal plants is very important since vast number of medicinal plants have been used for centuries as remedies for human diseases. Among them extracts from different parts of mangroves and mangrove associates are widely used throughout the world. For instance, stem of *Avicennia marina* is used for ulcers and bark of *Bruguiera sexangula* is used for antitumors. Mangrove and mangrove associates contain biologically active antiviral, antibacterial and antifungal. They provide a rich source of steroids, triterpenes, saponins, flavonoids, alkaloids and tannins [1]. Therefore, it is worth to screen mangrove plants for the presence of new antibacterial compounds to combat the normal pathogenic bacterial strains and hospital acquired antibiotic resistant bacterial strains.

Mangroves are one of the easiest tropical forest types to generate. They have the ability to grow where no other vascular plants can. The mangroves exist under stressful conditions such as violent environments, high concentration of moisture, high and low tides of water, and abundant living microorganisms and insects. They thrive in a very peculiar environment and serve as a bridging ecosystem between freshwater and marine systems. They possess an unusual morphology and physiognomy and the path of photosynthesis in mangroves is different from other glycophytes. They possess modifications to establish water and salt economy. There are modifications or alterations in other physiological processes such as carbohydrate metabolism or polyphenol synthesis and due to these reasons, they may have chemical compounds, which protect them from these destructive elements [2].

Mangroves from the coast of Teluk Payo, South Sumatra province were collected. Their extract of leaf, fruit, bark and root was scrutinized to determine their bioactive using *Escherichia coli* and *Staphylococcus aureus* and brine shrimp cytotoxic as target organism.

Materials and Methods

Collection and Extraction of Mangrove Bioactive

Four species of mangroves i.e. *Avicenna marina*, *A. alba*, *Rhizophora mucronata* and *Sonneratia caseolaris* were collected and identified from the mangrove forest in Teluk Payo, South Sumatra during December 2009 (**Figure 1**). Prior to the extraction, leaves, fruits, barks and roots of respective species were cleaned, shade dried in order to prevent photolysis and thermal degradation, then chopped into small pieces and ground coarsely in a mechanical grinder.




Figure 1. Map showing the study area, Teluk Payo, South Sumatra (Indonesia)

Extraction of Bioactive

Around 100 g of powdered mangrove material was extracted for 3 x 24 hour using methanol 80%, ethyl acetate 80% and hexane 80% as much 250 ml. The extracts were filtered using Whatman no. 1 filter paper. The fraction was evaporated at rotary evaporator at 40 - 50°C, then collected in airtight plastic vials and stored in the refrigerator for further studies.

Bioassays

Antimicrobial assay was carried out as described by Abesinghe and Wanigatunge [3] against *Escherichia coli* NBRC 13276 and *Staphylococcus aureus* NBRC 14237 pathogen. The cytotoxic activity of mangrove extracts was tested against freshly hatched free-swimming nauplii of *Artemia salina* Leach. The assay system was prepared with 3 ml of filtered seawater containing chosen concentration of mangroves extract in cavity blocks (embryo cup) and 10 nauplii each was transferred in experimental, vehicle control and negative control wells. Invariably the concentration of the experimental systems was determined on the basis of exploratory experiments. The percentage of mortality was determined by comparing the mean surviving larvae of the test and control tubes. The LC₅₀ value was determined using probit scale [4].

Column Chromatography and Thin Layer Chromatography (TLC)

The mangroves extract (2 ml) was loaded on a silica gel 60 Fe₂₅₄ (Merck) column packed with chloroform and eluted with chloroform and methanol (9:1 to 1:9) to yield fractions. The potential fractions were then examined by TLC on 25 TLC alluminium silica gel 60 Fe₂₅₄ (Merck) using same solvent with column chromatography. Individual fractions were collected and tested for bioactivity.

High Performance Liquid Chromatography (HPLC)

Mangrove extracts derived from column chromatography were injected to HPLC KNAUER, column Eurospher100-5C₁₈ (4.6 x 150 mm, 5 µm), injection volume 20 µl, PDA detector with 200-400 nm, flow rate of gradient 1 ml/minute, methanol:water (v/v) (at 0 minute 0% water, at 22 minute 100% methanol, at 30 minute 100% methanol, at 33 minute 100% water, at 40 minute 100% water).

Results and Discussion

Extraction of Component Bioactive

Methanol solvent produced more weight extract percentage than that of ethyl acetate and hexane solvent (**Table 1**).

Table 1. Mangrove Extracted with MeOH, EtOAc, and Hexane

Species of Mangrove	Part	Original weight (g)	Weight Extract (g)			Crude Extract (% b/v)		
			MeOH	EtOAc	Hexane	MeOH	EtOAc	Hexane
<i>A. marina</i>	Leaf	100	7.71	6.80	3.34	7.71	6.80	3.34
	Bark	100	4.22	4.08	2.21	4.22	4.08	2.21
	Fruit	100	11.8	10.89	2.23	11.80	10.89	2.23
	Root	100	5.22	4.72	2.56	5.22	4.72	2.56
<i>A. alba</i>	Leaf	100	7.93	7.23	4.23	7.93	7.23	4.23
	Bark	100	9.34	8.67	4.21	9.34	8.67	4.21
	Fruit	100	10.31	9.98	4.22	10.31	9.98	4.22
	Root	100	8.30	7.56	3.67	8.30	7.56	3.67
<i>R. mucronata</i>	Leaf	100	10.89	9.79	2.58	10.89	9.79	2.58
	Bark	100	6.73	5.56	2.78	6.73	5.56	2.78
	Root	100	6.75	6.34	3.53	6.75	6.34	3.53
<i>S. caseolaris</i>	Leaf	100	8.67	6.65	3.45	8.67	6.65	3.45
	Bark	100	6.89	6.34	3.12	6.89	6.34	3.12
	Fruit	100	8.90	7.54	3.45	8.90	7.54	3.45
	Root	100	9.78	8.67	4.34	9.78	8.67	4.34

Antibacterial Activity

Crude extracts of mangrove (leaf, fruit, bark and root) had the ability to inhibit bacterial growth of *E. coli* and *S. aureus*. The invitro antibacterial activity revealed that methanol extract of mangroves had remarkable antibacterial activity. Among four species tested, methanol extract of the leaf *S. caseolaris* exhibited wide spectrum of activity which suppress the growth of all tested bacteria, produced a mean zones of inhibition in *E. coli* culture (18 mm) and in *S. aureus* culture (19 mm) (**Table 2** and **Table 3**).

Table 2. Antibacterial activity screening of mangrove extracts towards *E. coli*

Species of Mangroves	Part	Diameter of inhibition (mm)*							
		MeOH (20µl/5µg)	EtOAc (20µl/5µg)	Hexane (20µl/5µg)	A (-)	B (-)	C (-)	D (+)	E (+)
<i>A.marina</i>	Leaf	10	8	0	0	0	0	10	15
	Bark	12	0	0	0	0	0	10	15
	Fruit	16	12	0	0	0	0	10	15
	Root	9	9	0	0	0	0	10	15
<i>A.alba</i>	Leaf	9	7	0	0	0	0	10	15
	Bark	10	8	0	0	0	0	10	15
	Fruit	13	9	0	0	0	0	10	15
	Root	9	8	0	0	0	0	10	15
<i>R.mucronata</i>	Leaf	11	9	0	0	0	0	10	15
	Bark	11	9	0	0	0	0	10	15
	Root	9	8	0	0	0	0	10	15
<i>S.caseolaris</i>	Leaf	18	17	0	0	0	0	10	15
	Bark	12	11	0	0	0	0	10	15
	Fruit	14	11	0	0	0	0	10	15
	Root	9	8	0	0	0	0	10	15

info: * optimization hour observation at the 16th, A (-) MeOH, B (-) EtOAc, C (-) Hexane, D (+) Penicillin 10 µg, E (+) Chloramphenicol 30 µg

Table 3. Antibacterial activity screening of mangrove extract towards *S. aureus*

Species of Mangroves	Part	Diameter of inhibition (mm)*							
		MeOH (20µl/5µg)	EtOAc (20µl/5µg)	Hexane (20µl/5µg)	A (-)	B (-)	C (-)	D (+)	E (+)
<i>A.marina</i>	Leaf	10	7	0	0	0	0	10	15
	Bark	12	11	0	0	0	0	10	15
	Fruit	12	9	0	0	0	0	10	15
	Root	7	10	0	0	0	0	10	15
<i>A.alba</i>	Leaf	9	7	0	0	0	0	10	15
	Bark	8	7	0	0	0	0	10	15
	Fruit	10	8	0	0	0	0	10	15
	Root	9	8	0	0	0	0	10	15
<i>R.mucronata</i>	Leaf	10	8	0	0	0	0	10	15
	Bark	10	8	0	0	0	0	10	15
	Root	9	8	0	0	0	0	10	15
<i>S.caseolaris</i>	Leaf	19	16	0	0	0	0	10	15
	Bark	16	11	0	0	0	0	10	15
	Fruit	11	11	0	0	0	0	10	15
	Root	10	8	0	0	0	0	10	15

info: * optimization hour observation at the 16th, A (-) MeOH, B (-) EtOAc, C (-) Hexane, D (+) Penicillin 10 µg, E (+) Chloramphenicol 30 µg

The result of bioactivity tests showed that mangrove extract bioactive compounds have strong power. According to Davidstout [5], that the power of antibacterial is pointed out by growth inhibition zone. The growth inhibition of >20 mm means very strong, inhibition zone of 10-20 mm means strong, inhibition zone of 5-10 mm means medium, and inhibition zone of <5 mm means weak.

The highest activity recorded was methanol extract of the leaf *S. caseolaris* inhibiting the growth of *E. coli* isolates (18 mm inhibition) and the growth of *S. aureus* isolates (19 mm inhibition).

The difference between the antibacterial activities of mangrove extract could be due to the type and quantity of antimicrobial substances present in each form.

Brine Shrimp Assay

The brine shrimp assay is considered as a reliable indicator for the preliminary assessment of toxicity [6]. This assay is widely employed in the screening process of botanical for the isolation of bioactive metabolites.

The extract of four mangrove species showed different mortality rate at different concentrations (**Table 4**). The mortality rate increased with the increase of concentration of each sample. The crude extracts of the leaf *S. caseolaris* indicated the highest LC₅₀ value of 34914,03 µg/ml meaning not toxic.

Table 4. *Artemia* cytotoxicity profile of mangrove extracts

Concentration µg/ml	Percentage of Mortality			
	Fruit of <i>A. marina</i>	Fruit of <i>A. alba</i>	Leaf of <i>R. mucronata</i>	Leaf of <i>S. caseolaris</i>
0 (Control)	0.00	0.00	0.00	0.00
10	3.33	3.33	3.33	13.33
100	3.33	6.67	10.00	16.67
200	3.33	30.00	13.33	23.33
500	6.67	30.00	23.33	33.33
1000	10.00	36.67	33.33	30.00
LC ₅₀ (µg/ml)	532108.08	6714.288	1043.019	34914.03
Category of toxicity [7]	Not toxic	Not toxic	Not toxic	Not toxic

Purification with Column Chromatography

First step of identification of bioactive compound of mangroves plant is by usage of column chromatography. Solvent applied is chloroform:methanol (9:1 to 1:9). For extract of the leaf *S. caseolaris* as much 20 fractions (yield extract) are produced. Then all fractions were antibacterial tested (*S. aureus* and *E. coli*).

Purification with TLC

Fraction showing bioactivity then underwent thin layer chromatography (TLC). Its R_f point (*Retardation fraction*) was measured. Extract of the leaf *S. caseolaris* eluted by chloroform:methanol (9:1 to 1:9) showed R_f points 0,78. (**Figure 2**). Spot on TLC is taken further and tested by antibacterial (*S. aureus* and *E. coli*). The growth inhibition was 7 mm.



Figure 2. Profile of bioactive fraction of *S. caseolaris* leaf on TLC

HPLC Characteristic of Bioactive Substance

Analysis of HPLC with detector photodiode array (PDA) ($\lambda = 200 - 400$ nm) showed that the dominant peak with its retention time and wavelength characteristic was similar with those of the standard compound characteristic stored in HPLC data base.

HPLC analysis of the leaf *S. caseolaris* showed have 4 peak dominant at retention time (Rt) 14,50 ($\lambda = 234$ nm), Rt 15,92 ($\lambda = 258$ nm), Rt 19,68 ($\lambda = 230$ nm), and Rt 23,13 ($\lambda = 223$ nm and 277

nm) (Figure 3).




Figure 3. HPLC Analysis of leaf *S. caseolaris* extract

Conclusions

Among four mangrove species screened, the broadest activity was showed by *S. caseolaris* extract, therefore this mangrove plant might posses potential source for further study of its bioactive compounds for the purpose of biopharmaceuticals.

Acknowledgements

We wish to thank to all parties supporting this research.

References

- [1] Bandaranayake, W. M. Traditional and medicinal uses of mangroves. *Mangroves and Salt Marshes*, 2: 133-148. 1998.
- [2] Bandaranayake, W. M. Bioactivities, bioactive compounds and chemical constituents of mangrove plants. *Wetlands Ecology and Management* 10: 421-452. 2002.
- [3] Abeysinghe, P, D and Wanigatunge, R. P. Evaluation of antibacterial activity of different mangrove plant extracts. *Ruhuna Journal of Science* 1:104-112. 2006.
- [4] Wardlaw, A.C. Practical statistics for experimental biologists. John Wiley and Sons, Chichester. 1985.
- [5] Davidstout. Disc plate method of microbiological antibiotic assay. *Journal of Microbiology* 4: 659-665. 1971.
- [6] Cyrus, W.G., Daniel, G.W., Nanyingi, M.O., Njonge, F.K. and Mbaria, J.M. Antibacterial and cytotoxic activity of Kenyan medicinal plants, *Memórias do Instituto Oswaldo Cruz*, 103(7): 650-652. 2008.
- [7] Meyer, B. N. Brine Shrimp: a convinient general bioassay for active plant constituents. *Plant Medica* 45: 31-34. 1982.
- [8] Haque, M. E., Islam, M. N., Rahman, M. H., Mohamad, A. U., Antimicrobial and Cytotoxic Activities of the Crude Extracts and Isolated Compounds of *Xylocarpus mollucensis*. *Dhaka Univ. J. Pharm. Sci.* 2: 109-112. 2007.
- [9] Kitamura, S., Anwar, C., Chaniago, A and Baba, S. *Handbook of Mangrove in Indonesia*. ISME. Japan. 1997.
- [10] Wu, J., Xiao, Q., Xu, J., Li, M.Y., Pana, J.Y. and Yang, M. Natural products from true mangrove flora: source, chemistry and bioactivities. *Natural Product Report*, 25: 955-981. 2008.