

# ANTI-INFLAMMATORY EFFECT OF BETEL QUID ON MUCOSAL WOUND OF MALE WISTAR (RATTUS NOVERGICUS) RATS

*by* Budi Santoso

---

**Submission date:** 04-Feb-2022 09:36AM (UTC+0700)

**Submission ID:** 1754571962

**File name:** EFFECT\_OF\_BETEL\_QUID\_ON\_MUCOSAL\_WOUND\_OF\_MALE\_WISTAR.pdf\_17.pdf (228.54K)

**Word count:** 4344

**Character count:** 24378

## ANTI-INFLAMMATORY EFFECT OF BETEL QUID ON MUCOSAL WOUND OF MALE WISTAR (*RATTUS NOVERGICUS*) RATS

RINDIT PAMBAYUN<sup>1\*</sup>, RAFIKA PUTRI<sup>2</sup>, BUDI SANTOSO<sup>1</sup>, TRI WARDANI WIDOWATI<sup>1</sup>, SITI RUSDIANA PUSPA DEWI<sup>3</sup>

<sup>1</sup>Departement of Food and Science, Faculty of Agriculture, Sriwijaya University, Palembang, Indonesia, <sup>2</sup>Student of Dentistry Study Program, Faculty of Medicine, Sriwijaya University, Palembang, Indonesia, <sup>3</sup>Departement of Oral Medicine, Dentistry Study Program, Faculty of Medicine, Sriwijaya University, Palembang, Indonesia  
Email: sitrus\_pd@yahoo.com

Received: 24 Jan 2019, Revised and Accepted: 26 May 2019

### ABSTRACT

**Objective:** True experimental *in vivo* with pre-test and post-test control group design was conducted at Animal House of Medical Faculty of Sriwijaya University, Palembang and Biomedical Science Laboratory of Palembang, South Sumatera.

**Methods:** Thirty rats were divided into five groups; three groups were treated with betel quid ointments at concentration of 5%, 10% and 20%, one group was positive control (hyaluronic acid) and last group was negative control (placebo). One-mm diameter of wound was made on lower lip mucosa of rats with cylinder diamond bur. Wound was induced with carrageenan. The number of neutrophils was counted on the first day and third day after treatment by using hematoanalyzer.

**Results:** There was a significant decrease of the number of neutrophils for respective groups ( $p < 0.05$ ). Betel quid ointment of 10% and 20% concentration had similar effect to 0.2% hyaluronic acid ( $p > 0.05$ ).

**Conclusion:** Betel quid has an anti-inflammatory effect on the mucosal wound of rats in a dose-dependent manner.

**Keywords:** Anti-inflammatory, Betel quid, Mucosal wound

© 2019 The Authors. Published by Innovare Academic Sciences Pvt Ltd. This is an open-access article under the CC BY license (<http://creativecommons.org/licenses/by/4.0/>) DOI: <http://dx.doi.org/10.22159/ijap.2019.v11s4.35298>

### INTRODUCTION

Inflammation is a biological response to tissue damage, caused by pathogens, chemicals, or physics. The main function of the inflammation process is to resist tissue damaged and keep it from spreading. Inflammation is characterized by heat (calor), redness (rubor), swelling (tumor), pain (dolor) and loss of function (functiolaesa) [2]. The sensation of heat is due to the increased movement of more blood in enlarged vessel on the wound area. The redness is caused by the increased number of erythrocytes to the affected area. Swelling is due to a buildup of fluid on permeable and dilated vessel into the surrounding tissue. Pain is because the stimulated nerve endings are released. Loss of function is related to immobility joint in the inflamed tissue and scar tissue replacement on the wound. This process is happened an hour after injury [3, 4].

Inflammation commonly found in oral mucosa is caused by a number of physical injury (i.e., bitten, collide, or instrumentation) or an infection [5]. Those injuries trigger the immune system as response, and show inflammation signs. The main processes occur during the inflammation are the increased supplying blood to the injury and lead increased blood movement in dilated arteries. The vessels become more permeable and easier for fluids and proteins to infiltrate [6]. After a few hours, neutrophils are released on wound area. The highest number of neutrophils are on first and second days after injury [7]. Neutrophils express as the primary defence against pathogens. On the site of injury, where it assembly activating signals to trigger bacterial killing, neutrophils are mobilized, then secrete antimicrobial agents such as cytokines, chemokines, and proteases. The rapid neutrophil migration from the circulation on to injury is regulated by vascular endothelium [8, 9]. Activation of neutrophils is initiated by cytokines, such as interleukins (IL-1 $\beta$ , IL-8), tumor necrosis factor (TNF $\alpha$ ), leukotrienes, pathogens/bacterial products and by hit stimulus [10]. Neutrophils eliminate microorganisms and dead cells through phagocytosis with different cytotoxic mechanism, the release of free radicals and deliver of proteolytic enzymes into phagosome [11]. When neutrophils confront with foreign particles, the cells will recognize the particles, swallow or phagocyte, then

destroy the foreign particles. Neutrophils perform the process of marginization, adhesion and migration before being ready for the function. The neutrophil's life cycle is regulated by granulocyte colony-stimulating factor (G-CSF) [7]. Along with the presence of neutrophils, macrophages are needed. Macrophages are formed by chemotaxis and migration. By day 3, the neutrophils are phagocytized by macrophages. Peripheral blood monocytes on the wound site differentiate to be macrophages [12]. Macrophages are formed by chemotaxis and migration. The number of macrophages reaches a maximum after 4-5 d. Macrophages play a role in the process of phagocytosis. Macrophages kill pathogenic microorganisms and initiate tissue regeneration processes [13].

Anti-inflammation agents are used to reduce the symptom and accelerate wound healing process. One of them is hyaluronic acid. Hyaluronic acid is considered one of anti-inflammatory agents [14]. Previous study mentioned that topical hyaluronic acid of 0.2% concentration is effective on the healing of gingivitis [15]. But hyaluronic acid has side effects, such as allergic or hypersensitivity reaction, edema, ecchymoses, hypercorrection, bluish discoloration, and indurated nodules [16]. Alternative medicine from natural product as anti-inflammatory agent with minimum side effect is needed to find. One of them is betel quid.

Betel quid is combination of betel (*Piper betle* Linn.), areca nut (*Areca catechu* Linn), gambier (*Uncaria gambir* Roxb.), and mineral slaked lime (calcium hydroxide) [17]. All components of betel quid have been proven to have anti-inflammatory effect. Betel has been used as anti-inflammation in oral medicine for long time. Alam *et al.* reported that *Piper betle* leaves had antioxidant, analgesic and anti-inflammatory activities [18]. The anti-inflammatory activities of *Piper betle* is due to its chemical constituents, such as eugenol, hydroxychavicol, quercetin,  $\beta$ -caryophyllene [19].

The betel quid is the dark red seed of *Areca catechu*. This plant has several therapeutic properties, including analgesics, anti-inflammatory and antioxidants [20]. Bhandare *et al.* revealed that hydroalcohol extract from *Areca catechu* nut inhibited extravasation of plasma protein and inflammatory progression in diameter [21].

The areca nut extract contains procyanidins, major condensed tannins that widely disseminates its phenolic compound and have pharmacological effect [22]. Procyanidin is effective in inhibiting the expression of pro inflammatory mediator and cytokines [23]. Kato et al reported that procyanidin suppressed TPA-induced inflammation of mouse, and its activity was stronger than indomethacin and glycyrrhetic acid [24].

Gambier (*Uncaria gambier* Roxb.) plays important role as an anti-inflammatory agent. Seto et al mentioned that gambier extract was effective in reducing neutrophil cells in inflammation phase of male Wistar's injury [25]. The anti-inflammatory activity of gambier is due to its chemical compound, catechins [26]. Catechins are polyphenol, and group of flavonoids that exhibit water soluble characteristics [27]. Trekli et al. reported that catechins suppressed cytokine-induced-IL6 and IL8 and showed anti-inflammatory activity [28].

Mineral slaked lime contained in betel quid is commonly used in dentistry. Mineral slaked lime has a formula of Ca(OH)<sub>2</sub> or calcium hydroxide. Calcium hydroxide has been proved has an antibacterial effect and anti-inflammation in the periapical lesion, so that it mostly used in root canal treatment [29]. Anti-inflammatory effect of calcium hydroxide is rely on alkalinity and calcium ion release. The high pH of calcium hydroxide encourages antibacterial effect, inflammatory reduction, and stimulate tissue repair [30]. Dixit et al stated that calcium hydroxide was successful resolution in treating periapical lesions [31].

The study about the anti-inflammatory effect of betel quid in male Wistar rats has not been done. The aim of this study was to investigate the anti-inflammatory effect of betel quid on the oral mucosal wound of male Wistar (*Rattus norvegicus* L.) rats.

## MATERIALS AND METHODS

### Material

This study was true experiment *in vivo* with pretest-posttest control group design. The research was conducted at Animal House of Medical Faculty of Sriwijaya University Palembang and Biomedical Science Laboratory of Palembang, South Sumatera. The protocol had been approved by Research Ethical Commission of Mohammad Hoesin General Hospital (RSMH) Palembang and Medical Faculty of Sriwijaya University with ethical certificate No. 391/kepkrsmhfksunri/2017.

### Animals

Thirty male white Wistar (*Rattus norvegicus* L.) rats (obtained from Pharmacy School of Bandung Institute of Technology) were divided into 5 groups. Group I was treated with 5% betel quid ointment, Group II was treated with 10% betel quid ointment, Group III was treated with 5% betel quid ointment, Group IV was positive control, treated with 0.2% hyaluronic acid ointment (purchased from Ricefarma Pharm. Co, Surabaya, Indonesia), and Group V was negative control, treated with placebo ointment. Rats (weighing 190-200 mg, aged 8-12 w old) were acclimated for 8 d at room temperature of 20-25 °C under 12:12 light-dark cycle, prior to the experimental period [32]. Samples were fed with standard pellet diet and water *ad libitum*.

### Preparation of betel quid ointment

Betel quid components consisted of betel leaf (*Piper betle* L.), areca nut (*Areca catechu* L.), gambier (*Uncaria gambier* Roxb.) and mineral slaked lime (calcium hydroxide) were collected from Babatoman Village, Sekayu Subdistrict, Musi Banyuasin District, South Sumatra

Province, Indonesia. All the constituents and material were identified and authenticated by the Faculty of Agriculture, Sriwijaya University, Indonesia. Betel quid was made in ointment preparats of 5 g. The composition of 5% betel quid ointment was as follows 250 mg betel quid mixture, 4.04 g vaseline album, 712.5 mg adeps lanae; while 10% betel quid was as follows 500 mg betel quid mixture, 3.82 g vaselin album, 675 mg adeps lanae; 20% betel quid was as follows 1 g betel quid mixture, 3.4 g vaselin album, 600 mg adeps lanae; and placebo ointment was as follows 4.25 g vaselin album and 750 mg adeps lanae. The ingredients of betel quid mixture were 8 g betel leaves, 2.5 g areca nut, 3.5 g gambier, and 2 g mineral slaked lime mixed homogeneously [33].

### Wound induction and treatment

Prior to wound induction, rats were anesthetized with 0.2 ml ketamine by i. m injection. Mandibular labial gingiva and lower lip mucosa of rats were swabbed with cotton wetted with 10% povidone iodine for sterilization. The lower lip of rats was withdrawn by using tweezer (Fisher brand™, Thermo fisher Co, UK) and labial gingiva was induced with 1% carrageenan by using spuit. One-mm cylinder diamond bur (Microdont, USA) was used to create wound at the depth of 1 mm. Blood was cleaned with wetted cotton sterile and dried. After 5 h, blood samples were performed on orbital sinus to count the number of segmented neutrophil. The blood samples were put in the pots filled with ethylene diamine tetra acid (EDTA) anti-coagulant.

After blood sample was taken on the first day, mucosal wound was treated with the ointments twice daily based on the groups. The treatment was done for 3 d. On the third day, orbital sinus blood samplings were taken to count the neutrophils after treatment. Segmented neutrophils were counted by using hematology analyzer (Mindray BC-2800, Shenzhen Mindray Bio-Medical Electronic Co., Ltd, China).

### Statistical analysis

Data were analyzed using SPSS ver. 22 (IBM® inc. pvt ltd, US) and Microsoft Excel (Microsoft inc®, Redmond). Shapiro Wilk's test and Levene's test were used to know the normality and homogeneity of samples, with  $p > 0.05$  as significance. Paired t-test was used to analyze the significant lowering number of neutrophils before and after treatment. The test was followed with one way Anova and Post-Hoc LSD. P value of  $< 0.05$  was indicated as significant statistically.

## RESULTS

The number of neutrophils was evaluated. Saphiro-Wilk and Levene's test showed  $p > 0.05$ , so that meant that data were normal and homogen. All groups showed the reduction of neutrophils before and after treatment significantly table 1. The lowest reduction number of neutrophils was showed on hyaluronic acid of 0.2%, followed by betel quid ointment of 20%, 10%, 5% concentration and placebo ointment.

The result of one-way Anova test exhibited  $p = 0.00$ . It meant that there was a significant difference in reducing neutrophils between and within groups after treatment. Data analysis was continued to Post-hoc LSD test fig. 2. The result showed that there was no significant difference between betel quid ointment at 10% and 20% concentration and positive control in reducing neutrophils. All groups presented significant effects to the negative control, except betel quid of 5% concentration. It meant that 10% and 20% betel quid ointment had a similar anti-inflammatory effect to 0.2% hyaluronic acid ointment.

Table 1: The number of neutrophils before and after treatment

	Means+SD		P value
	Before	After	
5% betel quid ointment	28.00+2.28	16.00+1.79	0.00*
10% betel quid ointment	28.50+4.89	11.17+2.04	0.00*
20% betel quid ointment	28.50+2.04	9.67+1.63	0.00*
0.2% hyaluronic acid	28.33+2.73	9.33+1.03	0.00*
Placebo	28.33+5.68	18.33+6.28	0.00*

\*significance =  $p < 0.05$ , paired t-test




Fig. 1: The number of neutrophils. Betel quid extract of 10% and 20% concentration displayed anti-inflammatory effect significantly and data were expressed as mean±SD, \* $p < 0.05$  versus positive control group; \*\* $p < 0.05$  versus the negative control group

## DISCUSSION

Chewing betel quid is done for many people all over the world since very ancient times. In Indonesia, the betel leaves, along with areca nut, gambier leaves and mineral slaked lime in wrapped package, are chewed together [34]. All of those components in betel quid consist of active compounds. The effectivity of anti-inflammatory is due to the synergic combination of pharmacological properties contained in betel quid. All constituents of the ingredients has been documented to have an anti-inflammatory effect. Eugenol, hydroxychavicol, terpenes (contained in betel leaf), procyanidin, alkaloids/*arecoline* (contained in areca nut), catechin, tannin (contained in gambier leaf), and calcium hydroxide have been showed in alleviating inflammation. Reddy *et al.* reported that *Piper betle* leaves had anti-inflammatory activity in experimental animals due to its active compounds [35]. Sarpongala *et al.* stated that *Areca catechu* L reduced paw edema in rats and its anti-inflammatory activity was similar to indomethacin [36]. Yimam *et al.* revealed that UP3005 (containing *Uncaria gambir* and *Morus* extract) inhibit inflammatory enzymatic activities of cyclooxygenase-1 (COX-1), COX-2 and lipoxygenase (5-LOX), respectively [37]. Louwakul and Lertchirakarn exhibited that calcium hydroxide was effective in repairing and treating inflamed dental pulp tissue [38].

Eugenol suppresses the expression of cyclooxygenase and decreases the production of proinflammatory cytokines [39, 40]. Hydroxychavicol has the ability in inhibiting platelet aggregation. The previous study reported that hydroxychavicol is effective to hamper COX-1 and COX-2 enzymes, ROS scavenger and platelet calcium signaling [41]. Eugenol and hydroxychavicol inhibit xanthine oxidase (XOD) and lipoxygenase (LOX) [42]. XOD plays an important role in neutrophil mediation. The activation of XOD is a response to proinflammatory and growth factor stimulation [43]. While LOX, group of oxidative enzymes, involves in the metabolism of pro-inflammatory mediators, i.e. prostaglandins and leukotrienes [44]. Terpenes block the oxidation of arachidonic acid and the release of inflammatory mediators [45].

Procyanidins modulate nitric oxide generation that plays an important role in acute and chronic inflammation and reduces COX-2 expression [46]. Alkaloid (*Arecoline*) contained in areca nut is capable to reduce pro-inflammatory mediators, such as prostaglandin E2 an interleukine-6 through the inhibition of COX and LOX metabolisms [47].

Catechin has been proved to have anti-inflammatory activity by inhibiting XO, IL-6, IL-8 production, COX-1/COX-2 expression, lipoxygenase (LOX) and phospholipase (PL) enzyme. The inhibition of COX and LOX disturbs synthesis of inflammatory mediators, such as prostaglandin and leukotriene. Inactivation of PL enzyme causes the inhibition of the release of arachidonic acid. As a result, the number of inflammatory mediators produced will decrease [48, 49]. Another anti-inflammatory mechanism of catechin is to eliminate the expression of pro-inflammatory cytokines, neutralize free

radical, inhibit NF-Kb activation so that inflammation reaction will be decreased [50].

Calcium hydroxide has ability in eliminating some bacteria in oral mucosa. The high alkalinity of calcium hydroxide promotes antimicrobial activities [51]. The presence of calcium hydroxide modify the environmental pH in inflammation areas, regulate inflammatory cell migration and proliferation, control the irritant agents, stimulate tissue mineralization [52]. The effect is based on the dissociation of calcium hydroxide into calcium and hydroxide ions [38, 53].

## CONCLUSION

In conclusion, the study shows that betel quid treatment has an anti-inflammatory effect on the mucosal wound of Wistar rats in a dose-dependent manner. The anti-inflammatory activities are due to the synergic pharmacological properties contained in betel quid.

## ACKNOWLEDGMENT

This research was financially supported by Research Institutes and Community Service of Sriwijaya University for Superior Research Profession (LP2M 2017, no. 1011/UN9.3.1/PP/2017).

## AUTHORS CONTRIBUTIONS

All authors have made substantial contribution to the work reported in the manuscript. Rindit Pambayun, Budi Santoso, Triwardhani developed the idea and designed study. Rafika Putri conducted the actual study, Siti Rusdiana Puspa Dewi involved in the designed study, administration process, the statistical analysis, and write up of the manuscript.

## CONFLICT OF INTERESTS

There are no conflict of interest in this study

## REFERENCES

1. Aramwit P, Towiwat P, Srichana T. An investigation of the anti-inflammatory potential of silk sericin. *Adv Sci Lett* 2013;19:3615-9.
2. Kumar V, Abbas AK, Aster JC. editors. *Robbins basic pathology*. Canada: Elsevier Saunders; 2013.
3. Puncard NA, Whelan CJ, Adcock I. *Journal of inflammation*. *J Inflamm (Lond)* 2004;1:1-4.
4. Gerdle B, Bijar G, Nazdar G, Emmanuel B, Torsten G. Signs of ongoing inflammation in female patients with chronic widespread pain: a multivariate, explorative, cross-sectional study of blood samples. *Medicine* 2017;96:e6130.
5. Hurlbutt, Michelle, Thomsen L. editors. *Dentifying recurrent oral ulcerations*. New York: PenWell; 2007.
6. Cawson RK, Odel EW. *Cawson's essentials of oral pathology and oral medicine*. China: Churchill Livingstone Elsevier; 2008.

7. Kruger P, Saffarzadeh M, Weber ANR, Rieber N, Radsak M, Bernuth HV, et al. Neutrophils: between host defence, immune modulation, and tissue injury. *Plos Pathol* 2015;11:1004651.
8. Lacy P. Mechanisms of degranulation in neutrophils. *Allergy Asthma Clin Immunol* 2006;2:98-108.
9. Leoni G, Neumann PA, Sumagin R, Denning TL, Nusraf A. Wound repair: role of immune-epithelial interactions. *Mucosal Immunol* 2015;8:959-68.
10. Wright HL, Moost RJ, Bucknall RC, Edwards SW. Neutrophil function in inflammation and inflammatory disease. *Rheumatol (Oxford)* 2010;49:1618-31.
11. Rosales C, Demaurex N, Lowell CA, E Uribe Querol E. Neutrophils: their role innate and adaptive immunity. *J Immunol Res* 2016. Doi:10.1155/2016/1469780
12. Rahat MA, Coffelt SB, Granot Z, Muthana M, Amedei A. Macrophages and neutrophils: regulation of the inflammatory microenvironment in autoimmunity and cancer. *Mediators Inflammation* 2016. Doi:10.1155/2016/5894347
13. Koh TJ, DiPietro LA. Inflammation and wound healing: the role of the macrophage. *Expert Rev Mol Med* 2011;13:e23.
14. Kapoor P, Sachdeva S, Sachdeva S. Topical hyaluronic acid in the management of oral ulcer. *Indian. J Derm* 2011;56:300-2.
15. Sapna N, Vandana KL. Evaluation of hyaluronan gel (Gengigel) as a topical applicant in the treatment of gingivitis. *J Investig Clin Dent* 2011;2:162-70.
16. Bitterman Deutsch O, Kogan L, Nasser F. Delayed immune mediated adverse effects of hyaluronic acid fillers: report of five cases and review of the literature. *Dermatol Reports* 2015;7:5851.
17. Marcelina, Samad R. Salivary profile of betel quid tobacco chewers in district of rembon, tana toraja. *J Dentomaxillofac Sci* 2013;12:103-13.
18. Alam B, Akter F, Parvin N, Pia RS, Akter S, Chowdhury J, et al. Antioxidant, analgesic, and anti-inflammatory activities of the methanolic extract of *Piper betle* leaves. *Avicenna J Phytomed* 2013;3:112-25.
19. Dwivendi V, Tripathi S. Review study on potential activity of piper betle. *J Pharmacog Phytochem* 2014;3:93.
20. Bhandare AM, Kshirsagar AG, Vyawahare NS, Thorve VS. Potential analgesic, anti-inflammatory and antioxidant activities of hydroalcoholic extract of *Areca catechu* L. *Nut Food Chem Toxicol* 2010;48:3412-7.
21. Bhandare A, Kshirsagar A, Vyawahare N, Sharma P, Mohite R. Evaluation of anti-migraine potentiation of *Areca catechu* to prevent nitroglycerin-induced delayed inflammation in rat meninges: possible involvement of NOS inhibition. *J Ethnopharmacol* 2011;136:267-70.
22. Pithayanukul P, Nithitanakool S, Bavovada R. Hepatoprotective potentiation of extracts from seeds of *Areca catechu* and nutgalls of *Quercus infectoria*. *Molecules* 2009;14:4987-5000.
23. Bak MJ, Truong VL, Kang HS, Jun M, Jeong WK. Anti-inflammatory effect of procyanidins from wild grape (*Vitis amurensis*). *Oxid Med Cell Longev* 2013. <http://dx.doi.org/10.1155/2013/409321>
24. Katoh M, Oizumi Y, Mohri Y, Hirota M, Makabe H. Synthesis of procyanidin B1, B2, and B4 and their anti-inflammatory activity: the effect of 4-alkoxy group of catechin and/or epicatechin electrophiles for condensation. *Lett Org Chem* 2012;9:233-8.
25. Wibowo IS, Dewi SRP, Yasmin U, Octaria F. FDI-IDA continuing education. Editors. The effectiveness of gambir extract (*Uncaria gambir* Roxb.) as an anti-inflammatory agents in wistar male rats. *Proceeding of FDI-PDGI Continuing Education*. Bandung, Indonesia; 2016. p. 11-2.
26. Anggraini T, Tai A, Yoshino T, Itani T. Antioxidative activity and catechin content of four kind of *Uncaria gambir* extracts from West Sumatra, Indonesia. *Afr J Biochem Res* 2011;5:33-8.
27. Mbaveng AT, Zhao Q, Kuete V. Harmful and protective effects of phenolic compounds from African medicinal plants. *Toxicol Survey African Med Plants* 2014;1:577-609.
28. Trekli M, Bettle D, Guesdon F. Anti-inflammatory actions of green tea catechins and ligands of peroxisome proliferator-activated receptors. *Int J Exp Pathol* 2004;85:A75.
29. Mustofa M, Saujanya KP, Jain D, Sajjanshetty S, Arun A, Uppin L, et al. Role of calcium hydroxide in endodontics: a review. *Global J Med Public Health* 2012;1:66-8.
30. De Freitas RP, Greatti VR, Alcalde MP, Cavenago BC, Vivan RR, Duarte MAH, et al. Effect of the association of nonsteroidal anti-inflammatory and antibiotic drugs on antibiogram activity and pH of calcium hydroxide pastes. *J Endo* 2017;43:131-4.
31. Dixit S, Dixit A, Kumar P. Unusual gingival enlargement: a rare case report. *Case Reports Dentistry* 2014. <http://dx.doi.org/10.1155/2014/536312>
32. Arts JMW, Kramer K, Arndt SS, Ohi F. Effects of transfer from breeding to research facility on the welfare of rats. *Animals (Basel)* 2014;4:712-28.
33. Wowor VN, Supit A, Marbun DR. Profile of chewing betel in Manado. *Unsrat Press* 2016;2:1-5.
34. Toprani R, Patel D. Betel leaf: revisiting the benefits of an ancient Indian herb. *South Asian J Cancer* 2013;2:140-1.
35. Reddy PS, Gupta RJ, Reddy SM. Analgesic and anti-inflammatory activity of hydroalcoholic extract of *Piper betle* leaves in experimental animals. *Int J Basic Clin Pharm* 2016;5:979-85.
36. Sarpangala KB, Ashwin D, Sarpangala M. Punarnava-a natural remedy by ayurveda. *Int J Ayurveda Pharm Res* 2016;4:78-83.
37. Yimam M, Lee YC, Kim TW, Moore B, Jiao P, Hong M, et al. Analgesic and anti-inflammatory effect of UP3005, a botanical composition containing two standardized extracts of *Uncaria gambir* and *Morus alba*. *Pharmacogn Res* 2015;7:S39-46.
38. Louwakul P, Lertchirakarn V. Incorporation of anti-inflammatory agent into calcium hydroxide pulp capping material: an *in vitro* study physical and mechanical properties. *Dent Mater J* 2012;31:32-9.
39. Huang X, Ma C. Anti-inflammatory effects of eugenol on lipopolysaccharide-induced inflammation reaction in acute lung injury via regulating inflammation and redox status. *Int Immunopharmacol* 2015;26:265-71.
40. Magalhaes CB, Riva DR, DePaula LJ, Brando-Lima A, Koatz VGL, Leal-Cardoso JH, et al. *In vivo* anti-inflammatory action of eugenol on lipopolysaccharide-induced lung injury. *J Appl Physiol* 2010;108:845-51.
41. Rintu D, Shinjini M, Kaustab M, Pramathadhip P, Umesh PS, Banarjee ER. Anti-oxidant and anti-inflammatory activities of different varieties of piper leaf extracts (*Piper betle* L.). *J Nutr Food Sci* 2015;5:415.
42. Pin KY, Chuah AL, Rashih AA, Mazura MP, Fadzureena J, Vimala S, et al. Antioxidant and anti-inflammatory activities of extracts of betel leaves (*Piper betle*) from solvents with different polarities. *J Trop Forest Sci* 2010;22:448-55.
43. Aboali M, Lall GS, Coughlan K, Lall HS, Gibbs BF, Sumbayev VV. Crucial involvement of xanthine oxidase in the intercellular signaling networks associated with human myeloid cell function. *Sci Rep* 2014;4:6307.
44. Wisastra R, Dekker FJ. Inflammation, cancer and oxidative lipoxygenase activity are intimately linked. *Cancer (Basel)* 2014;6:1500-21.
45. Theoduloz C, Delporte C, Valenzuela Barra G, Silva X, Cadiz S, Bustamante F, et al. Topical antiinflammatory activity of new hybrid molecules of terpenes and synthetic drugs. *Molecules* 2015;20:11219-35.
46. Lee KP, Sudjarwo GW, Kim JS, Dirgantara S, Maeng WJ, Hong H. The anti-inflammatory effect of Indonesian *Areca catechu* leaf extract *in vitro* and *in vivo*. *Nutr Res Prac* 2014;8:267-71.
47. Khan S, Mehmood MH, Ali ANA, Ahmed FS, Dar A, Gilani AH. Studies on anti-inflammatory and analgesic activities of betel nut in rodents. *J Ethnopharm* 2011;135:654-61.
48. Risma E, Ningsih S, Fachrudin F. *In vitro* study of xanthine oxidase inhibitory of gambir (*Uncaria gambir*) hunter roxb extracts. *Pharm J* 2017;9:862-5.
49. Morel A, Miller E, Bijak M, Saluk J. The increased level of COX-dependent arachidonic acid metabolism in blood platelets from secondary progressive multiple sclerosis patients. *Mol Cell Biochem* 2016;420:85-94.
50. Mota MAL, Landim JSP, Targino TSS, Silva SFR, Silva SL, Pereira MRP. Evaluation of the anti-inflammatory and analgesic effects of green tea (*Camellia sinensis*) in mice. *Acta Cirurgica Brasileira* 2015;30:242-6.
51. Mohammadi Z, Shalavi S, Yazdizadeh M. Antimicrobial activity of calcium hydroxide in endodontics: a review. *Chonnam Med J* 2012;48:133-40.

52. Subramaniam P, Konde S, Prashanth P. An *in vitro* evaluation of pH variations in calcium hydroxide liners. J Indian Soc Pedodontics Preventive Dentistry 2006; 24:144-5.
53. De Freitas RP, Greatti VR, Alcalde MP, Cavenago BC, Vivan RR. Effect of the association of nonsteroidal anti-inflammatory and antibiotic drugs on antibiofilm activity and pH of calcium hydroxide pastes. J Endo 2017;43:131-4.

# ANTI-INFLAMMATORY EFFECT OF BETEL QUID ON MUCOSAL WOUND OF MALE WISTAR (RATTUS NOVERGICUS) RATS

## ORIGINALITY REPORT

**21** %  
SIMILARITY INDEX

**16** %  
INTERNET SOURCES

**11** %  
PUBLICATIONS

**9** %  
STUDENT PAPERS

## PRIMARY SOURCES

**1** [jurnal.unissula.ac.id](http://jurnal.unissula.ac.id) 3 %  
Internet Source

**2** Submitted to Sriwijaya University 2 %  
Student Paper

**3** [repository.unair.ac.id](http://repository.unair.ac.id) 1 %  
Internet Source

**4** [worldwidescience.org](http://worldwidescience.org) 1 %  
Internet Source

**5** Submitted to Universitas Hasanuddin 1 %  
Student Paper

**6** Submitted to Udayana University 1 %  
Student Paper

**7** [jurnalmahasiswa.unesa.ac.id](http://jurnalmahasiswa.unesa.ac.id) 1 %  
Internet Source

**8** [jisppd.com](http://jisppd.com) 1 %  
Internet Source

**9** Amol Bhandare, Ajay Kshirsagar, Neeraj Vyawahare, Pushpendra Sharma, Reshma 1 %

Mohite. "Evaluation of anti-migraine potential of Areca catechu to prevent nitroglycerin-induced delayed inflammation in rat meninges: Possible involvement of NOS inhibition", Journal of Ethnopharmacology, 2011

Publication

10

[www.science.gov](http://www.science.gov)

Internet Source

1 %

11

[www.ncbi.nlm.nih.gov](http://www.ncbi.nlm.nih.gov)

Internet Source

1 %

12

"Abstracts", *Respirology*, 11/2011

Publication

<1 %

13

M. Katoh, Y. Oizumi, Y. Mohri, M. Hirota, H. Makabe. "Synthesis of Procyanidin B1, B2, and B4 and Their Anti-Inflammatory Activity: The Effect of 4-Alkoxy Group of Catechin and/or Epicatechin Electrophiles for Condensation", *Letters in Organic Chemistry*, 2012

Publication

<1 %

14

Submitted to New Jersey Institute of Technology

Student Paper

<1 %

15

Rafaela Pignatti de Freitas, Vanessa Raquel Greatti, Murilo Priori Alcalde, Bruno Cavalini Cavenago et al. "Effect of the Association of Nonsteroidal Anti-inflammatory and Antibiotic

<1 %



Drugs on Antibiofilm Activity and pH of Calcium Hydroxide Pastes", Journal of Endodontics, 2017

Publication

16

[www.koreascience.or.kr](http://www.koreascience.or.kr)

Internet Source

<1 %

17

Leoni, G, P-A Neumann, R Sumagin, T L Denning, and A Nusrat. "Wound repair: role of immune-epithelial interactions", Mucosal Immunology, 2015.

Publication

<1 %

18

Naresh Killi, Anil Tukaram Pawar, Rathna VN Gundloori. "Polyesteramide of Neem Oil and Its Blends as an Active Nanomaterial for Tissue Regeneration", ACS Applied Bio Materials, 2019

Publication

<1 %

19

[www.actanaturae.ru](http://www.actanaturae.ru)

Internet Source

<1 %

20

[www.tandfonline.com](http://www.tandfonline.com)

Internet Source

<1 %

21

[test.dovepress.com](http://test.dovepress.com)

Internet Source

<1 %

22

Pang Xinxin, Ya-ge Zhang, Li-jie Zhang, Qing Zhu, Hai-tao Shang, Zhen-tao Wang. "Autophagy Inhibits Renal Tubule Inflammation in Diabetic Kidney Disease Mice

<1 %

Via Inhibition of NLRP3 Inflammasome",  
Research Square Platform LLC, 2022

Publication

23

[pure.ulster.ac.uk](https://pure.ulster.ac.uk)

Internet Source

<1 %

24

[www.clonerresources.com](http://www.clonerresources.com)

Internet Source

<1 %

25

MS Firdaus, Nukman, Irsyadi Yani, Amir Arifin, Prana Arifita, Indra Surya. "The Effect of Heat Treatment on Fatigue Testing of Aluminum Cans", Journal of Physics: Conference Series, 2019

Publication

<1 %

26

Muhammad Umar Ijaz, Hussain Ahmed, Asma Ashraf, Sidra Aziz, K.A. Al-Ghanim, Mumtaz Akhtar, M. Nadeem Riaz, Shahid Mahboob. "Vitexin attenuates cisplatin-induced renal toxicity by reducing oxidative stress and inflammation", Journal of King Saud University - Science, 2021

Publication

<1 %

27

[journals.biologists.com](https://journals.biologists.com)

Internet Source

<1 %

28

[www.ijphrd.com](http://www.ijphrd.com)

Internet Source

<1 %

29

Submitted to Glasgow Caledonian University

Student Paper

<1 %

30

[www.panafrican-med-journal.com](http://www.panafrican-med-journal.com)

Internet Source

<1 %

31

"Medicinal Herbs and Fungi", Springer Science and Business Media LLC, 2021

Publication

<1 %

32

Jeong-Ran Park, Hanbyeol Lee, Seok-In Kim, Se-Ran Yang. "The tri-peptide GHK-Cu complex ameliorates lipopolysaccharide-induced acute lung injury in mice", *Oncotarget*, 2016

Publication

<1 %

33

Pornanong Aramwit, Pasarapa Towiwat, Teerapol Srichana. "Anti-inflammatory Potential of Silk Sericin", *Natural Product Communications*, 2013

Publication

<1 %

Exclude quotes  On

Exclude matches  Off

Exclude bibliography  On