

TROPICAL JOURNAL OF
**NATURAL
PRODUCT
RESEARCH**

TJNPR

www.tjnpr.org VOL. 1 (NO. 1) 2017



**Tropical Journal of
NATURAL PRODUCT RESEARCH**

This journal is indexed in Open J-Gate,
Crossref, Creative Commons, Index
Copernicus, Google Scholar.

All published papers are DOI assigned.

Official Journal of Natural Product Research Group
University of Benin

Available online at <https://www.tjnpr.org>*Original Research Article*

Bioactivity Endophytic Fungi Isolated from the Leaf Stalk of *Syzygium jambos* L. Alston

Kurratul Aini^{1,2*}, Elfita Elfita^{3,*}, Hary Widjajanti⁴, Arum Setiawan⁴¹Graduate School of Sciences, Faculty of Mathematics and Natural Sciences, Universitas Sriwijaya. Jl. Padang Selasa No. 524, Palembang 30139, South Sumatra, Indonesia²Universitas Islam Negeri Raden Fatah. Jl. Prof. K.H. Zainal Fikri, Palembang 30126, South Sumatra, Indonesia³Department of Chemistry, Faculty of Mathematics and Natural Sciences, Universitas Sriwijaya. Jl. Raya Palembang-Prabumulih Km 32, Indralaya, Ogan Ilir 30662, South Sumatra, Indonesia⁴Department of Biology, Faculty of Mathematics and Natural Sciences, Universitas Sriwijaya. Jl. Raya Palembang-Prabumulih Km 32, Indralaya, Ogan Ilir 30662, South Sumatra, Indonesia

ARTICLE INFO

Article history:

Received 22 May 2022

Revised 26 September 2022

Accepted 02 November 2022

Published online 01 December 2022

Copyright: © 2022 Aini *et al.* This is an open-access article distributed under the terms of the [Creative Commons Attribution License](https://creativecommons.org/licenses/by/4.0/), which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.

ABSTRACT

Antibiotic resistance and free radicals are a problem to be overcome by using antibiotics and antioxidants from natural ingredients. This study aimed to investigate the antibacterial and antioxidant activity of endophytic fungi (EF) isolated from the leaf stalks of jambu mawar (*Syzygium jambos* (L.) Alston). EF isolated from the leaf stalk of *S. jambos* was identified microscopically (shape of hyphae and spores) and macroscopically (growth patterns, colony color, texture, margin, and other characteristics). Antibacterial activity was tested on *Salmonella typhi*, *Escherichia coli*, *Staphylococcus aureus*, and *Bacillus subtilis* bacteria using the Kirby Bauer method. Antioxidant test using 2,2-diphenyl-1-picrylhydrazyl (DPPH) method. The molecular identified of EF with high antioxidant and antibacterial activity to determine the species and continued with the isolation of pure compounds. Identification of pure compounds was carried out by spectroscopic methods including 1D NMR (¹H-NMR and C-NMR). The results of the isolation of EF from the leaf stalk of *S. jambos* found four isolates, namely SJS1-SJS4. The results of the antibacterial test represented that the EF isolate SJS1 had strong activity against *S. aureus* and *B. subtilis* bacteria. The antioxidant activity test showed IC₅₀ value of 29.29 µg/mL. Molecular identification results showed that SJS1 was identified as *Lasiodiplodia theobromae*. Spectroscopic results of the pure compound identified as 3,5-dihydroxy-4-(4-hydroxyphenyl)tetrahydro-2H-pyran-2-one. The endophytic fungus *Lasiodiplodia theobromae* was isolated from the leaf stalk of *S. jambos* (L.) Alston has the potential as a source of antioxidants and antibacterial bioactive compounds that can be developed through further research, including in vitro and in vivo tests.

Keywords: Antibacterial, Antioxidants, *Lasiodiplodia theobromae*.

Introduction

The use of synthetic antibiotics with inappropriate doses causes pathogenic bacteria to become resistant and cause side effects. Therefore, research is needed to find alternative natural antibiotic ingredients. Endophytic fungi (EF) are reported to have metabolites that can inhibit bacterial growth. Some of EF such as *Fusarium*,¹ *Cladosporium*, *Chaetomium*, and *Ceratobasidium*² have antibacterial activity against *Escherichia coli* and *Staphylococcus aureus*. EF *Thanatephorus cucumeris* exhibits antibacterial activity.³ *Fusarium verticillioides* isolated from *Syzygium jambos* contains 3-hydroxy-4-(hydroxy(4-hydroxyphenyl)methyl)dihydrofuran-2-one compound which has antibacterial activity against *S. aureus* and *S. typhi*.⁴ Oxidation is a chemical reaction that results in the loss of electrons from atoms and produces free radicals.⁵ Free radicals are groups of atoms, molecules, or ions whose electrons are unpaired, unstable, and active in chemical reactions with other molecules.

*Corresponding author. E mail: kurratulaini_uin@radenfatah.ac.id; elfita.unsri.69@gmail.com

Tel: +62711-580056

Citation: Aini K, Elfita E, Widjajanti H, Setiawan A. Bioactivity Endophytic Fungi Isolated from the Leaf Stalk of *Syzygium jambos* L. Alston. Trop J Nat Prod Res. 2022; 6(11):1765-1772. <http://www.doi.org/10.26538/tjnpr/v6i11.4>

Official Journal of Natural Product Research Group, Faculty of Pharmacy, University of Benin, Benin City, Nigeria.

These unpaired electrons will seek and capture electrons from other substances to stabilize themselves. These free radical molecules are referred to as reactive oxygen species (ROS).⁶ Antioxidants play a role in reducing, preventing, and treating diseases caused by ROS. ROS are produced by metabolic pathways in cells, which play a role in the response to abiotic and biotic stresses.^{7,8} Antioxidant compounds have been widely used in the food, pharmaceutical, agricultural, and beauty product industries. The necessity for natural antioxidants is currently the main focus of several studies. EF produce bioactive compounds that have biological activity as antioxidant,^{5,9,10} such as *Fusarium oxysporum* from *S. jambos*.¹¹ To overcome synthetic antibiotic resistance and prevent free radicals, natural sources of antibiotics and antioxidants are required. Many studies have been carried out to find sources of medicinal crude materials and antioxidants derived from medicinal plants, one of which is jambu mawar (*Syzygium jambos*). *S. jambos* is known to contain active ingredients that have activity against pathogenic bacteria and as antioxidants.¹² The methanol extract of the *S. jambos* leaves and stem bark showed activity against *Staphylococcus aureus*.^{13,14} The leaves' ethanol extracts¹⁵⁻¹⁸ and fruit of the *S. jambos*¹⁹ have antioxidant activity. However, the existence of this jambu mawar is difficult to find and requires long cultivation periods to be used as raw material. Therefore, the alternative is to isolate the EF associated with *S. jambos* plants. Antioxidant activity is all about the elimination of free radicals and nascent oxygen [O]. On the other hand, many bacteria can also depend on this nascent oxygen to survive. The removal of free radicals or oxidants will affect antimicrobial activity. This could partially explained that antioxidant

properties correlate positively with antimicrobial activity. This correlation is the basis for finding new natural source that plays a role as antibacterial and antioxidant, such as the use of endophytic fungi.²⁰⁻²² EF associates and live in plant tissues without causing symptoms, disturbances, or diseases in their host plants, and is a promising source of bioactive agents.²³⁻²⁵ EF associated with medicinal plants have the potential to produce bioactive compounds. Recently, many researchers have conducted research on the potential of EF as a source of bioactive compounds. EF are important sources of bioactive compounds with unique chemical and biological activities.²⁶ EF produce metabolic compounds that are beneficial to humans as bioactive ingredients against pathogenic bacteria, viruses, cancer, treat diabetes, and antioxidants.^{27,28} The types of endophytic fungi obtained cannot be predicted due to environmental conditions that affect the diversity of endophytic fungi on the host plant. Therefore, this study is to determine the antibacterial activity, antioxidant, and compounds contained in endophytic fungi isolated from the *S. jambos* leaf stalk.

Materials and Methods

Sampling

A sampling of *Syzygium jambos* (L.) Alston leaf stalks was taken in August 2021 at the versatile ground of Perumahan Kencana Damai, Sukamaju, Sako, Palembang. The sample was obtained from the same tree, where the leaf stalks were from five leaves. The sample was identified morphologically in the Biosystematics Laboratory, Department of Biology, Faculty of Mathematics and Natural Science, Universitas Sriwijaya, Palembang with letter number 233/UN9.1.7/4/EP/2021.

Isolation and identification of endophytic fungi

Syzygium jambos (L.) Alston leaf stalks were washed with clean water and drained until the water dried. Surface sterilization began by immersing each sample in 3% sodium hypochlorite (NaOCl) for \pm 1 min. They were soaked in 70% alcohol for \pm 1 min, rinsed with sterile distilled water for \pm 1 min,^{29,30} placed in a petri dish containing potato dextrose agar (PDA, Merck®), which was added with chloramphenicol (0.2 g/L), and incubated at room temperature for 48 hr.¹¹ Pure isolates of EF were identified macroscopically and microscopically. Morphological characteristics were identified based on growth patterns, colony color, texture, margin, and other characteristics. Microscopic characteristics include the shape of hyphae and spores observed under a microscope (Hirox MXB-2500REZ), with the slide culture method. The results of microscopic and macroscopic observations will be compared with identification books for fungi^{29,30} and relevant journal articles.

Cultivation and extraction of endophytic fungi

Pure isolates of EF *S. jambos* leaf stalk were cultivated as much as 5 x 300 mL in potato dextrose broth by placing 6 blocks of pure culture in bottles. Then the cultures were incubated for four weeks (\pm 28 days) at room temperature under static conditions. After the incubation period, the endophytic fungal mycelia were separated from the liquid culture and partitioned in ethyl acetate at a ratio of 1:1 three times. The ethyl acetate extract was separated from the liquid culture, then evaporated using a rotary evaporator until a thick extract was obtained.³¹

Antibacterial activity test of endophytic fungi

Antibacterial activity was tested using the paper disc method with 400 μ g/mL by using Dimethylsulfoxide (DMSO) as solvent and positive control using tetracycline antibiotic 30 μ g/mL. The tested bacteria were represented by Gram-negative and Gram-positive bacteria, namely *Salmonella typhi* (ATCC 1408), *Escherichia coli* (InaCCB5), *Staphylococcus aureus* (InaCCB4), and *Bacillus subtilis* (InaCCB4) grown on Nutrient Agar media (NA). Antibacterial activity was indicated by the clear zone around the paper disc after being incubated for 24 hr at room temperature. The clear zone formed indicates the sensitivity of bacteria to antibacterial ingredients from EF extract. The criteria for antibacterial activity consist of strong, moderate, and weak which are determined by the following equation.³² With A = zone of

inhibition of the test sample (mm); B = zone of inhibition standard antibiotic (mm).

$$\text{Weak: } \frac{A}{B} \times 100\% < 50\%; \text{ Moderate: } 50\% < \frac{A}{B} \times 100\% < 70\%; \text{ Strong: } \frac{A}{B} \times 100\% > 70\%$$

Antioxidants activity assay of endophytic fungi

An antioxidant activity test was performed using the 2,2-diphenyl-1-picrylhydrazyl (DPPH) method. A total of 1.97 mg of DPPH was dissolved in 100 mL of 90% methanol solution so that a solution of 100 mL of 0.05 mM DPPH was obtained. The ethyl acetate extract of EF and ascorbic acid was dissolved to obtain concentrations of 50, 25, 12.5, 6.25, and 3.125 ppm. A total of 0.2 mL of EF extract of *S. jambos* leaf stalk was added with 3.8 mL of 0.05mM DPPH solution. The sample was placed in a dark place (without light) for 30 min. After that, the solution was analyzed using a UV-Vis spectrophotometer (Shimadzu UV-1900) $\lambda_{\text{maks}} = 517$ nm so that the absorbance value was obtained.³³ The absorbance results were obtained from the sample EF of *S. jambos* leaf stalk (a) and negative control (b), then the % inhibition was calculated using the formula:

$$\% \text{ inhibition} = \frac{\text{control absorbance} - \text{sample absorbance}}{\text{control absorbance}} \times 100 \%$$

The linear regression equation, $y = b + ax$, is obtained where the x -axis is the sample concentration and the y -axis is the % inhibition. The IC₅₀ value is the concentration value that can inhibit 50% of a test. Mathematically it can be seen as follows:

$$IC_{50} = \frac{50-b}{a}$$

Molecular identification of endophytic fungi

EF isolates that have strong antibacterial and antioxidant activity were identified molecularly to determine the species of the endophytic fungi. The molecular test of EF was carried out at the laboratory of Genetika Science Indonesia, Banten, Indonesia using method of Genomic DNA extraction with Quick-DNA Fungal Miniprep Kit (Zymo Research, D6005), polymerase chain reaction (PCR) amplification (Bioline, BIO-25048) with MyTaq HS Red Mix twice. Standard PCR primers using ITS1 dan ITS4.³⁴ Analysis of DNA structure using Molecular Evolution Genetics Analysis Version 11.³⁵

Isolation of chemical compound

Ethyl acetate extract of endophytic fungi exhibited strong antibacterial and antioxidant activity, then isolated the bioactive compounds. The bioactive compounds was analyzed using a thin-layer chromatography (TLC) silica gel G-60 F254 using *n*-hexane: EtOAc (5:5). The concentrated extract was separated by column chromatography over silica gel 60 (70–230 mesh) at the stationary phase (1:30) and eluent. The chosen eluent with increased polarity was *n*-hexane: EtOAc at a ratio of 10:0 to 0:10 (v/v). An eluate was collected and then tested using thin-layer chromatography with mixed eluent *n*-hexane: EtOAc (5:5). Similar chromatogram patterns were combined, evaporated, and rinsed with *n*-hexane to obtain the purified compound. The chemical structure of the compounds was determined using the following spectroscopy methods: 1DNMR (¹H-NMR and ¹³C-NMR).³⁶

Results and Discussion

Isolation and identification of endophytic fungi

The results of the isolation of EF from *S. jambos* leaf stalk found four isolates (SJS1 – SJS4). EF isolates were identified macroscopically by observing the front and bottom surfaces of the colonies, and colony color (Figure 1 and Table 1). Microscopic observations were carried out to determine the hyphae, shape, and type of spores (Figure 2 and Table 2). Four different genera were found: *Lasiodiplodia*, *Bipolaris*, *Aureobasidium*, and *Cladophialophora*.




Figure 1: EF 7 days on PDA of *S. jambos* leaf stalks; a) Isolate colony surface; b) Isolate colony reverse.




Figure 2: Microscopic characteristics (magnification: 1000x) of the EF 7 days on PDA of *S. jambos* leaf stalks

Antibacterial activity

The antibacterial activity of EF from *S. jambos* leaf stalk was evaluated using the paper disc diffusion method by measuring the zone of inhibition. The results of the antibacterial test are shown in Table 3. The results showed that EF SJS1 had strong activity against *S. aureus* and *S. typhi* bacteria, and moderate activity against *E. coli* and *B. subtilis*. The EF SJS4 had moderate activity against the four tested bacteria. SJS2 and SJS3 isolates had weak antibacterial activity against *E. coli* bacteria and moderate activity against *S. typhi*, *S. aureus* and *B. subtilis*.

Antioxidants activity

Antioxidant activity was tested using the DPPH method with parameter IC50. The antioxidant test results of the five isolates of EF showed that SJS1 isolates had a strong activity with an IC50 value of 29.29 µg/mL, while the other three isolates were very weak (Table 4 and Fig. 3). Many studies have been on the diversity of EF and their biological activities, especially in the genus *Syzygium*. Antibacterial activity was shown by EF isolated from the stem bark of *Syzygium jambos*,⁴ from leaves, bark, and root bark of *S. malaccense*³⁷ from root bark of *S. zeylanicum*,³⁸ from leaves and leaf stalks of *S. aqueum*.³³ Antioxidant activity was also demonstrated by EF isolated from *S. cummini* (*Eugenia jambolana*),³⁹ and from *S. aqueum*.⁴⁰ The antioxidant compound found in EF isolated from *S. aqueum* was 3-(hidroksil (3,4,5-trihidroksifenil)metil)-3,4-dihidro-2H-piran-4,5,6-triol, the antioxidant power was close to the activity of gallic acid with IC50 value of 11.4 µg/mL.⁴⁰ Alkaloids, phenols, flavonoids, saponins, and terpenes isolated from the EF *Eugenia jambolana* have antioxidant potential. These compounds exhibit antioxidant activity by breaking free radical chains through the donation of hydrogen atoms.³⁹ The isolate of SJS1 (*Lasiodiplodia*) exhibited strong antibacterial and antioxidant activity. It is known that *L. theobromae* has antioxidant^{41,42} and antibacterial activity.^{43,44} EF *L. theobromae* contain secondary metabolites in the form of Chloropreussomerin A – B and preussomerin K, H, G, F that have antibacterial activity against bacteria *S. aureus* and *B. subtilis*;⁴⁵ also contain compounds cyclohexenes and cyclohexenones, indoles, jasmonates, lactones, melleins, phenols, lasiodiplodins,⁴⁶ L-asparaginase compounds as anti-leukemia.⁴⁷

Table 1: Macroscopic characteristics of EF isolated from *S. jambos* leaf stalks

Isolate	Colony color	Reverse colony color	Texture	Topography	Pattern	Exudate drops	Radial line	Concentric circle
SJS1	Black and White	Black and White	Cottony	Umbonate	Zonate	-	-	√
SJS2	Dark brown and White	Dark brown	Velvety	Rugose	Zonate	-	√	-
SJS3	Light green and White	Light green and White	Cottony	Raised	Radiated	-	-	-
SJS4	White to gray	White to gray	Cottony	Raised	Radiated	-	-	√

Note: (-) = characteristic doesn't appear; (√) = characteristic appear

Table 2: Microscopic characteristics EF isolated from *S. jambos* leaf stalks

Isolate	Type of spore	Shape of spore	Hyphae	Specific characteristic	Genus / species
SJS1	Conidia	Globose	Septate	showing characteristic 2-celled, pigmented, longitudinally striate pycnidioconidia	<i>Lasiodiplodia</i> sp.
SJS2	Conidia	Globose	Septate	Conidia porosporous, solitary, pale brown, ellipsoidal, usually 4-celled.	<i>Bipolaris</i> sp.
SJS3	Conidia	Globose	Hyaline	Conidiophores lacking, hyaline, cylindrical, 1-celled.	<i>Aureobasidium</i> sp.
SJS4	Conidia	Subglobose	Aseptate	Conidial chains	<i>Cladophialophora</i>

Table 3: Antibacterial Activity EF of *S. jambos* leaf stalk

Sample (400 µg/disc)	<i>S. typhi</i>	<i>E. coli</i>	<i>S. aureus</i>	<i>B. subtilis</i>
<i>Tetracycline</i>	20.9 ± 0.29 100***	21.9 ± 0.21 100***	20.6 ± 0.34 100***	20.9 ± 0.29 100***
SJS1	14.2 ± 0.29 67.9**	14.8 ± 0.21 67.6**	14.5 ± 0.08 70.4***	14.8 ± 0.34 70.8***
SJS2	11.1 ± 0.25 53.1**	12.1 ± 0.16 55.3**	10.5 ± 0.46 51.0**	10.2 ± 0.25 48.8*
SJS 3	11.9 ± 0.29 56.9**	11.1 ± 0.17 50.7**	12.8 ± 0.21 62.1**	10.5 ± 0.16 50.2**
SJS4	11.2 ± 0.26 53.6**	11.0 ± 0.08 50.2**	10.9 ± 0.21 52.9**	7.8 ± 0.25 37.3*

Note: *) Weak; **) Moderate; ***) Strong

Table 4: Antioxidants Activity EF of *S. jambos* leaf Stalk

Test Sample	IC50 (µg/mL)
SJS1	29.29 ± 0.45
SJS2	166.42 ± 0.98
SJS3	199.21 ± 0.71
SJS4	149.17 ± 0.91
Ascorbic Acid	8.09 ± 0.09

Lasiodiplactone A has potential as an antidiabetic and anti-inflammatory agent.⁴⁸ Jasmonic acid produced by *L. theobromae* is the main ingredient in cosmetic products.⁴⁹ In this study, the compound produced by *L. theobromae* from *S. jambos* was different from the compound *L. theobromae* that had been published. It was found that compounds belonging to the phenolic group have antibacterial and antioxidant activity. Different host plants cause different secondary metabolites in fungi of the same species. This is caused by the presence of fungi on the host plant that can copy the compounds of host plant.

Bipolaris (code SJS2) found in *S. jambos* leaf stalk also had moderate antibacterial activity. *Bipolaris* is a fungal pathogen that causes a leaf spot on palms.⁵⁰ *Bipolaris* A–H compounds, derivatives of tetracyclic ophiobolin-type sesterterpenes produced by *Bipolaris* have antibacterial activity.⁵¹ Hybrid polyketide–terpenoid, 1-alkylated-3,5-dihydroxyphenyl derivative coupled to a farnesyl pyrophosphate unit (FPP), isolated from the fungus *Bipolaris zeicola* has immunosuppressive and cytotoxic activity,⁵² sesquiterpenoids and xanthenes compounds have anti-pathogenic activity of microorganisms.⁵³ *Bipolaris eleusines* produces chromone (S)-5-hydroxyl-2-(1-hydroxyethyl)-7-methylchromone compounds which have inhibitory power against *S. aureus*,⁵⁴ Ophiobolin A compounds have anticancer activity by inducing cell death.⁵⁵ *Aureobasidium* (code SJS3) has moderate activity against all four bacteria. Based on a literature study, *Aureobasidium* produces liamocins compounds that have antibacterial against *Streptococcus* spp.,⁵⁸ and yield volatile organic compounds that have antifungal against *Botrytis cinerea*.⁵⁶ *Aureobasidium pullulans* produce enzymes that are useful in industry, such as β-glucosidase, amylases, cellulases, lipases, proteases, xylanases and mannanases which have cytotoxic, antioxidant, and antibacterial properties.⁵⁷ It also produces pullulan (poly-α-1,6-maltotriose biopolymer) which is applied in the food, cosmetic and pharmaceutical industries,⁵⁸ poly(β-L-malic acid) (PMA) heavy oils and β-1,3-glucan,⁵⁹ hydroxydecanoic acid derivatives have antibacterial activity against *E. coli*, *S. aureus*, and *B. subtilis*.^{60,61}

Molecular identification

Molecular identification was carried out to determine the EF species that have vigorous antibacterial and antioxidant activity. DNA

sequences resulting from PCR amplification of SJS1 isolates were compared with DNA sequences available in the GenBank database, to analyze the phylogenetic relationship. The results showed the DNA sequence of SJS1 isolates was, GGAGCTCGAA AACTCGGTAA TGATCCTTCC GTAGGTGAAC CTGCGGAAGG ATCATTACCG AGTTTTTCGAG CTCCGGCTCG ACTCTCCCAC CCTTTGTGAA CGTACCTCTG TTGCTTTGGC GGCTCCGGCC GCCAAAGGAC CTTCAAACCTC CAGTCAGTAA ACGCAGACGT CTGATAAACA AGTTAATAAAA CTAAAACCTTT CAACAACGGA TCTCTGGTT CTGGCATCGA TGAAGAACGC AGCGAAATGC GATAAGTAAT GTGAATTGCA GAATTCAGTG AATCATCGAA TCTTTGAACG CACATTGCGC CCCTTGATAT TCCGGGGGGC ATGCCTGTTC GAGCGTCATT ACAACCCTCA AGCTCTGCTT GGAATTGGGC ACCGTCCTCA CTGCGGACGC GCCTCAAAGA CCTCGGCGGT GGCTGTTTCAAG CCCTCAAGCG TAGTAGAATA CACCTCGCTT TGGAGCGGTT GGCGTCGCC GCCGGACGAA CCTTCTGAAC TTTTCTCAAG GTTGACCTCG GATCAGGTAG GGATACCCGC TGAACTTAAAG CATATCAATA AGCGGA, with 566bp, GenBank code OM095454. Molecular characterization combined with morphological identification can identify fungi until to the species level. Molecular characterization was carried out on the area of rDNA Internal Transcribed Spacer (ITS). Currently, there are more than 90,000 fungi sequences in the ITS region, where this area is most widely used as a barcode area for fungi. PCR analysis used a pair of universal primers, namely ITS1 for forwarding primers and ITS4 for reverse primers. The amplification results of the ITS rDNA region varied by ±500 bp (Fig. 4). Sequence readings from each of the forward and reverse primers were processed by cutting the ends of the sequences with low peaks using the Bioedit program. Furthermore, the sequences are straightened so that the compilation sequences of the forward and reverse primers from the ITS rDNA area are obtained. Based on BLAST, the ITS isolate sequence SJS1 showed 98–100% similarity to the sequences of the *Lasiodiplodia theobromae*.

The construction of the EF SJS1 phylogeny tree is shown in Figure 3. The SJS1 phylogenetic tree showed that the isolate sequence was close to *Lasiodiplodia theobromae* with a bootstrap value of 100. The results of the phylogenetic analysis of the SJS1 isolate are in line with the phenotypic characters that have been discussed previously. In this study, EF with strong antibacterial and antioxidant activity were carried out by molecular tests and identified as *Lasiodiplodia theobromae* (code SJS1). *L. theobromae* is a member of the *Botryosphaeriaceae* as an EF and latent pathogen that survives in plant vascular tissues for a certain period of time without showing symptoms under suitable environmental conditions.^{62–67} With changes in environmental conditions inside or outside the host, some fungi can change their lifestyle from endophytes to pathogens.^{62,68} Pathogenic symptoms caused by *L. theobromae* such as cancer, dieback, and rotting of fruit and roots in nearly 500 plant species globally.^{69,70}




Figure 3: Phylogenetic analysis of endophytic fungi SJS1 (isolated from leaf stalk of *S. jambos*). This phylogenetic tree was constructed by using neighbor-joining method (1000 bootstrap replication) in MEGA11. (*: isolate target).




Figure 4: Compound 1 as 3,5-dihydroxy-4-(4-hydroxyphenyl) tetrahydro-2H-pyran-2-one.

Pure compound isolation

The $^1\text{H-NMR}$ spectrum of compound 1 (Figure 1 in Supplementary data) shows the presence of six proton signals of which four are signals for proton sp^3 which is a methylene signal at δ_{H} 3.61-3.81 ppm (2H, m), a proton methine signal at δ_{H} 4.14 ppm (1H, m), and two oxygenated methine signals at δ_{H} 6.23 (1H, s) and 5.16 (1H, d, $J=3$ Hz). Two other signals that appear on the spectrum are in the aromatic chemical shift region, namely at δ_{H} 7.64 and 8.17 ppm. In the spectrum of each signal appears doublet with the same plot constant, namely $J = 8.0$ Hz which has the integration of two protons. This indicates that compound 1 is a para-substituted aromatic compound, so it has two pairs of equivalent aromatic protons. Based on the analysis of the $^1\text{H-NMR}$ spectrum, compound 1 is a para-substituted aromatic compound with nine protons bonded to eight carbon atoms. The solvent used in this measurement is CDCl_3 so that protons bound to heteroatoms do not appear in the spectrum. The $^{13}\text{C-NMR}$ spectrum of compound 1 (Figure 2 in Supplementary data) showed the presence of nine signals. There are four sp^3 carbon signals, all of which are in the oxygenated carbon region, namely δ_{C} 58.6, 62.3, 67.5, and 71.4 ppm. Five other signals appear at $\delta_{\text{C}} > 100$ ppm. Two high-intensity signals indicate that compound 1 has two pairs of equivalent aromatic carbons (δ_{C} 124.2 and 128.4 ppm). Three carbon signals in the low field indicate the presence of oxyaryl carbon (δ_{C} 148.7 ppm), aromatic quaternary carbon (δ_{C} 151.7 ppm), and ester carbonyl carbon (δ_{C} 166.7 ppm). The analysis of the proton and carbon NMR spectra is reinforced by the data on the HMQC spectrum shown in Figure 2 (Supplementary data) and Table 5, namely the $^1\text{H-}^{13}\text{C}$ correlation through one bond. The HMQC spectrum showed six correlations consisting of two correlations on the aromatic ring, three correlations on oxygenated $^1\text{H-}^{13}\text{C}$, and one methylene proton correlation. The HMBC spectrum (Figure 3 in Supplementary data) showed a $^1\text{H-}^{13}\text{C}$ correlation through two or three bonds. The aromatic proton signal at δ_{H} 8.17 ppm indicates that there are three correlations, namely to carbon δ_{C} 124.2 ppm which is the carbon equivalent, δ_{C} 151.7 ppm which is aromatic quaternary carbon, and δ_{C} 148.7 ppm which is oxyaryl carbon. Another aromatic proton, namely at δ_{H} 7.64 ppm also has three correlations, namely to carbon at δ_{C} 128.4, 148.7, and 71.4 ppm which are carbon equivalents, oxyaryl carbon, and oxygenated carbon in their substituents, respectively. Furthermore, oxygenated methine protons at δ_{H} 5.16 ppm have four correlations, namely to carbon at δ_{C} 58.6, 62.3, 128.4, and 151.7 ppm, respectively. The correlation indicates that the oxygenated methine group is bound via three bonds with a para-substituted aromatic ring with a hydroxyl group. The proton methine at δ_{H} 4.14 ppm is triple-bonded with methylene carbon (δ_{C} 62.3 ppm) and carbonyl ester carbon (δ_{C} 166.7 ppm). The methylene proton at δ_{H} 3.61-3.81 ppm has two correlations, namely with oxygenated carbon at δ_{C} 71.4 ppm and methine carbon at δ_{C} 58.6 ppm. The proton hydroxyl signal does not appear on the spectrum because the pure compound is measured with the solvent CDCl_3 . Spectrum data for 1D NMR and 2D compounds are listed in Table 5.

Table 5: The NMR data of compound 1, recorded at $^1\text{H-}500$ MHz; $^{13}\text{C-}125$ MHz in CDCl_3

No. C	δ_{C} ppm	Type of C	δ_{H} ppm (ΣH , multiplicity, J (Hz))	HMBC
2	166.7	C		
3	67.5	CH	6.23 (1H, s)	166.7
4	58.6	CH	4.14 (1H, m)	62.3; 166.7
5	71.4	CH	5.16 (1H, d, $J=3$ Hz)	58.6; 62.3; 128.4; 151.7
6	62.3	CH_2	3.61-3.81 (2H, m)	58.6; 71.4
1'	148.7	C		
2'	128.4	CH	7.64 (1H, d, $J=8$ Hz)	71.4; 128.4; 148.7
3'	124.2	CH	8.17 (1H, d, $J=8$ Hz)	124.2; 148.7; 151.7
4'	151.7	C		
5'	124.2	CH	8.17 (1H, d, $J=8$ Hz)	124.2; 148.7; 151.7
6'	128.4	CH	7.64 (1H, d, $J=8$ Hz)	71.4; 128.4; 148.7

Based on the spectrum analysis of ¹H-NMR, ¹³C-NMR, HMQC, dan HMBC and HMBC, it can be explained that compound 1 has a substituted benzene ring at the para position respectively is a hydroxyl group and a pyran ring i.e. 3,5-dihydroxytetrahydro-2H-pyran-2-one. Thus, the proposed chemical structure of compound 1 is 3,5-dihydroxy-4-(4-hydroxyphenyl)tetrahydro-2H-pyran-2-one as shown in Figure. 4.

Conclusion

Four EF were found isolated from *Syzygium jambos* leaf stalk, namely *Lasiodiplodia*, *Bipolaris*, *Aureobasidium* dan *Cladophialophora*. These four EF were tested for antibacterial and antioxidant. The EF SJS1 -- *Lasiodiplodia theobromae* has high antibacterial activity against *S. aureus* and *B. subtilis* bacteria. Antioxidant activity of *L. theobromae* showed strong activity with an IC50 value of 29.29 µg/mL. Isolation of pure compound from *L. theobromae* obtained compound 3,5-dihydroxy-4-(4-hydroxyphenyl)tetrahydro-2H-pyran-2-one, this compound belongs to the class of phenolic.

Conflict of Interest

The authors declare no conflict of interest.

Authors' Declaration

The authors hereby declare that the work presented in this article is original and that any liability for claims relating to the content of this article will be borne by them.

Acknowledgments

The authors thank to the DRPM Kemenristek Republik Indonesia, which provided research funding through Hibah Disertasi Doktor 2022 as the third output, with contract no. 0166.05/UN9/SB3.LP2M.PT/2022.

References

- Toghueo RMK. Bioprospecting endophytic fungi from *Fusarium* genus as sources of bioactive metabolites. *Mycology*. 2020;11(1):1–21.
- Du W, Yao Z, Li J, Sun C, Xia J, Wang B, Shi D, Ren L. Diversity and Antimicrobial Activity of Endophytic Fungi Isolated from *Securinega suffruticosa* in the Yellow River Delta. *PLoS One*. 2020; 15(3):1–18.
- An C, Ma S, Shi X, Xue W, Liu C, Ding H. Diversity and Antimicrobial Activity of Endophytic Fungi Isolated from *Chloranthus japonicus* Sieb in Qinling Mountains, China. *Int J Mol Sci*. 2020; 21(17):1–15.
- Aini K, Elfita, Widjajanti H, Setiawan A, Kurniawati AR. Antibacterial activity of endophytic fungi isolated from the stem bark of jambu mawar (*Syzygium jambos*). *Biodiversitas*. 2022; 23(1):521–532.
- Manganyi MC and Ateba CN. Untapped Potentials of Endophytic Fungi: A Review of Novel Bioactive Compounds with Biological Applications. *Microorgan*. 2020; 8(12):1–25.
- Atta EM, Mohamed NH, Abdelgawad AAM. Antioxidants: an Overview on the Natural and Synthetic Types. *Eur Chem Bull*. 2017; 6(8):365–375.
- Mhamdi A and Van Breusegem F. Reactive oxygen species in plant development. *Dev*. 2018; 145(15):1–12.
- Das K and Roychoudhury A. Reactive oxygen species (ROS) and response of antioxidants as ROS-scavengers during environmental stress in plants. *Front Environ Sci*. 2014; 2(DEC):1–13.
- Ibrahim M, Oyebanji E, Fowora M, Aiyeolemi A, Orabuchi C, Akinnawo B, Adekunle AA. Extracts of endophytic fungi from leaves of selected Nigerian ethnomedicinal plants exhibited antioxidant activity. *BMC Complement Med Ther*. 2021; 21(1):1–13.
- Hidayat A, Turjaman M, Faulina SA, Ridwan F, Aryanto, Najmulah, Irawadi, TT, Iswanto AH. Antioxidant and Antifungal Activity of Endophytic Fungi Associated with Agarwood Trees. *J Korean Wood Sci Technol*. 2019; 47(4):459–471.
- Caicedo NH, Davalos AF, Puente PA, Rodriguez AY, Caicedo PA. Antioxidant Activity of Exo-Metabolites Produced by *Fusarium oxysporum*: An Endophytic Fungus Isolated from Leaves of *Otoba gracilipes*. *Microbiol Open*. 2019;8(10):1–7.
- Lim TK. *Syzygium jambos*. In: *Edible Medicinal And Non Medicinal Plants*. Springer Netherlands; 2012; 760–766.
- Wamba BEN, Nayim P, Mbaveng AT, Voukeng IK, Dzotam JK, Ngalani OJT, Kuete V. *Syzygium jambos* Displayed Antibacterial and Antibiotic-Modulating Activities against Resistant Phenotypes. Evidence-based Complement Altern Med. 2018;2018. doi:10.1155/2018/5124735
- Fathima SBN, Pandian R, Perumal PV. Antimicrobial Activity of Methanol and Ethyl Acetate Extracts of Three Medicinal Plants. *Int J Adv Sci Eng Technol*. 2017; 5(1):16–22.
- Jayasinghe ULB, Ratnayake RMS, Medawala MMWS, Fujimoto Y. Dihydrochalcones with Radical Scavenging Properties from the Leaves of *Syzygium jambos*. *Nat Prod Res*. 2007; 21(6):551–554.
- Hossain H, Rahman SE, Akbar PN, Khan TA, Rahman MM, Jahan IA. HPLC Profiling, Antioxidant and *In Vivo* Anti-Inflammatory Activity of The Ethanol Extract of *Syzygium jambos* Available in Bangladesh. 2016; 1–8.
- Devakumar J and Sudha SS. *In vitro* Phytochemical, Antioxidant and Cytotoxic Evaluation of *Syzygium jambos* L. (Alston). *J Pharm Res*. 2017; 11(3):235–238.
- Selvam NT, Venkatakrishnan V, Kumar SD. Tissue Level Antioxidant Activity of Leaf Extract of *Syzygium jambos* Linn. In Paracetamol Intoxicated Wistar rats. *J Intern Med*. 2014; 2(8):107–111.
- Reynertson KA, Kennelly EJ, Basile MJ. Antioxidant Potential of Seven Myrtaceae Fruits. 2005; 25–35.
- Borah SN, Goswami D, Sarma HK, Cameotra SS, Deka S. Rhamnolipid Biosurfactant against *Fusarium verticillioides* to Control Stalk and Ear Rot Disease of Maize. *Front Microbiol*. 2016; 7:1–10.
- Sen A and Imlay JA. How Microbes Defend Themselves From Incoming Hydrogen Peroxide. *Front Immunol*. 2021; 12(April):1–19.
- Nakai K and Tsuruta D. What are reactive oxygen species, free radicals, and oxidative stress in skin diseases? *Int J Mol Sci*. 2021; 22(19):1–16.
- Wilson D. Endophyte: The Evolution of a Term, and Clarification of Its Use and Definition. *Oikos*. 1995;73(2):274.
- Selim KA, El-Beih AA, Abdel-Rahman TM, El-Diwany AI. Biology of Endophytic Fungi. *Curr Res Environ Appl Mycol*. 2012; 2(1):31–82.
- Elfita, Munawar, Muharni, Sudrajat MA. Identification of New Lactone Derivatives Isolated from *Trichoderma* sp., An Endophytic Fungus of Brotowali (*Tinaspora crispa*). *HAYATI J Biosci*. 2014; 21(1):15–20.
- Gakuubi MM, Munusamy M, Liang ZX, Ng SB. Fungal Endophytes: A Promising Frontier for Discovery of Novel Bioactive Compounds. *J Fungi*. 2021;7(10):1–24.
- Adeleke BS and Babalola OO. Pharmacological Potential of Fungal Endophytes Associated with Medicinal Plants: A Review. *J Fungi*. 2021; 7(2):1–16.
- Rajamanikyam M and Vadlapudi V. Endophytic Fungi as Novel Resources of natural Therapeutics. *Brazilian Arch Biol Technol*. 2017; 60:1–26.
- Watanabe T. *Pictorial Atlas of Soil and Seed Fungi: Morphologies of Cultural Fungi and Key to Species*. Third Edit. New York: CRC Press: Taylor & Francis Group; 2010.
- Walsh TH, Hayden RT, Larone DH. *Larone's Medically Important Fungi: A Guide to Identification*. 6th Edition. ASM Press, Washington, DC; 2018.
- Nagarajan D. *In Vitro* Antioxidant Potential of Endophytic Fungi Isolated from *Enicostemma axillare* (Lam.) Raynal. and *Ormocarpum cochinchinense* (Lour.) Merr. *J Pharmacogn Phytochem*. 2019; 8(1):1356–1363.
- Elfita, Mardiyanto, Fitriya, Larasati JE, Julinar, Widjajanti H,

- Muharni. Antibacterial Activity of *Cordyline fruticosa* Leaf Extracts and Its Endophytic Fungi Extracts. *Biodiversitas*. 2019; 20(12):3804–3812.
33. Habisukan UH, Elfita, Widjajanti H, Setiawan A, Kurniawati AR. Antioxidant and Antimicrobial Activity of Endophytic Fungi Isolated from *Syzygium aqueum* Leaves. *J Phys Conf Ser*. 2021; 1751(1):1–8.
 34. Singha IM, Kakoty Y, Unni BG, Das J, Kalita MC. Identification and Characterization of *Fusarium* sp. Using ITS and RAPD causing Fusarium Wilt of Tomato Isolated from Assam, North East India. *J Genet Eng Biotechnol*. 2016; 14(1):99–105.
 35. Tamura K, Stecher G, Kumar S. MEGA11: Molecular Evolutionary Genetics Analysis Version 11. *Mol Biol Evol*. 2021; 38(7):3022–3027.
 36. Elfita E, Munawar M, Muharni M, Pratiwi G, Rahmadania R. A New Benzoyl Compound Isolated from the Endophytic Fungi of Kandis Gajah (*Garcinia griffithii*) and Asam Kandis (*Garcinia cowa*). *Makara J Sci*. 2016; 20(4):167–172.
 37. Hapida Y, Elfita, Widjajanti H, Salni. Biodiversity and antibacterial activity of endophytic fungi isolated from jambu bol (*Syzygium malaccense*). *Biodiversitas*. 2021;22(12):5668–5677.
 38. Elfita S, Widjajanti H, Setiawan A, Kurniawati AR. Diversity of Endophytic Fungi from The Root Bark of *Syzygium zeylanicum*, and The Antibacterial Activity of Fungal Extracts, and Secondary Metabolite. *Biodiversitas*. 2021; 22(10):4572–4582.
 39. Yadav M, Yadav A, Yadav JP. *In vitro* antioxidant activity and total phenolic content of endophytic fungi isolated from *Eugenia jambolana* Lam. *Asian Pac J Trop Med*. 2014; 7(S1):S256–S261.
 40. Habisukan UH, Elfita E, Widjajanti H, Setiawan A. Secondary Metabolite and Antioxidant Activity of Endophytic Fungi Isolated from *Syzygium aqueum* Leaves Stalk. *Biointerface Res Appl Chem*. 2022; 12(6):7584–7595.
 41. Jinu MV, Gini CK, Jayabaskaran C. *In vitro* antioxidant activity of cholestanol glucoside from an endophytic fungus *Lasiodiplodia theobromae* isolated from *Saraca asoca*. *J Chem Pharm Res*. 2015;7(1):952–962.
 42. Zhou J, Diao X, Wang T, Chen G, Lin Q, Yang X, Xu J. Phylogenetic diversity and antioxidant activities of culturable fungal endophytes associated with the mangrove species *Rhizophora stylosa* and *R. mucronata* in the South China Sea. *PLoS One*. 2018; 13(6):1–18.
 43. Taufiq MMJ and Darah I. Anti-Mrsa of the Ethyl Acetate Crude Extract From *Lasiodiplodia Pseudotheobromae* Ibrl Os-64, an Endophytic Fungus Isolated From Leaf of *Ocimum Sanctum* Linn. *Int J Pharm Pharm Sci*. 2018; 10(8):50.
 44. Setyaningsih R, Susilowati A, Prasetyani D. Antibacterial Activity of Endophytic Fungi in *Pometia pinnata* Against *Staphylococcus aureus* and *Methicillin-Resistant Staphylococcus aureus*. *Biodiversitas*. 2020;21(11):5408–5413.
 45. Chen S, Chen D, Cai R, Cui H, Long Y, Lu Y, Li C, She Z. Cytotoxic and Antibacterial Preussomerins from the Mangrove Endophytic Fungus *Lasiodiplodia theobromae* ZJ-HQ1. *J Nat Prod*. 2016; 79(9):2397–2402.
 46. Salvatore MM, Alves A, Andolfi A. Secondary metabolites of *Lasiodiplodia theobromae*: Distribution, chemical diversity, bioactivity, and implications of their occurrence. *Toxins (Basel)*. 2020; 12(7):1–29.
 47. Balbool BA, Abdel-azeem AM, Moubasher MH, Helmy EA. Production of L-Asparaginase (L-ASN) from endophytic *Lasiodiplodia theobromae* hosted *Teucrium polium* in Egypt. *Microb Biosyst*. 2018; 3(2):46–55.
 48. Chen S, Liu Z, Liu H, Long Y, Chen D, Lu Y, She Z. Lasiodiplactone A, a novel lactone from the mangrove endophytic fungus *Lasiodiplodia theobromae* ZJ-HQ1. *Org Biomol Chem*. 2017; 15(30):6338–6341.
 49. Eng F, Haroth S, Feussner K, Meldau D, Rehkter D, Ischebeck T, Brodhun F, Feussner I. Optimized Jasmonic Acid Production by *Lasiodiplodia theobromae* Reveals Formation of Valuable Plant Secondary Metabolites. *PLoS One*. 2016; 11(12):1–18.
 50. Dagher GM, Tomah AA, Alyaseri II. First Record the Fungus *Bipolaris australiensis* as a Cause of Leaf Blight Date Palm in Misan and Attempt to Control It in *In Vitro* Condition. *Plant Arch*. 2020; 20(July):966–970.
 51. Liu MT, He Y, Shen L, Hu ZX, Zhang YH. Bipolarins A–H, eight new ophiobolin-type sesterterpenes with antimicrobial activity from fungus *Bipolaris* sp. TJ403-B1. *Chin J Nat Med*. 2019; 17(12):935–944.
 52. Liu M, Zhang X, Shen L, Sun W, Lin S, Liu J, Cao F, Qi C, Wang J, Hu Z, Zhang Y. Bioactive Polyketide-Terpenoid Hybrids from a Soil-Derived Fungus *Bipolaris zeicola*. *J Org Chem*. 2021; 86(16):10962–10974.
 53. Yu JJ, Jin YX, Huang SS, He J. Sesquiterpenoids and xanthenes from the kiwifruit-associated fungus *Bipolaris* sp. And their anti-pathogenic microorganism activity. *J Fungi*. 2022; 8(9):1–17.
 54. He J, Li ZH, Ai HL, Feng T, Liu JK. Anti-bacterial chromones from cultures of the endophytic fungus *Bipolaris eleusines*. *Nat Prod Res*. 2019; 33(24):3515–3520.
 55. Maehara S, Yamane C, Kitamura C, Hinokuma M, Hata T. High ophiobolin A production in endophytic fungus *Bipolaris* sp. associated with *Datura metel*. *Nat Prod Res*. 2020;34(20):2990–2992. doi:10.1080/14786419.2019.1597352
 56. Di Francesco A, Zajc J, Gunde-Cimerman N, Aprea E, Gasperi F, Placi N, Caruso F, Baraldi E. Bioactivity of volatile organic compounds by *Aureobasidium* species against gray mold of tomato and table grape. *World J Microbiol Biotechnol*. 2020; 36(171):1–11.
 57. Botić T, Kralj-Kunčić M, Sepčić K, Batista U, Zalar P, Knez Z, Gunde-Cimerman N. Biological activities of organic extracts of four *Aureobasidium pullulans* varieties isolated from extreme marine and terrestrial habitats. *Nat Prod Res*. 2014; 28(12):874–882.
 58. Bozoudi D and Tsaltas D. The Multiple and Versatile Roles of *Aureobasidium pullulans* in the Vitivicultural Sector. *Fermentation*. 2018; 4(85):1–15.
 59. Li Y, Chi Z, Wang GY, Wang ZP, Liu GL, Lee CF, Ma ZC, Chi ZM. Taxonomy of *Aureobasidium* spp. and biosynthesis and regulation of their extracellular polymers. *Crit Rev Microbiol*. 2015; 41(2):1–10.
 60. Choi HG, Kim JW, Choi H, Kang KS, Shim SH. New Hydroxydecanoic Acid Derivatives Produced by an Endophytic Yeast *Aureobasidium pullulans* AJF1 from Flowers of *Aconitum carmichaeli*. *Molecules*. 2019; 24(22):1–11.
 61. Abdel-Lateff A, Elkhayat ES, Fouad MA, Okino T. Aureobasidin, New Antifouling Metabolite from Marine-Derived Fungus *Aureobasidium* sp. *Nat Prod Commun*. 2009; 4(3):389–394.
 62. Harishchandra DL, Zhang W, Li X, Chethana KWT, Hyde KD, Brooks S, Yan J, Peng J. A LysM Domain-Containing Protein LtLysM1 Is Important for Vegetative Growth and Pathogenesis in Woody Plant Pathogen *Lasiodiplodia theobromae*. *Plant Pathol J*. 2020; 36(4):323–334.
 63. Daengsuwan W, Wonglom P, Sunpapao A. First report of *Lasiodiplodia theobromae* causing spadix rot in *Anthurium andraeanum*. *J Phytopathol*. 2020; 168(2):129–133.
 64. Munirah MS, Azmi AR, Yong SYC, Nur Ain Izzati MZ. Characterization of *Lasiodiplodia theobromae* and *L. pseudotheobromae* causing fruit rot on pre-harvest mango in Malaysia. *Plant Pathol Quar*. 2017;7(2):202–213.
 65. Netto MSB, Assunção IP, Lima GSA, Marques MW, Lima WG, Monteiro JHA, Balbino VDQ, Michereff SJ, Phillips AJL, Camara MPS. Species of *Lasiodiplodia* associated with papaya stem-end rot in Brazil. *Fungal Divers*. 2014; 67(1):127–141.
 66. Netto MSB, Lima WG, Correia KC, da Silva CFB, Thon M, Martins RB, Miller RNG, Michereff SJ, Camara MPS. Analysis of phylogeny, distribution, and pathogenicity of Botryosphaeriaceae species associated with gummosis of *Anacardium* in Brazil, with a new species of *Lasiodiplodia*. *Fungal Biol*. 2017; 121(4):437–451.
 67. Sathya K, Parthasarathy S, Thiribhuvanamala G, Prabakar K. Morphological and Molecular Variability of *Lasiodiplodia theobromae* Causing Stem End Rot of Mango in Tamil Nadu, India. *Int J Pure Appl Biosci*. 2017;5(6):1024–1031.
 68. Peng J, Wu L, Zhang W, Zhang Q, Xing Q, Wang X, Li X, Yan J. Systemic Identification and Functional Characterization of

- Common in Fungal Extracellular Membrane Proteins in *Lasiodiplodia theobromae*. *Front Plant Sci.* 2021; 12(December):1–15.
69. Marsberg A, Kemler M, Jami F, Nagel JH, Postma-Smidt A, Naidoo S, Wingfield MJ, Crous PW, Spatafora J, Hesse CN, Robbertse B, Slippers B. *Botryosphaeria dothidea*: A latent pathogen of global importance to woody plant health. *Mol Plant Pathol.* 2016; 18(4):477–488.
70. Paolinelli-Alfonso M, Villalobos-Escobedo JM, Rolshausen P, Herrera-Estrella A, Galindo-Sanchez C, Lopez-Hernandez JF, Hernandez-Martinez R. Global transcriptional analysis suggests *Lasiodiplodia theobromae* pathogenicity factors involved in modulation of grapevine defensive response. *BMC Genomics.* 2016; 17(1):1–20.

Editorial Team

Editorial Board

Editor-in-Chief

Professor Abiodun Falodun

Professor of Pharmaceutical Chemistry

Natural product Research Group

University of Benin, Nigeria.

Email: editor.tjnpr@gmail.com; editor.tjnpr@uniben.edu

(<mailto:editor.tjnpr@gmail.com>; editor.tjnpr@uniben.edu)

Phone: +2348073184488

Associate Editors:

- **Professor Dr. Dr. Peter Langer**, Institute of Organic Chemistry, University of Rostock, (Germany)
- **Professor Frederick O. Ekhaïse**, Microbiology, University of Benin, Nigeria.
- **Professor Martins Emeje**: Professor of Drug Delivery/Nanomedicine in the Department of Pharmaceutical Technology and Raw Materials Development (PT&RMD) at the National Institute of Pharmaceutical Research and Development.

Editorial Assistant:

- **Erharuyi Osayemwenre**, Faculty of Pharmacy, University of Benin, Nigeria.

Board Members


- **Professor Ikhlas A. Khan**, National Center for Natural Product Research, Mississippi (USA)
- **Professor Nosa Egiebor**, College of Environmental Science & Forestry, State University of New York
- **Professor Samuel Qiu**, South China Botanical Gardens, Chinese Academy of Sciences (China)
- **Professor Xavier Barril**, De Fisicoquímica-Facultat De Farmàcia Universitat de Barcelona, (Spain)
- **Professor Abiodun Ogundiani**, Pharmaceutical Chemistry, OAU, Ile-Ife, (Nigeria)
- **Professor Thomas Kodadek**, The Scripps Research Institute, Scripps Florida (USA)
- **Professor Anthony B Ebeigbe**, Physiology, College of Medical Sciences, University of Benin, (Nigeria)
- **Professor Eric KI Omogbai**, Pharmacology and Toxicology, University of Benin, Nigeria
- **Professor Dr. Udo Kragl**, Institute of Organic Chemistry, University of Rostock, (Germany)

- **Professor Cyril O. Usifoh**, Faculty of Pharmacy, University of Benin, Nigeria.
- **Professor Azuka C Opara**, Clinical Pharmacy & Pharmacy Practice, University of Benin, (Nigeria).
- **Professor Ikhide G. Imumorin**, Biological Sciences, Georgia Institute of Technology, Atlanta (USA)
- **Professor Mark T. Hamann**, Medical University College, South Carolina (USA)
- **Professor Barbara Nebe**, University of Rostock, (Germany)
- **Professor Anthony I. Okoh**, University of Fort Hare, Alice (South Africa)
- **Professor Dr. M. Iqbal Choudhary**, HEJ, University of Karachi, (Pakistan)
- **Professor Omoanghe S. Isikhuemhen**, North Carolina A&T State University, (USA)
- **Professor Ezekiel Green**, University of Johannesburg, (South Africa)
- **Professor John Igoli**, Strathclyde Institute of Pharmacy and Biomedical Sciences, UK
- **Professor Peter Akah**, Pharmacology & Toxicology, University of Nigeria, Nigeria
- **Professor H.A.B. Coker**, Faculty of Pharmacy, University of Lagos (Nigeria)
- **Dr Kingsly Agho**, School of Science and Health, Western Sydney University (Australia)
- **E. Igbinosa**, Microbiology, University of Benin, Nigeria
- **Pius Fasinu**, School of Pharmacy, Campbell University (USA)
- **Professor Larry A Walker**, National Center for Natural Products Research, Mississippi, USA
- **Dr. Alireza Heidari**, Faculty of Chemistry, California South University (CSU), Irvine, California, USA
- **Professor Iyere O Onoagbe**, Faculty of Life Sciences, University of Benin, (Nigeria)
- **Professor Broderick Eribo**, Department of Biology, Howard University, Washington DC, (USA)
- **Professor Simon Gibbons**, School of Pharmacy, University College London, UK
- **Professor Masashi Mizuno**, Laboratory of Food & Nutritional Chemistry, Kobe University, Japan
- **Professor FBC Okoye**, Professor of Medicinal and Pharmaceutical Chemistry, Nnamdi Azikiwe University, Awka





(<https://tjnpr.org/index.php/home/about/submissions>)



More Graphical Abstracts (<https://tjnpr.dynetng.org/index.php/home/graphicalabstracts>)

Indexing & Abstracting



(<https://www.scopus.com/sourceid/21100933230>)



Ads by Google

Stop seeing this ad

Why this ad? ⓘ

Tropical Journal of Natural Product Research

COUNTRY

Nigeria



Universities and research institutions in Nigeria

SUBJECT AREA AND CATEGORY

Biochemistry, Genetics and Molecular Biology
 Biochemistry
 Molecular Medicine

Chemistry
 Analytical Chemistry

Medicine
 Complementary and Alternative Medicine

Pharmacology, Toxicology and Pharmaceutics
 Drug Discovery
 Pharmaceutical Science
 Pharmacology

PUBLISHER

Faculty of Pharmacy, University of Benin

H-INDEX

5

PUBLICATION TYPE

Journals

ISSN

26160684, 26160692

COVERAGE

2017-2021

INFORMATION

[Homepage](#)

[How to publish in this journal](#)

[Contact](#)

SCOPE


The Tropical Journal of Natural Product Research is open access, peer-reviewed international journal aimed at making important contributions in the field of Natural Product Research, Pharmaceutical and Natural Sciences. The journal covers all aspects of Pharmaceutical research, chemistry and biochemistry of naturally occurring compounds, ethnomedicine, ethnopharmacology, pharmacognosy, biomedical research, biotechnology and related disciplines. The journal welcome submissions from a broad spectrum of scientific endeavour involving biological evaluation of natural substances of plant, microbial and animal origin against different disease targets, processes or therapeutic strategies that can lead to or assist in the prevention and management of chronic and infectious diseases, clinical therapeutics, Isolation and characterization of metabolites, structure elucidation, synthesis and experimental biosynthesis of natural Product as well as developments of methods in these areas. Research papers in the fields of chemistry-biology boundary, e.g. fermentation chemistry, plant tissue culture investigations etc. are also welcomed. Although the journal focuses mainly on original research articles, timely, concise and focused reviews on recent progress in active areas of natural Product is also encouraged.







Join the conversation about this journal

Quartiles

FIND SIMILAR JOURNALS ?

<p>1 Pharmacognosy Journal</p> <p>IND</p> <p>76% similarity</p>	<p>2 Journal of Traditional and Complementary Medicine</p> <p>NLD</p> <p>76% similarity</p>	<p>3 Pharmacologyonline</p> <p>ITA</p> <p>75% similarity</p>	<p>4 Pharmacognosy</p> <p>IND</p> <p>73% similarity</p>
---	---	--	---





Tropical Journal of Natural Product Research

Q4 Analytical Chemistry

best quartile

SJR 2021
0.13

powered by scimagojr.com


← Show this widget in your own website

Just copy the code below and paste within your html code:

```
<a href="https://www.scimaç
```

SCImago Graphica

Explore, visually communicate and make sense of data with our **new data visualization tool.**



Home (<https://www.tjnpr.org/index.php/home/index>)

/ Archives (<https://www.tjnpr.org/index.php/home/issue/archive>)

/ Vol. 6 No. 11 (2022): Tropical Journal of Natural Product Research (TJNPR)

Published: 2022-12-02

Articles



(<https://www.tjnpr.org/index.php/home/article/view/1205class=>)


A Review of the Ethnobotanical, Phytochemical and Medicinal Values of *Phragmanthera* Hemi-Parasitized in Southwestern Nigeria
<http://www.doi.org/10.26538/tjnpr/v6i11.1>

(<https://www.tjnpr.org/index.php/home/article/view/1205class=>)

Maria M. Adeyemi, Olutayo S. Shokunbi, Oduwole O. Oluwalana
1743-1745

PDF (<https://www.tjnpr.org/index.php/home/article/view/1205class=>)

DOI (<https://www.tjnpr.org/index.php/home/article/view/1205class=>)



(<https://www.tjnpr.org/index.php/home/article/view/1206class=>)

BRAF Inhibitors in Carcinogenesis and Chemoprevention: Implications: A Review

<http://www.doi.org/10.26538/tjnpr/v6i11.2>

(<https://www.tjnpr.org/index.php/home/article/view/1206class=>)

Benedict C. Okoro, Titilope M. Dokunmu, Eziuno O. N. Iheagwam, Daniel I. Osaigbovo, Ibukunoluwa O. Oluwalana, J. Iweala
1746-1754

PDF (<https://www.tjnpr.org/index.php/home/article/view/1206class=>)

DOI (<https://www.tjnpr.org/index.php/home/article/view/1206class=>)



(<https://www.tjnpr.org/index.php/home/article/view/1208class=>)

Crossopteryx febrifuga (Afzel. ex G. Ethnobotany, Phytochemistry and P African Tree for Malaria and Beyond
<http://www.doi.org/10.26538/tjnpr/v6i11.3>

(<https://www.tjnpr.org/index.php/hon>

Nneka N. Ibekwe
1755-1764

PDF (<https://www.tjnpr.org/index.php/home/>

DOI (<https://www.tjnpr.org/index.php/home/æ>



(<https://www.tjnpr.org/index.php/home/article/view/1209class=>)


Bioactivity Endophytic Fungi Isolate Stalk of Syzygium jambos L. Alston
<http://www.doi.org/10.26538/tjnpr/v6i11.4>

(<https://www.tjnpr.org/index.php/hon>

Kurratul Aini, Elfita Elfita, Hary Widjajanti, Arur
1765-1772

PDF (<https://www.tjnpr.org/index.php/home/>

DOI (<https://www.tjnpr.org/index.php/home/æ>



(<https://www.tjnpr.org/index.php/home/article/view/1210class=>)

Detection of Herpes Simplex Virus 1 Affected by Conjunctivitis


<http://www.doi.org/10.26538/tjnpr/v6i11.5>

(<https://www.tjnpr.org/index.php/hon>

Ruqayh R. Mutter, Layla F. Ali
1773-1777

PDF (<https://www.tjnpr.org/index.php/home/>

DOI (<https://www.tjnpr.org/index.php/home/æ>



(<https://www.tjnpr.org/index.php/home/article/view/1211class=>)

Mapping of Global 100 Top-cited Art oleifera Lam. from Documents Inde Sciences Database

<http://www.doi.org/10.26538/tjnpr/v6i11.6>

(<https://www.tjnpr.org/index.php/hon>

Abdelaziz A. I Arbab, Taha H. Musa, Hassan F Elhady A. M. Omer, Olayinka K. Oderinde, Yin
1778-1789



(<https://www.tjnpr.org/index.php/home/article/view/1212class=>)

PDF (<https://www.tjnpr.org/index.php/home/>)

DOI (<https://www.tjnpr.org/index.php/home/>)

Photoabsorbent, Antioxidative, and Properties of Three Traditional Cosr
<http://www.doi.org/10.26538/tjnpr/v6i11.7>

(<https://www.tjnpr.org/index.php/hon>)

Soukaina Laaraj, Ibtissam Ouahidi, Lotfi Aarak
1790-1793

PDF (<https://www.tjnpr.org/index.php/home/>)

DOI (<https://www.tjnpr.org/index.php/home/>)



(<https://www.tjnpr.org/index.php/home/article/view/1221class=>)

Alterations in Biochemical Parameters Enzymes in Male Mice as Biomarker Pollution with Cadmium

<http://www.doi.org/10.26538/tjnpr/v6i11.8>

(<https://www.tjnpr.org/index.php/hon>)

Ahmed S. Aldhamin, Nuhad M. Hamid
1794-1797

PDF (<https://www.tjnpr.org/index.php/home/>)

DOI (<https://www.tjnpr.org/index.php/home/>)



(<https://www.tjnpr.org/index.php/home/article/view/1226class=>)

Antioxidant Activity and Inhibition of Reductase Enzyme by Bay Leaf (Sy Wight) Extract as a Treatment for H
<http://www.doi.org/10.26538/tjnpr/v6i11.9>

(<https://www.tjnpr.org/index.php/hon>)

Nanang Yunarto, Eva D. Helentina, Indah Suli Reswandaru, Sukmayati Alegantina, Ani Isnav
1798-1801

PDF (<https://www.tjnpr.org/index.php/home/>)

DOI (<https://www.tjnpr.org/index.php/home/>)



(<https://www.tjnpr.org/index.php/home/article/view/1225class=>)

Potential Role of Betel Leaf (*Piper b*
Extract as Antibacterial *Escherichia*
Inhibition of β -Ketoacyl-[Acyl Carrier
<http://www.doi.org/10.26538/tjnpr/v6i11.10>

(<https://www.tjnpr.org/index.php/hon>

Ayu T. Agustin, Eko Julianto, Julianus Julianus
1802-1808

PDF (<https://www.tjnpr.org/index.php/home/>

DOI (<https://www.tjnpr.org/index.php/home/>



(<https://www.tjnpr.org/index.php/home/article/view/1232class=>)

Effect of *Cnidoscolus aconitifolius* E
Injury Induced with Carbon Tetrachl
<http://www.doi.org/10.26538/tjnpr/v6i11.11>

(<https://www.tjnpr.org/index.php/hon>

Zamharira Muslim, Jon Farizal, Rahmad Abdil
1809-1813

PDF (<https://www.tjnpr.org/index.php/home/>

DOI (<https://www.tjnpr.org/index.php/home/>



(<https://www.tjnpr.org/index.php/home/article/view/1224class=>)

Ethnobotanical Study and Biodivers
Plants Used in the Province of Taza
Morocco

<http://www.doi.org/10.26538/tjnpr/v6i11.12>

(<https://www.tjnpr.org/index.php/hon>

Rajae El Brahimi, Azeddin El Barnossi, Mohar
1814-1831

PDF (<https://www.tjnpr.org/index.php/home/>

DOI (<https://www.tjnpr.org/index.php/home/>



(<https://www.tjnpr.org/index.php/home/article/view/1231class=>)

Antiplatelet Activity of *Piper cubeba*
Extract


<http://www.doi.org/10.26538/tjnpr/v6i11.13>

(<https://www.tjnpr.org/index.php/hon>

Kyky Herlyanti, Retno Murwanti, Wahyono, N
1832-1835

PDF (<https://www.tjnpr.org/index.php/home/>

DOI (<https://www.tjnpr.org/index.php/home/>



(<https://www.tjnpr.org/index.php/home/article/view/1234class=>)

Synthesis, Characterization and Bio

Some New Dinitrone Compounds

<http://www.doi.org/10.26538/tjnpr/v6i11.14>

(<https://www.tjnpr.org/index.php/home>)

Batool S. Haddad

1836-1839

PDF (<https://www.tjnpr.org/index.php/home/>)

DOI (<https://www.tjnpr.org/index.php/home/>)



(<https://www.tjnpr.org/index.php/home/article/view/1227class=>)

Assessment of Acute Toxicity, LD50

Histopathological Evaluation of Nige

Moringa oleifera Seed Extracts in W

<http://www.doi.org/10.26538/tjnpr/v6i11.15>

(<https://www.tjnpr.org/index.php/home>)

Marie U.E. Dibua, Daniel E. Garba, Esther C.

Emencheta, Emmanuel S. Okeke

1840-1843

PDF (<https://www.tjnpr.org/index.php/home/>)

DOI (<https://www.tjnpr.org/index.php/home/>)



(<https://www.tjnpr.org/index.php/home/article/view/1222class=>)

Comparative Antioxidant Activities o

Detarium microcarpum Guill & Perr

<http://www.doi.org/10.26538/tjnpr/v6i11.16>

(<https://www.tjnpr.org/index.php/home>)

Jane David, Francis M. Agwom, Stephen O. C

Taiwo E. Alemika

1844-1847

PDF (<https://www.tjnpr.org/index.php/home/>)

DOI (<https://www.tjnpr.org/index.php/home/>)



(<https://www.tjnpr.org/index.php/home/article/view/1233class=>)

Botanical and Bio-active Markers of

Honey

<http://www.doi.org/10.26538/tjnpr/v6i11.17>

(<https://www.tjnpr.org/index.php/home>)

Bayo O. Adeoye, Abolape A. Iyanda, Michael O

Adeoye, Ayodeji M. Oyerinde, Goodness O. C

1848-1853



(<https://www.tjnpr.org/index.php/home/article/view/1230class=>)

PDF (<https://www.tjnpr.org/index.php/home/>)

DOI (<https://www.tjnpr.org/index.php/home/>)

CTXM and OXA Genes in Foodborne Cooked Street Foods Sold in the South of Nigeria

<http://www.doi.org/10.26538/tjnpr/v6i11.18>

(<https://www.tjnpr.org/index.php/home>)

Olufeyikemi A. Adeleke, Abimbola A. Owoseni
1854-1857

PDF (<https://www.tjnpr.org/index.php/home/>)

DOI (<https://www.tjnpr.org/index.php/home/>)



(<https://www.tjnpr.org/index.php/home/article/view/1239class=>)

In vitro Antioxidant and Antidiabetic Extracts of the Stem Bark of Cylicoc (Harms) Mimosaceae

<http://www.doi.org/10.26538/tjnpr/v6i11.19>

(<https://www.tjnpr.org/index.php/home>)

Vincent O. Imieje, Chika F. Onochie, Bridget N
1858-1863

PDF (<https://www.tjnpr.org/index.php/home/>)

DOI (<https://www.tjnpr.org/index.php/home/>)



(<https://www.tjnpr.org/index.php/home/article/view/1240class=>)

Comparative Study of the Influence and Hydroxypropyl Methylcellulose Profile of Ibuprofen from Subcutaneous Formulations


<http://www.doi.org/10.26538/tjnpr/v6i11.20>

(<https://www.tjnpr.org/index.php/home>)

Collins O. Airemwen, Jude E. Isesele, Johnbu
Michael U. Uhumwangho
1864-1869

PDF (<https://www.tjnpr.org/index.php/home/>)

DOI (<https://www.tjnpr.org/index.php/home/>)



(<https://www.tjnpr.org/index.php/home/article/view/1241>class=)

In silico Investigation on Isatin (1H-i Derivatives as Potential AntiTumor I Alpha

<http://www.doi.org/10.26538/tjnpr/v6i11.21>

(<https://www.tjnpr.org/index.php/home/article/view/1241>class=)

Abel K. Oyebamiji, Emmanuel T. Akintayo, Ba Odelade, Babatunde O. Adetuyi, Amin Hafiz, C 1870-1875

PDF (<https://www.tjnpr.org/index.php/home/article/view/1241>class=)

DOI (<https://www.doi.org/10.26538/tjnpr/v6i11.21>)



(<https://www.tjnpr.org/index.php/home/article/view/1242>class=)

Rice and Non-Rice Staple Consumption among Urban and Rural Communities in Nigeria


<http://www.doi.org/10.26538/tjnpr/v6i11.22>

(<https://www.tjnpr.org/index.php/home/article/view/1242>class=)

Raphael A. Omolehin, Segun B. Fakayode, Sa Olajumoke O. Alabi 1876-1881

PDF (<https://www.tjnpr.org/index.php/home/article/view/1242>class=)

DOI (<https://www.doi.org/10.26538/tjnpr/v6i11.22>)



(<https://www.tjnpr.org/index.php/home/article/view/1243>class=)

Antioxidant, Anti-hyperlipidemic and Potentials of Diet Supplemented with Roasted Seeds on Isoproterenol-Induced Oxidative Stress in Wistar rats

<http://www.doi.org/10.26538/tjnpr/v6i11.23>

(<https://www.tjnpr.org/index.php/home/article/view/1243>class=)

Adetoun E. Morakinyo, Femi E. Omoniyi, Sam Agbabiaka, Paula A. Odole, Abel K. Oyebamiji O. Bello-Ogunesan, Emmanuel T. Akintayo, C Sunday A. Akintelu, Temitope A. Oyedepo, Jor 1882-1887

PDF (<https://www.tjnpr.org/index.php/home/article/view/1243>class=)

DOI (<https://www.doi.org/10.26538/tjnpr/v6i11.23>)



(<https://www.tjnpr.org/index.php/home/article/view/1244class=>)

Evaluation of Gallus gallus domesticus Glass as an Alternative to Dental Plaster

<http://www.doi.org/10.26538/tjnpr/v6i11.24>

(<https://www.tjnpr.org/index.php/home>)

Joshua C. Onyia, Eze E. Ajaegbu, Benneth M. Ikuesan, Abdulrasheed M. Bello, Ijeoma O. Okeadaobi J. Dieke, Adaora L. Onuora, Flora N. E. Juliana O. Ndubuisi, Ethel E. Adimora, Juliet C. Iloabuchi, Nnyeneime U. Bassey, Florence O. 1888-1892

PDF (<https://www.tjnpr.org/index.php/home/>)

DOI (<https://www.tjnpr.org/index.php/home/>)



(<https://www.tjnpr.org/index.php/home/article/view/1245class=>)

Aphrodisiac Properties and Reproductive Concentrations of Ethanol Leaf Extract of *Conoclinium chinensis* in Male Wistar Rats

<http://www.doi.org/10.26538/tjnpr/v6i11.25>

(<https://www.tjnpr.org/index.php/home>)

Grace S. Effiong, Othuke B. Odeghe, Nse U. I. Chinonso C. Obu, Emmanuel O. Nwuzor, Ifeanyi O. Oviru 1893-1900

PDF (<https://www.tjnpr.org/index.php/home/>)

DOI (<https://www.tjnpr.org/index.php/home/>)



(<https://www.tjnpr.org/index.php/home/article/view/1246class=>)

Evaluation of Ultrasonically Assisted Thai Polyherbal Formulation, Mahajam, and Anti-Inflammation Activities

<http://www.doi.org/10.26538/tjnpr/v6i11.26>

(<https://www.tjnpr.org/index.php/home>)

Adisak Thomudtha, Pornpun Laovachirasuwan 1900-1905

PDF (<https://www.tjnpr.org/index.php/home/>)

DOI (<https://www.tjnpr.org/index.php/home/>)

Erratum: In vitro Antioxidant Activities of Some Re(I) Metal Carbonyls Synthesized from Isatin Derivatives

<http://www.doi.org/10.26538/tjnpr/v6i10.28>

(<https://www.tjnpr.org/index.php/home/article/view/1278>)

Adebomi A. Ikotun, Esther E. Babajide, Tolulope O. Omolekan, Chijioke J. Ajaelu
1723-1726

PDF (<https://www.tjnpr.org/index.php/home/article/view/1278/2078>)

DOI (<https://www.tjnpr.org/index.php/home/article/view/1278/2079>)

Erratum: Classification of Dysmenorrhea among Students at Ubon Ratchathani Rajabhat University, Thailand According to the Māhaachortàrat Scripture
<http://www.doi.org/10.26538/tjnpr/v6i6.12>

(<https://www.tjnpr.org/index.php/home/article/view/1279>)

Phanida Kamuttachat, Pitchanan Thiantongin
900-905

PDF (<https://www.tjnpr.org/index.php/home/article/view/1279/2080>)

DOI (<https://www.tjnpr.org/index.php/home/article/view/1279/2081>)

Erratum: Artemether-Lumefantrine Combination for Non-Comorbid Falciparum Malaria: A Clinico-Parasitological Efficacy Study
<http://www.doi.org/10.26538/tjnpr/v6i10.26>

(<https://www.tjnpr.org/index.php/home/article/view/1280>)

Chinelo K. Ezejiegu, Angus N. Oli, Chinwendu N. Ezomike, Christian C. Uba, Uchenna C. Ogwaluonye,
Charles O. Esimone
1715-1718

PDF (<https://www.tjnpr.org/index.php/home/article/view/1280/2082>)

DOI (<https://www.tjnpr.org/index.php/home/article/view/1280/2083>)

Erratum: Economic Activities of Rice Production Practice and the Outcome in Southern Guinea Savanna of Nigeria
[. http://www.doi.org/10.26538/tjnpr/v6i9.31](http://www.doi.org/10.26538/tjnpr/v6i9.31)

(<https://www.tjnpr.org/index.php/home/article/view/1281>)


Tanko P. Sani, Raphael A. Omolehin, Shaffi Abullahi, Usman K. Abubakar
1523-1533

PDF (<https://www.tjnpr.org/index.php/home/article/view/1281/2084>)

DOI (<https://www.tjnpr.org/index.php/home/article/view/1281/2085>)



(<https://tjnpr.org/index.php/home/about/submissions>)



More Graphical Abstracts (<https://tjnpr.dynetng.org/index.php/home/graphicalabstracts>)

Indexing & Abstracting





(<https://www.scopus.com/sourceid/21100933230>)



(<https://www.scopus.com/sourceid/21100933230>)

Keywords

