

# Is TRMM product good proxy for gauge precipitation over peat land area of the South Sumatera?

*by* Muhammad Irfan

---

**Submission date:** 19-Jul-2020 04:49PM (UTC+0700)

**Submission ID:** 1359288824

**File name:** SICBAS\_2019.pdf (1.26M)

**Word count:** 2549

**Character count:** 12753

PAPER

## Is TRMM product good proxy for gauge precipitation over peat land area of the South Sumatera?

To cite this article: Muhammad Irfan *et al* 2019 *J. Phys.: Conf. Ser.* **1282** 012021



View the [article online](#) for updates and enhancements.



**IOP | ebooks™**

Bringing you innovative digital publishing with leading voices to create your essential collection of books in STEM research.

Start exploring the collection - download the first chapter of every title for free.

## Is TRMM product good proxy for gauge precipitation over peat land area of the South Sumatera?

Muhammad Irfan<sup>1,2a</sup>, Wijaya Mardiansyah<sup>1</sup>, Menik Ariani<sup>1</sup>, Albert Sulaiman<sup>3</sup>  
and Iskhaq Iskandar<sup>1,2b</sup>

<sup>1</sup>Department of Physics, Faculty of Mathematics and Natural Sciences, Sriwijaya University, Indonesia;

<sup>2</sup>Graduate School of Sciences, Faculty of Mathematics and Natural Sciences, Sriwijaya University, Indonesia;

<sup>3</sup>Agency for Assessment and Application Technology, Jakarta, Indonesia

Email: <sup>a</sup> irfplg@yahoo.com, <sup>b</sup> iskhaq.iskandar@gmail.com,

**Abstract.** The Tropical Precipitation Measuring Mission (TRMM) provides daily precipitation data covering the tropical region. In this study, data from the TRMM were compared with those obtained from direct measurement using an integrated observation system namely *SEnsory data transmission Service Assisted by Midori Engineering* laboratory (SESAME). The SESAME directly measures and records the ground water level, soil moisture, skin temperature and precipitation in the peatland area. There are four SESAME stations used in this study, two stations located at the Peatland Hydrological Unit (PHU) of the Saleh River and the other two are located at the PHU of the Lumpur River for validating the TRMM precipitation data. The analysis indicates that the precipitation data obtained from the TRMM have significant correlation with those recorded by the SESAME. The coefficient correlations at those PHU Saleh River I, PHU Saleh River II, PHU Lumpur River I and PHU Lumpur River II are 0.52, 0.48, 0.53 and 0.52, respectively. Meanwhile, the root mean square errors (RMSEs) of the TRMM relative to SESAME data at those four stations are 1.88 mm/d, 2.62 mm/d, 1.53 mm/d and 2.31 mm/d, respectively. Therefore, the TRMM product is a good proxy for gauge precipitation over the peatland area of the South Sumatera.

### 1. Introduction

Precipitation is defined as products of condensation of atmospheric water vapor that falls from the earth, for example rain, snow and hail. It is one of the most important hydrology parameters and has a large influence in the supply of fresh water. In addition, precipitation also has a significant impact on the economic and environmental conditions of an area [1-4].

Rain gauge is one of a common tool for measuring precipitation. However, its spatial distribution is sparse and it cannot cover a wide area. From the last few decades, satellite remote sensing has been used for the measurement of precipitation, including the TRMM satellites. Although satellite measurement covers a wide area, its spatial and temporal accuracy might not be significant. Therefore, evaluation of the satellite remote sensing data using ground station is required [5-7].

In order to obtain ground-based measurement data, an in-situ measurement system for hydrological and climatological parameters has been developed. This system is called *SEnsory data transmission Service Assisted by Midori Engineering* laboratory (SESAME) system. In the South Sumatra, this system was installed at 8 locations in the Peatland Hydrological Unit (PHU) [8].



<sup>1</sup> Content from this work may be used under the terms of the [Creative Commons Attribution 3.0 licence](https://creativecommons.org/licenses/by/3.0/). Any further distribution of this work must maintain attribution to the author(s) and the title of the work, journal citation and DOI.

Published under licence by IOP Publishing Ltd

This study is designed to evaluate the TRMM precipitation data by comparing with the SESAME data obtained in the peatland area of South Sumatera. In particular, this study is intended to address the following question:

- What is the correlation between TRMM precipitation data and the SESAME precipitation data?
- Is the TRMM results good for precipitation measuring over peat land area of the South Sumatera?

## 2. Data

This study used the TRMM 3 hours precipitation data for a period of 1 July 2017 to 31 March 2018. In addition, the SESAME data observed from 4 locations: PHU Sungai Lumpur 1, PHU Sungai Lumpur 2, PHU Sungai Saleh 1, and PHU Sungai Saleh 2 for a period of 1 July 2017 to 31 March 2018. The locations of this study shown in Figure 1.

PHU Lumpur River 1	PHU Lumpur River 2	PHU Saleh River 1	PHU Saleh River 2
			
-3.1436443, 105.1843584	-2.9106579, 105.0824855	-3.4583709, 104.9209707	-2.6768524, 105.1434387

Figure 1. Locations of the SESAME stations used in this study.

### 2.1. SESAME (Sensory data transmission Service Assisted by Midori Engineering laboratory)

The SESAME is a system for measuring hydrological and climatological parameters and it was designed for the peatland area in Indonesia. The parameters measured are surface temperature, ground water level (GWL), soil moisture, and precipitation. The system measures data using sensors, records and disseminates them through mobile communication networks, then processes and transmits data, and finally provides output that can be analyzed and transmitted to the users [8]. In Indonesia, this system has been installed in 17 locations of peatland area, 8 of which are in the South Sumatera.

### 2.2. TRMM (Tropical Precipitation Measuring Mission)

The TRMM satellite was developed by the National Aeronautics and Space Administration (NASA) and the Japan Aerospace Exploration Agency (JAXA). The TRMM satellite has started the measurements in 1997 [9]. The satellite brings 5 channels of PR (Precipitation Radar), TMI (TRMM Microwave Channel), VIRS (Visible and Infrared Channel), CERES (Clouds and the Earth's Radiant Energy System), and LIS (Lightning Imaging Sensor). In particular the PR and TMI channels have missions in measuring the precipitation. Both channels are able to observe rain structures and play an important role in knowing the mechanisms of global climate change and monitoring of environmental variations.

The TRMM precipitation data have been available since 1998 [9,10]. The data have 3 levels [10]. Level 1 provides raw data but it has been calibrated and corrected geometrically. Level 2 contains images of original precipitation at the same spatial resolution when the satellite passes through the recorded area. Level 3 provides processed data which are ready to use. In this study, we used the TRMM data level 3, with a spatial resolution of  $0.25^\circ \times 0.25^\circ$  and temporal resolution of 3 hours.

## 3. Methodology

This research was conducted at 4 (four) stations, namely the PHU Sungai Lumpur 1, the PHU Sungai Lumpur 2, the PHU Sungai Saleh 1 and the PHU Sungai Saleh 2. In order to evaluate the TRMM

precipitation data, a statistical analysis has been conducted, namely the linear regression, linear correlation and standard errors analyses [11-13].

### 3.1. Linier regression

In this study, a simple linier regression analysis was used to find are lationship between two continuous variables, namely the precipitation data from the SESAME and those from the TRMM satellite. The SESAME data are as the dependent variables, while the independent variable is the TRMM data. The general form of linier regressionis:

$$y = a + bx, \quad (1)$$

$$a = \frac{(\sum y \sum x^2) - (\sum x \sum xy)}{N(\sum x^2)(\sum x)^2}, \quad (2)$$

$$b = \frac{N(\sum xy) - (\sum x \sum y)}{N(\sum x^2)(\sum x)^2}, \quad (3)$$

where y is a dependent variable, x is a independent variable, a is intercept and b is slope.

### 3.2. Linier correlation

The linear correlation is used to determine how closely the relationship between two or more variables. The calculation of correlation coefficient for two variablesis determined by:

$$r_{xy} = \frac{1}{N-1} \sum_{i=1}^N \frac{(x_i - \bar{x})(y_i - \bar{y})}{s_x s_y}, \quad (4)$$

where  $s_x$  and  $s_y$  are standard deviation for each variables ( $x, y$ ). If the correlation coefficient  $r = \pm 1$ , then the two variables ( $x, y$ ) clusters along a straight line and they have a perfect correlation. Note that the positive (+) value means that the two variables are “in-phase”. Meanwhile, the negative (-) value means that the two variables are “out-of-phase”. For  $r = 0$ , the two variables are scattered randomly on the graph and there is little or no relationship between the variables.

### 3.3. Standard errors

3.3.1. *Mean Bias Error (MBE)*. The MBE value indicates a systematic error estimation. A positive (negative) value means that the estimation is greater (smaller) than the real measurement. The MBE is calculated by:

$$MBE = \frac{1}{N} \sum_{i=1}^N (x_i - y_i) \quad (5)$$

3.3.2. *Root mean square error (RMSE)*. RMSE values indicate random errors from satellite data.

$$RMSE = \sqrt{\left(\frac{1}{N} \sum_{i=1}^N \{x_i - (MBE) - y_i\}^2\right)} \quad (6)$$


3.3.3. *Mean absolute error (MAE)*. MAE value indicate the accuracy of estimate with the average error in unit of the same size as the original.

$$MAE = \frac{1}{N} \sum_{i=1}^N |x_i - y_i| \quad (7)$$


Where  $x_i$  is the estimated value (TRMM data),  $y_i$  is the value of in-situ measurement (SESAME data), and N is the number of data used.

## 4. Result and discussion

A graph of precipitation data recorded by SESAME and TRMM over a period of 1 July 2017 to 31 March 2018 is shown in Figure 2 below:



**Figure 2.** The graph comparison of precipitation SESAME data and TRMM satellite data on four location of research.



**Figure 3.** The graph liner regression of precipitation SESAME and TRMM satellite onfour location of research.



It was calculated the precipitation correlation and errors between the SESAME data and the TRMM data using equation 1 until 7. Figure 3 shows the scatter plot of daily precipitation correlation between the SESAME data and TRMM data at the PHU Lumpur River 1, PHU Lumpur River 2, PHU Saleh River 1, and PHU Lumpur Saleh 2 for a period 1 July 2017 to 31 March 2018.

Table 1 shows the correlation coefficient and error of precipitation SESAME and TRMM satellite on location of research: PHU Lumpur River 1, PHU Lumpur River 2, PHU Saleh River 1, and PHU Saleh River 2. The results shows that the coefficient correlations of TRMM precipitation and SESAME precipitation for 4 research locations were 0.52, 0.48, 0.53 and 0.52, respectively. The root mean square errors (RMSEs) of the TRMM relative to SESAME data were 1.88 mm/d, 2.62 mm/d, 1.53 mm/d and 2.31 mm/d, respectively. So that, the TRMM precipitation data and SESAME precipitation data has a strong correlation.

**Table 1.** The correlation coefficient and error of precipitation SESAME and TRMM satellite on four location of research.

Parameter	PHU Lumpur River 1	PHU Lumpur River 2	PHU Saleh River 1	PHU Saleh River 2
r	0.52	0.48	0.53	0.52
MBE	-0.027	-0.95037	-0.04744	-0.00212
RSME	1.883058	2.617391	1.534115	2.305494
MAE	3.846557	3.7637	3.592344	3.686813

## 5. Conclusion

The analysis shows that the precipitation data from TRMM weather satellite is significantly correlated with the observed precipitation data from the SESAME. Therefore, it could be concluded that TRMM product is good proxy for gauge precipitation over peat land area of the South Sumatera.

## Acknowledgements

We thank the Peatland Restoration Agency for providing us the SESAME data. This study is supported by the Universitas Sriwijaya through the *Hibah Unggulan Kompetitif* 2018 for the first author and the *Hibah Unggulan Profesi* for the last author.

## References

- [1] Fanin T and van der Werf G R 2017 Precipitation – fire linkages in Indonesia ( 1997 – 2015 ) *Biogeosciences* **14** 3995–4008.
- [2] Liu H -W, Yu J -Y, and Chen C-A 2018 Changes of tropical precipitation and convective structure under global warming projected by CMIP5 model simulations *Terr. Atmos. Ocean. Sci.*, **29** (4) 429–440.
- [3] Jiang S 2014 *et al* Improvement of Multi-Satellite Real-Time Precipitation Products for Ensemble Streamflow Simulation in a Middle Latitude Basin in South China,” *Water Resour. Manag.*, **28** (8) 2259–2278.
- [4] Wang Y, Yang J, Chen Y, De Maeyer P, Li Z and Duan W 2018 Detecting the Causal Effect of Soil Moisture on Precipitation Using Convergent Cross Mapping, *Sci. Rep.*, **8**(1) 12171.
- [5] Cao Y, Zhang W and W. Wang 2018 Evaluation of TRMM 3B43 data over the Yangtze River Delta of China, *Sci. Rep.*, **8** (1) 1–13.
- [6] Wang W *et al* 2016 Modelling hydrologic processes in the Mekong River basin using a distributed model driven by satellite precipitation and rain gauge observations, *PLoS One*, **11**, (3) 1–20
- [7] Cai Y *et al* 2015 Spatio-temporal analysis of the accuracy of tropical multisatellite precipitation analysis 3b42 precipitation data in mid-high latitudes of China, *PLoS One*, **10** (4) 1–23.
- [8] Shigenaga Y, Takahashi H, Teguh R, Kencana W, Yokoyama S, and Jaya A, Field Data Transmission System, SESAME-SATREPS, by using Cell-phones Digital telecommunications network.

- [9] Kummerow C, et al 2000 The status of the Tropical Precipitation Measuring Mission (TRMM) after two years in orbit. *J. Appl. Meteorol.*, **39** 1965–1982.
- [10] Cao Y, Zhang W, and Wang W 2018 Evaluation of TRMM 3B43 Data over The Yangtze River Delta of China, *Scientific Reports* 8, article number 5290.
- [11] Emery W J, and Thomson R E 2004 Data Analysis Method in Physical Oceanography, 2nd Edition, Elsevier B V, Amsterdam, The Netherlands. 638.
- [12] Iskandar I, Irfan M, Syamsuddin F, Johan A and Poerwono P 2012 Trend in Precipitation over Sumatra under the Warming Earth *Int. Jour. of Remote Sensing and Earth Sci.*, **8** 19-24.
- [13] Agustin A, Mardiansyah W, Setiabudidaya D, and Iskandar I, Wind Satand RAMA Buoy: acomparison of ocean-atmosphere data, MATEC Web of conference 101, 04005, 2017



# Is TRMM product good proxy for gauge precipitation over peat land area of the South Sumatera?

## ORIGINALITY REPORT

14%

SIMILARITY INDEX

%

INTERNET SOURCES

14%

PUBLICATIONS

%

STUDENT PAPERS

## PRIMARY SOURCES

- 1 Chandan Kumar Chakrabarty, Abd Halim Bin Shamsuddin, Ibrahim Bin Ahmad, Mohamed Nor Bin Mohamed Desa et al. "4th International Conference on Energy and Environment 2013 (ICEE 2013)", IOP Conference Series: Earth and Environmental Science, 2013

Publication

3%
- 2 Manh-Hung Le, Venkataraman Lakshmi, John Bolten, Duong Du Bui. "Adequacy of Satellite-derived Precipitation Estimate for Hydrological Modeling in Vietnam Basins", Journal of Hydrology, 2020

Publication

1%
- 3 Rifki Nurfaiz, Bintang Wibowo, Gladhi Guarddin, Petrus Mursanto. "Low power wireless network for efficient peatland monitoring system", 2017 International Workshop on Big Data and Information Security (IW BIS), 2017

Publication

1%

Meijing Zhang, Conrado de Leon, Kati

4

Migliaccio. "Evaluation and comparison of interpolated gauge rainfall data and gridded rainfall data in Florida, USA", Hydrological Sciences Journal, 2018

Publication

---

1%

5

Hsiao-Wei Liu, Jia-Yuh Yu, Chao-An Chen. "Changes of tropical precipitation and convective structure under global warming projected by CMIP5 model simulations", Terrestrial, Atmospheric and Oceanic Sciences, 2018

Publication

---

1%

6

Yan Jiang, Changming Liu, Xuyong Li. "Hydrological Impacts of Climate Change Simulated by HIMS Models in the Luanhe River Basin, North China", Water Resources Management, 2014

Publication

---

1%

7

Frédéric Satgé, Denis Ruelland, Marie-Paule Bonnet, Jorge Molina, Ramiro Pillco. "Consistency of satellite-based precipitation products in space and over time compared with gauge observations and snow- hydrological modelling in the Lake Titicaca region", Hydrology and Earth System Sciences, 2019

Publication

---

1%

8

"Tropical Peatland Ecosystems", Springer

1%

9

Claudia Stickler, Olivia David, Charlotta Chan, Juan Pablo Ardila, Tathiana Bezerra. "The Rio Branco Declaration: Assessing Progress Toward a Near-Term Voluntary Deforestation Reduction Target in Subnational Jurisdictions Across the Tropics", *Frontiers in Forests and Global Change*, 2020

Publication

1%

10

Liu, Q.. "Comparison of radiative signals between precipitating and non-precipitating clouds in frontal and typhoon domains over East Asia", *Atmospheric Research*, 201005

Publication

1%

11

Zachary Fasnacht, Alexander Vasilkov, David Haffner, Wenhan Qin, Joanna Joiner, Nickolay Krotkov, Andrew M. Sayer, Robert Spurr. "A geometry-dependent surface Lambertian-equivalent reflectivity product for UV/Vis retrievals: Part II. Evaluation over open ocean", *Copernicus GmbH*, 2019

Publication

1%

12

Wei Wang, Chen Ji, Fangye Lin, Jun Zou, S. Dorbolo. "Water drops bouncing off vertically vibrating textured surfaces", *Journal of Fluid Mechanics*, 2019

Publication

<1%

---

13

Tao Zhang. "Response of water vapor and clouds to El Niño warming in three National Center for Atmospheric Research atmospheric models", Journal of Geophysical Research, 2006

Publication

---

<1%

14

G. P. Matthews\*, V. G. Brook, G. S. Elliot, T. H. Khan-Gandapur. "Cognitive structure determination as a tool in science teaching", European Journal of Science Education, 2007

Publication

---

<1%

15

J. C. Lin. "Effects of biomass-burning-derived aerosols on precipitation and clouds in the Amazon Basin: a satellite-based empirical study", Journal of Geophysical Research, 10/12/2006

Publication

---

<1%

---

Exclude quotes      On

Exclude matches      Off

Exclude bibliography      On