

Phytochemical and Larvicidal Activity of Pineapple (*Annanas comusus* L.) Peel Extract Against *Aedes aegypti* of Dengue Vector

by Rico Januar Sitorus

Submission date: 07-Mar-2023 12:16PM (UTC+0700)

Submission ID: 2030961919

File name: jpnr-2023-01-006_28-32.pdf (364.29K)

Word count: 3383

Character count: 18268

Phytochemical and Larvicidal Activity of Pineapple (Ananas comusus L.) Peel Extract Against *Aedes aegypti* of Dengue Vector

Krisna Delita^{1,2}, Nurhayati Damiri^{1,3}, Rico Januar Sitorus³, Poedji Loekitowati Hariani⁴

¹Department of Environmental Science, Doctoral Program, Sriwijaya University, South Sumatera, Indonesia.

²Sriwigama College of Agricultural Sciences, Palembang, Indonesia

³Department of Plant pests and diseases, Agricultural Faculty, Sriwijaya University, Indralaya Ogan Ilir District, South Sumatera, Indonesia

⁴Faculty of Public Health Diseases, Sriwijaya University, South Sumatera Indonesia

Abstract

Control of the *Aedes aegypti* DHF vector with synthetic insecticides has an impact on resistance. As an alternative is the use of plant materials, one of which is pineapple skin waste which is widely available and has not been utilized. This study aims to determine the phytochemical compounds contained and biolarvicidal activity of pineapple peel against *Aedes aegypti*. Held. Laboratory experimental study with randomized block design effect of concentration on mortality of *Aedes aegypti* larvae. Based on the research results, the ethanol extract of pineapple peel contains flavonoids, tannins, saponins, alkaloids and terpenoids and these compounds have biolarvicidal activity. Ethanol extract at concentrations of 400 ppm and 500 ppm and at concentrations of 100 to 300 ppm has very low activity. Letal Konsentrasi ekstrak ethanol kulit nanas adalah Pineapple peel ethanol extract has potential in the future as an environmentally friendly biolarvicidal.

Keywords: Pineapple Pell , Biolarvasida, *Aedes aegypti*.

INTRODUCTION

Dengue hemorrhagic fever is a problem in many regions of the world, especially in tropical areas, including Indonesia. Palembang City is located in the Province of South Sumatra Indonesia, is one of the endemic cities for dengue fever in Indonesia. Dengue Hemorrhagic Fever (DHF) In 2021 the cumulative DHF in Indonesia was recorded at 71,044 cases with a cumulative number of 690 deaths [1]. Dengue fever is an arbovirus disease yang carried by the *Aedes aegypti* mosquito is the primary vector in the transmission of DHF [2], [3].

The people and government of Indonesia have tried to overcome dengue fever by controlling vectors because there is no DHF vaccine yet. Control is carried out in various ways, but the most preferred and widely practiced method is chemical control using insecticides.

The use of insecticides for a long time and with improper management can have a negative impact on the environment and dengue hemorrhagic fever vectors, which can cause resistance to *Aedes aegypti* dengue hemorrhagic vectors.. The results of molecular research on *Aedes aegypti* in Palembang found mutations in the VGSC gene at the Val1016Ile point, as proof of the resistance mechanism which is a target site for synthetic pyrethroid insecticides [4].

Address for correspondence: Nurhayati Damiri

Department of Environmental Science, Doctoral Program, Sriwijaya University,
South Sumatera, Indonesia.

Email: nurhayati@fp.unsri.ac.id

Access this article online

Quick Response Code:



Website:

www.pnrjournal.com

DOI:

10.47750/pnr.2023.14.01.006

For reprints contact: pnjournal@gmail.com

How to cite this article: Krisna D, Nurhayati D, Rico J S, Poedji L H, Phytochemical and Larvicidal Activity of Pineapple (Ananas comusus L.) Peel Extract Against *Aedes aegypti* of Dengue Vector, J PHARM NEGATIVE RESULTS 2023;14(1): 28-32.

There is a need for efforts to control the dengue vector using environmentally friendly insecticides. One of the plants that is widely available in the city of Palembang is the Queen variety pineapple plant which has been adapted for a long time in the province of South Sumatra, known as the Palembang pineapple. Pineapple produces solid waste in the form of peel and weevil. Pineapple waste can reach 48.6% of the total weight of the fruit, consisting of pineapple peel and hump [5]. Pineapple peel is an untapped waste, even though pineapple peel contains vitamin C, carotenoids and flavonoids [17].

Pineapple skin is thought to contain phytochemicals that act as insecticides, anti-bacterial and anti-fungal. Pineapple peel from Rayalseema region of Andhra Pradesh, India contains phytochemical compounds, namely flavonoids, tannins, saponins, sterols, terpenoids and alkaloids, these compounds can be used as raw materials for insecticides. According to [7, 8], Ethanol extract of pineapple fruit contains saponins, tannins, flavonoids and terpenoids that act to inhibit the growth of *Escherichia coli*, *Pseudomonas aeruginosa*, *Staphylococcus aureus* and *Bacillus amyloliquefaciens* bacteria with an inhibition zone of 15-17 mm. Pineapple peel waste originating from fruit traders contains flavonoids, alkaloids, anthocyanins and tannins and pineapple peel has a high inhibitory power against the activity of *E. coli* and *S. Aureus* bacteria. Pineapple peel extract at a concentration of 4% was able to kill *Culex* mosquitoes up to 97.5% [9- 10- 12].

This study aims to determine the phytochemical content of the ethanol extract of pineapple peel which acts as a biolarvicidal against *Aedes aegypti* as a vector of dengue hemorrhagic fever.

MATERIALS AND METHODS

Study Area

This study is a laboratory experimental study with a randomized block design. To test the concentration of pineapple peel ethanol extract on larval mortality and determine the phytochemical content. The research was carried out in the Special Integrated Laboratory of the Sriwijaya University Palembang Postgraduate Laboratory and Baturaja Health Research and Development Center Laboratory. It will be held from March to June 2022.

The materials used in this study are pineapple peel used in this study was collected from pineapple planting centers, East Prabumulih District, Prabumulih City, South Sumatra Province Indonesia Latitude south side 3°-4°, Longitude Eastern 104°-105°, Altitude 48 m, Annual Raining Average 2.723 mm/year, Daily Temperature average 20.13 °C, Comperative humidity average 79,03% and type of soil alluvial, clay and sandy [13]. Egg *Aedes aegypti* Strain Liver Pool from Health Research and Development entomology laboratory. While the chemicals used in this study were aluminum foil, filter paper, parchment paper, ethanol p.a (Merck™), distilled water, AlCl₃ (Merck™),

and potassium acetate (Merck™), HCl 2 N, Mg, FeCl₂, reaction Meyer, Wagner, Dragendorf. Chloroform, H₂SO₄, Lieberman Burchard reagent. The tools used are, blender (Philips™), ayakan no.60 dan 40, rotary evaporator (IKA™ HB 10), vacuum pump (B-One™), waterbath (MEMMERT™), perangkat alat gelas, mikro pipet, cawan porselen, inkubator/dry box (PH050 A™), dan timbangan analitik (BEL™).

Preparasi Sample

Pineapple peel that has been cleaned cut into small pieces and then dried in the sun not directly covered with a black cloth covered. Then weigh the dry weight. As much as 500 g was crushed using a blender and sieved with a size of 40 – 60 mesh and crushed to obtain simplicia powder. then the pineapple skin simplicia was extracted using 2 L ethanol pa 96% as a solvent by the maceration method. Furthermore, it was evaporated using a rotary evaporator at 60 °C to obtain a thick extract

Phytochemical Test.

Flavonoid Test. 2 ml of the solution was put into a test tube, then heated for 15 minutes, after which 1 ml of concentrated HCl and 0.05 mg Mg powder were added. If a red, yellow or orange color is formed, this indicates the presence of flavonoids [10]. **Tannin test.** A total of 1 ml of the solution was put into a test tube, then 2-3 drops of 1% FeCl₃ solution were added. If a dark blue or blackish green color is formed, it indicates the presence of tannins [14]. **Saponin Test.** 2 ml of the test solution is put into a test tube, then added with 10 ml of hot water, after that it is cooled and shaken for 10 seconds, if there is foam about 10 minutes with a height of 1 to 10 cm and when 2N HCl is added, the foam does not disappear, indicating the presence of saponins [15]. **Alkaloid Test.** 0.5 g of thick pineapple peel extract was weighed, dissolved with 1 ml of 2N HCl and 9 ml of aquadest. Then put into a test tube and then heated on a water bath for 2 minutes, then cooled and filtered. The resulting filtrate was then divided into 3 test tubes where each of Mayer, Wagner and Dragendorf reagents was added. If a white precipitate is formed in Mayer's reagent, an orange precipitate in Dragendorf's reagent and a brown precipitate in Wagner's reagent, indicates the presence of alkaloids [17]. **Terpenoid Test.** The test extract was dissolved in chloroform and then filtered. The filtrate is then added with a few drops of concentrated sulfate. If a brownish red color is formed at the boundary between the two phases, this indicates that terpenoids are present [17].

Bioassay Larvasida

Eggs of *Aedes aegypti* Strain liverpool F 146 were maintained until instar. Each treatment consisted of five replications. A total of 100 ml of pineapple peel extract from its concentrations of 100 ppm, 200 ppm, 300 ppm, 400 ppm and 500 ppm were put into a 250 ml disposable plastic cup, 25 *Aedes aegypti* larvae were added. Observations were made on dead larvae for 24 hours. Observation of the percentage of larval mortality is carried out by

$$\% \text{ mortality of tested larvae} = \frac{\sum \text{dead tested larvae}}{\sum \text{tested larvae}} \times 100$$

Data analysis

The larval mortality percentage data were tested for normality and homogeneity. If the data is normally distributed and homogeneous, then it is continued with Analysis of Variance (ANOVA), if the ANOVA test results show a significant effect, then it is continued with the Tukey test. To get the value of LD 50 is done by probit test.

RESULTS

Results of the phytochemical test of the Palembang pineapple pell contained phytochemical compounds flavonoids, tannins, saponins, alkaloids, terpenoids, but did not contain steroids contain steroids.

Table Pineapple Pell Phytochemical Compounds Derived from 5 production center in South Sumatra.

| Phytochemical compounds | content | Reagent | Color indicators |
|-------------------------|---------|-----------|------------------|
| Flavanoid | +++ | HCl | Black, Reddish |
| Tanin | ++ | FeCl3 10% | Dark blue |
| Saponin | +++ | Aquadest | Stable foam |

Table 1. ANOVA Effect of pineapple peel ethanol extract concentration on the death of Aedss aegypti.

| | Sum of Squares | df | Mean Square | F | Sig. |
|----------------|----------------|----|-------------|---------|-------|
| Between Groups | 776,300 | 4 | 194,075 | 130,837 | 0,000 |
| Within Groups | 22,250 | 15 | 1,483 | | |
| Total | 798,550 | 19 | | | |

9
 Based on the results of the ANOVA test, Tukey's follow-up test was carried out to see the differences between 10 concentrations of the pineapple peel ethanol extract, the results of the Tukey test are presented in the following table 2.

8
 Table 2. Effect of concentration of ethanol extract of pineapple peel on the mortality of Aedes aegypti larvae

| Concentration (ppm) | Larvvae Mortality | Percentage of Larval Mortality |
|---------------------|-------------------------|--------------------------------|
| 100 | 7,75 (4,22 - 11,28) a | 31 (16,88 - 45,12) a |
| 200 | 17,25 (15,73 - 18,77) b | 69 (62,92 - 72,08) b |
| 300 | 20,00 (18,70 - 21,30) c | 80 (74,80 - 85,20) c |
| 400 | 24,25 (22,73 - 25,00) d | 97 (90,92 - 100,0) d |
| 500 | 25,00 (25,00 -25,00) d | 100 (100,0 - 100,0) d |

16
 Values represent the mean ±standard deviation (SD), n=5

Values in the same column with different superscript letters are significantly different Tukey test (p < 0,005)

8
 The effect of the concentration of the ethanol extract of pineapple peel at a concentration of 100 ppm showed a low action as an Aedes aegypti biolarvicidal, but with an increase in concentration the biolarvicidal activity of the extract increased and at a concentration of 500 ppm it could kill 100 Aedes aegypti larvae (Figure 1).

| | | + HCl | |
|-----------|-----|--------------------|------------------------|
| Alkaloid | +++ | Dragendorff | Orange red precipitate |
| Alkaloid | +++ | Wagner | Chocolate Deposit |
| Alkaloid | +++ | Mayer | White precipitate |
| Terpenoid | ++ | Lieberman Buercard | brown ring |

Note:: +++ (strong)

++ (medium)

+ (weak)

4
 The results of the data normality test showed that the data was normally distributed with p 0.05 = 0.725. 0.05 and the results of the homogeneity test with p 0.05 = .007 showed homogeneous data, then continued with analysis of variance (ANOVA) to see the effect of the concentration of ethanol extract as a biolarvicide on the death of Aedes aegypti. Based on ANOVA, it showed that the concentration of ethanol extract had a significant effect with an F-value of 130.837 greater than F-table at P0.05 = 0.000 <0.05, indicating that bioactive had a significant effect on the death of Aedes aegypti. The results of the ANOVA test can be seen in the following table 1.

12
 The results of the Tukey test showed a significant difference for all concentrations, except between 400 pp and 500 ppm, there was no significant difference in the death of Aedes aegypti larvae.




Figure 1. Effect of Ethanol Extract Concentration on Total Larval Mortality

Table 3. Lethal Concentration of Pineapple Peel Ethanol Extract Against Aedes aegypti Larvae.

| LC 50 (ppm) | LC 95 (ppm) | r | Persamaan regresi |
|-----------------------|-----------------------|------|--------------------|
| 144,04 (89.70-187.19) | 408,93(297,91-856,31) | 0.94 | Y = 12.5 + 0.175 x |

DISCUSSION

Pineapple peel extract has the potential as a biolarvicidal based on the results of analysis containing phytochemical compounds of flavonoids, saponins, tannins, alkaloids and terpenoids. Hasil penelitian yang dilakukan oleh [18]., ekstrak etanol kulit nanas mengandung senyawa fitokimia flavonoid, tanin, saponin dan alkaloid. The content of different secondary metabolites in a plant will cause a distinctive insecticidal effect on insects [18]. The toxicity of pineapple peel to Aedes aegypti larvae is due to the phytochemical compounds it contains, where flavonoids can inhibit breathing and cause nerve wilting, saponins can reduce mucous membranes, alkaloids can act as stomach poisons, tannins can damage tissues and terpenoids can damage breathing [18, 19].

Concentrations of 100 ppm to 300 ppm larvicidal activity of the ethanol extract of pineapple peels are low, but with increasing concentrations of 400 ppm and 500 ppm the activity of the ethanol extract of pineapple peels is very good. Higher concentrations disrupt the respiration and stomach of the larvae. The larval stage represents the most delicate stage in the life cycle of insect vectors. Therefore, efforts to control these vectors are usually directed at the larval stage for disrupting their biological cycle by different strategies [20].

The lethal concentration (LC 50) of the ethanol extract of pineapple peel was 144.04 ppm and the maximum killing power of LC 95 was 408.93 ppm. This shows that the ethanol extract of pineapple peel is toxic to Aedes aegypti larvae. LC50 1000 mg/L = Toxic and LC 50 500 ppm and is said to be active [21]

The results of the probit analysis Lethal Concentration 50 (LC 50) of the ethanol extract of pineapple peel was 144.04 ppm and the maximum killing power was 408.93 ppm (LC 95). The correlation between extract concentration and activated larvicide is very close with a value of $r = 0.94$ and the regression equation can be seen in the table 3.

CONCLUSION

Pineapple peel extract has the potential to be developed as an environmentally friendly Aedes aegypti biolarvicid that can replace synthetic insecticides in controlling Aedes aegypti as a vector of dengue hemorrhagic fever in its larval stage. Concentrations up to 300 ppm very low activity and 500 ppm concentration 100 percent mortality of Aedes aegypti larvae.

ACKNOWLEDGEMENTS

This paper is part of an Environmental Science Doctoral Dissertation. The author would like to coaching team whose help and support made the research possible.

REFERENCES

- [1] Indonesian Ministry of Health. 2022. The situation of DHF (dengue) in Indonesia. Indonesian Ministry of Health. Directorate of Prevention and Control of Vector and Zoonotic Communicable Diseases.
- [2] T. A. Garjito et al., "Stegomyia Indices and Risk of Dengue Transmission: A Lack of Correlation," Front. Public Heal., vol. 8, no. July, hal. 1–13, 2020, doi: 10.3389/fpubh.2020.00328.
- [3] Wagner, C.E. et al., "Climatological, virological and sociological drivers of current and projected dengue fever outbreak dynamics in Sri Lanka," J. R. Soc. Interface, vol. 17, no. 167, 2020, doi: 10.1098/rsif.2020.0075.
- [4] Ghifari A 2013. Detection of insecticide synthetic pyrethroid resistance on dengue vector Aedes aegypti (L.) in Palembang using polymerase chain reaction. Aspirato 5(2):34-44.
- [5] Inawan DWP, Kamo, Jayadi H (2014). Effect of cayenne pineapple peel extract in reducing the number of Coli bacteria in clean water disinfection process. Journal of Widya Warta : 96-107.
- [6] Putri RMA, Yuanita T, Roelianto M (2016). Antibacterial power of pineapple skin extract (Ananas comosus) on the growth of enterococcus faecalis bacteria. Conservative Densistry Journal 6(2):1-6.
- [7] Yeraganreddy PR, Peraman R, Routhu H, Babu N (2013). In vitro

- antitubercular and antibacterial activities of isolated constituents and column fractions from leaves of *Cassia occidentalis*, *Camellia sinensis* and *Ananas comosus*. *African Journal of Pharmacology and Therapeutics* 2(4):116-123.
- [8] Kaushik J, Kundu (2018). Phytochemical screening, anti-oxidant and anti-microbial activity of polyphenolic flavonoids isolated from fruit of *Ananas comosus* in various solvents. *International Journal of Scientific and Publication* 8(2):31-55.
- [9] Dabesor AP, Asowata-Ayodele AM, Umoitte P (2017). Phytochemical compositions and antimicrobial activities of *Ananas comosus* pell (M) and *Cocos nucifera* Kernel (L) on selected food borne pathogens. *American Journal of Plant Biology* 2(2):73-76. Doi:10.11648/j.ajpb.20170202.15.
- [10] Tomini VI, Haiti M (2018). Analysis of pineapple (*Ananas comosus*) on the death of *Aedes aegypti* mosquito larvae. *Journal of Health* 9(3):412-428.
- [11] Juariah S and Irawan MP (2017). Biolarvacide ethanol extract of pineapple skin (*Ananas comosus* L. Merr) against *Culex* sp. Mosquito larvae.
- [12] Sharma G, Vivek I, Gupta AK, Ganjewala D, Gupta C, Prakash D (2017). Phytochemical composition, antioxidant and antibacterial potential of under utilized parts of some fruits. *International Food Research Journal* 24(3):1167-1173.
- [13] Central Bureau of Statistics for Prabumulih City. Prabumulih in Figures 2021.
- [14] Kumaradewi DAP, Subaidah WA, Andayani WA, Al-mokaram A (2021). Phytochemical screening and activity test antioxidant ethanol extract on *Buni* leaves (*Antidesma bunius* L. Spreng) using DPPH method. *Journal of Research in Science Education* 7(2):275-280..
- [15] Harborne JB 1998. *Phytochemical methods: A guide to modern technologies of plant analysis*. 3rd Edition. London, Chapman and Hall, 302 p.
- [16] Rabima and Marshall 2017. Stability test of antioxidant cream preparations ethanol extract 70% from melinjo (*Gnetum gnemon* L.) seeds. *Indonesian Natural Research Pharmaceutical Journal* 2(1):107-121.
- [17] Sinurat JP, Krisdianilo V, Karo RM, Berut R 2020. Analysis of total terpenoids from *Maniltoa grandiflora* (A. Gray.) Scheff leaves using TLC and HPLC methods. *Journal of Science and Applied Chemistry* 2(2):40-44.
- [18] Reiza, I.A., Laode Rijai, L. and Mahmudah, F.2019. *Skrining Fitokimia Ekstrak Etanol Kulit Nanas (Ananas comosus (L.) Merr)*. Proceeding of Mulawarman Pharmaceuticals Conferences. Samarinda, 16-17 Oktober 2019, pp.104 -108. DOI: <https://doi.org/10.25026/mpc.v10i1.371>
- [19] Nursal and Sari Yeanny. M.S. The Egg Hatchability and The Development of *Aedes aegypti* Mosquitoes in Ethanol Extracts of The Leaves of Bitter Melon (*Momordica charantia* L.) and Basil (*Ocimum basilicum* L.). *IOP Conf. Series: Earth and Environmental Science* 305 (2019) 012048. Doi:10.1088/1755-1315/305/1/012048.
- [20] Perumalsamy H, Jang MJ, Kim JR, Murugan K, Ahn YJ (2015). Larvicidal activity and possible mode of action of four flavonoids and thw fatty acids identified in *Milletia pinnata* seed toward three mosquito species. *Parasite and Vector* 8:237.
- [21] Cui C, Yang Y, Zhao T, Zou K, Peng C, Wan HCX, How R (2019). Insecticidal activity and insecticidal mechanism of total saponins from *Camellia oleifera*. *Molecules* 24(4518):1-12
- [22] Carneiro, VCS, Lucena, L.B, Ronaldo Figueiró, R. and el Victório, C.P. Larvicidal activity of plants from Myrtaceae against *Aedes aegypti* L. and *Simulium pertinax* Kollar (Diptera). 2021. *Journal of the Brazilian Society of Tropical Medicine*. Vol.:54:(e00922020): 2021.
- [23] Sumihe G, Runtuwene MRJ, Rorong JA (2014). Phytochemical analysis and determination of LC 50 liwas leaf methanol extract. *Scientific Journal of Science* 14(2):125-128.

Phytochemical and Larvicidal Activity of Pineapple (Annanas comusus L.) Peel Extract Against Aedes aegypti of Dengue Vector

ORIGINALITY REPORT

17%

SIMILARITY INDEX

11%

INTERNET SOURCES

12%

PUBLICATIONS

3%

STUDENT PAPERS

PRIMARY SOURCES

- 1 N F Dhenge, P Pakan, K Lidia. "Larvacide effectiveness of Papaya leaf extract (Carica papaya) on the mortality of larvae vector of Dengue hemorrhagic fever caused by Aedes aegypti", IOP Conference Series: Earth and Environmental Science, 2021
Publication 2%
- 2 gotomydoctor.com
Internet Source 2%
- 3 www.scielo.br
Internet Source 1%
- 4 Endang Sri Sunarsih, Ebta Narasukma Anggraeny, Patricia Sanggita Listyoputri Wibowo, Novi Elisa. "Phytochemical Screening and Antioxidant Activity of Strawberry Juice (Fragaria ananassa Duchesse) Against Ureum Level, Creatinin, and Enzyme Catalase Activity In Isoniazid-Induced Wistar Male Rats", STRADA Jurnal Ilmiah Kesehatan, 2020
Publication 1%

| | | |
|-----------------|---|-----|
| 5 | <p>Begum Fauziyah, Asna Defi Batrisyia, Alif Firman Firdausy, Burhan Ma'arif, Siti Maimunah. "Analysis of Bioactive Compounds in Nabeez Water", Biomedical and Pharmacology Journal, 2022</p> | 1 % |
| Publication | | |
| 6 | <p>Khadijah, Nunuk Hariani Soekamto, Siti Maisuri Tadjuddin Chalid, Nur Fatin Rafidah. " Total Phenol Content and Activities of Antioxidant Extracts Methanol Limes By Uv-Vis Spectrophotometry ", E3S Web of Conferences, 2021</p> | 1 % |
| Publication | | |
| 7 | <p>Submitted to West Coast University</p> | 1 % |
| Student Paper | | |
| 8 | <p>www.scitepress.org</p> | 1 % |
| Internet Source | | |
| 9 | <p>Muntasir, Yohannes Mau Abanit, Pius Weraman, Ahmad Fudholi. "Natural larvicide mixed lime leaf extract and galangal rhizome to mortality Aedes aegypti larvae", Biocatalysis and Agricultural Biotechnology, 2022</p> | 1 % |
| Publication | | |
| 10 | <p>Nurani Hasanela, Fensia Analda Souhoka. "Effect of Heating Coarse Extract of Brown Macroalgae (Padina australis) from Tial</p> | 1 % |

Waters, Salahutu District, Central Maluku
Regency on Antioxidant Activity", Indo. J.
Chem. Res., 2022

Publication

11 ejournal2.litbang.kemkes.go.id 1 %
Internet Source

12 cyberleninka.org 1 %
Internet Source

13 jurnal.ugm.ac.id 1 %
Internet Source

14 Nursal, Mayang Sari Yeanny. " The Egg
Hatchability and The Development of
Mosquitoes in Ethanol Extracts of The Leaves
of Bitter Melon (L.) and Basil (L.) ", IOP
Conference Series: Earth and Environmental
Science, 2019 1 %
Publication

15 journal2.uad.ac.id 1 %
Internet Source

16 www.mdpi.com 1 %
Internet Source

17 journal.ubb.ac.id 1 %
Internet Source

Exclude quotes On

Exclude bibliography On

Exclude matches < 1%