

Assesment of Antioxidant Activity Test of Kersen Leaf (The phenolic.) and Epiphyte with DPPH (2.2-Diphenyl-1-Picrylhidrazyl)

by Nita Parisa

Submission date: 18-Jul-2022 01:53PM (UTC+0700)

Submission ID: 1872040392

File name: 9-Article_Text-819-1-10-20220411.pdf (249.99K)

Word count: 3430

Character count: 18588



Assesment of Antioxidant Activity Test of Kersen Leaf (*The phenolic.*) and Epiphyte with DPPH (2,2-Diphenyl-1-Picrylhidrazyl)

Nurlutfiyah Aini¹, Nita Parisa^{2*}, Fatmawati³

¹ Undergraduate Student Medical Faculty, Universitas Sriwijaya, Palembang, Indonesia

² Department of Pharmacology, Medical Faculty, Universitas Sriwijaya, Palembang, Indonesia

³ Department of Biochemistry, Medical Faculty, Universitas Sriwijaya, Palembang, Indonesia

ARTICLE INFO

Keywords:

Antioxidant
Kersen Leaf
Epiphyte DPPH
IC₅₀

*Corresponding author:

Nita Parisa

E-mail address:

nitaparisa@unsri.ac.id

All authors have reviewed and approved the final version of the manuscript.

<https://doi.org/10.37275/AMCR.v1i2.9>

ABSTRACT

Antioxidant is very important to give protection against free radical activity and highly reactive molecules that could lead in slowing the progression of degenerative disease. In case of degenerative disease, internal antioxidant cannot neutralize the increasing concentration of free radical. Because of that, human needs external antioxidant. Kersen (*Muntingia calabura* L.) is a plant that is known for its antioxidant content. Plants containing antioxidant experience is kersen (*Muntingia calabura* L.). Research study to determine the antioxidant activity of Kersen plant and knows the difference of antioxidant activity, based on the process of extract and infusion. Research was done by experimental study which was oriented in testing antioxidant activity in (*Morinda citrifolia* L.) extract and infusion. Extraction was done by using 96% ethanol as solvent meanwhile infusion was made by using aquadest. Extract and infusion were divided into group of concentration and antioxidant activity was tested by DPPH(2,2-Diphenyl-1-Picrylhidrazyl) method by measuring the absorbance using spectrophotometer at 520 nm wavelength. Percentage of DPPH inhibition and IC₅₀ then analyze d using linear regression analysis. Ethanolic extract of kersen leaf and epiphyte had IC₅₀ value of 113,801 ppm and 98,7802 ppm, respectively. Kersen leaf infusion showed 191,7624 ppm IC₅₀ values, besides its epiphyte had 131,6750 ppm. Antioxidant activity of *Muntingia calabura* L. in the order from kersen leaf an epiphyte and epiphyte extract has a higher antioxidant content than others.

1. Introduction

Indonesia is one of the countries which has high-level biodiversity in the world, 3rd ranked after Brazil and Zaire. The biodiversity includes plants of flora and fauna which are spread throughout Indonesia.¹ There are 40,000 species of flora that grow in the world,

30.000 species found in Indonesia with 1.845 species of plants have the potential as traditional medicine.² According to the POM Agency (2006), there are 283 types of plants that have been registered for the use of traditional medicines. Only 13 of the 283 types of medicinal plants have been cultivated. Traditional medicine is widely used to treat several diseases such as urolithiasis, diabetes, high blood

pressure and others.

The ability of a plant as a drug is caused by the content of chemical compounds or active compounds that have the working power of treatment. One of the chemical constituents that have the working power of treatment is antioxidant.³ The emergence of free radicals (hydroxyl) in biochemical mechanisms in the body are the causes of degenerative diseases.⁴ Degenerative diseases, such as osteoporosis, cardiovascular, cancer, diabetes mellitus, and others can be reduced by consuming antioxidants. This is related to the work system of antioxidants which can inhibit oxidation reactions, by binding free radicals and



molecules which are very reactive.⁵

One potential source of natural antioxidants is plants because they contain flavonoid compounds, chlorophyll and tannin.⁶ Kersen (*Muntingia calabura* L.) is a plant that has the potential to be a natural antioxidant.⁷ Antioxidants kersen (*Muntingia calabura* L.) are found in all parts of flowers, fruit and leaves, and the highest activity on the part of the leaf. Various studies show that kersen leaves contain active components of saponins, flavonoids, and tannins when extracted using methanol and ethanol solvents.⁸

The administration of the ethanol extract of the leaves has an effect on the *B.carambolae* fruit fly. The higher the concentration of extracts, the lower the number of pupae and imago fruit fly that appear (Putri, 2016), and the greater the total flavonoid content, the higher the antibacterial activity.^{9,10} Kersen leaves are believed to have the ability to be an antibacterial to *Streptococcus mutans* which have glucosyltransferase enzymes (GTF).¹¹

Another study stated that the antioxidant activity of extract of noni leaf was smaller with IC_{50} value of 98.68 $\mu\text{g/mL}$ than noni leaf infusion with IC_{50} value of 75.65 $\mu\text{g/mL}$.¹² There were differences in the results of antioxidant activity using cold (extraction) and heat (infusa) on medicinal plants. As we know, people use medicinal plants by boiling the plants. Even though the effects of infusion or heating can damage the secondary metabolites in the plant.¹³

The epiphyte has a chemical compound similar to the host plant it occupies. In another study, ethanol extract of kepel parasite leaves has higher antioxidant activity (IC_{50} value 6.43 $\mu\text{g/mL}$) than ethanol extract of leaves of kepel (IC_{50} value $12.57 \pm 0.7 \mu\text{g/mL}$).^{14,15} It is also expected that the associated epiphyte will contain antioxidant activity. In general, Kersen leaf and epiphyte of the truth contain flavonoids which have antioxidant power.

2. Methods

This study is an experimental analytic study with post-test only control group design to determine the

ratio of antioxidant levels to leaves and epiphyte kersen (*Muntingia calabura* L.). The study was conducted from October to November 2018 in the Laboratory.

Biochemistry, Faculty of Medicine, Sriwijaya University. The object of this research is a green plant (*Muntingia calabura* L.) which will be extracted in cold (extraction) and hot (infusa). The parts of the plant to be sampled are kersen leaves and epiphyte. The criteria for this research object are fresh, perfectly shaped and clean, dark green leaves and parasitic leaves.

Data analysis was performed using the Statistical Package for Social Science (SPSS) program and Linear Regression Test to determine the direction and relationship between the independent variables and the dependent variable and to predict the value of the dependent variable if it increases or decreases.

3. Results

Table 1 below presents the absorbance values and percent inhibition of kersen leaf extract. From the 6 concentrations of the kersen leaves, the highest inhibition was obtained at 49.17279% at a concentration of 100 ppm and the lowest was 43.060666% at a concentration of 10 ppm.

Table 2 below shows the absorbance and percent inhibition values of the parasite leaf extract. Of the 6 concentrations of parasitic leaves, the highest percentage inhibition was 49.17279% at a concentration of 100 ppm and the lowest was 39.52206% at a concentration of 10 ppm.

Table 3 below shows the absorbance value and the percentage of inhibition in the infusion of kersen leaves. From the 6 concentrations of the kersen leaves, the highest inhibition was obtained at 47.61029% at a concentration of 100 ppm and the lowest was 45.18382% at a concentration of 10 ppm.

Table 4 below shows the absorbance value and percent inhibition of epiphyte infusion. From the 6 concentrations of epiphyte leaf, the highest inhibition was obtained at 48.71324% at a concentration of 100 ppm and the lowest was 44.71507% at a concentration of 10 ppm.



The measurement of antioxidant activity was carried out using linear regression analysis in SPSS. Linear regression analysis was used to see how much influence x (concentration) has on y (% inhibition) so that the results of the linear regression equation can be seen in the value of x as IC₅₀ (Inhibitory Concentration 50) by replacing the y value to 50. In table 5 below shows that leaf extract cherry leaf has moderate antioxidant activity with an IC₅₀ value of 113.801 ppm.

5 Table 6 below, shows that epiphyte extract has a strong antioxidant activity with an IC₅₀ value of 98.7802 ppm.

Table 7 below, shows that the infusion of kersen leaves has a weak antioxidant activity, that is, with an IC₅₀ value of 191.7624 ppm.

Table 8 below, shows that the parasite leaf infusion

has moderate antioxidant activity with an IC₅₀ value of 98.7802 ppm.

In table 9 below, the results of antioxidant extracts of ethanol extract of leaves and epiphyte plants (*Muntingia calabura* L.) are presented. From the sample, the highest antioxidant activity was found in the epiphyte extract with an IC₅₀ value of 98,7802 ppm, compared to kersen leaf extract with an IC₅₀ value of 113.801 ppm.

In table 10 below, the results of antioxidant activity of leaf infusion and epiphyte are obtained from plants (*Muntingia calabura* L.). From the sample, the highest antioxidant activity was obtained in epiphyte infusion with an IC₅₀ value 131.6750 ppm, compared to kersen leaf infusion with an IC₅₀ value of 191.7624 ppm.

Table 1. Absorbance and percent inhibition of kersen leaf extract

Extract	Concentration (ppm)	Absorbance	Absorbance Of DPPH	% Inhibition
Kersen Leaf	10	0.619	1.088	43.06066
	20	0.609	1.088	44.02574
	30	0.598	1.088	44.99081
	50	0.588	1.088	45.95588
	70	0.577	1.088	46.92096
	100	0.553	1.088	49.17279

Table 2. Absorbance and percent inhibition of epiphyte extract

Extract	Concentration (ppm)	Absorbance	Absorbance DPPH	% Inhibition
Epiphyte	10	0.658	1.088	39.52206
	20	0.6335	1.088	41.77390
	30	0.623	1.088	42.73897
	50	0.595	1.088	45.31250
	70	0.567	1.088	47.88603
	100	0.553	1.088	49.17279

Table 3. Absorbance and percent inhibit kersen leaf infusion

Infusion	Concentration (ppm)	Absorbance Sample	Absorbance DPPH	% Inhibition
Kersen Leaf	10	0.5964	1.088	45.18382
	20	0.5928	1.088	45.51471
	30	0.5892	1.088	45.84559
	50	0.5832	1.088	46.39706



Table 4. Absorbance and Percent of inhibition of Epiphyte Infusion

Infusion	Concentration (ppm)	Absorbance Sample	Absorbance DPPH	% Inhibition
	10	0.6015	1.088	44.71507
	20	0.5895	1.088	45.81801
Epiphyte	30	0.585	1.088	46.23162
	50	0.5805	1.088	46.64522
	70	0.5715	1.088	47.47243
	100	0.558	1.088	48.71324

Table 5. Antioxidant activity of kersen leaf extract

Sample	Regression Value	Equation IC ₅₀	Antioxidant Activity
Kersen Leaf Extract	$Y = 0.0642 x + 42.649$	113.801 ppm	Moderate

Table 6. Antioxidant activity of epiphyte extract

Sample	Regression Value	Equation IC ₅₀	Antioxidant Activity
Epiphyte Extract	$Y = 0.107 x + 39.391$	98.7802 ppm	Strong

Table 7. Antioxidant activity of kersen leaf infusion

Sample	Regression Value	Equation IC ₅₀	Antioxidant Activity
Kersen Leaf Infusion	$Y = 0.0261 x + 44.995$	191.7624 ppm	Weak

Table 8. Antioxidant Activity of Epiphyte Infusion

Sample	Regression Value	Equation IC ₅₀	Antioxidant Activity
Kersen Leaf Infusion	$Y = 0.0261 x + 44.995$	191.7624 ppm	Weak

Table 9. Antioxidant Activity of Ethanol Extract of Kersen Plant (*Muntingia calabura* L.)

Sample	IC ₅₀ Value
Kersen Leaf Extract	113.801 ppm
Epiphyte Leaf Extract	98.7802 ppm

Table 10. Antioxidant Activity of Kersen Infusion (*Muntingia calabura* L.)

Sample	IC ₅₀ Value
Kersen Leaf Infusion	191.7624 ppm
Epiphyte Leaf Infusion	131.6750ppm



4. Discussion

An increase in% inhibition of ethanol extract indicated that an increase in extract concentration would affect the ability of extracts to soak free radicals.

This result is supported by research that states the percentage inhibition (percent inhibition) of free radical activity will also increase along with concentration increasing.¹⁶

The results of the existence of ethanol extract of kersen leaves have an IC₅₀ value of 113.801 ppm, classified as a moderate antioxidant (100-150 ppm). While the ethanol extract of the epiphyte has an IC₅₀ value of 98.7802 ppm, classified as a very strong antioxidant (50-100 ppm). This shows that the leaves and epiphyte in kersen plants have antioxidant activity that is classified as strong with the highest antioxidant activity found in epiphyte because the smaller the IC₅₀ value the greater the antioxidant activity.

The results of this study are in accordance with previous studies using different objects namely kepel plants, where ethanol extract of kepel parasite leaves had higher antioxidant activity (IC₅₀ value 6.43 ppm) than ethanol extract of kepel leaves (IC₅₀ value 12.57±0.7 ppm).^{14,17}

Another theory that supports ethanol extract of the epiphyte has antioxidant activity that is higher than the leaves of the stemcell because the epiphyte contains more antioxidant compounds than the stem cell. Antioxidant compounds found in epiphyte are amino acids, carbohydrates, alkaloids, saponins, flavonoids, alkaloids, terpenoids, and tannins.^{18,19} While the antioxidant compounds found in kersen leaves are active components of saponins, flavonoids, and tannins.²⁰

In this study, infusion (heating) was carried out by soaking the plants with 500 ml of water to boil so that the kersen leaves infusion were obtained with an IC₅₀ value of 191.7624 ppm, classified as a weak antioxidant (150-200 ppm). While the parasite leaf infusion has an IC₅₀ value of 131.675 ppm, classified as a moderate antioxidant (100-150 ppm).

The results of this study are in accordance with previous studies using different objects, namely on the

soursop plant, where the infusion of leaves of soursop parasites has a higher antioxidant activity (IC₅₀ value 33.5 ppm) than the infusion of soursop leaves (IC₅₀ value 45 ppm).¹⁵

When compared between ethanol extract and infusion from each part of the plant, **the results showed that the preparation of ethanol extract had a stronger antioxidant activity** than infusion. The results of this study are in accordance with previous studies that kersen leaf infusion has a higher antioxidant activity of 196.80 ppm, whereas in ethanol extract, kersen leaves have antioxidant activity of 164.12 ppm.^{21,22}

The heating process of the infusa which can cause a reduction or damage to the content of antioxidant compounds found in the leaves and leaves of the parasite, the plant seeds (*Muntingia calabura* L.). Cersen plant has phenolic compounds. Phenolic compound is one of the constituents of green plants which act as antioxidants. Extraction at temperatures above 60°C can reduce phenolic compounds found in plants²³ and in this study it has heated at the temperature of 100°C. The use of heating at a high enough temperature will damage the active compounds contained in simplicia, especially antioxidant compounds.²⁴ In addition, compounds flavonoids and tannins found in plants are heat resistant.²⁵ Antioxidant activity decreases with increasing temperature and length of heating time.²⁶ This is because the longer the oven drying process (heating), the more free water content evaporates so that the mass of the dry material produced will also decrease.²⁷

5. Conclusion

Chrysanthemum plants (*Muntingia calabura* L.) have antioxidant activity in the leaves and epiphyte. Epiphyte extract has stronger antioxidant activity than kersen leaf extract. Epiphyte extract has stronger antioxidant activity than the kersen leaves. The extract of extract has a higher antioxidant activity than infusion.



6. References

1. Nugroho A, Anis T, and Ulfah M. Analysis of the diversity of fruiting plants in Surokoto protected forest, Kendal, Central Java and their potential as a bird conservation area. 2015; 1(2): 472-476.
2. Zuraida, Lelana A and Nuronia HS., Development of Forestry Biopharmaca. Indonesian Forestry Biopharmaca Flowers from Forest Plants for the Excellence of Nations and Countries. Center for Plantation Forestry Research and Development. Forestry Research and Development Agency. Ministry of Forestry. 2009; 3-13.
3. Hernani and Tri Marwati. Post-Harvest Technology of Medicinal Plants. BB Post Harvest: Bogor. 2012
4. Oeinitan, J. 2013. Antioxidant Power of Ethanol Extract of Mangosteen Fruit Skin (*Garcinia mangostana* Linn.) Results of Stirring and Reflux. Surabaya Student Scientific Journal. 1 (2): 1-2.
5. Winarsi, H. Natural Antioxidants and Free Radicals. Yogyakarta: Kansius. 2007.
6. Lee S, Ahn K, Paik HY, Chung Sang-Jin et al. Serum immunoglobulin E (IgE) levels and dietary intake of Korean infants and young children with atopic dermatitis. Nutrition research and practice; 2012; 6(5): 429-35.
7. Khan MAY, Ramadas D, Mundasada SC, Kumar SN, Chikkanna D. Antioxidant and free radical scavenging activities from root extracts of *Muntingia Calabura*. Scholar Journal of Applied Medical Sciences (SJAMS). 2015; 3(6B): 2309-2312
8. Surjowardojo. P, Sarwiyono, Thohari. I, Ridhowi. A, Quantitative and Qualitative Phytochemical Analysis of *Muntingia Calabura* L. Journal of Biology, Agriculture and Healthcare. 2014; 4(16).
9. Putri, D. A. Effect of Giving Kersen Leaf Extract (*Muntingia calabura* L.) on *Bactrocera carambolae* Fruit Flies. Ahmad Dahlan University. 2016; 9(2): 139-143.
10. Manik KM, Islam MAS, Wahid K. Md, Morshed, MM, Kamal, S., et al. Investigation of In Vitro Antioxidant, Antimicrobial and Thrombolytic Activity of the Exocarp of *Spondias pinnata* (Anacardiaceae). Borderless Science Publishing. 2013; 1(3): 191-201.
11. Isnarianti R, Wahyudi IA and Puspita RM. *Muntingia calabura* L. Leaves Extract Inhibits Activity Glucosyltransferase of *Streptococcus mutans*. Journal of Dentistry Indonesia. 2013; 20(3): 59-63.
12. Irianti T, Puspitasari ALM. and HR R. Catching Activities of 2-2-Diphenil-1- Pycrilhydrazil (DPPH) Radical Ethanolic Extract of Menkudu Leaves (*Morinda citrifolia* L.), and Brotowali Stem (*Tinospora crispa* L.), Water Fraction and Hydrolyzed Water action. Faculty of Pharmacy, Gaja Mada University, Yogyakarta. 2015; 20(3). 140-148.
13. Hudha, et al. Effervescent Powder Based on Beluntas Leaf Extract (*Pluchea indica* less) as a Source of Natural Antioxidants. Food Journal and Agroindustry. 2015; 3(4): 1412-1422.
14. Sunami T, Pramono S, and Asmah R. Antioxidant-free Radical Scavenging of Flavonoids from The Leaves of *Stelechocarpus burahol* (Bl.) Hook. F. and Th., Indonesian Pharmacy Magazine, 2007; 18(3): 111-116.
15. Patria, W. and Soegihardjo, C. Antioxidant Activity Test Using 1,1-Difeny-2- Pikrilhidrazil (DPPH) Radical and Determination of Total Phenolic Content of Ethyl Acetate Fraction of Leaf Ethanolic Extract of *Dendrophthoe pentandra* L. Miq. which grows on the Kepel tree (*Stelechocarpus burahol* (Bl) Hook. f.). Science and Community Pharmacy. 2013; 10(1): 1693-5683.
16. Hanani, E., Mun'im, A., and Sekarini, R. 2005. Identification of Antioxidant Compounds in Sponge *Callyspongia* SP of the Thousand Islands. Science Magazine.
17. Patria W. and Soegihardjo, C. Antioxidant



- Activity Test Using 1,1-Difenyl-2- Pikrilhidrazi (DPPH) Radical and Determination of Total Phenolic Content of Ethyl Acetate Fraction of Ethanolic Leaves of Parental Leaves *Dendrophthoe pentandra* L. Miq. which grows on the Kepel tree (*Stelechocarpus burahol* (Bl) Hook. f.). *Science and Community Pharmacy*. 2013; 10(1): 1693-5683.
18. Ikawati, M., Wibowo, A.E., Navista, S.O.U., & Adelina, R. Utilization of parasites as anticancer agents, International Seminar of Indonesia - Malaysia Update 2008, Gadjah Mada University and Universiti Sains Malaysia. 2008.
 19. Nirwana, A. P., Astrin, O. P. and Widiyani, T. Phytochemical Screening Ethanol Extract of Kersen Parasite Leaves (*Dendrophthoe pentandra* (L.) Miq.). *University Journal Eleven March*. 2015; 3(2): 9-15.
 20. Surjowardojo. P, Sarwiyono, Thohari. I, Ridhowi. A, Quantitative and Qualitative Phytochemical Analysis of *Muntingia Calabura* L. *Journal of Biology, Agriculture and Healthcare*. 2014; 4(16).
 21. Marjoni, M. R., Afrinaldi & A. D. Novita. Phenol Total Content and Antioxidant Activity of Kersen Leaf Water Extract (*Muntingia calabura* L.). *Yarsi Medical Journal*. 2015; 23(3): 187-196.
 22. Hasanah M, Andriani N, Noprizon. Comparison of antioxidant activity of ethanol extract of kersen leaves (*Muntingia calabura* L.) as a result of maceration and re-extraction. *Scientia*. 2016; 6 (2); 84-90.
 23. Lantah, Montolalu and Reo. Phytochemical content and antioxidant activity of *Kappaphycus alvarezii* Seaweed Methanol Extract. *Journal of Fishery Product Technology Media* 2017; 5(3): 167-172.
 24. Pratiwi, Dina. Wahdaningsih S. Isnindar. Restianti, Eka. Test Antioxidant Activity of Shallot Leaf (*Eleutherine Americana* Merr.) With DPPH 2,2-Diphenyl-1- Picrylhydrazyl Method). *Traditional Medicine*. 2013; 18(1): 9-16. ISSN: 1410-5918.
 25. Gupita, C. Effect of Various pHs of Fruit Juice and Temperature of Pasteurization on Antioxidant Activities and Acceptance Levels of Mangosteen Fruit Skin. *Journal of Nutrition College*. 2012. 1(1): 67-79.
 26. Husni, A., Deffy R.P, and Iwan Y.B.L. Antioxidant Activity of *Padina* sp. at various temperatures and drying times. *Journal of Fisheries*. 2014; 9(2): 165-173.
 27. Sudarmadji, S., Haryono, B., & Suhardi. *Food and Agriculture Material Analysis*. Liberty. Yogyakarta. 2007.



Assesment of Antioxidant Activity Test of Kersen Leaf (The phenolic.) and Epiphyte with DPPH (2.2-Diphenyl-1-Picrylhidrazyl)

ORIGINALITY REPORT

20%
SIMILARITY INDEX

19%
INTERNET SOURCES

6%
PUBLICATIONS

7%
STUDENT PAPERS

PRIMARY SOURCES

1	hmpublisher.com Internet Source	10%
2	Submitted to ESADE Student Paper	3%
3	Submitted to Sriwijaya University Student Paper	2%
4	www.bioscmed.com Internet Source	1%
5	journal.uinjkt.ac.id Internet Source	1%
6	"Bioactive Compounds in Underutilized Vegetables and Legumes", Springer Science and Business Media LLC, 2021 Publication	1%
7	I Ketut Budaraga, Eva Susanti, Asnurita Asnurita, Elliza Nurdin, Ramaiyulis Ramaiyulis. "The Antioxidant Characteristics of The Liquid Smoke of Cocoa Shell (Theobroma cacao, I)	1%

In Different Water Content Variations", Journal of Applied Agricultural Science and Technology, 2019

Publication

8	www.real-statistics.com Internet Source	1 %
9	Aparicio-Soto, Marina, Susana Sánchez-Fidalgo, Alejandro Gonzalez-Benjumea et al. "Naturally Occurring Hydroxytyrosol Derivatives: Hydroxytyrosyl Acetate and 3,4-Dihydroxyphenylglycol Modulate Inflammatory Response in Murine Peritoneal Macrophages. Potential Utility as New Dietary Supplements", Journal of Agricultural and Food Chemistry Publication	1 %
10	jurnal.iseibandung.or.id Internet Source	1 %
11	media.neliti.com Internet Source	1 %

Exclude quotes On

Exclude matches < 1%

Exclude bibliography On