

ASSOCIATION OF APPLIED BIOLOGISTS

ASPECTS OF APPLIED BIOLOGY 113

**Making Crop Rotations
Fit for the Future**

**Great North Museum at the Hancock,
Newcastle-upon-Tyne, UK**

20-21 December 2011

Produced by the Association of Applied Biologists

**Copies available from: The AAB Office
Warwick Enterprise Park, Wellesbourne,
Warwick CV35 9EF, UK
<http://www.aab.org.uk>**

REFERENCES

The correct form of reference for this publication, which is based on a meeting of the Association of Applied Biologists, is:

Aspects of Applied Biology **113**, *Making Crop Rotations Fit for the Future*, pp. xx–xx

Papers are included herein without any liability for loss or damage suffered as a result of their application or use. Reference herein to trade names and proprietary products without special acknowledgement does not imply that such names, as defined by the relevant protection laws, may be regarded as unprotected and thus free for general use. No endorsement of named products is intended nor is any criticism implied of similar products which are not mentioned. *Please note* certain names of chemicals featured in this publication are Registered Trademarks.

This publication is copyright under the Berne Convention and the Universal Copyright Convention. All rights reserved. Apart from any relaxation permitted under national copyright laws, no part of this publication may be reproduced, stored in a retrieval system or transmitted in any form or by any means without prior permission of the copyright owners. Permission is not, however, required to copy Abstracts of papers on condition that a full reference to the source* is shown. Multiple copying of the contents of the publication without permission from both The Association of Applied Biologists through the Executive Officer and separately from the author, or other holder of the unilateral copyright, is always illegal.

The Association of Applied Biologists and Editors cannot be held responsible for errors or any consequences arising from the use of information contained in this publication; the views and opinions expressed do not necessarily reflect those of the Association of Applied Biologists and Editors.

ENQUIRIES

Enquiries concerning the technical content of chapters should be addressed directly to the authors but other matters should be directed to the Executive Officer, AAB Office, Warwick Enterprise Park, Wellesbourne, Warwick CV35 9EF, UK.

Printed in UK

Published by the Association of Applied Biologists
Warwick Enterprise Park, Wellesbourne, Warwick CV35 9EF, UK

© 2011 The Association of Applied Biologists

ISSN 0265-1491

*Full reference: *Aspects of Applied Biology* **113**, *Making Crop Rotations Fit for the Future*, published by the Association of Applied Biologists, Warwick Enterprise Park, Wellesbourne, Warwick CV35 9EF, UK.



Legume Futures is an EU funded research programme that aims to deliver more-sustainable legume supported cropping systems.

For more information please visit:
www.legumefutures.eu

INTRODUCTION

Crop rotations play an important role in maintaining crop productivity and soil health in a wide range of farming systems. They are important for sustaining crop yield (feed, food and fuel crops) and fertility building, as well as weed, pest and disease control. UK farming systems have tended to adopt simplified rotations; however, more crops are now available to farmers and there is support to diversify rotations through the inclusion of cover crops as well as new market opportunities for non-food crops. Increased crop diversity is also known to support increased biodiversity on farm. It has also been suggested that increased crop diversity will give added resilience to farming systems in the light of climate change. This inter-disciplinary conference will allow consideration of the impacts of rotations on crop yield (feed and food crops), nutrient cycling, weed, pest and disease control, farm biodiversity and environmental impact. Not all rotations will work everywhere and thus there is a need to consider the drivers for rotation design and the trade-offs between the different ecosystem services that rotations can provide for future agriculture. This conference brings together scientists and advisors from a range of disciplines and environments in Europe and beyond. We look forward to fruitful discussion and exchange of information which can be applied to the design of crop rotations in future agriculture.

Elizabeth Stockdale
Christine Watson

CONTENTS

	<i>Page</i>
Below-ground interactions for sustainable cropping systems MICHAEL J GOSS, ISABEL BRITO, MÁRIO CARVALHO, SABARUDDIN KADIR & AMARILIS DE VARENNES	1–8
Agri-Environment Stewardship; improving performance and agronomic management through manipulating the rotation of in-field options A R LEAKE, P G L THOMPSON & P E JARVIS	9–14
New Farming Systems Research (NFS) project: long term research seeking to improve the sustainability and resilience of conventional farming systems R M STOBART & N L MORRIS	15–23
The use of crop yields and gross margins to optimise arable rotations and farm financial output J MACKAY	25–28
The impact of break-crop and cereal rotations on crop performance and profit margins P D FORRISTAL & J GRANT	29–36
Effects of diverse and cereal-intensive crop rotations on crop yields and soil fertility - reflections after the first cycle of the Nafferton long-term trials J M COOPER, M BARANSKI, A CARMICHAEL, R CHAPMAN, M EYRE, G HALL, L LUECK, L REMPELOS, C SCHMIDT, D SREDNICKA-TOBER, P N SHOTTON, C TETARD-JONES, S WILCOCKSON & C LEIFERT	37–43
Delivering the barebones for designing more weed suppressive crop rotations L BASTIAANS & H N C BERGHUIJS	45–52
An investigation of crop choice through crop rotation WEIQI LUO & NIGEL BOATMAN	53–60
The design of a new cropping system experiment to be used as a research platform – maize and winter wheat in monoculture and rotations G BERGKVIST, B BÅTH & I ÖBORN	61–66
Sustainability Trial in Arable Rotations (STAR project): a long term farming systems study looking at rotation and cultivation practice R M STOBART & N L MORRIS	67–73
Long-term field experiments in Sweden – what are they designed to study and what could they be used for? G BERGKVIST & I ÖBORN	75–85
Using legumes to drive crop rotations and sustainable farming systems F L STODDARD	87–88
Age of grass-clover sward: does it affect the predictions N fixed for subsequent crops? C F E TOPP, R M REES & C A WATSON	89–92

	<i>Page</i>
Benefits and limitations of white clover N₂ fixation as the main N input into mixed cropping rotations and dairy pastures: a case study of the recent agricultural intensification in the Canterbury Plains, New Zealand M ANDREWS, B A McKENZIE, S HODGE & J L MOIR	93–96
Fertility building strategies in organic crop rotations F RAYNS, A ROSENFELD & U SCHMUTZ	97–100
Implication of green manure crop cultivation with rock phosphate in organic rotations O G G KNOX, R L WALKER, A C EDWARDS, A MCVITTIE & P MASKEL	101–106
Trace elements in agricultural systems C F E TOPP, C A WATSON & I ÖBORN	107–110
The long-term effects of widely differing soil pH on the yields of an eight course crop rotation established in 1961 R L WALKER, C A WATSON & A C EDWARDS	111–115
Effects of crop management regimes and potato genotype on Nitrogen Use Efficiency L REMPELOS, J COOPER, C TETARD-JONES, S WILCOCKSON, M EYRE, G HALL, P SHOTTON, A M R GATEHOUSE & C LEIFERT	117–118
Can we make crop rotations fit for a multi-functional future? E A STOCKDALE & C A WATSON	119–126
 <i>Posters</i>	
Assessing the economic value of cereal-legume intercrops in low-input rotational cropping systems JOHN A BADDELEY, VALENTINI A PAPPA, ROBIN L WALKER, CHRISTINE A WATSON & ROBERT M REES	127–132
The importance of beetle banks for small mammals B BOAG & B FENTON	133–135
The importance of no-till in the development of cropping systems to maximise benefits of arbuscular mycorrhiza symbiosis ISABEL BRITO, MÁRIO CARVALHO & MICHAEL J GOSS	137–141
The use of intercrops in rotations V A PAPPA, R M REES, J A BADDELEY, A PRISTERI, HERWART BÖHM, F L STODDARD & C A WATSON	143–148
The effects of crop rotation and soil management on worm activity and soil physical quality R L WALKER, C A WATSON, G ARMSTRONG & B BALL	149–154
Improving barley yields in organic stockless farming systems through innovations in the management of green manure swards S HANSEN, R B FRØSETH, A K BAKKEN, H RILEY, K T KRISTENSEN & M A BLEKEN	155–157

See discussions, stats, and author profiles for this publication at: <http://www.researchgate.net/publication/278021106>

Below-ground interactions for sustainable cropping systems

ARTICLE · JANUARY 2011

READS

8

5 AUTHORS, INCLUDING:



Michael J Goss

University of Guelph

118 PUBLICATIONS 2,519 CITATIONS

SEE PROFILE



Isabel Brito

Universidade de Évora

11 PUBLICATIONS 43 CITATIONS

SEE PROFILE



Mário Carvalho

Universidade de Évora

70 PUBLICATIONS 183 CITATIONS

SEE PROFILE



Sabaruddin Kadir

Universitas Sriwijaya

14 PUBLICATIONS 7 CITATIONS

SEE PROFILE

Below-ground interactions for sustainable cropping systems

By MICHAEL J GOSS¹, ISABEL BRITO², MÁRIO CARVALHO², SABARUDDIN KADIR³
and AMARILIS DE VARENNES⁴

¹*University of Guelph, Kemptville Campus, Kemptville K0G 1J0, Ontario, Canada*

²*University of Évora, ICAAM, Apartado 94, 7002 - 540 Évora, Portugal*

³*University of Sriwijaya, Faculty of Agriculture, Jl. Padang Selasa No. 524, Bukit Besar,
Palembang, South Sumatra 30139, Indonesia*

⁴*Technical University of Lisbon, Instituto Superior de Agronomia, Tapada da Ajuda,
1349-017 Lisbon, Portugal*

Summary

Key aspects of cropping systems are described in terms of their impacts on soil, water and air resources. The importance of microbial symbiotic relations with crop plants are briefly considered in the context of nutrient resource use efficiency and the resilience of plants to biotic and abiotic stress. It is argued that cropping systems need to include crops with fibrous root systems and legumes in the rotation to ensure sustainable use of resources. Three series of experiments are discussed that considered how the efficacy of rhizobia and indigenous arbuscular mycorrhizal fungi could be enhanced in sustainable cropping systems. Evidence is presented to demonstrate that there are potential benefits to crop production from improved management of soil resources through the adoption of reduced tillage practices coupled with suitable crop rotation and weed control.

Key words: Cropping systems, mycorrhizal fungi, rhizobium, nutrient acquisition, soil productivity, plant resilience

Introduction

Cropping systems describe the temporal and spatial arrangements of crops together with the associated management of soil, water and vegetation. Although the aim of a cropping system may be to enhance the productivity of land per unit of input (Lal, 2003), it must be evaluated over several crop cycles to ensure sustainability of the resource base. Soil resource needs to be resilient to the forces of wind and water erosion and less susceptible to the loss of plant nutrients or transport of contaminants by leaching or in surface runoff. Water resources need to be protected from excessive exploitation and contamination, while volatilisation of ammonia and the release of gases, such as nitrous oxide and methane, to the atmosphere must be minimised. The total amount and the dynamics of soil organic matter, normally expressed in terms of carbon (soil organic carbon, SOC), are critical for these requirements and to maintain biological activity.

Plant roots contribute to soil organic matter both by their death and decay and the release of mucilage from their apical regions. During their existence roots undergo extension and turnover; the latter occurring particularly within the various categories of branch roots (Goss & Watson, 2003). Fibrous root systems, typically associated with grasses, tend to result in greater root-length densities (the length of root per unit volume of soil) than do tap roots, which in turn can be

faster growing. Larger values of root-length density help to enmesh soil particles and enhance the resistance of aggregates to disruptive forces. Extraction of water by roots helps increase the strength of aggregates as cementing agents, present in or derived from root-secreted mucilage or decaying plant material, are brought into closer contact with soil particles as water menisci retreat from larger pores of the soil into finer ones (Goss & Kay, 2005).

Over the last century and a half, huge increases in agricultural production have come from the breeding of crop varieties that perform better, particularly being more resistant to disease, the development of more efficient and reliable systems of drainage and irrigation to meet the need for water and aeration, and the use of synthetic fertilisers to provide sufficient nitrogen (N) and phosphorus (P). Leguminous crops, particularly forage species, can obtain more than 80% of their N requirement from the atmosphere by biological N fixation through their symbiosis with rhizobia (Table 1). Some of the N fixed can be passed to non-fixing crops, for example to grass in a mixed stand with alfalfa or clover (Ta & Faris, 1987*a,b*), and after the legume crop has been harvested or ploughed under (Ebelhar *et al.*, 1984; Sheaffer & Seguin, 2003).

Table 1. *Estimates of crop N derived from N₂ fixation by legumes*

Plant	N in crop (kg N ha ⁻¹)	Plant N derived from the atmosphere (%)
Peanut	37–206	22–92
Soybean	17–450	14–98
Cowpea	9–39	12–70
Alfalfa	51–386	46–92
Annual medics	100–200	79–86
White clover	45–291	62–93
Red clover	15–373	35–87

The effectiveness of biological N fixation in legumes can be greatly enhanced if arbuscular mycorrhizal fungi (AMF) also colonise the roots; with the impact being greater if that occurs at an early stage of seedling development (Goss & de Varennes, 2002; de Varennes & Goss, 2007). There is evidence that some N may pass directly between fixing and non-fixing plants in a mixed stand through AMF (Haystead *et al.*, 1988; Frey & Schüepp, 1993). Uptake of mineral N released by the turnover of roots and nodules of legumes appears to be the more important mechanism (Goss *et al.*, 2002; Haystead & Marriott, 1979). The role of AMF in supplying to their hosts nutrients that have limited mobility in soil, such as P and Zn, is well established. Direct uptake of P by roots from the soil can be so down-regulated in the presence of AMF that the only supply is through the fungal hyphae (Smith *et al.*, 2004). Govindarajulu *et al.* (2005) showed that N can also move as ammonium ions (NH₄⁺) from the fungi into host root cells.

Mycorrhizas confer further benefits to the host crop, including reduced uptake of toxic metals (Arines & Vilariño 1989; Bethlenfalvai & Franson 1989), improved tolerance to drought (Augé 2004; Cho *et al.*, 2006) and greater resistance to soil-borne pathogens (Harrier & Watson, 2004). Mycorrhizal fungi also contribute importantly to development and maintenance of soil structure (Goss & Kay, 2005). Cropping systems need to capture and exploit all the advantages offered by both microbial symbioses. Ensuring the optimum rate of colonisation of a new crop by both AMF and rhizobia should be an important objective of soil management options within the cropping system. Enhancing mycorrhizal associations can be achieved by inoculating with exotic strains or by promoting the activity of indigenous AMF (Bagyaraj, 1992). However, inoculation at the field scale requires such a large amount of material that it is considered impractical (Fitter *et al.*, 2011). The success of inoculation with AMF is also determined by their ability to compete with indigenous fungi (Gianinazzi-Pearson & Diem, 1982). Practices that could enhance the efficacy of indigenous AMF would be of considerable benefit.

Colonisation can be initiated by three types of AMF propagule: spores, extraradical hyphae and hyphae from colonised roots fragments. Runner hyphae from a well-developed extraradical mycelium are quicker to initiate colonisation in a new host than other sources of inoculum (Martins & Read, 1997), particularly when the number of viable spores is limited (Read *et al.*, 1976) or soil temperature is not optimal (Entry *et al.*, 2002). Intensive tillage of soil will disrupt extraradical mycelium networks and limit the opportunities for colonisation (Jasper *et al.*, 1989). In contrast, adoption of tillage systems that minimise soil disturbance below the depth of seed placement can encourage persistence of the extraradical mycelium. In temperate climates better protection of the soil surface is afforded by autumn sowing of crops that require vernalisation before reproductive growth can begin. However, such crops are harvested much earlier than the corresponding spring-sown varieties so increasing the period between components of a rotation. Under Mediterranean conditions varieties that have no requirement for vernalisation are commonly grown over winter, resulting in an even longer period between successive crops as sufficient precipitation is required before sowing to ensure plant establishment. But is the longevity of the extraradical mycelium adequate under hot dry conditions in the absence of living host plants? Can the roots of the weeds, which germinate with the first rains, form a living bridge for mycelium networks between successive crops in the rotation? The effect soil disturbance has on the colonisation is also important in developing strategies to optimise the viability of the extraradical mycelium.

Methodology

In the controlled environment pot studies that are discussed below, the underlying common experimental approach was based on that of Fairchild & Miller (1988). Essentially two different inoculum potentials of indigenous AMF were established by growing a base crop, firstly in a 'pre-treatment' cycle, then in two or more 'treatment' cycles. Pots were filled with field soil that had been fully disrupted in passing through a 4–8 mm sieve. The soil was tamped to a bulk density of about 1.2 Mg m⁻³. At the end of the pre-treatment cycle, pots were selected at random and for half the soil was removed as two or three layers, which were kept separate, and again passed through a sieve before being repacked into the same layers within the pot. Root material that had not passed through the sieve was cut into lengths of approximately 1 cm and returned to the soil layer from which they had been separated. These pots formed the 'Disturbed' treatment. The remaining pots formed the 'Undisturbed' treatment. At the end of each treatment cycle soil in the pots of the Disturbed treatment was again disrupted as described above. Establishment of contrasting AMF inoculum potentials was defined by better growth of the host crop in at least the last of these treatment cycles. Further consequences of the differential inoculum potential were then investigated on the growth and development of a 'test' crop, usually over a single cycle. The water content of the soil was maintained at the value of that from a well-drained field. Nutrients were applied to ensure that plants of the last treatment cycle and the test crop showed no deficiency symptoms.

In the first series, clay pots of soil were buried in the field with their tops level with the soil surface in the Alentejo region of Portugal and left without water for the period from the harvest of wheat (*Triticum aestivum* L., cv. Coa) of the last treatment cycle (July) until October. Wheat was used as the test crop. Disruption of the soil in the disturbed treatment only occurred in the pre-treatment and treatment cycles and was not carried out after the last of those cycles or as part of the preparation for planting the test crop. The initial cycles were each of 21 days and the test crop was sampled after 10, 21 and 35 days growth.

In the second study, maize (*Zea mays* L.) was sown for the pre-treatment and treatment cycles and soybean (*Glycine max* (L.) Merr.) was used as the test crop. Peat-based Rhizobium (*Bradyrhizobium japonicum* strain 532C) inoculum was placed at the bottom of a small hole formed by a piece of dowel and the pre-germinated seed of the soybean was placed on top and the soil eased back to

cover the seed. To investigate early colonisation the test crop (*Glycine max* (L.) Merr. cv. Korada) was sampled after 10, 23 and 49 days. To investigate the significance of soil P, $\text{Ca}(\text{H}_2\text{PO}_4)_2 \cdot \text{H}_2\text{O}$ was added to pots at the start of the treatment cycles to achieve four levels of amendment, 0, 20, 40, and 80 mg P kg soil⁻¹. These soybean plants (*Glycine max* (L.) Merr. cv. Evans) were sampled only at podfill (49 days).

For the third study, a mixture of Mediterranean weeds was grown in each pot during the pre-treatment cycle for at least one month. The weeds selected were Persian ryegrass (*Lolium rigidum* Gaudin), wild oat (*Avena sterilis* L.) and littleseed canarygrass (*Phalaris minor* Retz.). For the ‘Undisturbed’ treatment the weeds were killed using herbicide while in the ‘Disturbed’ treatment shoots and roots were chopped and incorporated during soil disturbance. The wheat was sampled after 21 and 28 days.

Results

Despite a period of 2 months exposure to severe drying coupled with hot air temperatures and the absence of supporting plants, indigenous AMF were able to colonise the following wheat crop. The greater colonisation was associated with the establishment of the crop in the absence of significant soil disturbance below the depth of seeding during the treatment cycles (Table 2).

Table 2. *Effect of soil disturbance during treatment cycles on arbuscular and hyphal colonisation of roots in a wheat test crop, between 10 and 35 days after planting*

Previous soil treatment	Days after emergence	Proportional AM colonisation	
		Hyphal	Arbuscular
Undisturbed	10	0.10 ^c	0.04 ^d
Disturbed		0.07 ^c	0.02 ^d
Undisturbed	21	0.19 ^b	0.11 ^c
Disturbed		0.10 ^c	0.06 ^d
Undisturbed	35	0.29 ^a	0.21 ^a
Disturbed		0.21 ^b	0.15 ^b

Values with the same letter within columns are not significantly different from each other ($P = 0.05$).

Assuming development of an intact extraradical mycelium in the disturbed treatment was restricted to the last treatment cycle, the difference in colonisation must have been the result of a greater resilience of the mycelium that had been developed over the three or four cycles in the undisturbed treatment.

Table 3. *Early effect of soil disturbance on AMF colonisation and nodule formation in soybean*

Soil treatment	Days after emergence	AMF colonization parameters			Nodules plant ⁻¹	
		Hyphae (%)	Arbuscules (%)	Vesicles (%)	Number	Dry weight (mg)
Undisturbed	10	70 ^a	56 ^a	0.8 ^a	14 ^a	-
Disturbed		17 ^b	14 ^b	0.0 ^b	8 ^b	-
Undisturbed	23	85 ^a	79 ^a	5.0 ^a	25 ^a	18 ^a
Disturbed		43 ^b	42 ^b	0.3 ^b	21 ^a	4 ^b

For each date, values in a column followed by the same letter are not significantly different as estimated by the *t*-test at $P = 0.05$.

The greater AMF inoculum potential associated with the undisturbed treatment resulted in faster formation of mycorrhizas in soybean (Table 3). However, by podfill roots were colonised by hyphae or arbuscules to about 80% or more and differences induced by soil disturbance in these parameters were small or not statistically different (Table 4). In contrast, the frequency of vesicles in roots from the disturbed treatment was only about half that of plants grown in undisturbed soil (Table 4); consistent with there being faster colonisation by AMF in the absence of disturbance. The addition of large applications of P fertiliser tended to reduce colonisation rates (Table 4).

Table 4. *The effect of P fertiliser application and soil disturbance on AMF colonisation in soybean roots at podfill (R₅)*

Treatment	Soil disturbance		Phosphorus applied (P mg kg ⁻¹ soil)			
	Undisturbed	Disturbed	0	20	40	80
Hyphal	94 ^a	93 ^a	96 ^a	95 ^a	92 ^a	89 ^b
Arbuscular	79 ^a	78 ^b	85 ^a	81 ^{ab}	77 ^{bc}	72 ^c
Vesicular	15 ^a	8.9 ^b	14 ^a	13 ^a	12 ^a	7.9 ^b

Values followed by the same letter in the same row and under the same treatment are not significant at $P = 0.05$.

The faster colonisation of the soybean roots by AMF was accompanied by earlier nodule formation (Table 3). When P levels in the soil were small the total weight of nodules from plants in disturbed soil was less than those from undisturbed soil although the number of nodules was about the same (Table 3). As P levels increased the number of nodules tended to increase, especially in undisturbed soil but differences in weight of nodules were greatly reduced (data not shown). The proportion of N in the plant that was derived from the atmosphere was greater in undisturbed than disturbed soil (Tables 5 and 6) and was enhanced by the addition of P (Table 6).

Table 5. *Effect of soil disturbance on the N content, N₂ fixation, and use of soil N by soybeans*

Soil treatment	Days after emergence	N concentration (g kg ⁻¹ air-dry soil)	N content (mg plant ⁻¹)	Ndfa (%)
Undisturbed	10	75 ^a	15 ^a	-
Disturbed		74 ^a	16 ^a	-
Undisturbed	23	49 ^a	28 ^a	-
Disturbed		46 ^b	25 ^a	-
Undisturbed	49	32 ^b	89 ^a	32 ^a
Disturbed		39 ^a	76 ^b	12 ^b

For each date, values in a column followed by the same letter are not significantly different as estimated by the t -test at $P = 0.05$. Ndfa – N derived from the atmosphere.

The rate of N fixation measured at podfill in disturbed soil (Table 6) was consistent with the values obtained from the differences between N in nodulating and non-nodulating soybean isolines, 2.4 and 0.5 mg N per plant day⁻¹ (Kadir, 1994).

Arbuscular colonisation rate of wheat was enhanced 21 days after planting, in the treatment where weeds were controlled by herbicide rather than by soil disturbance. Enhanced AMF colonisation promoted early P acquisition and growth of the crop (Table 7). The method of weed control significantly affected wheat AM colonisation parameters after 14 and 21 days, with soil disturbance resulting in poorer AM colonisation (Table 8).

Table 6. Proportion of N derived from the atmosphere at podfill of soybean as determined by ^{15}N dilution and the daily rate of fixation measured as plant N derived from ^{15}N gas applied to roots

P applied mg kg^{-1}	Percentage of N derived from the atmosphere (by ^{15}N Dilution)			
	0	20	40	80
Undisturbed	34 ^{cA}	79 ^{bA}	86 ^{aA}	81 ^{aA}
Disturbed	28 ^{cB}	59 ^{bB}	76 ^{aB}	82 ^{aA}
Rate of N fixation in disturbed soil ($\text{mg N per plant day}^{-1}$)	0.5 ^b	ND	ND	2.6 ^a

Values followed by the same letter (lower case) in the same row or under the same P treatment (upper case) are not significant at $P = 0.05$. ND – Not determined.

Table 7. Effect of weed control method on wheat growth and AM colonisation parameters after 21 days

Pretreatment	Shoot weight (g pot^{-1})	P uptake (mg pot^{-1})	Hyphal colonisation	Arbuscular colonisation
No Weeds	1.32 ^a	2.84 ^a	0.32 ^a	0.21 ^b
Weeds				
- Systemic herbicide	1.22 ^{ab}	2.70 ^a	0.39 ^a	0.28 ^a
- Soil disturbed	1.02 ^b	2.03 ^b	0.22 ^b	0.18 ^b

For each measured parameter means followed by the same letter are not significantly different ($P = 0.05$).

Comparison of results for the ‘No weeds’ and ‘Systemic Herbicide treatments in Tables 7 and 8 suggest that the benefits of AM colonisation from extraradical mycelium associated with weed roots increased as that from other inoculum types decreased (difference between ‘No weeds treatments). The type of herbicide (contact or systemic) had no impact on colonisation of the wheat crop (Table 8).

Table 8. Effect of weed control method on AM colonisation in wheat

Days after planting	With weeds prior to wheat planting			‘No weeds’	
	Method of weed control				
	Systemic Herbicide	Contact Herbicide	Disturbance		
Hyphal colonisation	14	0.68 ^b	0.75 ^a	0.48 ^c	0.07 ^d
	21	0.84 ^a	0.77 ^a	0.46 ^b	0.09 ^c
Arbuscular colonisation	14	0.35 ^a	0.36 ^a	0.08 ^b	0.02 ^b
	21	0.40 ^a	0.33 ^a	0.14 ^b	0.02 ^c

For each measured parameter means followed by the same letter are not significantly different ($P \leq 0.05$).

Conclusions

Cropping systems need to include crops with fibrous root systems and legumes in the rotation to ensure sustainable use of resources. Coupled with tillage practices that limits soil disturbance to the depth of seed placement, crops can establish more effective symbioses with indigenous AMF. This can have important consequences in terms of greater resistance to drought and to

soil-borne pathogens, improved utilisation of P and other nutrients with slow mobility in soil but also the reduced availability of toxic elements. In addition, biological N fixation can be enhanced through interactions between AMF and rhizobia. Weeds may be a useful means of enhancing extraradical mycelium as a viable means of colonising crops separated in time, particularly in warm dry periods.

References

- Arines J, Vilariño A. 1989.** Use of nutrient: phosphorus ratios to evaluate the effects of vesicular arbuscular mycorrhiza on nutrient uptake in unsterilized soils. *Biology and Fertility of Soils* **8**:293–297.
- Augé R M. 2004.** Arbuscular mycorrhizae and soil/plant water relations. *Canadian Journal of Soil Science* **84**:373–381.
- Bagyaraj D J. 1992.** Vesicular-arbuscular mycorrhiza: application in agriculture. *Methods in Microbiology* **24**:359–373.
- Bethlenfalvy G J, Franson R L. 1989.** Manganese toxicity alleviated by mycorrhizae in soybean. *Journal of Plant Nutrition* **12**:953–970.
- Cho K, Toler H D, Lee J, Ownley B H, Stutz J C, Moore J L, Augé R M. 2006.** Mycorrhizal symbiosis and response of sorghum plants to combined drought and salinity stresses. *Journal of Plant Physiology* **163**:517–528.
- Ebelhar S A, Frye W W, Blevins R L. 1984.** Nitrogen from legume cover crops for no-tillage corn. *Agronomy Journal* **76**:51–55.
- Entry J A, Rygielwicz P T, Watrud L S, Donnelly P K. 2002.** Influence of adverse soil conditions on the formation and function of arbuscular mycorrhizas. *Advances in Environmental Research* **7**:123–138.
- Fairchild G L, Miller M H. 1988.** Vesicular-arbuscular mycorrhizas and the soil-disturbance induced reduction of nutrient absorption in maize. II. Development of the effect. *New Phytologist* **110**:75–84.
- Fitter A H, Helgason T, Hodge A. 2011.** Nutritional exchanges in the arbuscular mycorrhizal symbiosis: implications for sustainable agriculture. *Fungal Biology Reviews* **25**:68–72.
- Frey B, Schüepp H. 1993.** A role of vesicular-arbuscular (va) mycorrhizal fungi in facilitating interplant nitrogen transfer. *Soil Biology & Biochemistry* **25**:651–658.
- Gianinazzi-Pearson V, Diem H G. 1982.** Endomycorrhizae in the tropics. In *Microbiology of Tropical Soils and Plant Productivity*, pp. 209–251. Eds Y R Dommergues and H G Diem. The Hague: Martinus Nijhoff/Dr W Junk Publishers.
- Goss M J, de Varennes A. 2002.** Soil disturbance reduces the efficacy of mycorrhizal associations for early soybean growth and N₂ fixation. *Soil Biology & Biochemistry* **34**:1167–1173.
- Goss M J, Kay B D. 2005.** Soil aggregation. In *Roots and Soil Management: Interactions between Root and the Soil. Agronomy Monograph No. 48*, Chapter 9. Eds S F Wright and R W Zobel. Madison: ASA, CSSA and SSSA.
- Goss M J, Watson C A. 2003.** The importance of root dynamics in cropping systems research. *Journal of Crop Production* **8**:127–156.
- Goss M J, de Varennes A, Smith P S, Ferguson J A. 2002.** N₂ fixation by soybeans grown with different levels of mineral nitrogen, and the fertilizer replacement value for a following crop. *Canadian Journal of Soil Science* **82**:139–145.
- Govindarajulu M, Pfeffer P E, Jin H, Abubaker J, Douuds D D, Allen J W, Bücking H, Lammers P J, Shachar-Hill Y. 2005.** Nitrogen transfer in the arbuscular mycorrhizal Symbiosis. *Nature* **435**:819–835.
- Harrier L A, Watson C A. 2004.** The potential role of arbuscular mycorrhizal (AM) fungi in the bioprotection of plants against soil-borne pathogens in organic and/or other sustainable farming systems. *Pest Management Science* **60**:149–157.

- Haystead A, Marriott C. 1979.** Transfer of legume nitrogen to associated grass. *Soil Biology & Biochemistry* **11**:99–104.
- Haystead A, Malajczuk N, Grove T S. 1988.** Underground transfer of nitrogen between pasture plants infected with vesicular arbuscular mycorrhizal fungi. *New Phytologist* **108**:417–423.
- Jasper D A, Abbott L K, Robson A D. 1989.** Soil disturbance reduces the infectivity of external hyphae of vesicular- arbuscular mycorrhizal fungi. *New Phytologist* **112**:93–99.
- Kadir S. 1994.** *Interaction between infection with rhizobium and indigenous mycorrhizas on nitrogen fixation and phosphorus nutrition in soybean.* Ph.D. Thesis, University of Guelph.
- Lal R. 2003.** Cropping systems and soil quality. *Journal of Crop Production* **8**:33–52.
- Martins M A, Read D J. 1997.** The effects of disturbance on the external mycelium of arbuscular mycorrhizal fungi on plant growth. *Pesqui Agropecu Brasil* **32**:1183–1189.
- Read D J, Koveheri H K, Hodson J. 1976.** Vesicular-arbuscular mycorrhizal in natural vegetation systems. *New Phytologist* **77**:641–653.
- Sheaffer C C, Seguin P. 2003.** Forage legumes for sustainable cropping systems. *Journal of Crop Production* **8**:187–216.
- Smith S E, Smith F A, Jakobsen I. 2004.** Functional diversity in arbuscular mycorrhizal (AM) symbioses: the contribution of the mycorrhizal P uptake pathway is not correlated with mycorrhizal responses in growth or total P uptake. *New Phytologist* **112**:93–99.
- Ta T C, Faris M A. 1987a.** Effects of alfalfa proportions and clipping frequencies on timothy-alfalfa mixtures. I. Competition and yield advantages. *Agronomy Journal* **79**:817–820.
- Ta T C, Faris M A. 1987b.** Effects of alfalfa proportions and clipping frequencies on timothy-alfalfa mixtures. II. Nitrogen fixation and transfer. *Agronomy Journal* **79**:821–824.
- de Varennes A, Goss M J. 2007.** The tripartite symbiosis between legumes, rhizobia and indigenous mycorrhizal fungi is more efficient in undisturbed soil. *Soil Biology & Biochemistry* **39**:2603–2607.