

BOT14_Zulfikhar.pdf

WORD COUNT

2946

TIME SUBMITTED

08-APR-2020 07:00PM

PAPER ID

57315885

Diversity and Composition of Tree Species of the Secondary Tropical Lowland Forest as a Response to Structure Change of the Meranti-Dangku Landscape, South Sumatra, Indonesia

Zulfikhar¹, Hilda Zulkifli², Sa²ruddin Kadir³, Iskhaq Iskandar⁴

¹Graduate School of Environmental Sciences, Sriwijaya University, Palembang, South Sumatra, Indonesia. 14

²Department of Biology, Faculty of Mathematics and Natural Sciences, Sriwijaya University, Ogan Komering Ilir, South Sumatra, Indonesia. 2

³Department of Soil Science, Faculty of Agriculture, Sriwijaya University, Palembang, South Sumatra, Indonesia.

⁴Department of Physics, Faculty of Mathematics and Natural Sciences, Sriwijaya University, Ogan Komering Ilir, South Sumatra, Indonesia.

* Corresponding authors: [zulfikhar.zl@gmail.com]

Abstract – The diversity and composition of tree species of the tropical lowland forest will change, when there are fragmentation and changes of landscape structure of natural forest, and the dynamics of changes in the composition and diversity of trees species occur in the landscape [1]. However, there are variations among the study areas, which can be determined by tracing the linkage and interaction among indicators of changes in the composition and diversity of tree species [2].

The research objectives are to examine the the trees species diversity forest, and interactions between species on the landscape structure change in the Meranti-Dangku forest, to better understand the effects of landscape modification against individuals and groups of species. We expect that the research findings can be used to enhance the efforts on the forest restoration and biodiversity conservation.

We applied a tree vegetation analysis of fragmented natural secondary forest, and conducted observations in 32 units of sample plots, whose size is 20m x 50m. Ecological analysis of all tree species with dbh \geq 10 cm was conducted by identifying the local name, latin name of the tree species, and its typical characteristics, and then making the sample of herbarium. Hill's Diversity Number produced a family of number related to diversity, effective species richness, a rare species and the abundance of common species in the population [3]. We also calculated the importance value index of tree species and the Jaccard similarity index, to differentiate the level of the forest succession and the gradation among sub-landscape.

The results showed significance of some parameters, which has played important roles in the detecting changes in biodiversity, namely index of species richness and species evenness, and basal area of a tree stand. Three other parameters are not significant. The changes of the tree species composition of the natural forest succession, as respond to the changes of landscape structure, and the gradation among sub-landscape, can be used as a reference on species selection in rehabilitation and restoration of the fragmented natural forest, as a part of the planning on human-modified landscape.

Keyword: Diversity indices, Importance Value Index, Similarity Index, Inter-correlation.

1. INTRODUCTION

The diversity and composition of tree species of the tropical lowland forest will change when there are fragmentation and changes of landscape structure of natural forest, and the dynamics of changes in the composition and diversity of trees species occur in the landscape [1]. However, there are variations among the study areas, which can be determined by tracing the linkage and interaction among indicators of changes in the composition and diversity of tree species [2].

The research objectives are to examine the trees species diversity forest and interactions between species on the landscape structure change in the Meranti-Dangku forest, to better understand the effects of landscape modification against individuals and groups of species. We expect that the research findings can be used to enhance the efforts on the forest restoration and biodiversity conservation.

2. METHODS

2.1 Materials

Work map of field survey in the form of an Indonesia topographical map of 1:50,000 scale, a vegetation cover type map as resulted from interpretation of Landsat TM Imagery acquisition in 2013, GPS handheld, magnetic compass, clinometer Suunto, meter types, phi band, Haga, camera, and materials needed to make herbarium.

2.2 Procedures

We applied a tree vegetation analysis of fragmented natural secondary forest and conducted observations in 32 units of sample plots, whose size 20m x 50m. We measure diameters of all tree species with dbh \geq 10 cm, identified the local name, latin name of the tree species and its typical characteristics, and next made the sample of herbarium for identification in Herbarium Bogoriensis, LIPI.

To analysis of trees species diversity and composition, we use Hill's Diversity Number, which produced a family of number related to diversity, effective species richness, effective number of species, evenness of species, and the abundance of common species in the population [4; 3]). We also calculating the tree density and basal areal of tree stand [5], Importance Value Index of tree species [6], Shannon-Wiener Diversity Index, Simpson Dominance Index, and Simpson Evenness Index [7]. We also calculate the Jaccard Similarity Index [8; 7] to differentiate the structur of forest, level of the forest succession and the gradation among sub-landscape.

3. RESULTS AND DISCUSSION

Species Accumulation Curve

Within 32 sample plots, which covers 3.2 ha, we found a total of 774 species and 1,598 trees, and we processed data of species into a species - accumulation curve. We found 341 species and two unidentified species. The species-accumulation curve showed that it has reached the asymptotic number, where the addition of sample plot did not significantly increase the number of species.

Composition and Diversity of Trees Species

We use Hill's Number of Diversity [9] to determine the richness of species, effective number of species (true diversities), abundance distribution of species, and the evenness of species, as well as the structure of the forest trees, i.e. the density of trees and basal area of trees stand (Table 1). We calculate the important value index of tree species, Jaccard's index for measuring the similarity between communities, and Shannon and Wiener Index to differentiate the level of forest succession, entropy, and gradien between the sub-landscape.

Indicators	Total Sample Plot	Mean	Std. Deviation	Indicators	Total Sample Plot	Mean	Std. Deviation
1. Kapas				3. Meranti Ilir			
N ₀	5	27.00	14.71	N ₀	7	16.57	10.34
N ₁	5	22.93	12.63	N ₁	7	13.82	8.99
N ₂	5	19.44	11.05	N ₂	7	11.76	7.90
E _{0,1} = N ₀ /N ₁	5	1.15	.11	E _{0,1} = N ₀ /N ₁	7	1.21	.188
N/ha	5	562	169	N/ha	7	420	198
BA (m ² /ha)	5	20.47	14.29	BA (m ² /ha)	7	15.84	13.84
2. Meranti Ulu				4. Dangku			
N ₀	13	26.62	8.43	N ₀	7	20.29	4.54
N ₁	13	20.98	9.28	N ₁	7	15.00	5.16
N ₂	13	16.87	9.68	N ₂	7	11.44	5.06
E _{0,1} = N ₀ /N ₁	13	1.37	.296	E _{0,1} = N ₀ /N ₁	7	1.47	.469
N/ha	13	588	309	N/ha	7	479	206
BA (m ² /ha)	13	22.54	7.77	BA (m ² /ha)	7	13.24	5.52

Table 1. The results of the calculation of the indices of biodiversity of forest vegetation based on Hill's Diversity Number and stands structure on each sub-landscape

The results of the study showed that the increase in the effective number of species (true diversities) correlated with a decrease in evenness of species. This is the identifier of the process of secondary succession of forest ecosystems, which indicates the occurrence of a grouping of the new tree species population distribution (clumpiness).

Importance Value Index of Tree Species

Based on the highest order of the important value index, the main species composition in the sub-landscape of Kapas, i.e. Palaquium gutta-percha

(Hook) Ball, Litsea sp, Xanthophyllum rufum, Benn, Shorea parvifolia Dyer, and Castanopsis acuminatissima DC. In the sub-landscapes of Meranti Ulu (specially of young trees class), Meranti Ilir and Dangku are dominated by secondary species or pioneers species, such as Bellucia pentamera Naudin, Bellucia axinantha Trian, Macaranga gigantea (Rchb. f. & Zoll.) Mull. Arg., Endospermum diadenum (Miq.) Airy Shaw, and Ficus variegata Blume. The degraded natural forest has a small average of an important value index, none of dominant species, and the species composition is similar to the natural forest (old growth dipterocarp forest).

Tree diversity Index	Kapas		Meranti Ulu		Meranti Ilir		Dangku	
	DBH >10-30 (cm)	DBH >30 (cm)	DBH >10-30 (cm)	DBH >30 (cm)	DBH >10-30 (cm)	DBH >30 (cm)	DBH >10-30 (cm)	DBH >30 (cm)
1. Shannon's and Wiener Index	4.51	3.04	4.60	5.22	3.59	3.32	3.46	3.24
2. Simpson Dominance Index	0.01	0.06	0.02	0.03	0.10	0.04	0.04	0.07
3. Simpson Evenness Index	0.01	0.04	0.01	0.01	0.01	0.03	0.03	0.01

Table 2. Index of tree diversity of young tree (dbh \geq 10-30 cm) and mature tree (dbh >30 cm) at four locations of the Meranti-Dangku landscape.

Ecological analysis showed significance of some parameters, which have played significant roles in detecting changes in biodiversity, namely index of species richness, species evenness, and basal area of a tree

stand. Three other parameters are not significant. The changes of the tree species composition of the natural forest succession, as respond to the changes of landscape structure, and the gradation among sub-landscape, can be used as a indicator in species selection in rehabilitation and restoration of the fragmented natural forest, as a part of the human-modified landscape.

Barbour states [8] that the value of Shannon's and Wiener Diversity Index can range between 0-7, with criteria: 0-2 (low), 2 (medium), and > 3 (high). Thus, the level of species diversity index of young and mature trees at four locations are classified as high.

Simpson's Dominance Index of Dangku forest was the highest at both young and mature trees. However, for the forest at Dangku had Shannon's and Wiener (Table 2.) Diversity Index is lower, at both young and mature trees, compared to a third of other locations. Specifically, the young trees, with a diameter of ≥ 10 –30 cm, at Dangku had the highest Simpson's Evenness Index., compared to an other three locations.

Discussion

Correlation between Tree Species Diversity Index

The calculation of the value of the correlation between the vegetation of the forest biodiversity indices, was conducted based on Hill's Diversity Number, and structure of the forest stand, performed in Kapas, Meranti Ulu and Meranti Ilir, indicating there is a very significant positive correlation ($p=0.01$) between the number of species in the plot or the richness of species, and the number of effective species (true diversities), and the abundance of species distribution, and as well as between the number of effective species with an abundance of species distribution.

In addition, in Kapas, there was a significant positive correlation ($p = 0.01$) between a richness of tree species and tree density of forest stands, as well as a highly significant positive correlation ($p = 0.01$) between the abundance of tree species distribution with a basal area of forest tree stands.

In Meranti Ulu, shows that the richness of species and the effective number of species (true diversities) Meranti Ulu, both negatively correlated in a very significant ($p=0.01$) against species evenness. This means that if the total number of species in the plots and the effective number of species increased, so this is precisely indicated that species evenness will decrease. Species diversity of ensembles in undisturbed primary forest was distinctly higher than in disturbed or secondary forest. Disturbed forest with tall trees remaining appeared to represent impoverished subsets of the undisturbed primary forest community [10].

Importance Value Index of Tree Species

The analysis to the top ten IVI showed that each group of forest on each location, and a different diameter class was dominated by different species, thus the ability of species to live in a place is very depending on its ability to adapt to environmental conditions of the habitat. Therefore, the environmental conditions are very instrumental in the selection of species to be able to survive in a habitat.

There is no dominant young tree species, where the average IVI is low. This means that in the sub-landscape Kapas has a high species richness, especially for young trees. Old dipterocarp forests in Kapas and Meranti Ulu sub-landscape are managed using silviculture system of the restoration of natural forest ecosystem, the species of young trees and mature trees, have low IVI compared to IVI in secondary forest in sub-landscape of Meranti Ilir and Dangku.

The Implication against Island Biogeography Theory

Every ecologist know [11], that, Island Biogeography Theory (IBT) states that the number of species in a habitat Island is governed by a balance of dependence distance colonization and dependence of extinction area, and it predicts that the habitat of the smaller and more isolated will be supported by fewer species.

A fragmented landscape, whether in the dissection phase, dissipation or break apart, then in accordance with the IBT will happen to lose and decline in species richness, species-area curve as that hypothesized by IBT [12].

Deviations against the IBT as stated by [13] are, that, the dynamics of populations and communities are often heavily enforced by the relatively fragmentary habitat against the natural condition. Likewise, other discoveries are that habitat fragmentation affects different species in different ways. Some species are declining sharply, or disappeared in fragments, remain stable in other area, and increases dramatically, elsewhere [14].

The results of this research also show, that, the young growth tree species in the sub-landscape of Meranti Ulu and Meranti Ilir, and the young growth and mature growth of tree species in sub-landscape of Dangku, were dominated by a secondary tree species, which controlled the growth phases of a tree in the process of forest succession. The symptoms are so common on the type of vegetation that lead to the climatic conditions and to stability. The species composition of the natural forest, that, has developed in the long run, will show physiognomy, fenologi, and low power regeneration, and tends to be steady. Thus, the new dynamics of the forest community are manifestly and less conspicuously, where the turning generations, or species regeneration, does not seem as if, as a result of certain rare species are dominant, since all species have to adapt in a prolonged period of time [15].

At the stage of forest climax, then, the IBT is likely to be re-occur [15]. It was observed during this research results that the reduction the number of populations and species of trees occurred, when the phase of forest fragmentation is Attrition, and an increase in the number of species of trees and evenness index of tree species on the secondary growth forest, during the phase of Dissipation and Dissection [16].

4. CONCLUSIONS

The effect of the change of landscape metrics, especially on the regrowth and regeneration of the forest succession, is significantly to increase the richness of species, the evenness index of tree species, and basal area of tree stand. The increase in the effective number of species (true diversities) was followed by a decline in the evenness index of tree species, and the grouping of the new tree species population (clumpiness). This is an indicator of the occurrence of a succession process of the secondary forest ecosystems.

Dangku sub-landscape has an average value of Shannon's and Wiener entropy index, an index of tree species richness, a tree density, and basal area, which are higher than in Meranti Ilir sub-landscape. This indicates that the entropy condition for the succession process of secondary forest ecosystems in Dangku sub-landscape relatively more steady than on Meranti Ilir sub-landscape.

Secondary forest of the adjacent sub-landscape of Meranti Ilir and Dangku are experiencing the high pressure from the deforestation and forest degradation. An average value of density of trees stand is almost the same, and has a very significant positive correlation between density of trees stand and index of tree species evenness. However, the Jaccard's similarity index both sub-landscapes is very low.

The change of landscape metrics led to the decolonization of the composition and diversity of tree species in the succession of secondary natural forest, and significantly cause a rise in the index of tree species richness, and the index of tree species evenness. This phenomenon is deviation from the Island Biogeography Theory (IBT). However, it is temporal, and will follow the dynamics of the forest succession process until a climax of forest succession is achieved, as the old growth dipterocarp forest. This phenomenon occurs due to the edge effects, shade-intolerant in gap dynamics of forest canopies, connection between patches, as well as seed dispersal factors.

REFERENCES

- [1] L. Fahrig, "Effects of habitat fragmentation on biodiversity," *Annu Rev Ecol Evol Syst*, vol. 34, pp. 487–515, 2003.
- [2] M. Martínez-Ramos, I. A. Ortiz-Rodríguez, D. Piñero, R. Dirzo, and J. Sarukhán, "Anthropogenic disturbances jeopardize biodiversity conservation within tropical rainforest reserves," *Proc. Natl. Acad. Sci.*, vol. 113, no. 19, pp. 5323–5328, 2016.
- [3] A. Chao, N. J. Gotelli, T. C. Hsieh, E. L. Sander, K. H. Ma, R. K. Colwell, and A. M. Ellison, "Rarefaction and extrapolation with Hill numbers: A framework for sampling and estimation in species diversity studies," *Ecol. Monogr.*, vol. 84, no. 1, pp. 45–67, 2014.
- [4] M. O. Hill, "Diversity and evenness: a unifying notation and its consequences," *Ecology*, vol. 54, no. 2, pp. 427–432, 1973.
- [5] J. A. Husch, B., Beers, T.W. and Kershaw Jr, *Forest mensuration*. John Wiley & Sons., 2002.
- [6] A. I. Soerianegara, I., *Ekologi Hutan Indonesia*. Edisi Pert. Fakultas Kehutanan - Institut Pertanian Bogor., 2005.
- [7] D. Setiadi, "Keanekaragaman Spesies Tingkat Pohon di Taman Wisata Alam Ruteng , Nusa Tenggara Timur," *Biodiversitas*, vol. 6, no. April, pp. 118–122, 2005.
- [8] J. K. B. and W. D. P. Barbour, G.M., *Terrestrial Plant Ecology*. New York: Cummings Publishing Company, Inc., 1987.
- [9] A. Chao, C.-H. Chiu, and L. Jost, "Unifying species diversity, phylogenetic diversity, functional diversity, and related similarity and differentiation measures through Hill N_j numbers," *Annu. Rev. Ecol. Evol. Syst.*, vol. 45, no. 1, pp. 297–324, 2014.
- [10] K. Fiedler and C. H. Schulze, "Forest modification affects diversity (but not dynamics) of speciose tropical pyraloid moth communities," *Biotropica*, vol. 36, no. 4, pp. 615–627, 2004.
- [11] R. MacArthur and E. Wilson, "Island biogeography," *Monogr. Popul. Biol.*, vol. 1, no. 1, pp. 1–203, 1967.
- [12] J. Fischer and D. B. Lindenmayer, "Landscape modification and habitat fragmentation: A synthesis," *Global Ecology and Biogeography*, vol. 16, no. 3, pp. 265–280, 2007.
- [13] W. F. Laurance, T. E. Lovejoy, H. L. Vasconcelos, E. M. Bruna, R. K. Didham, P. C. Stouffer, C. Gascon, R. O. Bierregaard, S. G. Laurance, and E. Sampaio, "Ecosystem Decay of Amazonian Forest Fragments: a 22-Year Investigation," *Conserv. Biol.*, vol. 16, no. 3, pp. 605–618, Jun. 2002.
- [14] R. E. Ricklefs and I. J. Lovette, "The roles of island area per se and habitat diversity in the species-area relationships of four Lesser Antillean faunal groups," *J. Anim. Ecol.*, vol. 68, no. 6, pp. 1142–1160, 1999.
- [15] C. Echeverría, A. C. Newton, A. Lara, J. M. R. Benayas, and D. A. Coomes, "Impacts of forest fragmentation on species composition and forest structure in the temperate landscape of southern Chile," *Glob. Ecol. Biogeogr.*, vol. 16, no. 4, pp. 426–439, 2007.
- [16] Zulfikhar, Zulkifli H. Kadir S. and Iskandar I., "The Rapid Changes of the Landscape Structure of the Meranti-Dangku Tropical Lowland Forest in the South Sumatra Province, Indonesia," *Sriwij. J. Environ.*, vol. 2, no. 1, pp. 8–18, 2017.

7%

SIMILARITY INDEX

PRIMARY SOURCES

- 1 Konrad Fiedler. "Forest Modification Affects Diversity (But Not Dynamics) of Speciose Tropical Pyraloid Moth Communities¹", *Biotropica*, 2004 31 words — 1%

Crossref
- 2 Ady Mara, Karna Wijaya, Wega Trisunaryati Mudasir. "Effect of Sulfuric Acid Treatment and Calcination on Natural Zeolites of Indonesia", *Asian Journal of Chemistry*, 2016 29 words — 1%

Crossref
- 3 Nurlita Pertiwi, Nur Anny Suryaningsih Taufieq, St. Fatmah Hiola. "The Diversity of Riparian Trees Vegetation at Around The Lawo River, South Sulawesi, Indonesia", *Journal of Physics: Conference Series*, 2019 19 words — 1%

Crossref
- 4 John S. Gray. "The measurement of marine species diversity, with an application to the benthic fauna of the Norwegian continental shelf", *Journal of Experimental Marine Biology and Ecology*, 2000 18 words — 1%

Crossref
- 5 Laurance, W.F.. "Theory meets reality: How habitat fragmentation research has transcended island biogeographic theory", *Biological Conservation*, 2008⁰⁷ 17 words — 1%

Crossref
- 6 Martins, Inês Santos, Vânia Proença, and Henrique Miguel Pereira. "The unusual suspect: 10 words — < 1%

Land use is a key predictor of biodiversity patterns in the Iberian Peninsula", *Acta Oecologica*, 2014.

[Crossref](#)

7 Alan Feest, Ian Merrill, Philippa Aukett. "Does Botanical Diversity in Sewage Treatment Reed-Bed Sites Enhance Invertebrate Biodiversity?", *International Journal of Ecology*, 2012

9 words — < 1%

[Crossref](#)

8 Ermias T. Azeria, Isabel Sanmartín, Stefan Ås, Allan Carlson, Neil Burgess. "Biogeographic patterns of the East African coastal forest vertebrate fauna", *Biodiversity and Conservation*, 2007

9 words — < 1%

[Crossref](#)

9 Pardini, R.. "The challenge of maintaining Atlantic forest biodiversity: A multi-taxa conservation assessment of specialist and generalist species in an agro-forestry mosaic in southern Bahia", *Biological Conservation*, 200906

8 words — < 1%

[Crossref](#)

10 R. Bozorgipour, J. W. Snape. "The relationship between in vitro performance of haploid embryos and the agronomic performance of the derived doubled haploid lines in barley", *Theoretical and Applied Genetics*, 1992

8 words — < 1%

[Crossref](#)

11 Rocha-Santos, Larissa, Michaele S. Pessoa, Camila R. Cassano, Daniela C. Talora, Rodrigo L.L. Orihuela, Eduardo Mariano-Neto, José C. Morante-Filho, Deborah Faria, and Eliana Cazetta. "The shrinkage of a forest: Landscape-scale deforestation leading to overall changes in local forest structure", *Biological Conservation*, 2016.

8 words — < 1%

[Crossref](#)

12 A. Khaled, A. El Nemr, A. El Sikaily. "An assessment of heavy-metal contamination in surface sediments of the Suez Gulf using geoaccumulation indexes and statistical analysis", Chemistry and Ecology, 2006 7 words — < 1%

Crossref

13 "Stability of Tropical Rainforest Margins", Springer Science and Business Media LLC, 2007 6 words — < 1%

Crossref

14 Bakri Rio Rahayu, Tarmizi Taher, Poedji Loekitowati Hariani, Aldes Lesbani. "Congo red and direct yellow dye removal from aqueous solution by Zn/Cr layered double hydroxides", AIP Publishing, 2018 6 words — < 1%

Crossref

EXCLUDE QUOTES OFF

EXCLUDE SOURCES OFF

EXCLUDE BIBLIOGRAPHY ON

EXCLUDE MATCHES OFF